



HAL
open science

Erythrocyte glutathione peroxidase (GSH-Px) polymorphism in Finnsheep

F. Atroshi, S. Sankari, S. Oeterberg, M. Sandholm

► **To cite this version:**

F. Atroshi, S. Sankari, S. Oeterberg, M. Sandholm. Erythrocyte glutathione peroxidase (GSH-Px) polymorphism in Finnsheep. *Annales de génétique et de sélection animale*, 1982, 14 (1), pp.100-100. <hal-00893384>

HAL Id: hal-00893384

<https://hal.science/hal-00893384v1>

Submitted on 11 May 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire HAL, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



HAL Authorization

In vitro testing of blast transformation, immune rosette formation, and IF-detectable IgG-positive cell counts indicated a 40-50 p. 100 decrease of immunological responsiveness relative to the control. Toxin neutralizing titres to necrotic enteritis/NE/vaccine were in fact significantly lower in the sera of T₂-treated pigs.

In the given conditions of experiment, T₂ developed a distinct immunosuppressive action in the early phase of immune induction in growing pigs by interfering with the function of both T- and B-lymphocytes.

Erythrocyte glutathione peroxidase (GSH-Px) polymorphism in Finnsheep

F. ATROSHI *, S. SANKARI, S. OETERBERG and M. SANDHOLM

Department of Pharmacology and Toxicology () / Department of Biochemistry
College of Veterinary Medicine, Helsinki 55, Finland
and Department of Animal Breeding, Agricultural Research Centre, Vantaa, Finland*

Erythrocyte glutathione peroxidase (GSH-Px) was studied in 200 *Finnsheep*. The concentration of GSH-Px varied between 500 to 1 800 μ kat/l red cells and the distribution revealed a bimodality despite the same diet and selenium intake. The peroxidase activity of the sheep erythrocytes was classified high if it was $> 1\,000$ μ kat/l or low if it was $< 1\,000$ μ kat/l. Inheritance studies of GSH-Px types revealed that they are controlled by a pair of autosomal genes, the genes for GSH-Px high being dominant to the genes for GSH-Px low. These two groups of sheep were then compared for other blood parameters and production characteristics. It is proposed that the low GSH-Px may represent an adaptation to low selenium intake.

SESSION II

PROBLÈMES D'ÉLEVAGE SOUS LES TROPIQUES

Cattle production in the tropics and improvement through breeding

J. RENDEL

*Animal Production and Health Division
Food and Agriculture Organization of the United Nations
Via delle Terme di Caracalle, 00100 Rome, Italy*

Results from breeding and improving dairy and beef cattle in the tropics are reviewed. It is concluded that for improved dairy production, climate *per se* is no serious constraint to the use of European-type cattle in tropical highlands or in dry, hot areas provided that management is good. In hot, humid areas, European-type cattle do not adapt well, and local well-adapted breeds or crosses should be used. Suitable breeding policies are outlined. Under conditions of smallholder production, the extensive system with crossbred cattle, once-a-day milking and calf suckling has a role to play. Beef production is generally carried