



HAL
open science

**FTIR spectroscopy and taxonomic purpose:
Contribution to the classification of lactic acid bacteria**

Caroline Amiel, Laurence Mariey, Catherine Denis, Patricia Pichon, Josette
Travert

► **To cite this version:**

Caroline Amiel, Laurence Mariey, Catherine Denis, Patricia Pichon, Josette Travert. FTIR spectroscopy and taxonomic purpose: Contribution to the classification of lactic acid bacteria. *Le Lait*, 2001, 81 (1-2), pp.249-255. 10.1051/lait:2001128 . hal-00895477

HAL Id: hal-00895477

<https://hal.science/hal-00895477>

Submitted on 11 May 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Evolution, biodiversity, taxonomy

FTIR spectroscopy and taxonomic purpose: Contribution to the classification of lactic acid bacteria

Caroline AMIEL^{a*}, Laurence MARIEY^a, Catherine DENIS^b,
Patricia PICHON^b, Josette TRAVERT^a

^a Laboratoire de Physico-Chimie et Biotechnologies, IUT de Caen,
Université de Caen Basse Normandie, 14032 Caen Cedex, France

^b ADRIA Normandie, boulevard du 13 juin 1944, BP 2, 14310 Villers-Bocage, France

Abstract — The capacity of FTIR spectroscopy for identification and classification of lactic acid bacteria has been proved on brewery strains and more recently by ourselves on dairy strains. During our study, we underlined that IR spectra give very informative data which can be used for old or more recent taxonomical purposes. We tried to illustrate this aspect with two examples: the distinction between *Streptococcus thermophilus* and *Streptococcus salivarius*, the taxonomic range of *Lactobacillus casei*, *paracasei*, *zeae* and *rhamnosus*. The interest of this tool for taxonomic studies will be discussed.

taxonomy / FTIR spectroscopy / *Streptococcus* / *Lactobacillus*

Résumé — Apport de la spectroscopie infrarouge à transformée de Fourier en taxonomie : contribution à la classification de bactéries lactiques. La capacité de la spectroscopie IRTF pour l'identification et la classification de bactéries lactiques a été démontrée pour des souches de lactobacilles brassicoles et plus récemment par notre équipe en matière de souches laitières. Au cours de notre étude, il nous est apparu que les données spectrales obtenues, étaient riches en informations pouvant être utilisées pour étayer des discussions taxonomiques anciennes ou plus récentes. Nous illustrons notre propos par deux exemples : la distinction entre les souches types de *Streptococcus thermophilus* et *Streptococcus salivarius* et la position taxonomique des espèces *Lactobacillus casei*, *paracasei*, *zeae* et *rhamnosus*. L'intérêt et la place de cet outil en taxonomie bactérienne est présenté.

taxonomie / spectroscopie IRTF / *Streptococcus* / *Lactobacillus*

* Correspondence and reprints
Tel.: (33) 2 31 56 71 19; fax: (33) 2 31 56 71 65; e-mail: C.amiel@iutcaen.unicaen.fr

1. INTRODUCTION

Bacteria FTIR spectra give a global picture of whole cellular components (fatty acids, intracellular and membrane proteins, polysaccharides, nucleic acids) [5, 16] and are considered to be a phenotypic method. Moreover, spectra give information relative to nucleic acids and in this way it may be proposed as a link between the both genomic and phenotypic approaches. In the last twenty years, genomic data has led to many taxonomic changes, and species usually confused can now be distinguished. This is the case for *S. salivarius* and *S. thermophilus*, and for *Lb. casei*, *Lb. zaeae* and *Lb. rhamnosus* species involved in the dairy industry that we have studied by FTIR spectroscopy [1, 2]. By using more and more efficient methods a lot of new species have been created each year, and systematic bacteriology is more and more complex. The interest of these new species is not always very clear for lack of easy methods to identify them routinely. The comparison of FTIR spectra using statistical analysis makes it possible to classify them. In a previous study, we recorded the spectra of 12 species of *Lactobacillus* involved in the dairy industry [1, 2]. In this paper, we point out the capacity of FTIR spectroscopy for taxonomic discussions about some species and we present the screening of wild strains belonging to these species.

2. MATERIALS AND METHODS

2.1. Bacteria growth and sample preparation

The strains come from international collections ATCC, CNRZ, LMG and CIP and were kept frozen ($-80\text{ }^{\circ}\text{C}$) in Adria-Normandie. Wild strains were isolated from "Pont-L'Evêque" cheese (strains p64, p135) by Adria-Normandie and from "Camembert" cheese by LMA, University of Caen

Basse-Normandie. All strains were grown for $24\text{ h} \pm 2\text{ h}$ in optimal conditions: MRS, $37\text{ }^{\circ}\text{C}$ for lactobacilli, M 17, $37\text{ }^{\circ}\text{C}$ for streptococci. Cells are pelleted by centrifugation $4\text{ }000\text{ rpm}$ for 10 min and washed twice with saline solution as previously described [1]. Five microlitres of the concentrated bacteria were put on a ZnSe spectral window and dried for an hour at $50\text{ }^{\circ}\text{C}$.

2.2. Spectroscopic measurements

Bacterial spectra were recorded between $4\text{ }000$ and 700 cm^{-1} using a FTIR spectrometer (Nicolet 250, Nicolet Instrument, Thermo-Optek, Montigny le Bretonneux, France) equipped with a KBr beamsplitter and a DTGS detector. Sixty-four scans were averaged per spectrum at a resolution of 4 cm^{-1} . All spectra were submitted to a "quality test" adapted from Helm et al. [10]. After this test, validated spectra were normalized to one absorbance unit using the amide I spectral band located at about $1\text{ }640\text{ cm}^{-1}$.

2.3. Statistical analysis

Three different methods were employed to compare spectra.

2.3.1. Discriminant analysis

Using Omnic TQ analyst software 1.2 (Nicolet instrument, Thermo-Optek, Montigny-le-Bretonneux, France), discriminant analysis is performed by the calculation of Mahalanobis distances between spectra [12].

2.3.2. Search standard method

Using the same software, we can obtain the percentage of similitude between an unknown spectrum and spectra of a database. Results are noted from 1 to 100 (100 is the best correlation between spectra) [1, 2].

2.3.3. Hierarchical cluster analysis

Using Omnic software (Nicolet instrument, Thermo-Optek, Montigny-le-Bretonneux, France), spectra were first smoothed and the first derivative was calculated. Then, using SPSS 8.0 for Windows software, each spectra was considered as an observation, and each point of the spectra as a variable. Euclidian spectral distances were calculated and Ward algorithm was performed to obtain the dendrogram.

3. RESULTS AND DISCUSSION

3.1. Discrimination between *S. salivarius* and *S. thermophilus*

Normally only one species of the genus *Streptococcus* (*S. thermophilus*) is found in the dairy industry [7, 13] but the species *S. salivarius*, very close to *S. thermophilus*, is an opportunistic bacteria in man and that is why it is so important to separate them. The taxonomic border between the two species has been discussed for years: in 1984, on the basis of DNA hybridization and fatty acid results, Farrow and Collins [9] proposed to classify them in the same species. Later, Schleifer et al. [17]

underlining a DNA homology of only 60% in optimal conditions and 30% in stringent conditions between the two species, refiled them as distinctive species. This last taxonomic position was validated in 1980 by Skerman [18]. However, in practice, strains belonging to these two species are difficult to separate by phenotypic tests.

Using FTIR spectroscopy, both type strains (NCDO 573^T and ATCC 7073^T) show spectral differences visible to the naked eye, particularly between 3 100–2 800 cm⁻¹, a region corresponding to fatty acids vibrations, and around 1 200–900 cm⁻¹ (polysaccharides vibrations) (Fig. 1). Calculation of Mahalanobis distances between the spectra of the two strains (over 100 versus 0.67 for distances between spectra of each strain) in the region 1 500–700 cm⁻¹, translates a good discrimination of the two strains (Fig. 2, Tab. D). Likewise, the percentage of homology between the two strains is less than 90%, even though in our experience the limit for the same species is 95% [1, 2]. These three arguments, obtained from the IR spectrum alone, are sufficient to confirm the argument of Schleifer et al. rather than the one of Farrow et al. and to propose FTIR spectroscopy as a complementary tool for the discrimination of the two species.




Figure 1. FTIR spectra of *S. thermophilus* (—) and *S. salivarius* (.....) type strains.




Figure 2. Discriminant analysis of spectra from *S. thermophilus* and *S. salivarius* type strains.

Table I. Mahalanobis distances between type strains of *S. thermophilus* and *S. salivarius*.

Strain		Distance to class	
		<i>S. thermophilus</i>	<i>S. salivarius</i>
<i>S. salivarius</i>	spectra 1	101.74	0.67
	spectra 2	115.61	0.67
ATCC 7073 ^T	spectra 3	131.03	0.67
	spectra 1	0.67	112.73
<i>S. thermophilus</i>	spectra 2	0.67	120.56
	spectra 3	0.67	115.09

3.2. Contribution to the discussion about the taxonomic position of *Lb. casei* and related taxa

Taxonomic discussion about *Lb. casei* species and related taxa is old and always uncertain: five subspecies admitted to the approved list of Bacterial Names in 1980 [18], four in the last edition of the Bergey's Manual of Systematic Bacteriology [11], and, more recently, three separate species (*Lb. casei*, *Lb. paracasei* and *Lb. rhamnosus*), on the basis of molecular arguments proposed by Collins et al. [4]. At the same time, since 1971, the proximity of the type strains ATCC 393^T (*Lb. casei*) and ATCC 15820^T (*Lactobacterium zeae*, reclassified as *Lb. rhamnosus*) has been underlined by different research teams [3, 6, 8, 14, 15]. Most

of the authors proposed the combination of ATCC 393^T and ATCC 15820 in the species *Lb. zeae* (type strain ATCC 15820). Moreover, Dellaglio's team proposes the rejection of the species *Lb. paracasei* and the recognition of ATCC 334 as the new type strain of *Lb. casei*. However, these changes have been rejected by the Juridical Commission of the International Committee on Systematic Bacteriology [19] for fear of confusion in the existing taxonomy.

Comparison of FTIR spectra of the type strains considered and some other collection strains does not show many differences between them except for *Lb. rhamnosus* CNRZ 212^T (Fig. 3). Similarly, the hierarchical cluster obtained from spectral data, using the Ward method, confirms the




Figure 3. FTIR spectra of *Lb. zeae* (CIP 103253: ---), *Lb. casei* (CNRZ 313^T ----), *Lb. paracasei* subsp *paracasei* (CNRZ 62^T - - - -), *Lb. paracasei* subsp *tolerans* (CIP 102309^T) and *Lb. rhamnosus* (CNRZ 212^T ———).

proximity between CNRZ 313^T and CIP 103253, the singularity of ATCC 334, and the real distance of CNRZ 212^T from other strains (Fig. 4). Yet, contrary to the results of Dicks et al. [8], strains CNRZ 62^T and CIP 103024^T are closer to the casei-zeae cluster

than to strain ATCC 334. Since our results are partly in accordance with those of Dellaglio's team we decided to consider as *Lb. zeae* all strains closely related to ATCC 103253 and/or CNRZ 313^T; to consider as *Lb. paracasei* all strains related to CNRZ 62^T,




Figure 4. Hierarchical cluster analysis using Ward algorithm on Euclidian spectral distances.

and CIP 103402^T; and to keep the strains close to CNRZ 212^T as *Lb. rhamnosus*. The remaining strains, including ATCC 334, are kept as *Lb. casei*.

In this way, when we try to identify wild strains by comparing their spectra with those of type strains, we can easily distinguish the *Lb. rhamnosus* strains from the *zeae* and the *casei* strains. Table II gives the results of FTIR identification in comparison with the results of phenotypic tests (Api 50 CHL) and for some strains with the RAPD method. The correlation between RAPD and Api methods was good for five strains among the six tested. The correlation between the RAPD and FTIR methods was good for the six strains. The differences observed between the FTIR and Api methods can be explained insofar as phenotypic tests do not

take into account the species *Lb. zeae* and consider as *Lb. rhamnosus* the strains which ferment rhamnose. In fact, ATCC 15820 and related strains ferment rhamnose too.

With this precision, FTIR spectroscopy allows a more precise classification of the casei-group than phenotypic tests. This classification is as efficient as those obtained with genomic methods such as RAPD. In both cases, FTIR spectroscopy is more rapid.

4. CONCLUSION

Concerning the taxonomic subject, our results confirm only in part those of Dellaglio et al. but don't allow the grouping together of *Lb. casei* and *Lb. paracasei* strains around the type strain ATCC 334,

Table II. Comparison of identifications with different methods.

Strain	Identification by		
	Api50CHL	RAPD	IRTF
435	<i>paracasei</i>	<i>paracasei</i>	<i>paracasei</i>
468	<i>paracasei</i>	<i>paracasei</i>	<i>paracasei</i>
733	<i>paracasei</i>	<i>paracasei</i>	<i>paracasei</i>
736	<i>paracasei</i>	<i>paracasei</i>	<i>paracasei</i>
833	<i>paracasei</i>	<i>paracasei</i>	<i>paracasei</i>
936	<i>paracasei</i>	ND	"casei 334"
p64	ND	ND	"casei 334"
102	<i>paracasei</i>	ND	"casei 334"
103	<i>paracasei</i>	ND	"casei 334"
196	<i>paracasei</i>	ND	"casei 334"
82	<i>paracasei</i>	ND	"casei 334"
83	<i>paracasei</i>	ND	"casei 334"
84	<i>paracasei</i>	ND	"casei 334"
62	<i>rhamnosus</i>	ND	<i>rhamnosus</i>
80	douteux (rhamnose+)	ND	<i>rhamnosus</i>
81	<i>rhamnosus</i>	ND	<i>rhamnosus</i>
272	ND	ND	<i>rhamnosus</i>
85	<i>rhamnosus</i>	ND	<i>casei/zeae</i>
79	<i>rhamnosus</i>	<i>casei</i>	<i>casei/zeae</i>
p135	<i>paracasei</i>	ND	<i>casei/zeae</i>
837	<i>paracasei</i>	ND	<i>casei/zeae</i>

since two distinct clusters are present. Nevertheless, the method allows one to set apart easily *Lb. zeae* strains from *Lb. rhamnosus* strains, contrary to Api 50CHL which cannot distinguish them. Moreover in the case of *streptococci*, FTIR spectroscopy allows one to set apart type strains of *S. thermophilus* and *S. salivarius* so we can propose to study these two species more precisely with this method.

The information given by FTIR spectra is complementary to genomic information and can be introduced in a polyphasic taxonomic approach.

REFERENCES

- [1] Amiel C., Mariey L., Curk-Daubié M.C., Pichon P., Travert J., Potentiality of Fourier Transform Infrared Spectroscopy (FTIR) for discrimination and identification of dairy lactic acid bacteria, *Lait* 80 (2000) 1–15.
- [2] Amiel C., Potentialités de la spectroscopie infrarouge à transformée de Fourier (IRTF) pour l'identification et la caractérisation de bactéries d'implication laitière, thèse de l'université de Caen, 2000, pp. 71–95.
- [3] Chen H., Lim C.K., Lee Y.K., Chan Y.N., Comparative analysis of the gene encoding 23s-5s-rRNA intergenic spacer regions of *Lactobacillus casei* related strains, *Int. J. Evol. Microbiol.* 50 (2000) 471–478.
- [4] Collins M.D., Philips B.A., Zanoni P., Deoxyribonucleic acid homology studies of *Lactobacillus casei*, *Lactobacillus paracasei* sp. nov., subsp. *paracasei* and subsp. *tolerans*, and *Lactobacillus rhamnosus* sp. nov. comb. nov., *Int. J. Syst. Bacteriol.* 39 (1989) 105–108.
- [5] Curk M.-C., Peladan F., Hubert J.-C., Identification bactérienne par spectrométrie infrarouge à transformée de Fourier, *Bull. Soc. Fr. Microbiol.* 9 (1994) 278–283.
- [6] Dellaglio F., Dicks L.M.T., Dutoit M., Torriani S., Designation of ATCC 334 in place of ATCC 393 (NCDO 161) as the neotype strain of *Lactobacillus casei* subsp. *casei* and rejection of the name *Lactobacillus paracasei*, *Int. J. Syst. Bacteriol.* 41 (1991) 340–342.
- [7] Dellaglio F., de Roissart H., Torriani S., Curk M.C., Janssens D., Caractéristiques générales des bactéries lactiques, in: Roissart H., Luquet F.M. (Eds.), *Bactéries Lactiques*, Loriga, Uriage, France, 1994, pp. 25–70.
- [8] Dicks L.M.T., Du Plessis E.M., Dellaglio F., Lauer E., Reclassification of *Lactobacillus casei* subsp. *casei* ATCC 393 and *Lactobacillus rhamnosus* ATCC 15820 as *Lactobacillus zeae* nom. rev., designation of ATCC 334 as the neotype of *L. casei* subsp. *casei*, and rejection of the name *Lactobacillus paracasei*, *Int. J. Syst. Bacteriol.* 46 (1996) 337–340.
- [9] Farrow J.A.E., Collins M.D., DNA base composition, DNA-DNA homology and long-chain fatty acid studies on *Streptococcus thermophilus* and *Streptococcus salivarius*, *J. Gen. Microbiol.* 130 (1984) 357–362.
- [10] Helm D., Labishinski H., Naumann D., Elaboration of a procedure for identification of bacteria using Fourier-Transform IR spectral libraries: a septwise correlation approach, *J. Microbiol. Methods* 14 (1991) 127–142.
- [11] Kandler O., Weiss N., Regular, nonsparing Gram-positive rods, in: Sneath P.H.A., Mair N.A., Sharpe M.E., Holt J.G. (Eds.), *Bergey's Manual of System. Bacteriol.*, Vol. 2, Williams and Wilkins, Baltimore MD, 1986, pp. 1209–1234.
- [12] Lefebvre J., Introduction au Analyses Statistiques Multidimensionnelles, Masson, Paris, France, 1980.
- [13] Leveau J.Y., Bouix M., La Flore Lactique, in: Bourgeois C.M., Leveau J.Y. (Eds.), *Techniques d'analyses et de contrôle dans les industries agro-alimentaires*, Vol. 3, Tec et Doc Lavoisier, 1980, pp. 106–129.
- [14] Mills C.K., Lassel E.F. *Lactobacterium zeae* Kuznetsov, a later subjective synonym of *Lactobacillus casei* (Orla-Jensen) Hansen and Lassel, *Int. J. Syst. Bacteriol.* 23 (1973) 430–432.
- [15] Mori K., Yamazaki K., Ishiyama T., Katsumata M., Kobayashi K., Kawai Y., Inoue N., Shinano H., Comparative sequence analyses of the genes coding for 16S rRNA of *Lactobacillus casei*-related taxa, *Int. J. Syst. Bacteriol.* 47 (1997) 54–57.
- [16] Naumann D., Schultz P.C., Helm D., What can infrared spectroscopy tell us about the structure and composition of intact bacterial cells?, in: Mantsch H.H., Chapman D. (Eds.), *Infrared Spectroscopy of Biomolecules*, Wiley-Liss, 1996, pp. 279–310.
- [17] Schleifer K.H., Ehrmann M., Krusch U., Neve H., Revival of the species *Streptococcus thermophilus*, *Syst. Appl. Microbiol.* 14 (1991) 386–388.
- [18] Skerman V.B.D., McGowan V., Sneath P.H.A., Approved lists of bacterial names, *Int. J. Syst. Bacteriol.* 30 (1980) 225–420.
- [19] Wayne L.G., Action of the Judicial Commission of the International Committee on Systematic Bacteriology on requests for opinions published between January 1985 and July 1993, *Int. J. Syst. Bacteriol.* 44 (1994) 177–178.