



HAL
open science

Interaction of salmon gonadotropin subunits : spectroscopic studies

R. Salesse, Josselin Garnier, Bernard Breton

► **To cite this version:**

R. Salesse, Josselin Garnier, Bernard Breton. Interaction of salmon gonadotropin subunits : spectroscopic studies. *Annales de biologie animale, biochimie, biophysique*, 1978, 18 (4), pp.767-771. 10.1051/rnd:19780502 . hal-00897344

HAL Id: hal-00897344

<https://hal.science/hal-00897344>

Submitted on 11 May 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Interaction of salmon gonadotropin subunits : spectroscopic studies

par R. SALESSE, J. GARNIER, B. BRETON *

Laboratoire de Biochimie physique, I. N. R. A.
Département de Biochimie et de Biophysique,
Université de Paris-Sud 91405 Orsay Cedex

* Laboratoire de Physiologie des Poissons, I. N. R. A.
78350 Jouy en Josas, France

Summary. Pituitary gonadotropins of female and male pacific salmon *Oncorhynchus tshawytscha* were prepared separately. The two preparations exhibited different sedimentation coefficients (2.8 and 2.3, respectively) but similar circular dichroism spectra indicative of low α helix and high β sheet contents. Both hormones gave a difference spectrum ($1\ 750\ \text{M}^{-1}\ \text{cm}^{-1}$ at 287 nm) characteristic of perturbed tyrosine and phenylalanine residues when dissociated at acid pH. These results suggest that fish and mammal gonadotropins exhibit the same general folding of their polypeptide chains and undergo the same conformational transition when their subunits associate ; however, fish gonadotropin subunits reassociate at considerably faster rates.

Introduction.

The presence of two distinct gonadotropins in fish is still an open and interesting question. A single gonadotropin having both lutropin and follitropin activities was found in carp (Billard *et al.*, 1970 ; Burzawa-Gérard, 1971, 1974a ; Sundararaj *et al.*, 1976) and in salmon (Donaldson *et al.*, 1972), although physicochemical, biological or immunological evidence for two distinct gonadotropins has been presented (Idler *et al.*, 1975a ; Pierce *et al.*, 1976).

As those of mammalian origin, fish gonadotropins are composed of two dissimilar subunits (Donaldson *et al.*, 1972 ; Burzawa-Gérard, 1974b ; Burzawa-Gérard *et al.*, 1975) which can be dissociated in acid condition with loss of biological activity. This activity loss in mammalian hormones is accompanied by a specific conformational change of the subunits which is association-dependent (see review in Garnier, 1978). From experiments reported below, it is shown that two gonadotropin preparations, one from female salmon (s GTH₁), the other from male salmon (s GTH₂), also undergo the same reversible conformational transition during the association-dissociation process of their two subunits.

Material and methods.

The gonadotropins were prepared from separate female and male pituitary glands of pacific salmon *Oncorhynchus tshawytscha*, according to Idler *et al.* (1975a). Assayed *in vitro* for trout oocyte maturation (Jalabert *et al.*, 1974), both preparations exhibited the same potency and were two times more active than a sample obtained from Idler and prepared according to his method (Idler *et al.*, 1975b).

Protein concentrations were determined by a Lowry assay with bovine serum albumin as standard.

Ultracentrifuge experiments were carried out in a Spinco Beckman ultracentrifuge model E.




FIG. 1. — CD spectra of *sGTH*₁ (bottom) and *sGTH*₂ (above) assuming a mean molecular weight per residue of 141.

— : native hormone ;
 - - - : dissociated hormone (pH 2.2).

Circular dichroism spectra were recorded with a Jouan Dichrograph III ; light paths were 0.5 to 0.01 cm. Mean molecular weight per residue was assumed as 141.

Optical densities and U. V. difference spectra were recorded in a Cary 118 spectrophotometer with a 1 cm light path cuvette in a thermostated holder. Kinetics of dissociation and reassociation of hormones were followed at 287 nm.

Results and discussion.

The two preparations were found to differ essentially by their sedimentation coefficient ($S_{20,w}$) of 2.8 for s GTH₁ (0.5 mg/ml) and 2.3 for s GTH₂ (0.7 mg/ml) at neutral pH, 0.1 M NaCl. Both exhibited the same absorbance spectra with a maximum at 275-276 nm characteristic of tyrosine residue with no tryptophane contribution.

They had similar circular dichroism (CD) spectra (fig. 1) with minima at 280 nm, 210 nm (s GTH₂) or 208 nm (s GTH₁) and 193 nm; s GTH₁ had another small band at 234 nm. All these bands were already reported for mammalian gonadotropins and they are indicative of low α helix and high β sheet contents. These CD spectra suggest that the general gonadotropin folding has been conserved through evolution from fish to mammal.

Other interesting common features are the blue shift of the far U. V. CD, the decrease of the CD band at 280 nm (fig. 1) and the appearance of difference spectra ($1750 \text{ M}^{-1} \text{ cm}^{-1}$ at 287 nm) characteristic of perturbed tyrosine and phenylalanine residues (fig. 2) when the hormones are dissociated at acid pH. By raising the pH again to neutral, the two subunits reassociated with recovery of most of the native CD and loss of the difference spectrum. This suggests that salmon gonadotropins undergo the same conformational transition as mammalian gonadotropins when their two subunits associate.




FIG. 2. — UV difference spectrum of s-GTH₁ upon dissociation at pH 2.2
Reference : s-GTH₁ at pH 5.6.

Association-dissociation processes were followed with time (fig. 3); contrary to mammalian hormones, they were found to be much more rapid processes. For example, reassociation of sGTH at neutral pH from acid pH at 37 °C was 5 to 10 times faster than association of o-lutropin subunits or 20 to 50 times faster than h-choriogonadotropin reassociation (Pernollet *et al.*, 1976).




FIG. 3. — Change with time of the absorbance at 287 nm of s-GTH₁ during dissociation of the subunits at pH 2 (curve 1) and re-association at pH 6.4 (curve 2). Temperature 37 °C, 0.1 M NaCl. Dashed parts correspond to extrapolated absorbance change to zero time.

These observations strengthen the hypothesis that subunit association and subsequent conformational change are necessary steps *in vivo* for the formation of active gonadotropin.

*Symposium sur la Reproduction des Poissons
Paimpont, France, 19-21 septembre 1977.*

Acknowledgments. — We are very grateful to Dr. Idler for providing us with a sample of salmon gonadotropin for bioassays.

Résumé. A partir d'hypophyses du saumon *Oncorhynchus tshawytscha* femelle et mâle deux préparations de gonadotropine ont été obtenues. Elles ne diffèrent que par leur coefficient de sédimentation respectivement 2,8 et 2,3. Leurs spectres de dichroïsme circulaire sont semblables et indiquent une faible teneur en hélice α et une forte teneur en structure β . A pH acide, ces deux hormones se dissocient en leurs sous-unités, donnant naissance à un spectre de différence ($1750 \text{ M}^{-1} \text{ cm}^{-1}$ à 287 nm) caractéristique de résidus tyrosines et phénylalanines perturbées. Ces résultats suggèrent que les gonadotropines de poisson et de mammifère présentent la même conformation générale de leurs chaînes polypeptidiques et qu'elles subissent la même transition conformationnelle quand leurs sous-unités s'associent à la différence que les sous-unités de gonadotropine de poisson s'associent à des vitesses beaucoup plus grandes.

References

- BILLARD R., BURZAWA-GÉRARD E., BRETON B., 1970. Régénération de la spermatogenèse du cyprin hypophysectomisé (*Carassius auratus*) par un facteur gonadotrope hautement purifié de carpe. *C. R. Acad. Sci. Paris, Sér. D*, **271**, 1896-1899.
- BURZAWA-GÉRARD E., 1971. Purification d'une hormone gonadotrope hypophysaire de poisson téléostéen, la carpe (*Cyprinus carpio* L.). *Biochimie*, **53**, 545-552.
- BURZAWA-GÉRARD E., 1974a. Etude biologique et biochimique de l'hormone gonadotrope d'un poisson téléostéen, la carpe (*Cyprinus carpio* L.). *Mém. Mus. nat. Hist. nat., sér. A, Zoologie*, **86**, 1-77.
- BURZAWA-GÉRARD E., 1974b. Séparation et réassociation des sous-unités de l'hormone gonadotrope d'un poisson téléostéen, la carpe (*Cyprinus carpio* L.). *C. R. Acad. Sci. Paris, sér. D*, **279**, 1681-1684.
- BURZAWA-GÉRARD E., GONCHAROV B. F., FONTAINE Y. A., 1975. L'hormone gonadotrope hypophysaire d'un poisson chondrostéen, l'esturgeon (*Acipenser stellatus*). II. Propriétés biochimiques. *Gen. comp. Endocrinol.*, **27**, 296-304.
- DONALDSON E. M., YAMAZAKI F., DYE H. M., PHILLEO N. W., 1972. Preparation of gonadotropin from salmon (*Oncorhynchus tshawytscha*) pituitary glands. *Gen. comp. Endocrinol.*, **18**, 469-481.
- GARNIER J., 1978. Molecular aspects of the subunit assembly of the glycoprotein hormones. In McKERNS K. W., *The gonadotropins : structure and function*, Plenum Publ. Co., New York (in press).
- IDLER D. R., BAZAR L. S., HWANG S. J., 1975a. Fish gonadotropin(s). III. Evidence for more than one gonadotropin in chum salmon pituitary glands. *Endocr. Res. Commun.*, **2**, 237-249.
- IDLER D. R., BAZAR L. S., HWANG S. J., 1975b. Fish gonadotropin(s). II. Isolation of gonadotropin(s) from chum salmon pituitary glands using affinity chromatography. *Endocr. Res. Commun.*, **2**, 215-235.
- JALABERT B., BRETON B., BILLARD R., 1974. Dosage biologique des hormones gonadotropes de poissons par le test de maturation *in vitro* des ovocytes de truite. *Ann. Biol. anim. Bioch. Biophys.*, **14**, 217-228.
- PERNOLLET J.-C., GARNIER J., PIERCE J. G., SALESSE R., 1976. *In vitro* activation of glycoprotein hormones. Hybridization of subunits from thyrotropin, lutropin and human choriogonadotropin. *Biochim. biophys. Acta*, **446**, 262-276.
- PIERCE J. G., FAITH M. R., DONALDSON E. M., 1976. Antibodies to reduced S-carboxymethylated alpha subunit of bovine luteinizing hormone and their application to study the purification of gonadotropin from salmon (*Oncorhynchus tshawytscha*) pituitary glands. *Gen. comp. Endocrinol.*, **30**, 47-60.
- SUNDARARAJ B. I., NAYYAR S. K., BURZAWA-GÉRARD E., FONTAINE Y. A., 1976. Effect of carp gonadotropin on ovarian maintenance maturation and ovulation in hypophysectomized catfish, *Heteropneustes fossilis* (Bloch). *Gen. comp. Endocrinol.*, **30**, 472-476.