



**HAL**  
open science

## Peripheral plasma progesterone concentration in zebu (*Bos indicus*) cows during pregnancy

E. Mukasa-Mugerwa, Azage Tegegne

► **To cite this version:**

E. Mukasa-Mugerwa, Azage Tegegne. Peripheral plasma progesterone concentration in zebu (*Bos indicus*) cows during pregnancy. *Reproduction Nutrition Development*, 1989, 29 (3), pp.303-306. <hal-00899054>

**HAL Id: hal-00899054**

**<https://hal.science/hal-00899054v1>**

Submitted on 11 May 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire HAL, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



HAL Authorization

## Peripheral plasma progesterone concentration in zebu (*Bos indicus*) cows during pregnancy

E. Mukasa-Mugerwa and Azage Tegegne

*Animal Reproduction and Health Section, International Livestock Center for Africa (ILCA) P. O. Box 5689, Addis Ababa, Ethiopia*

(received 13 August 1988, accepted 21 April 1989)

**Summary** — Plasma progesterone concentrations were determined weekly during gestation averaging  $283 \pm 2$  d in Ethiopian zebu (*Bos indicus*) cows. Mean progesterone levels increased from  $0.2 \pm 0.1$  ng/ml at oestrus (service) to  $3.1 \pm 1.6$  ng/ml on d 7 and  $8.1 \pm 2.1$  ng/ml on d 21. Progesterone levels remained elevated throughout pregnancy. Hormone concentration differed significantly between cows ( $P < 0.001$ ) and with the wk of pregnancy ( $P < 0.05$ ); it tended to be higher during the last trimester of pregnancy. Mean levels dropped sharply to below 1 ng/ml during the last wk of pregnancy with considerable variation (C.V. = 39 to 63%) among cows. These data indicate that pregnancy in Ethiopian zebu cows can be reliably diagnosed by determining circulatory plasma progesterone levels.

**progesterone — zebu cow — pregnancy**

**Résumé** — *Evolution du taux de progestérone dans le plasma sanguin périphérique chez la vache zébu au cours de la gestation.* Les concentrations de progestérone plasmatique ont été déterminées chaque semaine au cours de la gestation ( $283 \pm 2$  jours) chez 17 vaches zébu éthiopiennes. Le niveau moyen de la progestérone s'élève depuis  $0,2 \pm 0,1$  ng/ml le jour de l'oestrus (J0) jusqu'à  $3,1 \pm 1,6$  ng/ml à J7 et  $8,1 \pm 2,1$  ng/ml à J21. Le niveau de la progestérone reste élevé pendant toute la gestation. Les concentrations d'hormone diffèrent significativement entre vaches ( $P < 0,001$ ) et selon la semaine de gestation ( $P < 0,05$ ); elles ont tendance à augmenter durant le dernier tiers de la gestation. Au cours de la dernière semaine de gestation les niveaux moyens chutent brutalement en-dessous de 1 ng/ml, avec de fortes variations (c.v. = 39 à 63%) entre vaches. Ces résultats indiquent que la gestation chez les vaches zébu éthiopiennes peut être correctement diagnostiquée par la mesure des niveaux de progestérone dans le plasma sanguin périphérique.

**progestérone — vache zébu — gestation**

## Introduction

Cattle in the tropics are predominantly of the zebu (*Bos indicus*) type. Most are raised traditionally and experience nutritional disease and thermal stresses that affect their reproductive efficiency. Although calving rates exceeding 70% have been reported in Zebu cows under modern and improved management (Rennie *et al.*, 1976; Eversbush, 1978), the fertility rates of animals raised traditionally seldom exceed 60% (Reed *et al.*, 1974; Rennie *et al.*, 1976; Butterworth and McNitt, 1984; Wilson, 1985). These disparities probably arise from prenatal reproductive wastage.

The concentration of progesterone in blood or milk can be determined by radioimmunoassay (RIA) or the enzyme-linked immunosorbent assay (ELISA) technique (Arnstadt and Cleere, 1981; Arnstadt and Schmidt-Adamopolou, 1982; Dinar and Sreenan, 1988). ELISA can be as accurate and reliable as RIA (Meyer, 1986; Sauer *et al.*, 1986; Magnin *et al.*, 1988) and is more accessible, cheaper and safer than RIA (Blake and Gould, 1984), and thus, well suited to use in developing countries. This study was undertaken to determine the progesterone concentration in the peripheral plasma of pregnant Ethiopian zebu cows using the ELISA technique in order to know whether it would help to monitor their reproductive efficiency.

## Materials and Methods

### *Animals and location of study*

The study used 17 pluriparous pregnant Ethiopian Highland zebu cows of the Small

East African zebu type. The cows were bred between April and July, 1987, and calved from January to April, 1988. Animals were located at the ILCA Debre Berhan experiment station, 120 km north of Addis Ababa, at an altitude of about 2 800 metres. Total station rainfall was 710 mm in 1987, of which 450 mm fell during the heavy rains season from May to October (August : 289 mm). The remainder fell as small showers from February to April (November, December, January : 0 mm). Mean monthly minimum temperature was 5.9 °C (range 2.1–8.5 °C). Mean monthly maximum temperature was 20.7 °C (range 18.5–22.1 °C). Cows grazed in fenced native pastures of *Andropogon*, *Festuca* and *Pennisetum* spp. grasses mixed with *Trifolium semense* legume during the day and had access to water *ad libitum*. They were penned overnight and supplemented to maintain body condition scores of 5 to 7 (Nicholson and Butterworth, 1986). Cows were mated naturally with a 3/4 Friesian x 1/4 Boran bull.

### *Blood collection and progesterone analysis*

Blood was taken weekly from the jugular vein of all cows until they were palpated *per rectum* to determine pregnancy 8 wk after copulation. Five of the pregnant cows were thereafter bled weekly until wk 38 or 39 of gestation. All cows were bled twice during the last wk of pregnancy and first wk after calving. Blood was collected in heparinised vacutainer tubes (Becton-Dickinson) and centrifuged within 15 min of collection to separate the plasma which was stored at -20 °C for subsequent progesterone assay. Plasma progesterone concentration was determined by the ELISA technique (Foulkes *et al.*, 1982; Parker *et al.*, 1988) using Ovucheck kits (Cambridge Veterinary Sciences). Procedural details of the technique were as *per* manufacturer's recommendation and have been reported previously (Mukasa-Mugerwa *et al.*, 1989). For plasma containing 8 ng/ml of progesterone, the intra- and inter-assay coefficients of variation for kits delivered in different batches have been assessed as 0.18 and 0.27, respectively. Data were analysed by least squares procedures (Harvey, 1977) using a model in which animal and calf sex were regarded as fixed effects and wk of gestation as a continuous variable. All the results are expressed as mean  $\pm$  SEM.

## Results

Mean length of gestation was  $283.3 \pm 1.8$  d ( $n = 17$ ). The associated average plasma progesterone is shown in Figure 1. Progesterone levels were elevated throughout gestation but differed significantly between cows (range :  $12.4 \pm 2.1$ — $7.4 \pm 1.9$  ng/ml,  $P < 0.001$ ).

Progesterone concentrations increased from  $0.2 \pm 0.1$  ng/ml at copulation (d 0) to  $3.1 \pm 1.6$  ng/ml on d 7 and  $7.1 \pm 1.9$  ng/ml on d14. Progesterone concentrations remained high ( $8.1 \pm 2.1$  ng/ml) until d 21, the day of the next expected oestrus, instead of declining as in non-pregnant animals. After conception, progesterone levels remained high and were significantly ( $P < 0.001$ ) elevated by the stage of gestation, with a tendency for higher levels in the last trimester of pregnancy.

Mean progesterone levels declined sharply prior to parturition, from  $8.8 \pm 3.49$  ng/ml 2 wks before calving to  $6.9 \pm 4.34$  ng/ml 1 wk before. Hormone values then fell to less than 1.0 ng/ml after calving, with considerable variation between cows (C.V. = 39 to 63%). Mean levels remained below 1.0 ng/ml during the first wk postpartum when sampling stopped. In the wk after calving there was little variability in progesterone levels between cows.

## Discussion

The 283 d average duration of gestation obtained was within the range of 270—292 d already established for zebu cows (Plasse *et al.*, 1968; Chandramohan and




Fig. 1. Mean ( $\pm$  SEM) progesterone concentrations in Ethiopian highland (zebu) cattle during pregnancy. Cows bred between April and July 1987.  $n = 17$  week 0—8 and 38—39;  $n = 5$  week 8—38.

Bhat, 1981) and taurine cows (Hunter, 1980).

The observation that, up to 15 d after copulation, all cows had progesterone values similar to those observed in the luteal phase of the oestrous cycle, agrees with Donaldson *et al.* (1970), Henricks *et al.* (1972) and Robertson (1972). Details of mean plasma progesterone levels during the spontaneous and induced oestrous cycle of Ethiopian cows are reported by Mukasa-Mugerwa *et al.* (1989). Our observation that progesterone levels did not decline at 15 d after oestrus but remained elevated 21 d after successful copulation, and throughout gestation, is consistent with data on Haryana (Agarwal *et al.*, 1980), Brahman (Shelton and Summers, 1983) and taurine (Melampy *et al.*, 1959; Henricks *et al.*, 1972; Robertson, 1972) cows. The observation that progesterone levels vary significantly with stage of gestation is nonetheless contrary to that of Agarwal *et al.* (1980) in Haryana cows. On the other hand, our observation of rapid declines in progesterone levels towards parturition agrees with previous reports (Donaldson *et al.*, 1970; Henricks *et al.*, 1972; Edquist *et al.*, 1973; Smith *et al.*, 1973). Progesterone levels under 1 ng/ml during the week after delivery were also observed by Coetzer *et al.* (1978) in Africander cattle. However, we failed to demonstrate the temporary rise in progesterone values which these authors reported during and shortly after parturition.

The corpus luteum is the major source of progesterone during pregnancy in the cow (Shelton and Summers, 1983). Bilateral ovariectomy prior to 200 d of gestation terminates pregnancy (Tanabe, 1966; Estergreen *et al.*, 1967). However, Estergreen *et al.* (1967) and Hoffmann *et al.* (1979) found that pregnancy was maintained if ovariectomy was performed

after 200 d of gestation. Wickersham and Tanabe (1967) found that the functional activity of the corpus luteum remained relatively constant during gestation and our results show that average progesterone levels were higher during the last trimester than before. It seems that 200 d after conception, an extraovarian source of progesterone exists in taurine cattle (Wendorf *et al.*, 1983), as well as in zebu cattle (our data). The placenta and adrenal glands, which are known to produce progesterone (Melampy *et al.*, 1959; Wendorf *et al.*, 1983), may be the additional source of this hormone.

We conclude that determination of circulatory progesterone can be used to diagnose pregnancy status in Ethiopian zebu cows. This would assist studies on the timing and magnitude of prenatal reproductive losses in these tropical cattle.

### Acknowledgements

The authors wish to thank Girma Abebe and Tekeste Tibebu for technical assistance, Solomon Zewde for data analysis and Paul Neate for editing the manuscript.

### References

- Agarwal S.P., Agarwal V.K. & Ahmad A. (1980) Studies on steroid hormones. III. Serum progesterone concentration in zebu cows during pregnancy. *Indian J. Anim. Sci.*, 50, 706-709
- Arnstadt K.I. & Cleere N.F. (1981) Enzymeimmunoassay for determination of progesterone in milk from cows. *J. Reprod. Fert.* 62, 173-180
- Arnstadt K.I. & Schmidt-Adamopolou B. (1982) Direct enzymeimmunoassay for determination

- of progesterone in milk from cows. *Br. Vet. J.* 138, 436-438
- Blake C. & Gould B.J. (1984) Use of enzyme immunoassay techniques : a review. *Analyst* 109, 533-547
- Butterworth M.H. & McNitt J.I. (1984) The Malawi zebu. *Wild Anim. Rev.* 49, 6-12
- Chandramohan K.K. & Bhat P.N. (1981) Note on factors affecting first gestation length in Sahiwal cattle. *Indian J. Anim. Sci.* 51, 788-790
- Coetzer W.A., van Niekerk C.H. & Morgenthal J.C. (1978) Hormone levels in peripheral plasma of the Afrikaner cow. II. Progesterone and total unconjugated oestrogen levels during late pregnancy and after parturition. *South Afr. J. Anim. Sci.* 8, 7-10
- Dinar A. & Sreenan J.M. (1988) The use of a rapid, high throughput enzymeimmunoassay for progesterone in a cow synchronisation programme. In : Proc. 11th Inter. Cong. Anim. Reprod. and A.I. 26-30 June 1988, Dublin, Ireland. No 137.
- Donaldson L.E., Bassett J.M. & Thorburn G.D. (1970) Peripheral plasma progesterone concentration of cows during puberty, oestrous cycles, pregnancy and lactation, and the effects of undernutrition or exogenous oxytocin on progesterone concentration. *J. Endocrinol.* 48, 599-614
- Edquist L., Ekman L., Gustafsson B. & Johansson E.D. (1973) Peripheral plasma levels of oestrogen and progesterone during late bovine pregnancy. *Acta Endocrinol.* 72, 81-88
- Estergreen V.L., Frost O.L., Gomes W.R., Erb R.E. & Bullard J.F. (1967) Effect of ovariectomy on pregnancy maintenance and parturition in dairy cows. *J. Dairy Sci.*, 50, 1293-1295
- Eversbusch G.C.J. (1978) Reproductive performance in a herd of Charolais cattle compared with that of the zebu (the Brahman breed) and its crosses in the tropics of Mexico. *Veterinaria Mex. City* 9, 225-226
- Foulkes J.A., Cookson A.D. & Sauer M.J. (1982) Artificial insemination of cattle based on daily enzyme immunoassay of progesterone in whole milk. *Vet. Rec.* 111, 302-303
- Harvey W.R. (1977) *User's guide for mixed model least-squares and maximum likelihood computer program (LSML 76)*. Ohio State University, OH
- Henricks D.M., Dickey J.F., Hill J.R. & Johnson W.E. (1972) Plasma oestrogen and progesterone levels after mating, and during late pregnancy and postpartum cows. *Endocrinology* 90, 1336-1342
- Hoffmann B., Wagner W.C., Hixon J.E. & Bahr J. (1979) Determination concerning the functional status of the corpus luteum and the placenta around parturition in the cow. *Anim. Reprod. Sci.* 2, 253-266
- Hunter R.H.F. (1980) *Physiology and Technology of Reproduction in Female Domestic Animals*. Academic Press. London. p. 393
- Magnin M., Humblot P., Staples M., Khalil M., Thibier M. & Somme G. (1988) Accuracies of pregnancy diagnosis by a progesterone RIA and direct enzymeimmunoassay (Bovitest). Proc. of the 11th Inter. Cong. Anim. Reprod. and A.I., 26-30 June 1988, Dublin, Ireland. No 104
- Melampy R.M., Hearn W.R. & Rakes J.M. (1959) Progesterone content of bovine reproductive organs and blood during pregnancy. *J. Anim. Sci.* 18, 307-313
- Meyer H.H.D. (1986) Possible ways of improving enzymeimmunoassay (EIA) techniques and their application in animal production. Proc. Symp. on Nuclear and related techniques in animal production and health. 17-21 March 1986. Int. Atomic Energy Agency, Vienna. pp. 255-262
- Mukasa-Mugerwa E., Azage T., Mattoni M. & Cechini G. (1989) Effect of oestrous synchronisation with prostaglandin F<sub>2</sub> alpha in Ethiopian highland zebu (*Bos indicus*) cows. *Anim. Prod.* 48, 367-373
- Nicholson M.J. & Butterworth M.H. (1986) *A guide to condition scoring of Zebu cattle*. ILCA. Addis ababa, Ethiopia. p. 29
- Parker B.N.J., Foulkes J.A., Jones P.C., Dexter I. & Stephens H. (1988) Prediction of calving times from plasma progesterone concentration. *Vet. Rec.* 122, 88-89
- Plasse D., Warnick A.C., Reese R.E. & Koger M. (1968) Reproductive behaviour of *Bos indicus* females in a subtropical environment. II. Gestation length in Brahman cattle. *J. Anim. Sci.* 27, 101-104
- Reed J.B.H., Doxey D.L., Forbles A.B., Finlay R.S., Geering I.W., Smith S.D. & Wright J.D. (1974) Productive performance of cattle in Botswana. *Trop. Anim. Health Prod.* 6, 1-21
- Rennie T., Light D., Rutherford A., Miller M., Fischer I., Pratchett D., Capper B., Buck N. & Trail J.C.M. (1976) Beef cattle productivity under traditional and improved management in Botswana. *Trop. Anim. Health Prod.* 9, 1-6

- Robertson H.A. (1972) Sequential changes in plasma progesterone in the cow during the oestrous cycle, pregnancy, at parturition and postpartum. *Can. J. Anim. Sci.* 52, 645-658
- Sauer M.J., Foulkes J.A., Worsfold A. & Morris B.A. (1986) Use of progesterone 11-glucuronide-alkaline phosphatase conjugate in a sensitive microtitre-plate enzyme immunoassay of progesterone in milk and its application to pregnancy testing in dairy cattle. *J. Reprod. Fert.* 76, 375-391
- Shelton J.N. & Summers P.M. (1983) Progesterone and oestradiol-17beta in the peripheral plasma of Brahman cows with single and twin pregnancies. *Theriogenology* 20, 47-52
- Smith V.G., Edgerton L.A., Hafs H.D. & Convey E. M. (1973) Bovine serum oestrogens, progestins and glucocorticoids during late pregnancy, parturition and early lactation. *J. Anim. Sci.* 36, 391-396
- Tanabe T.Y. (1966) Essentiality of the corpus luteum for maintenance of pregnancy in dairy cows. *J. Dairy Sci.* 49, 731, abstr.
- Wendorf G.L., Lawyer M.S. & First N.L. (1983) Role of the adrenals in the maintenance of pregnancy in cows. *J. Reprod. Fert.* 68, 281-287
- Wickersham E.W. & Tanabe T.Y. (1967) Functional status of bovine corpora lutea of pregnancy. *J. Dairy Sci.* 50, 1001, abstr.
- Wilson R.T. (1985) Livestock production in central Mali : reproductive aspects of sedentary cows. *Anim. Reprod. Sci.* 9, 1-9