



**HAL**  
open science

# Serum lipid and protein profiles in eutrophic or hypotrophic newborn at term and in their mothers. H Merzouk 1

H Merzouk, M Meghelli-Bouchenak, N Meghelli, N Korso, K Kerzabi, J Belleville

► **To cite this version:**

H Merzouk, M Meghelli-Bouchenak, N Meghelli, N Korso, K Kerzabi, et al.. Serum lipid and protein profiles in eutrophic or hypotrophic newborn at term and in their mothers. H Merzouk 1. *Reproduction Nutrition Development*, 1992, 32 (5-6), pp.509-509. hal-00899565

**HAL Id: hal-00899565**

**<https://hal.science/hal-00899565v1>**

Submitted on 11 May 2020

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.


L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

### Serum lipid and protein profiles in eutrophic or hypotrophic newborn at term and in their mothers.

H Merzouk <sup>1</sup>, M Meghelli-Bouchenak <sup>2</sup>, N Meghelli <sup>2</sup>, N Korso <sup>3</sup>, K Kerzabi <sup>3</sup>, J Belleville <sup>4</sup> (<sup>1</sup> *Université de Tlemcen, Laboratoire de Physiologie Animale*; <sup>2</sup> *Institut des Sciences de la Nature, Université d'Oran, Laboratoire de Physiologie Animale et de la Nutrition, Oran*; <sup>3</sup> *Centre Hospitalo-Universitaire de Tlemcen, Service de Maternité, Tlemcen, Algeria*; <sup>4</sup> *Université de Bourgogne, Faculté des Sciences Mirande, Unité de Recherches de Nutrition Cellulaire et Métabolique, Dijon, France*)

This study was conducted in the Tlemcen region (Algeria) on 38 hypotrophic newborn ( $2290 \pm 30$  g), 20 eutrophic newborn ( $3570 \pm 93$  g) and their mothers. The effects of hypotrophy on serum lipid and protein amounts were studied in the newborn. The same investigation was carried out in mothers at delivery. Blood samples were taken from the umbilical vein immediately after delivery and cutting the umbilical cord. Blood was obtained from the mothers just after placental expulsion by venous puncture of the elbow pleat.

Cause analysis showed that primiparity, malnutrition in the mother and pregnancy were independent factors responsible for the increased hypotrophic risk in the Tlemcen population. In mothers of hypotrophic newborn compared with those of eutrophic newborn, lower levels of total



**Fig 1.** Each value represents mean  $\pm$  SEM. Comparison of eutrophics ( $n = 20$ ) versus hypotrophics ( $n = 38$ ) was performed using Student's test; \*  $P < 0.01$ ; \*\*  $P < 0.001$ ; \*\*\*  $P < 0.0001$ .

lipids, triacylglycerol, phospholipid, total protein and total globulin were observed as well as an increase in  $\gamma$  globulins.

In hypotrophic compared with eutrophic newborn a significant increase in triacylglycerol, apo-B and VLDL levels was noted. On the other hand, hypotrophy induced a decrease in cholesterol ester, phospholipid, protein, apo A1 amounts, relative values of HDL and LDL and the albumin/globulin ratio.

In conclusion, hypotrophy involved lipid and protein profiles similar to those of marasmic children. The hypotrophic newborn at term should be considered as a previously marasmic child.