



**HAL**  
open science

# PATHOGENESIS AND BIOLOGY OF ANOPLOCEPHALINE CESTODES OF DOMESTIC ANIMALS

Vs Narsapur

► **To cite this version:**

Vs Narsapur. PATHOGENESIS AND BIOLOGY OF ANOPLOCEPHALINE CESTODES OF DOMESTIC ANIMALS. *Annales de Recherches Vétérinaires*, 1988, 19 (1), pp.1-17. hal-00901779

**HAL Id: hal-00901779**

**<https://hal.science/hal-00901779>**

Submitted on 11 May 2020

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

## **PATHOGENESIS AND BIOLOGY OF ANOPLOCEPHALINE CESTODES OF DOMESTIC ANIMALS**

VS NARSAPUR

Department of Parasitology, Bombay Veterinary College, Parel, Bombay-400 012, India

### **Plan**

#### Introduction

#### *Pathogenesis of Anoplocephaline cestodes*

#### *Biology of Anoplocephaline cestodes*

#### Developmental stages in oribatid hosts

#### Conditions of development

#### *Oribatid intermediate hosts*

#### Oribatid species as intermediate hosts

#### Oribatid host specificity

#### Alternative intermediate hosts

#### *Biological phenomena in nature*

#### Conclusions

Adult tapeworms parasitic in domestic herbivorous animals, exclusively belong to family Anoplocephalidae and are common in many countries. Yamaguti (1959) has recognized 23 species under seven genera in these hosts. This group has received good attention of helminthologists all over the world. Considerable literature has been accumulated since last few years and the need for a review article on the topic is felt necessary. In the present paper, literature on pathogenicity and biology of Anoplocephaline cestodes of domestic animals is reviewed and up-to-date bibliography available to the author is presented. Most of the Russian work was available only in the form of abstracts in English but original references are cited in the bibliography for the better use of readers.

### **Pathogenesis of Anoplocephaline cestodes**

There is a wide divergence of opinion regarding the pathogenic effects of these tapeworms in cattle, sheep and goats. Hawkins (1946), Kates and

Goldberg (1951) did not notice any observable injurious effects nor any significant retardation of growth in lambs heavily infected, experimentally with *Moniezia expansa*. Haematological studies by Deshpande *et al* (1980b) did not show any alteration in the values of haemoglobin, packed cell volumes and erythrocyte counts during prepatency of experimental monieziasis. But many of the Russian workers have noted high degree of pathogenicity, and adverse effects on weight gains and on yields of meat and wool. In lambs, Tableman (1946) recorded cases of convulsions and death and Hansen *et al* (1950) retarded weight gains and anaemia due to pure *Moniezia* infections.

Stampa (1967) found that in the lambs treated for *Moniezia sp* the weight gain was 109.8 g per day during first four weeks of age as against 69.4 g per day in untreated group. Afonina (1967) found weight gain of 10.2 kg in three months in treated lambs as against 4.2 kg in untreated ones.

Lyashenko and Teplov (1974) found that in sheep infected for three months, the wool yield per animal was less by 0.84 kg and meat yield by 4.11 kg as compared to uninfected sheep. They also recorded 43.5 % mortality in lambs due to monieziasis. Efner (1974) also found that the carcass yield in the infected Polish long wool sheep was less by 8.10 kg per carcass than in non-infected group.

In trials of anthelmintic treatment for monieziasis on several farms, Shakiev (1973) found that mortality rate was completely checked in treated farms while it remained at 3 and 0.8 % in lambs and ewes respectively of untreated farms. He further, noticed that wool yield was better and more lambs were born in treated farms.

In very extensive trials on 600 000 sheep, Vibe (1976) found that anthelmintic dosing against *Avitellina sp*, and *Moniezia sp*, brought down the mortality by 12.4 times. This also reduced mortality due to enterotoxaemia by 11.7 times thereby showing that high incidence of enterotoxaemia is associated with tapeworm infections in lambs.

The exact modes of pathogenicity in Anoplocephalines is still not fully known. The possibility of toxic effect of excretions and secretions of the parasite is indicated by Shakurova (1974) who showed that 50 to 150 mg of *M expansa* antigen injected intra-peritoneally into mice resulted in toxicity and death.

The injury caused to the intestinal mucosa also contributes to the pathogenicity as scolex of *M expansa* is buried deep into the mucosa (Casarosa 1964) and proliferative inflammation, nodulations, deeply inserted scolices, disquamation of epithelium and cellular infiltration in the jejunum is described in massive infections of *Stilesia globipunctata* (Bankov 1971, Amjadi 1971). Yet another explanation for the pathogenicity could be absorption of enormous amount of food by the parasite. Narsapur (1976c) found that during the period of 17 days (30th to 47th days) of prepatency in lambs *M expansa* grows from 2.2 to 195.0 cm in length and opined that for achieving such gigantic growth the parasite must be utilising enormous amount of food available in the intestine which otherwise would have been utilized by the host.

Soulsby (1982) concluded that light infections are of little importance and that as a rule only lambs, kids and calves under six months are substantially affected. However, from the evidence available it can reasonably be assumed that Anoplocephaline infections are of considerable economic importance for the animal industry.

**Biology of Anoplocephaline cestodes**

**1.- Developmental stages in oribatid hosts**

*Moniezia expansa* is known since 1810 as a common tapeworm parasite of cattle, sheep and goats, but the knowledge regarding its life history and mode of transmission remained a mystery for a very long time. In 1890, Curtice, initiated serious studies in this direction and after a series of speculative experiments conducted by several research workers over a period of 47 years, the opinion had centered on the possibility of involvement of an intermediate host living on the pastures. The number of observations on epidemiology of *M expansa* made by Stoll (1935a, 1935b, 1936, 1937a, 1937b) contributed substantially to this belief although he himself did not appear to have been convinced then. It was Stunkard (1937a, 1937b) who made the epoch making discovery that an oribatid mite was the long searched intermediate host of *M expansa*, an anoplocephalid tapeworm.

The developmental process of *M expansa* in oribatid hosts together with morphology of stages has first been described by Stunkard (1938) follo-

wed by many workers. A general account that emerges out from these studies indicates that the pattern of development and morphology of larval stages is similar in all the species of Anolanoplocephaline cestodes. Oncosphere which is found in body cavity of mites within 48 h after their exposure to tapeworm eggs, undergoes a continuous




Fig 1. - Camera lucida drawings of developmental stages of *Moniezia expansa* in *Schelorbates laevigatus*.

- 1 : oncosphere
- 2, 3 : oblong stage
- 4, 5 : elongate stage
- 6 : stage with body and cercomere
- 7, 8 : stage with scolex, body and cercomere
- 9 : cysticercoid

process of development without any abrupt changes in their morphology demarcating different stages. However, following six morphological stages can be recognized before the infective stage (ie cysticeroid) is reached (Narsapur 1976c) (fig 1) :

1. Oncosphere :  $22.9 \mu \times 21.3 \mu$  in size with well arranged six embryonal hooks each  $7.18 \mu$  in length.
2. Oblong stage :  $103 \mu \times 79.33 \mu$  in size with embryonal hooks irregularly arranged.
3. Elongate stage :  $158.3 \mu$  in length, wider at ( $84.6 \mu$ ) one end and narrow at the other.
4. Stage with body and cercomere : body is sub-spherical, measures  $134.6 \mu \times 100 \mu$  and bears tail like cercomere  $100 \mu$  in length. Few embryonal hooks present on the tail.
5. Stage with scolex body and cercomere : the body is further divided by transverse constriction to anterior and middle parts with cercomere remaining as posterior part. At the anterior part there is accumulation of calcareous bodies and aggregation of cells indicating developing suckers. Few embryonal hooks may still be present on the tail.
6. Cysticeroid : the stage results by retraction of anterior part (scolex) into the middle part with sides closing on to from cyst wall. After slight shrinkage, the cyst wall becomes rigid. Cercomere persists as an appendage with few embryonal hooks still present. The scolex exhibits active movements.

The size of cysticeroid is subject to great variation depending upon the number of cysticeroids in the body cavity of the mite. Thus Narsapur (1976c) found smaller ( $110 \times 100 \mu$ ) cysticeroids of *M expansa* in mites bearing many number and larger ( $175 \times 140 \mu$ ) ones in mites bearing a single cysticeroid.

The position of embryonal hooks on different stages shows that there is a reversal of antero-posterior axis of the larvae as it transforms from oncosphere to cysticeroid (Stunkard 1938, 1941, Narsapur 1971).

Narsapur (1976c) further observed that *M expansa* oncospheres remained dormant in the body cavity of the mites up to 12 days after infection and the development then on started at a very uneven pace, so that during this phase larvae under various stages of development were present in the body cavity of the same mite. He cited that in a mite dissected after 57 days of the infection, all the stages from oncosphere to cysticeroid were found. Similar observations have been made earlier by Stunkard (1938, 1940, 1941), Anantaraman (1951) and Freeman (1952) with regard to different species of Anoplocephalines. Stunkard (1941) opined that this was due to influencing factors such as mite vector, number of larvae in

haemocoel, age and development of larvae. Freeman (1952) stated that as most of the mites in experimental condition pick up the infection within 24 to 48 h, the above variations cannot be attributed to delayed pick up of the infection by mites.

It therefore appears, that more than anything else, larvae in the body cavity themselves exert a suppressive effect on each other, which triggers uneven development and it further accentuates as larvae in advanced stage exert more and more suppressive effect on less advanced larvae. Some of these suppressed larvae might reach cysticeroid stage belatedly and many of them die out as is shown in experimental infection with *M expansa* by Narsapur (1976c) that mites dissected on 30 th day revealed 0.2 to 0.3 cysticeroids per mite along with many earlier stages while the same batches dissected on 57th day revealed 2.2 to 7.0 cysticeroids per mite unaccompanied by earlier stages.

## 2. Developmental conditions

The developmental period in oribatid hosts varies greatly and this appears to be mainly, inversely influenced by temperature conditions, although, Stunkard (1938) mentions the mite species as an additional factor. The development of oncospheres in cysticeroids is performed in the best conditions at  $25^\circ\text{C}$ . At this optimum the formation of infesting larvae is achieved in three months (Euzeby 1966). In warmer countries (between  $27$  and  $33^\circ\text{C}$ ) the development is accelerated, that is about from one to three months. In colder countries (between  $18$  and  $20^\circ\text{C}$ ) evolution in oribatids is slower and needs about from three to five months (refer tables 1a, 1b). However, the prepatent period in vertebrate hosts appears to be rather uniform as  $45 \pm 7$  days in *Moniezia sp* is reported by many workers (Shorb 1939, Anantaraman 1951, Potemkina 1951, Mehra and Srivastava 1955 a b, Kuznetsov 1968, Davydov and Smirnov 1972, Alkov 1971, 1972, Gapon 1974, Worley *et al* 1974, Narsapur 1976c, Kramnoi 1980).

A patent period of 154 days (Kuznetsov 1968), 132 to 256 days (Davydov and Smirnov 1972) have been reported in case of *M expansa* but spontaneous expulsion of the infection due to acquisition of immunity (self-cure) takes place within two to three months of patency which explains why infection in lambs occur at two to five months and is rare after six months of age (Worley *et al* 1974, Gapon 1974, Lin *et al* 1975).

## Oribatid intermediate hosts

### 1. Species of oribatids as intermediate host

There are many reports published from different

Tableau 1a – Summary of literature of relationship between temperature and developmental period of *Moniezia expansa*

Intermediate host Period of development (days)	Temperature (°C)	Reference
<i>Scheloribates laevigatus</i>		
27	28 a	Narsapur and Prokopic (1979)
45	20	Skorski <i>et al</i> (1984)
49-57	27-33	Narsapur (1976c)
75	27	Kuznetsov (1970)
97	18-20 a	Narsapur and Prokopic (1979)
150	18-20 b	Jurasek (1962)
<i>Scheloribates latipes</i>		
27	28 a	Narsapur and Prokopic (1979)
77-79	27	Nazarova (1970)
85-98	23	Nazarova (1970)
97	18-20 a	Narsapur and Prokopic (1979)
<i>Scheloribates fimbriatus</i>		
49-57	27-33	Narsapur (1976c)
<i>Scheloribates semidesertus</i>		
77-79	27	Nazarova (1970)
85-98	23	Nazarova (1970)
<i>Galumna sp</i>		
150	18-20 b	Jurasek (1962)
<i>Pilogalumna tenuiclavus</i>		
92	20	Skorski (1984)
<i>Archipteria sp</i>		
150	18-20 b	Jurasek (1962)
<i>Platinothrus peltifer</i>		
27	28 a	Narsapur and Prokopic (1979)
97	18-20 a	Narsapur and Prokopic (1979)
<i>Liacarus sp</i>		
150	18-20 b	Jurasek (1962)
<i>Liacarus coracinus</i>		
27	28 b	Narsapur and Prokopic (1979)
97	18-20 b	Narsapur and Prokopic (1979)
<i>Leibstadia similis</i>		
35	20	Skorski <i>et al</i> (1984)
65	27	Skorski <i>et al</i> (1984)

a: at 85 % of humidity  
b: at 100 % of humidity

countries on the oribatid intermediate hosts of Anaplocephalids and the list is growing continuously. The present position is that 53 species of oribatids are listed as intermediate hosts of *M. expansa* and 31 for *M. benedeni*.

Information on *Avitellina sp* is very scanty because many experimental infections have ended in failure due to peculiarities in the morphology of the uterus of this cestode (Matevosyan 1978). However, Kuznetsov (1965, 1966) experimentally infected adults and larvae of insects of the order

*Psocoptera* viz, *Lachesilla pedicularia* and *L. pedicularia* var *brevipennis* with *Avitellina sp* and obtained cysticercoids. Nadakal (1960a, 1960b) exposed thousands of *Protoschelobates sp* to infection of *Avitellina centripunctata* and obtained oncospheres in only five of them. Narsapur (1974b) successfully infected *Scheloribates laevigatus* and *S. fimbriatus* to eggs of *Avitellina lahorea* and found cysticercoids, on 50th and 57th days. In *S. laevigatus*, 3 of 91 mites and in *S. fimbriatus* 3 of 158 mites showed larval development, with one cysticercoid per mite.

Tableau 1b – Summary of literature of relationship between temperature and developmental period of *Moniezia benedeni*

Intermediate host Period of development (days)	Temperature (°C)	Reference
<i>Scheloribates laevigatus</i>		
87	12-26	Kramnoi (1973)
150	18-20	Prokopic (1962a)
<i>Scheloribates latipes</i>		
34	28 a	Prokopic and Narsapur (1981)
77-79	27	Nazarova (1970)
85-98	23	Nazarova (1970)
87	12-26	Kramnoi (1973)
<i>Scheloribates semidesertus</i>		
77-79	27	Nazarova (1970)
85-98	23	Nazarova (1970)
<i>Galumna eliminata</i>		
150	18-20	Prokopic (1962a)
<i>Trichoribates trimaculatus</i>		
150	18-20	Prokopic (1962a)
<i>Trichoribates novus</i>		
34	28 a	Prokopic and Narsapur (1981)
<i>Ceratozetes sp</i>		
87	12-26	Kramnoi (1973)
<i>Punctoribates punctum</i>		
87	12-26	Kramnoi (1973)
<i>Archipteria coleoptera</i>		
150	18-20	Prokopic (1962b)
<i>Zygoribatula skrjabini</i>		
77-79	27	Nazarova (1970)
85-98	23	Nazarova (1970)
<i>Patinothrus peltifer</i>		
34	28 a	Narsapur and Prokopic (1981)
<i>Oribatula sp</i>		
162	17-20	Cai and Jin (1984)

a: at 85 % of humidity

Psocopteran insects are intermediate hosts also for *Thysaniezia sp* (Allen 1959, Kuznetsov 1965, 1966) in addition to seven species of oribatids, seven species of oribatid mites are reported intermediate hosts of *Stilesia globipuncta* two species of *Anoplocephala magna*, 18 of *A perfoliata*, 6 of *Paranoplocephala variabilis*, 4 of *P ryjkovi* and 2 of *P manilana*, in many of the papers the parasites and their hosts are mentioned only by generic name or by family name and these have not been included in the present list.

Apart from the adult stages of oribatids, the possibility of involvement of their juvenile stages in the transmission of tapeworms was investigated

by Narsapur (1979). He found that the nymphal stage of oribatids pick up the tapeworm infection in the same way as adults and that the tapeworm larvae survive the moult in the body of juvenile oribatids reaching cysticeroid stage by the time juvenile become adult oribatids. This has significance in the epizootology since juvenile oribatids occur abundantly on pastures during prepeak seasons of oribatid mite populations.

## 2. Oribatid host specificity

In view of the growing list of oribatid species reported as intermediate hosts of anoplocephalids, two divergent views existed on the oribatid host

Table 2a – Summary of literature on *Scheloribates* as intermediate hosts of *Moniezia expansa* in different countries

Intermediate host Country	References
<i>Scheloribates laevigatus</i>	
USSR	Potemkina (1941)
USA	Kates and Runkel (1948)
Canada	Rao and Choquette (1951)
USSR	Orhekov (1960)
USSR	Sokolova and Panin (1960)
Czechoslovakia	Prokopic (1962a)
Czechoslovakia	Jurasek (1962)
Bulgaria	Bankov (1965)
USSR	Alkov (1971)
China	Lin <i>et al</i> (1975)
India	Narsapur (1976c)
Czechoslovakia	Narsapur and Prokopic (1979)
GFR	Skorski <i>et al</i> (1984)
USSR	Kuznetsov (1970)
<i>Scheloribates latipes</i>	
USSR	Potemkina (1948)
USSR	Orhekov (1960)
Czechoslovakia	Prokopic (1962)
USSR	Nazarova (1970)
USSR	Alkov (1971)
Czechoslovakia	Narsapur and Prokopic (1979)
<i>Scheloribates pallidulus</i>	
Bulgaria	Bankov (1965)
<i>Scheloribates fimbriatus africanus</i>	
Chad	Graber and Gruvel (1969)
<i>Scheloribates fimbriatus</i>	
India	Narsapur (1976c)
<i>Scheloribates perforatus</i>	
Chad	Graber and Gruvel (1969)
<i>Scheloribates madrasensis</i>	
India	Anantaraman (1951)
<i>Scheloribates semidesertus</i>	
USSR	Nazarova (1970)
<i>Scheloribates chauhani</i>	
China	Lin <i>et al</i> (1975)
<i>Scheloribates sp</i>	
Hungary	Kassai and Mahunka (1965)

specificity since long time. Kates and Runkel (1948), Potemkina (1951) considered that susceptibility of different species of mites varied greatly but Kassai and Mahunka (1964, 1965) believed that there was no high degree of host specificity in different oribatids and that any species larger than 300 to 400 microns in length had the ability of acting as intermediate host. However, the scanning of world literature (table 2a, 2b, 2c, 3a, 3b, 3c) shows that *Scheloribates sp* followed by *Galumna sp* have been more often incriminated as intermediate hosts of *M expansa* and *M benedeni* in several countries and by this fact alone they could be more specific interme-

mediate hosts of the two tapeworms than other species of oribatid mites. Intermediate hosts of *Avitellina sp*, *Stilesia globipunctata* and *Thysaniezia* reported in several countries have been summarized in tables 4 and 5.

Intermediate hosts of *Anoplocephala perfoliata*, observed in USSR, are : *Galumna obivius*, *Galumna nervosus*, *Scheloribates laevigatus*, *Scheloribates latipes*, *Carbodidae*, *Archipteria sp*, *Liacaridae*, *Platinothrus pelfiter*, *Hermaniella granulata*, *Eremaeus oblongus*, *Parachipteria punctata*, *Ceratozetes bulanovae*, *Trichoribates incisellus*, *Galumna diforma*, *Urubambates schachtachtiskoi*, *Zygoribatula*

Table 2b – Summary of literature on *Protoschelobates* and *Galumna* as intermediate hosts of *Moniezia expansa* in different countries

Intermediate host Country	References
<i>Protoschelobates</i> sp	
India	Nadakal (1960)
<i>Protoschelobates seghettii</i>	
USA	Runkel and Kates (1947)
<i>Protoschelobates curtipes</i>	
USA	Runkel and Kates (1947)
<i>Galumna</i> sp	
USA	Stoll (1938)
USA	Stunkard (1937, 1938)
India	Anantaraman (1951)
Czechoslovakia	Jurasek (1962)
Bulgaria	Bankov (1965)
USSR	Alkov (1971)
<i>Galumna elimata</i>	
Czechoslovakia	Prokopic (1962a)
<i>Galumna nigra</i>	
USA	Stoll (1938)
<i>Galumna obvia</i>	
USSR	Potemkina (1941)
USSR	Alkov (1971)
<i>Galumna longipluma</i>	
China	Lin <i>et al</i> (1975)
<i>Galumna virginienensis</i>	
USA	Kates and Runkel (1948)
USA	Edney and Kelly (1953)
China	Lin <i>et al</i> (1975)
<i>Galumna curvum</i>	
China	Lin <i>et al</i> (1975)
<i>Galumna emarginata</i>	
USA	Krull (1939)
USA	Kates and Runkel (1948)
<i>Galumna flagellata</i>	
USSR	Orekhov (1960)
<i>Galumna type minor</i>	
USSR	Orekhov (1960)

*tenicola*, *Zygoribatula microporosa*, *Leibstadia similis*, and four unspecified oribatid mites (Bashkirova 1941, Kuliev 1963, Urmanbetova 1979).

Intermediate hosts of *Anoplocephala magna*, observed in USSR, are : *Scheloribates laevigatus*, *Scheloribates latipes* and four unspecified oribatid mites (Bashkirova 1941, Urmanbetova 1979).

Intermediate hosts of *Paranoplocephala mamillana*, observed in URSS, are : *Galumna obvia* and *Allogalumna longipluma* (Bashkirova 1941).

Intermediate hosts of *Paranoplocephala variabilis* are : *Galumna* sp, *Galumna curvum*, *Galumna virginienensis*, *Scheloribates laevigatus*, *Oribatella quadricarnata* and *Platynothrus peltifer* (Gleason and Buckner 1979).

Intermediate hosts of *Paranoplocephala ryjikovi*, observed in China, are : *Scheloribates chauhani*, *Galumna virginienensis*, *Scheloribates* sp and *Paralumna lydia* (Lin *et al* 1982).

Two criteria viz percentage of mites infected and



Table 2c – Summary of literature on *Pergalumna*, *Pilogalumna*, *Scutovetex*, *Oribatula*, *Ceratozetes*, *Trichoribates*, *Punctoribates* and *Protoribates* as intermediate hosts of *Moniezia expansa* in different countries.

Intermediate host Country	References
<i>Pergalumna nervosa</i>	
Bulgaria	Bankov (1961)
<i>Pergalumna sp</i>	
Hungary	Kassai and Mahunka (1965)
<i>Pilogalumna tenuiclavus</i>	
GFR	Skorski (1984)
<i>Scutovetex minutus</i>	
UK	Morgan (1947)
UK	Rayski (1947)
<i>Oribatula minuta</i>	
USA	Kates and Runkel (1948)
<i>Ceratozetes mediocris</i>	
Hungary	Kassai and Mahunka (1965)
<i>Trichoribates trimaculatus</i>	
Czechoslovakia	Prokopic (1962)
USSR	Alkov (1971)
<i>Trichoribates novus</i>	
Bulgaria	Bankov (1965)
<i>Trichoribates incisellus</i>	
USSR	Alkov (1971)
<i>Punctoribates sp</i>	
USSR	Potemkina (1951)
<i>Punctoribates punctum</i>	
Bulgaria	Bankov (1965)
USSR	Alkov (1971)
<i>Punctoribates hexagonus</i>	
USSR	Sokolova and Panin (1960)
<i>Protoribates lophotrichus</i>	
USSR	Rukarina <i>et al</i> (1960)

average number of cysticercoids per mite could be applied to either naturally infected mites or to the experimentally exposed ones to determine relative host specificity. Narsapur (1976a) compared the performances of *S laevigatus* and *S fimbriatus* by these criteria after exposing them to uniform suspension of the eggs of *M expansa* and *M benedeni* and found that for *M expansa*, *S laevigatus* was more specific intermediate host than *S fimbriatus* and for *M benedeni* it was reverse.

Thus, in the epidemiological consideration of Anoplocephaline infections, it is necessary to recognize the variations in the host specificity of oribatid mites and determine the predominant and

highly host specific oribatid species occurring in the area. In general, it may be said that wherever, *Scheloribates sp* and *Galumna sp* predominate in the soil, the area is vulnerable for outbreaks of Anoplocephaline infections.

### 3. Alternative intermediate hosts

Since the discovery of life cycle of *Moniezia expansa* by Stunkard in 1937a, a large number of papers have been published on the biology of Anoplocephalid tapeworms from which two generalizations can be made viz

Table 2d – Summary of literature on *Archipteria*, *Adoristus*, *Platynothrus*, *Eremaeus*, *Ceratoppia*, *Liacarus*, *Peloptulus*, *Unguizetes*, *Zygoribatula*, *Leibstadia*, *Hypozetes* and *Spatiodaemaeus* as intermediate hosts of *Moniezia expansa* in different countries

Intermediate host Country	References
<i>Archipteria</i> sp	
USSR	Potemkina (1951)
Czechoslovakia	Jurasek (1962)
<i>Archipteria coleoptera</i>	
Czechoslovakia	Prokopic (1962b)
<i>Adoristus ovatus</i>	
USSR	Potemkina (1951)
<i>Platynothrus peltifer</i>	
USSR	Shaldibina (1953)
Czechoslovakia	Narsapur and Prokopic (1979)
<i>Eremaeus hepaticus</i>	
USSR	Shaldibina (1953)
<i>Ceratoppia bipitis</i>	
USSR	Shaldibina (1953)
<i>Liacarus</i> sp	
Czechoslovakia	Jurasek (1962)
<i>Liacarus coracinus</i>	
Czechoslovakia	Prokopic (1967)
Czechoslovakia	Nasarpur and Prokopic (1979)
<i>Peloptulus phaenotus</i>	
Bulgaria	Bankov (1965)
<i>Unguizetes reticularis</i>	
Chad	Graber and Gruvel (1969)
<i>Zygoribatula exarata</i>	
Hungary	Kassai and Mahunka (1965)
<i>Zygoribatula lata</i>	
Argentina	Yanavella (1971)
<i>Zygoribatula skrjabini</i>	
USSR	Ilyasov (1970)
USSR	Nazarova (1970)
<i>Leibstadia similis</i>	
GFR	Skorski <i>et al</i> (1984)
<i>Hypozetes</i> sp	
India	Deshpande <i>et al</i> (1980)
<i>Spatiodaemaeus subverticulipes</i>	
USSR	Alkov (1972)

Table 3a – Summary of literature on *Scheloribates* and *Protoschelobates* sp as intermediate hosts of *Moniezia benedeni* in different countries

Intermediate host country	References
<i>Scheloribates laevigatus</i>	
USSR	Potemkina (1944a)
Czechoslovakia	Prokopic (1962a)
Bulgaria	Bankov (1965)
USSR	Kuznetsov (1970)
USSR	Alkov (1971)
USSR	Kramnoi (1973)
India	Narsapur (1976b)
Czechoslovakia	Prokopic and Narsapur (1978)
<i>Scheloribates latipes</i>	
USSR	Potemkina (1951)
Czechoslovakia	Prokopic (1962a)
USSR	Nazarova (1970)
USSR	Alkov (1971)
USSR	Kramnoi (1973)
USSR	Abkev (1976)
Czechoslovakia	Prokopic and Narsapur (1978)
<i>Scheloribates semidesertus</i>	
USSR	Nazarova (1970)
<i>Scheloribates fimbriatus</i>	
USSR	Nazarova (1970)
India	Narsapur (1976b)
<i>Scheloribates madrasensis</i>	
India	Anantaraman (1951)
<i>Scheloribates</i> sp	
China	Cai and Jin (1984)
<i>Protoschelobates</i>	
India	Nadakal (1960)

(1) oribatid mites are the intermediate hosts of only Anoplocephalids and

(2) Anoplocephalids are almost exclusively transmitted by oribatid mites.

However, certain exceptions are reported for both. Prokopic (1962b) showed that *Rodentolepis straminea* (Hymenolepidoidea) can also develop in *Achipteria coleoptera*, an oribatid mite.

Involvement of insects of the order Psocoptera in the life cycle of an anoplocephalid tapeworm viz *Thysaniezia actinoides* was reported for the first time by Allen (1959) which was later confirmed in respect of *Thysaniezia* sp and *Avitellina* sp by Kuznetsov (1963, 1965, 1966)).

In case of *Oochoristica osheroffi*, an anoplocephalid tapeworm of rattle snakes, flour beetles (*Tribolium confusum*) migratory grass hoppers (*Conozoa wallula*), and two striped grass hoppers (*Melonoplus bivittatus*) have been shown to be intermediate hosts (Widmer and Oken 1967).

Recently Fritz (1985) again went into the question of alternative intermediate hosts of *M expansa* and

reported an interesting finding that larvae of fire ants *Solenopsis invicta* readily fed on eggs of *M expansa* and showed oncospheres in the gut. It therefore appears that although the above made generalizations hold good in much part, the question of alternative intermediate hosts of Anoplocephalines in different biotopes is still open for curious investigations.

### Biological phenomena in nature

Some interesting observations have been made regarding mode of infection of mites in nature. According to Krull (1939), mites cannot ingest the tapeworm eggs entirely but make hole in the shell and suck the contents during which hexacanth embryo is taken in.

Stunkard (1938) considered that mites feed on tapeworm eggs purely incidentally or accidentally. The eggs of *Moniezia* survive only for few hours (4 to 12) on pasture and few (nine) days in sheep faeces as noted by Kuznetsov (1959) in Volga

Table 3b – Summary of literature on *Galumna*, *Trichoribates*, *Ceratozetes*, *Punctoribates* and *Archipteria* as intermediate hosts of *Moniezia benedeni* in different countries

Intermediate host Country	References
<i>Galumna</i> sp	
India	Anantaraman (1951)
USSR	Alkov (1971)
<i>Galumna elimata</i>	
Czechoslovakia	Prokopic (1962a)
<i>Galumna obvia</i>	
USSR	Potemkina (1944)
USSR	Alkov (1971)
<i>Galumna nervosus</i>	
USSR	Potemkina (1951)
<i>Trichoribates</i> sp	
USSR	Potemkina (1951)
<i>Trichoribates trimaculatus</i>	
Czechoslovakia	Prokopic (1962a)
USSR	Alkov (1971)
<i>Trichoribates incisellus</i>	
USSR	Alkov (1971)
<i>Trichoribates novus</i>	
Czechoslovakia	Prokopic and Narsapur (1981)
<i>Ceratozetes mediocris</i>	
Hungary	Kassai and Mahunka (1965)
<i>Ceratozetes</i> sp	
USSR	Kramnoi (1973)
<i>Punctoribates</i> sp	
USSR	Potemkina (1951)
<i>Punctoribates punctum</i>	
USSR	Alkov (1971)
USSR	Kramnoi (1973)
<i>Archipteria</i> sp	
USSR	Potemkina (1951)
USSR	Akbev (1976)
<i>Archipteria coleoptera</i>	
Czechoslovakia	Prokopic (1962b)

region of USSR. This period might be still shorter in tropics.

Kozlov and Tikhomirova (1979) state that oribatid mites in fact avoid fresh cattle faeces for 1.5 to 2 weeks which is best time for infection with Anoplocephalines. However, they explain, coprophagous beetles invade fresh faeces almost immediately and proceed to bury it. The infective material is taken into the ground where it is protected from

rapid dessication and where it is brought in contact with oribatid host. The observations by Fritz (1985) indicate that fire ants *Solenopsis invicta* also help in disseminating and preserving the tapeworm eggs in the soil.

Under natural conditions, on heavily contaminated pasture the mites are exposed to heavy concentration of tapeworm eggs, but the percentage of infected mites in nature as reported by different

Table 3c – Summary of literature on *Adoristus*, *Liacarus*, *Spatiodaemeus*, *Platinothrus*, *Oribatula* and *Zygoribatula* as intermediate hosts of *Moniezia benedeni* in different countries

Intermediate host Country	References
<i>Adoristus ovatus</i>	
USSR	Potemkina (1951)
<i>Liacarus coracinus</i>	
Czechoslovakia	Prokopic (1967)
<i>Spatiodaemeus subverticillipes</i>	
USSR	Alkov (1971)
<i>Platinothrus peltifer</i>	
USSR	Alkov (1971)
Czechoslovakia	Prokopic and Narsapur (1981)
<i>Oribatula sp</i>	
China	Cai and Jin (1984)
<i>Zygoribatula longiporosa</i>	
Australia	Roberts (1953)
USSR	Sokolova and Panin (1960)
<i>Zygoribatula frisiae</i>	
USSR	Sokolova (1958)
USSR	Sokolova and Panin (1960)
<i>Zygoribatula cognata</i>	
USSR	Sokolova and Panin (1960)
<i>Zygoribatula exarata</i>	
Hungary	Kassai and Mahunka (1965)
<i>Zygoribatula skrjabini</i>	
USSR	Ilyasov (1970)
USSR	Nazarova (1970)

workers is between 0.05 % to 2.0 % and rarely up to 5 % (Prokopic 1962a, Lin *et al* 1975, 1983). This might be due to heavy mortality in mites resulting from infection of tapeworm eggs as noted by Potemkina (1944a) and Prokopic (1962a). Interestingly enough the hexacanth embryos arrest the reproduction of mites and make them sterile (Kramnoi 1978). Thus, to some extent, tapeworm infection in grazing animals paradoxically limits the oribatid mite population on the pastures.

In the final host, only a few number (3 to 4 %) of the ingested cysticercoids reach the adult stage (Narsapur 1976c). Thus, it is likely that heavy tapeworm infections encountered in animals are the result not of a single heavy exposure but of continuous pick up over a longer period. In the latter process the percentage of cysticercoids developing to adult parasite in final host is likely to be still low as immunological responses of the host are coming into play. It therefore stands to reason that,

heavy infections with Anoplocephaline cestodes occur only in certain circumstances viz (1) in the seasons when oribatid mite population, particularly of *Schelorbates sp* and *Galumna sp* is high in nature and (2) over crowding of the pastures with infected animals in this season. Hence, anthelmintic treatment of the herd or flock at the time corresponding to the beginning of prepaek periods of oribatid mite population would therefore be a suitable strategy to control Anoplocephaline tapeworm infections as postulated by Narsapur (1977).

## Conclusions

Anoplocephaline cestodes have been conclusively shown to adversely affect the growth, production and resistance of the animals. Since 1938, very extensive literature has appeared on the biology of these tapeworms. The larval development in the

Table 4 – Summary of literature on intermediate hosts of *Avitellina* and *Stilesia* in different countries

Intermediate host Country	References
<i>Avitellina</i> sp	
<i>Lachisella pedicularia</i> (Psocoptera : Insecta)	
USSR	Kuznetsov (1965, 1966)
<i>Lachisella pedicularia</i> var <i>brevipenis</i> (Psocoptera : Insecta)	
USSR	Kuznetsov (1965, 1966)
<i>Scheloribates laevigatus</i>	
India	Narsapur (1976)
<i>Scheloribates fimbriatus</i>	
India	Narsapur (1976)
<i>Stilesia globipunctata</i>	
<i>Scheloribates indica</i>	
India	Tandon (1963)
<i>Scheloribates perforatus</i>	
Chad	Graber and Gruvel (1967)
<i>Scheloribates conglobatus</i>	
Chad	Graber and Gruvel (1967)
<i>Schelolibates fimbriatus africanus</i>	
Chad	Graber and Gruvel (1967)
<i>Galumna pellucida</i>	
Chad	Graber and Gruvel (1967)
<i>Africacarus calcaratus</i>	
Chad	Graber and Gruvel (1967)
<i>Erythraeus</i> sp	
India	Tandon (1963)

body cavities of oribatid mite hosts is influenced by the species of the mite, intensity of infection and the environmental temperature.

All over the world, 53 species of oribatid mites have been shown to be intermediate hosts of *Moniezia expansa*, 31 of *M benedeni*, 2 of *Avitellina* sp, 7 of *Thysaniezia* sp, 7 of *Stilesia globipunctata*, 2 of *Anoplocephala magna*, 4 of *Pryjukovi* and 2 of *P mamillana*.

A degree of host specificity has been demonstrated between the species of mites and tapeworms. *Scheloribates* sp and *Galumna* sp have been more often incriminated in several countries as intermediate hosts of common Anoplocephalines and hence can be considered highly specific intermediate hosts.

Nymphal stages of oribatids can also be infected and the tapeworm larvae survive moults in their

body cavities which adds to the dimensions of tapeworm epidemiology.

The question of alternate intermediate hosts is still fascinating as it is known that in case of some species, intermediate hosts other than oribatid mites are involved. The direct or indirect role of other creatures like beetles, ants, grasshoppers, etc... in the spread of Anoplocephalines has once again drawn the attention of researchers.

The immune expulsion phenomenon is another area that needs more elucidation and strategic treatment for control, more field trials.

**Acknowledgements**

The author wishes to record sincere thanks to Dr SR Rao (retired), Dr SM Niphadkar and Dr PD Sar-

deshpande (Bombay Veterinary College, Bombay-400 012, India) and Dr J Prokopic (Institute of Para-

sitology, Praha, Czechoslovakia) for encouragement and help in his work cited in this review.

Table 5 – Summary of literature on intermediate hosts of *Thysaniezia* in different countries

Intermediate host Country	References
<i>Thysaniezia</i> sp	
<i>Lachisella pedicularia</i> (Psocoptera : Insecta)	
USSR	Kuznetsov (1965, 1966)
<i>Lachisella Pedicularia var brevipennis</i>	
USSR	Kuznetsov (1965, 1966)
<i>Thysaniezia giardi</i>	
<i>Zygoribatula Skrjabini</i>	
USSR	Iliasov (1970)
<i>Thysaniezia ovilla</i>	
<i>Scheloribates laevigatus</i>	
USSR	Potemkina (1944b, 1951)
<i>Scheloribates latipes</i>	
USSR	Potemkina (1944b, 1951)
<i>Trichoribates sp</i>	
USSR	Potemkina (1944b, 1951)
<i>Galumna obvious</i>	
USSR	Potemkina (1944b, 1951)
<i>Adoristus ovatus</i>	
USSR	Potemkina (1944b, 1951)
<i>Phthiracarus</i>	
USSR	Potemkina (1944b, 1951)

## References

- Afonina YA, 1967. Effects of copper sulphate and phenothiazine salt mixtures on monieziasis in lambs. *Veterinariya* (Mosc) 44:49-50
- Akbev MSH, 1976. Infection of oribatid mites with *Moniezia ova*. *Sb Nauchnykh Tr Mosk Vet Akad* 86:86-89
- Alkov MV, 1971. Epizootology of monieziasis of ruminants. *Veterinariya* (Mosc) 48:60-61
- Alkov MV 1972. Epizootology of monieziasis of ruminants. *Mater Nauchnykh Issled Chlenov Vses Ova Gel'mintol* 24:3-5
- Allen RW, 1959. A preliminary note on the larval development of finged tapeworm of sheep, *Thysanosoma actinoides* (Diessing 1934) in Psocids (*Psocoptera corrodentia*). *J Parasitol* 45:537-538
- Amjadi AR, 1971. Studies on histopathology of *Stilesia globipunctata* infections in Iran. *Vet Rec* 88:486-488
- Anantaraman M, 1951. The development of *Moniezia*, the large tapeworm of domestic ruminants. *Sci Cul* 17:155-157
- Bankov DE, 1965. Research into the biology of *Moniezia* and the epizootology of monieziosis in Bulgaria. *Vet Sci* 2:238-294
- Bankov DE, 1971. Klinik Pathologie und Chemotherapie der Stilesiose beim Schaf. *Angew Parasitol* 12:90-96
- Bashkirova EJ, 1941. Contribution to the study of the biology of the tape worm *Anoplocephala perfoliata* (Goeze 1782) parasitic in horses. *Compt Rew (Doklady) Acad Sci USSR* 30:576-578
- Cai XP, Jin JS, 1984. Study on the life cycle of *Monezia benedeni*. *J Vet Sci Tech (Shougi Keji Zazhi)* 12:26-30

- Casarosa L, 1964. Indagini isopatologiche sulle modalita d'attacco dello scolice di *Moniezia expansa* (Rudolphi 1810) nella parete intestinale degli ovini. *Parasitologia* (Rome) 6:39-43
- Curtice C, 1890. The animal parasites of sheep. Washington DC 222:36
- Davydov AS, Smirnov NF, 1972. (Life span of *Moniezia expansa* during experimental infection of sheep). *Mater Nauchnykh Issled Chlenov vses ova Gel'mintol* 24:47-49
- Deshpande PD, Shastri UV, Ghafoor MA, 1980a. Role of *Hypozetes sp* an oribatid mite from India as intermediate host of *Moniezia expansa*. *Proc III India Congr Parasitol* 3:75
- Deshpande PD, Anantwar LG, Shastri UV, Ghafoor MA, 1980b. Effect of development of *Moniezia expansa* in the kids on some haematological and biochemical values. *Proc 3rd India Congr Parasitol* 3:96
- Edney JM, Kelly GW, 1953. Some studies on *Galumna virginienensis* and *Moniezia expansa* (Acarina : Oribatoidea-Cestoda : Anoplocephalidae). *J Tennessee Acad Sci* 28:287-296
- Efner T, 1974. The influence of intestinal tape worms on weight gains and the composition of carcass halves of wethers. *Med Weter* 30:81-83
- Euzeby J, 1966. Maladies et affections déterminées par le parasitisme des cestodes dans les maladies vermineuses des animaux domestiques et leurs incidences sur la pathologie humaine, vol 2, maladies dues aux plathelminthes, fasc 1, cestodoses, 208-234. Vigot, Paris
- Freeman RS, 1952. The biology and life history of *Monoecocestus*, Beddard 1914 (Cestoda : Anoplocephalidae) from porcupines. *J Parasitol* 38:111-129
- Fritz GN, 1985. A consideration of alternative intermediate hosts for *Moniezia expansa* (Cestoda : Anoplocephalidae). *Proc Helminthol Soc Wash* 52:51-53
- Gapon NM, 1974. Post larval development of *Moniezia* in calves. *Helminthol Abstr* 48:246
- Gleason LN, Buckner RL, 1979. Oribatid mites as intermediate hosts of certain anoplocephalid cestodes. *Trans Ky Acad Sci* 40:27-32
- Graber M, Gruvel J, 1967. Les vecteurs de *Stilesia globipunctata* (Rivolta 1874) du mouton. *Rev Elev Med Vet Pays Trop* 20:261-271
- Graber M, Gruvel J, 1969. Oribates vecteurs de *Moniezia expansa* (Rudolphi 1810) du mouton dans la région de Fort Lamy. *Rev Elev Med Vet Pays Trop* 22:521-527
- Hansen MF, Todd AC, Kelly GW, Cawein M, 1950. Effects of pure infections of tapeworm *Moniezia expansa* on lambs. *Bull Kent Agric Exp Stn* 556:11
- Hawkins PA, 1946. Studies on sheep parasites. 7. *Moniezia expansa* infections. *J Parasitol* 32
- Ilyasov IN, 1970. In Oribatids and their role in the process of soil formation. Ed Bulanova, Zakhvatkina EM *et al* *Akademiya Nauk Litovskoi, USSR Pp*:235-238
- Jurasek V, 1962. Development of Monieziasis in experimentally infected intermediate hosts – oribatid mites – from South Slovakia. *Folia Vet* 6:93-101
- Kassai T, Mahunka S, 1964. Vizsgalatok a Monieziak Koztigazdairol. *Magyar Allatorvosok Lapja* 17:531-558. *Helminthol Abstr* 34:307
- Kassai T, Mahunka S, 1965. Studies on the tape worms in ruminants. 2 Oribatids as intermediate hosts of *Moniezia sp*. *Acta Vet Hung* 15:227-249
- Kates KC, Goldberg A, 1951. The pathogenicity of the common sheep tape worm *Moniezia expansa*. *Proc Helminthol Soc Wash* 18:87-101
- Kates KC, Runkel CE, 1948. Observations on oribatid mite vectors of *Moniezia expansa* on pastures with a report of several new vectors from United States. *Proc Helminthol Soc Wash* 15:10-33
- Kozlov DP, Tikhomirova VA, 1979. *Tr Gel'mintol Lab* 29:67-70
- Kramnoi VYA, 1973. Development times of *Moniezia benedeni* from cattle in the intermediate host in Annur region USSR. *Helminthol Abstr* 45:3832
- Kramnoi VYA, 1978. The effect of Anoplocephalate cysticercoids on the reproductive functions of oribatids. In *Ekologiya Zhivotnykhi Faunistika. Tyumen USSR*:19-23
- Kramnoi VYA, 1980. Biology of bovine *Moniezia* in the priamure. *Tr Dal'nevost Zon Nauchno-Issled Vet Inst* 11:34-35
- Krull WH, 1939. On the life history of *Moniezia expansa* and *Cittotaenia sp* (cestoda : Anoplocephalidae). *Proc Helminthol Soc Wash* 6:10-11
- Kuliev KA, 1963. Observations on the development of *Anoplocephala perfoliata* (Goeze 1782) in an intermediate host (oribatid mites). *Problemy Parazitologii* (Trudy IV Nauchnoi Konferentsii Parazitologov UKR SSR). pp 210-211
- Kuznetsov MI, 1959. Survival of *Moniezia* eggs in pastures in lower Volga steppes. *Byull Nauchno – Tekh Inf V Ses Inst Gel'mintol in KI Skryabina* 5:48-51
- Kuznetsov MI, 1963. Intermediate hosts of *Thysaniezia* and *Avitellina*. *Tr Vses Inst Gel'mintol* 10:5-8
- Kuznetsov MI, 1965. Study of the biology of the causative agent of Avitellinosis in sheep. *Tr Uzb Nauchno Issled Inst Vet* 17:118-125
- Kuznetsov MI, 1966. On the intermediate hosts of the species of *Avitellina* and *Thysaniezia* of sheep. *Temat Sb Rabs Gel'mintol* (Trudy UI GIS) 12:25-37



- Kuznetsov MI, 1968. The susceptibility of sheep of different ages to *Moniezia expansa* and *M benedeni*. Tr Vses Inst Gel'minthol 14:210-222
- Kuznetsov MI, 1970. Development times of *Moniezia cysticercoids* in *Schelorbates laevigatus* under natural conditions. In Oribatid and their role in the process of soil formation. Vilnius : Akademiya Nauk Litovskoi SSR pp 223-227
- Lin YK, Ho YC, Sung YL, 1975. Studies on the epidemiology of Monieziasis (*M expansa*) and the biology of its natural vectors. Acta Zool Sin 21:141-154
- Lin YK, Guan J, Wang P, Yan W, 1982. Studies on the development cycle of *Paranoplocephala ryjikovi* (Spassky 1950) in the intermediate host. Acta Zool Sin 28:262-270
- Lin YK, Jian X, Guan J, Wang PP, Zhang C, Yang W, 1983. Studies on the anoplocephala cestodes and oribatid mite vectors of *Moniezia* in Ar-Bar country Sichuan province China. Acta Zoologica Sin 29:323-332
- Lyashenko IS, Teplov OV, 1974. Effect of Monieziasis on meat and wool yields of sheep. Byull Vses Inst Gel'minthol CM Kl. Skryabina 14:34-41
- Matevosyan EM, 1978. On the life cycle of *Avitellina centripunctata* (Rivolta 1874) cestoda. Biol Zh Arm 31:979-982
- Mehra KN, Srivastava HD, 1955a. Studies on the life history of *Moniezia expansa* (Rudolphi 1810) a broad tape worm of ruminants. Proc 42nd Indian Sc Cong III p 352
- Mehra KN, Srivastava HD, 1955b. Studies on the life history of *Moniezia benedeni* (Moniez 1879) a tape worm of ruminants. Proc 42nd Indian Sc Cong III p 352
- Morgan DO, 1947. Parasitic helminths of sheep. Vet Rec 59:494
- Nadakal AM, 1960a. Observations on the life cycle of *Avitellina centripunctata* (Rivolta 1874) an anoplocephalinae cestode from sheep and goat. J Parasitol 46:12
- Nadakal AM, 1960b. *Proteoschelobates* sp, an oribatid mite from India as a potential vector of the sheep tape worm *Moniezia expansa* and *Moniezia benedeni*. J Parasitol 46:817
- Narsapur VS, 1971. Ecological and biological studies on oribatid fauna of India (Bombay region) together with observations on the life cycle of common anoplocephalid tape worms. A thesis submitted to MPKV, Rahuri, India
- Narsapur VS, 1974a. Ecological and biological studies on oribatid fauna of India (Bombay region) together with observations on the life cycle of common anoplocephalid tape worms. Indian Vet J 50:165-166
- Narsapur VS, 1974b. A note on the vectors of *Avitellina lahorea* (Woodland 1927) in India. Indian Vet J 51:54-56
- Narsapur VS, 1976a. Laboratory infections of *Schelorbates* sp (Oribatid mites) with *Moniezia benedeni* and *Moniezia expansa*. J Helminthol 50:153-156
- Narsapur VS, 1976b. Intermediate hosts and larval development of *Moniezia benedeni* in India. J Parasitol 62:720
- Narsapur VS, 1976c. Observations on the biology of sheep tape worm *Moniezia expansa*. Indian J Anim Sci 46:603-609
- Narsapur VS, 1977. Strategic anthelmintic treatment for the control of Monieziasis in sheep, goat and cattle. Indian Vet J 54:856-858
- Narsapur VS, 1979. First report on the development of *Moniezia expansa* in juvenile stages of oribatid mites. Folia Parasitol (Praha) 25:210
- Narsapur VS, Prokopic J, 1979. The influence of temperature on the development of *Moniezia expansa* (Rudolphi 1810) in oribatid mites. Folia Parasitol (Praha) 26:239-243
- Nazarova SA, 1970. In Oribatids and their role in the process of soil formation. Ed Bulanova, Zackhvatkina EM *et al*, Vilnius Akademiya Nauk Litovskoi SSR pp 229-233
- Orekhov MD, 1960. Tr Turkm Nauchno Issled. Inst Zhivotnovod Vet 2:267-288. Helminthol Abstr 30:446
- Petrovic Z, Bordsoski A, Popovic B, 1970. Prevention of *Moniezia* infection in sheep and its economic importance. Vet Glas 24:991-994
- Potemkina VA, 1941. Contribution to the biology of *Moniezia expansa* (Rudolphi 1810) a tape worm parasitic in sheep and goats. CR (Doklady) Acad Sci URSS 30:474-476
- Potemkina VA, 1944a. On the decipherment of the biological cycle in *Moniezia benedeni* (Moniez 1874) tape worm parasite of cattle and other domestic animals. CR (Doklady) Acad Sci URSS 42:146-148
- Potemkina VA, 1944b. Contribution to the study of development of *Thysaniezia ovilla* (Rivolta 1878) a tape worm parasitic in ruminants. CR (Doklady) Acad Sci URSS 43:43-44
- Potemkina VA, 1948. On the biology of *Moniezia expansa*. Collected papers on Helminthology, dedicated to KI Kryabin on 40th Anniversary of the Union Inst Helminthology, Moscow pp 177-184
- Potemkina VA, 1951. Monieziasis in calves. Tr Gel'minthol Lab. Akad Nauk SSR 5:299-302
- Prokopic J, 1962a. Seasonal dynamics of occurrence of tape worms of the genus *Moniezia* and questions of their development cycle in the pasture regions of Sumara mountains. Cesk Parasit 9:355-364
- Prokopic J, 1962b. Roztoc *Achipteria coleoptera* (Ill). Mezihostitel tasmenice *Rodentolepis straminea* (Goeze 1782). Zool Listy 11:183-187
- Prokopic J, 1967. Bionomische Studien üben Bandwürmer der Gattung *Moniezia*. Angew Parasitol 8:200-209
- Propokic J, Narsapur VS, 1981. Experimental study on the life cycle of *Moniezia benedeni* (Moniez 1879). Folia Parasitol Praha 28:54
- Rao NSK, Choquette LPE, 1951. On the findings of an intermediate host for *Moniezia expansa* (Rudolphi 1810) in Eastern Quebec. Can J Comp Merl 15:12-14

- Rayski C, 1947. Sheep tape worms and their intermediate hosts. *Scott Farmer* 55:284
- Roberts FHS, 1953. *Zygoribatula longiporosa* (oribatei Acarina), an intermediate host of *Moniezia benedeni* (Moniez 1879) Anoplocephalidae in Australia. *Aust J Zool* 1:239-241
- Rukarina *et al*, 1960. *Veterinaria* (Sarajevo) 9:497-514. *Helminthol Abstr* 30:172
- Runkel CE, Kates KC, 1947. A new intermediate host (*Protoschelobates seghattii* nsp Acarina : Schelioribatidae) of the sheep tape worm *Moniezia expansa*. *Proc Helminthol Soc Wash* 14:64-67
- Shakiev ESH, 1973. Anthelmintic and therapeutic efficacy of compounds against *Moniezia* in sheep. *Tr Uzb Nauchno-Issled Vet Inst* 20:193-200
- Shakurova FM, 1974. The toxicity of *Moniezia expansa* antigen. *Uch Zap Kaz Vet Inst im NE Bauman* 115:268-270
- Shaldibina ES, 1953. Papers on Helminthology presented to Academic KI Skryabin on his 75th birthday Moscow Izadatelsteio Akademii Nauk SSR 747-755
- Shorb DA, 1939. Preliminary observations of the effect on the sheep of pure infestation with tape worm *Moniezia expansa*. *Proc Helminthol Soc Wash* 6:77-79
- Skorski P, 1984. Investigations into the maintenance of *Schelioribates laevigatus*, *Leibstadia similis* and *Pilo galumna tenuiclavus* and their infection with *Moniezia expansa*. Inaugural Dissertation Ludwig – Maximilians Universität Munich GFR p 69
- Skorski P, Barutski B, Boch J, 1984. Investigations on the prevalence of oribatid mites on sheep pastures and their role as intermediate hosts of *Moniezia expansa*. *Berl Muench Tierarztl Wochenschr* 97:291-295
- Sokolova IB, 1958. *Zygoribatula frisiae* intermediate host of *Moniezia benedeni* in South Kazakhstan. *Tr Inst Zool Akad Nauk Kaz SSR* 9:242-243
- Sokolova IB, Panin VY, 1960. *Tr Inst Zool Akad Nauk Kaz. SSR* 12:145-148
- Soulsby E.J.L, 1982. In *Helminth, Arthropods and Protozoa of domesticated animals*. 7th Ed, Bailliere Tindall, London
- Stampa S, 1967. A contribution towards influence of tape worms on live weights of lambs. *Vet Med Rev Leverkusen* 1:81-85
- Stoll NR, 1935a. Tape worm studies 1 – Restricted pasture sources of *Moniezia* infection. *Am J Hyg* 21:628-646
- Stoll NR, 1935b. Tape worm studies 2 – Persistence of pasture stage of *Moniezia expansa*. *Am J Hyg* 22:683-703
- Stoll NR, 1936. Tape worm studies 3 – Sheep parasitized with *Moniezia expansa*. *J Parasitol* 22:161-179
- Stoll NR, 1937a. Rates of acquisition of grazing sheep of *Moniezia expansa* and what they reveal of the available pasture infestation. *J Parasitol* 23:568-569
- Stoll NR, 1937b. Tape worm studies 5 – Absence of *Moniezia expansa* from sheep intestine early after infection. *Am J Hyg* 26:148-161
- Stoll NR, 1938. Tape worm studies 7 – Variations in pasture infestation with *Moniezia expansa*. *J Parasitol* 24:527-545
- Stunkard HW, 1937a. The life cycle of *Moniezia expansa*. *Science* 86:312
- Stunkard HW, 1937b. Studies on *cittoteania* sp and *Moniezia expansa*. *J Parasitol* 23:569
- Stunkard HW, 1938. Development of *Moniezia expansa* in the intermediate host. *Parasitology* 30:491-501
- Stunkard HW, 1940. The morphology and life history of the cestode *Bartiella studeri*. *Am J Trop Med Hyg* 20:305-327
- Stunkard HW, 1941. Studies on the life history of Anoplocephaline cestodes of hares and rabbits. *J Parasitol* 27:299-325
- Tableman HG, 1946. Taeniasis in lambs. *Vet Med* 41:455-456
- Tandon RS, 1963. Observations on the life history of the Anoplocephaline cestode, *Stilesia globipunctata* (Rivolta 1974) (Subfamily Thysanosominae) a common parasite of ruminants in India. *Parasitologia* 5:183-188
- Urmanbetova CL, 1979. Anoplocephalid infections in horses in Northern Krizia and their control. *Helminthol Abstr* 52:362
- Vibe PP, 1976. Parasitic coenoses in sheep. *Veterinariya* (Mosc) 6:58-60
- Widmer EA, Oken OW, 1967. The life history of *Oochoristica osheroffi*, Meggit 1934 (Cyclophyllidea : Anoplocephalidae). *J Parasitol* 53:343-349
- Worley DE, Jacobson RH, Barret RE, 1974. The chronology of tape worm (*Moniezia expansa*) acquisition by sheep on summer ranges in Montana and Idaho. *Proc Helminth Soc Wash* 41:19-22
- Yamaguti S, 1959. In *Systema helminthum*. Vol 2 The cestodes of vertebrates. Inter Science New York pp 369-401
- Yanavella FG, 1971. An ecological focus of parasitism of sheep by *Moniezia* and detection of the intermediate host. *Analecta Vet* 3:21-28