

Assessing the effect of a single dose florfenicol treatment in feedlot cattle on the antimicrobial resistance patterns in faecal Escherichia coli

Anna Catharina B. Berge, William Epperson, Robbi Pritchard

▶ To cite this version:

Anna Catharina B. Berge, William Epperson, Robbi Pritchard. Assessing the effect of a single dose florfenicol treatment in feedlot cattle on the antimicrobial resistance patterns in faecal Escherichia coli. Veterinary Research, 2005, 36 (5-6), pp.723-734. 10.1051/vetres:2005027. hal-00902995

HAL Id: hal-00902995 https://hal.science/hal-00902995

Submitted on 11 May 2020 $\,$

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés. Vet. Res. 36 (2005) 723-734 © INRA, EDP Sciences, 2005 DOI: 10.1051/vetres:2005027

Original article

Assessing the effect of a single dose florfenicol treatment in feedlot cattle on the antimicrobial resistance patterns in faecal Escherichia coli

Anna Catharina B. BERGE^a, William B. EPPERSON^{b,c*}, Robbi H. PRITCHARD^d

^a Veterinary Medicine Teaching and Research Center, University of California Davis, 18830 road 112, Tulare, CA, 93274, USA

^b Department of Veterinary Science, College of Agriculture and Biological Sciences, South Dakota State University, Brookings, South Dakota, 57007, USA

^c Present address: Department of Veterinary Preventive Medicine, College of Veterinary Medicine, The Ohio State University, 1920 Coffey road, Columbus, Ohio, USA ^d Department of Animal and Range Science, College of Agriculture and Biological Sciences,

South Dakota State University, North Campus drive, Brookings, South Dakota, 57007, USA

(Received 10 August 2004; accepted 21 February 2005)

Abstract – The objective of this clinical trial was to examine the effect of a single dose of florfenicol on antimicrobial resistance patterns in faecal E. coli of feedlot steers. Steers (n = 370), were purchased from two sources and housed in outdoor concrete floored pens. Two cattle from each pen (n = 42 pens, 84 cattle) were randomly selected for faecal sampling at study day 1, 14, 28, and 42. One sampled animal from each of 21 pens was randomly selected to receive a single 39.6 mg/kg dose of florfenicol subcutaneously at study day 11. Ten lactose positive colonies were isolated from faecal swabs and tested for antimicrobial resistance to 11 antimicrobials using the disk diffusion method. Zones of inhibition were grouped using cluster analysis and clusters were ordered by increasing multiple resistance. A cumulative logistic regression model using generalized estimating equations was used to assess factors associated with increasing levels of multiple resistance. Immediately post-treatment, all isolates obtained from treated cattle belonged to multiple resistant clusters containing chloramphenicol resistance. Though less pronounced in later sampling, resistance to chloramphenicol and other antimicrobials persisted. Antimicrobial treatment, sampling time and animal source, as well as interactions between these variables, were important predictors of the odds of E. coli belonging to a more resistant cluster. A very clear but transitory shift to increasingly multiple resistant faecal E. coli in response to florfenicol treatment was observed. There was no indication of horizontal transfer of resistant E. coli between steers. Level of resistance was influenced by complex interaction of animal source and previous management.

antimicrobial resistance / food animal / florfenicol / Escherichia coli

1. INTRODUCTION

There has been ongoing debate concerning the effect of therapeutic and non-therapeutic veterinary antimicrobials on human health [25, 28]. Animal agriculture has been implicated as a major user of antimicrobials, and, by extension, a major contributor to antimicrobial resistance. Traditionally, antimicrobial resistance was studied through

^{*} Corresponding author: Epperson.1@osu.edu

use of a specific organism, and through monitoring antimicrobial resistance in that organism over time [26]. Often these specific organisms were pathogens, and intensive study allowed identification of emerging trends, estimation of prevalence in animals or animal products, and investigation of antimicrobial resistance mechanisms. However, for population assessment, use of a pathogenic organism may be problematic since prevalence may be low [24]. The objective of population monitoring may be more toward determining predominate antimicrobial resistance, rather than documenting low prevalence antimicrobial resistance.

Antimicrobial resistance assessment of commensal organisms has been used to study gastrointestinal tract ecology and track environmental organisms to a presumed source [14, 16, 23]. Commensal E. coli have been used as a monitor of antimicrobial resistance in cattle and swine populations [5, 10, 30, 31]. Use of commensal organisms is convenient and allows measurement of the resistance gene pool that may be transmitted to pathogenic bacteria. Several studies have indicated that interspecies resistance transfer occur in bacteria, and increased recovery of antimicrobial resistant faecal E. coli was seen in dairy farms affected with resistant salmonellosis [6, 8, 11]. The level of resistance in commensal enteric bacteria has been correlated with resistance in pathogenic bacteria [11].

In cattle, factors apart from antimicrobial administration affect antimicrobial resistance, including age, environment, and nutrition [9, 16–18]. Long-term use of antimicrobials may select for antimicrobial resistance that persists over time, though duration of antimicrobial resistance in faecal *E. coli* in feedlot cattle following therapeutic antimicrobial usage appears to be short [20, 29].

Florfenicol is a veterinary fluorinated analogue of tiamphenicol approved in 1996 in the United States for bovine respiratory disease pathogens [13]. Florfenicol is a bacteriostatic antimicrobial with similar mode of action to chloramphenicol, binding to the 50S subunit of bacterial ribosomes. The *flo* gene confers resistance to chloramphenicol and florfenicol, and is most commonly located on a large transferable plasmid in bovine enteric *E. coli* [27, 33].

The objective of this study was to examine the effect of a single therapeutic dose of florfenicol on antimicrobial susceptibility patterns in faecal *E. coli* of steers within one month post treatment.

2. MATERIALS AND METHODS

2.1. Animal sampling and treatment

The clinical trial was performed on animals included in nutrition experiments at the South Dakota State University Ruminant Nutrition Research Center in the autumn of 2001. Angus and Angus crossbred steers (n = 370), weighing approximately 270 kg were purchased directly from source A (n = 196) and B (n = 174) and arrived on October 25 and October 31, 2001 respectively. Calves and dams were rounded up on the range and steer calves were weaned and shipped directly from the ranch. Sixty steers from source B had been weaned and held in a dirt lot for approximately one month prior to shipment to the research center. Cattle were used in two nutrition experiments, apart from the study here. Source A steers were assigned to one of two treatments in a study involving dietary trace mineral inclusion rate. Source B steers were placed into one of four treatments to study effect of various grain byproducts on health and performance. Cattle were housed in 42 open concrete floor pens (8-10 head/pen). Prior to the study, pens had been scraped free of physical debris and were unoccupied for 30 days. The ionophore monensin (Rumensin, Elanco Animal Health, Indianapolis, Indiana, USA), was incorporated into the diet at a rate of 20 grams/ton of feed, and was the only

Antimicrobial Code Concentration (µg) Amoxicillin/clavulanic acid 20/10 AMC Ampicillin AMP 10 Cephalothin 30 CEF Chloramphenicol CHL 30 Gentamicin GEN 10 Nalidixic acid NAL 30 Spectinomycin SPT 100 Streptomycin STR 10 Sulfisoxazole SULF 250 Tetracycline TET 30 Sulfamethoxazole/trimethoprim SXT 23.75/1.25

Table I. Antimicrobials and concentrations used for susceptibility testing of faecal *E. coli* isolates using the disk diffusion method.

antimicrobial in feed or water. Two cattle from each pen (n = 84) were randomly enrolled into the study. Serial rectal faecal samples were obtained using cotton tipped swabs from all study animals on days 1, 14, 28, and 42. At study day 11, one study animal from half the pens (n = 21) was randomly selected to receive a single 39.6 mg/ kg dose of florfenicol (Nuflor - Schering-Plough Animal Health, Kenilworth, New Jersey, USA) by subcutaneous injection. The use of this treatment regime was within label indications. Animals selected for antimicrobial treatment came from both source A (n = 12, 25% of cattle) and source B (n = 9, 12)25% of cattle). A total of 48 steers in 24 pens from source A and 36 steers in 18 pens from source B were sampled.

2.2. Sampling and isolation of E. coli

Swabs were plated onto MacConkey agar within 3 h of acquisition and incubated 24 h at 35 °C. Ten well isolated lactose positive colonies (or as many that were present, if less than 10) were randomly selected from each plate for antimicrobial resistance testing. Biochemical confirmation was performed on 49 randomly selected isolates using triple sugar iron agar (TSI), motility, and indole media.

Antimicrobial susceptibility tests were performed using the disk diffusion assay [4]. The selected colonies were inoculated into NaCl sterile solution (0.85%) to achieve an optical density corresponding to 0.5 McFarland units. A sterile swab was dipped into the adjusted suspension and streaked onto Mueller-Hinton (MH) agar plates (150 \times 15 mm) to form a uniform lawn of bacterial growth. Eleven drug-impregnated disks were placed on the surface of the agar using a disk dispenser (Tab. I). The plates were incubated at 37 °C for 16 to 18 h and zones of inhibition around each disc measured to the nearest mm. The quality control strain E. coli ATCC 25922 (ATTC, Manassas, Virginia, USA) was used for each batch of isolates tested [1].

2.3. Quantitative analysis

Data from *E. coli* antibiograms were a series of continuous measurements representing inhibition zone size to each of 11 antimicrobials. Descriptive statistics were calculated on isolates to determine the zone

A.C.B. Berge et al.

Table II. Antimicrobial susceptibility clusters of faecal *E. coli* isolates (total number = 3145) in a trial evaluating single-dose florfenicol treatment of steers. Mean disk diffusion zone (mm) is given for each cluster-antimicrobial combination, as well as number of isolates (Frequency) in each cluster*.

Cluster	Frequency	AMC	AMP	CEF	CHL	GEN	NAL	SPT	STR	SXT	SULF	TET
А	2518	25	23	22	25	23	27	24	18	31	25	26
В	263	25	23	22	25	23	26	24	17	30	25	11
С	23	22	6	21	26	23	27	25	18	31	24	6
D	5	27	26	24	6	22	29	23	6	31	24	6
E	54	24	22	21	25	23	27	24	11	24	6	6
F	31	25	23	22	26	23	28	15	12	26	6	6
G	71	26	24	22	6	23	26	23	6	22	6	7
Н	12	23	6	21	26	22	28	15	9	25	6	6
Ι	12	20	6	20	6	24	26	24	7	21	6	6
J	8	20	6	20	6	22	29	16	6	6	6	6
Κ	31	11	6	6	6	23	27	25	6	19	6	6
L	107	13	6	6	6	23	28	15	6	23	6	7
М	10	16	6	12	6	12	28	6	6	21	6	6

* Zone sizes in dark shading indicate resistance to the antimicrobial and zone sizes in light shading indicate intermediate resistance, according to National Committee for Clinical Laboratory Standards (NCCLS) guidelines for human *E. coli*. Zone sizes without shading indicate sensitivity according to same standard.

size distribution for each antimicrobial. Antimicrobials with a bimodal distribution of inhibition zones were used in grouping the strains using cluster analysis. Clusters were determined using Ward's minimum variance method and squared Euclidean distance, as described [5, 12, 32]. Clusters were considered ordinal outcomes, and ordered according to increasing resistance, based on a decreasing sum of mean inhibition zone sizes to the eleven antimicrobials. The statistical software package SAS version 8 (SAS Institute, Cary, North Carolina, USA) was used for data analysis.

Contingency tables of effects of source (A or B), sampling (1, 2, 3 or 4), and florfenicol treatment on *E. coli* antimicrobial cluster membership were developed. The effect of florfenicol treatment on the distribution of chloramphenicol resistance was evaluated with the Chi-square statistic. The Wilcoxon-Mann-Whitney test was used to determine differences in resistance between source groups. A comparison of antimicrobial resistance clusters between steers being housed in the same pen as treated steers and those steers in pens without any treated steers was performed using stratified analysis, and post-antibiotic treatment cluster distribution was tested using the Wilcoxon-Mann-Whitney test.

A mixed effects cumulative logistic regression model using generalized estimating equations was developed, using cluster membership as the outcome of interest [3, 19]. The model predicted the odds of the *E. coli* belonging to a more resistant cluster, given a set of explanatory variables including source ("source", consisting of two categories [A or B]), early weaning (termed "wean", consisting of two categories to account for steers from source B that were weaned one month prior to shipment and other), sampling time ("time", 1–4), florfenicol treatment ("abtx", yes/no categorical), pen mate to treated steer (yes/no categorical), and nutrition treatment (two categories). Second and third order interactions were included in the model. A stepwise forward selection procedure was used in a population average model for variable selection. Since a large number of observations were available, a *p* value of ≤ 0.1 was used for entry and retention in the model. The variables that were retained in the population average model were thereafter introduced into a model using generalized estimating equations (GEE) to control for multiple isolates from each faecal sample [15]. This was performed with a repeated measure on calf at each sampling occasion with an independent correlation matrix design. The proportional odds assumption was assessed by the score statistic and by graphing odds ratios for sampling and treatment group at each level in the cluster hierarchy and assessing trends. The final model was described by the following equation: Logit(cluster) = $\beta_0 + \beta_1$ (abtx) + β_2 (source) + β_3 (wean) + β_4 (time) + β_5^2 $(abtx \times time) + \beta_6 (source \times time) + \beta_7$ $(abtx \times source)/(repeated sampling within$ calf × time).

3. RESULTS

3.1. Overview

During the study, no animal in the sampling cohort became ill or necessitated antimicrobial treatment. Of the total 370 cattle received, only 1 animal (not sampled in this study) required treatment for illness between arrival and study end.

In total, 3179 isolates were obtained from the steers. Of the 49 isolates biochemically tested, 48 (98%) produced reactions typical of *E. coli*. The inhibition zones for most antimicrobials demonstrated a bimodal distribution. All isolates were sensitive to nalidixic acid. Since antimicrobial susceptibility tests to nalidixic acid exhibited no bimodal distribution, they were not included as variables to determine antimicrobial resistance cluster membership, but are included in description of clusters.

3.2. Antimicrobial resistance clusters

The E. coli antimicrobial resistance cluster characteristics and number of isolates belonging to each cluster are shown in Table II. Clusters were optimized to contain minimal intra-cluster variability to the eleven antimicrobials. The optimal number of clusters was based upon having isolates with disk diffusion zones only in one zone of the bimodal distribution. Thirteen clusters adequately described the major antimicrobial resistance patterns and were ordered in increasing levels of resistance from A to M. Of the 3179 isolates, 34 (about 1% of the dataset) were trimmed during the clustering algorithm, leaving 3145 isolates for analysis. The majority of isolates (82%) belonged to cluster A, containing bacteria susceptible to the 11 antimicrobials. Tetracycline resistance was seen in the remaining clusters (B-M) combined with other resistance to one or several antimicrobials. Chloramphenicol resistance, which is present in E. coli when florfenicol resistance is present, was seen in cluster D, G and I–M.

3.3. Statistical analysis

A contingency table of E. coli cluster membership by sampling time and source is shown (Tab. III). The influence of florfenicol treatment on E. coli cluster membership was assessed (Tab. IV). Treated steers displayed a significantly greater proportion of isolates with chloramphenicol resistance than non-treated steers at sampling days 14, 28 and 42. On the second sampling, occurring 3 days post-treatment (day 14), all isolates obtained from treated cattle belonged to clusters containing chloramphenicol resistance, compared to only 1% of isolates from non-treated steers. Though less pronounced in the third and fourth sampling, treated cattle had significantly greater proportion of

Table III. Percentage of faecal *E. coli* isolates (total number = 3145) by source group (A and B), sampling occasion (day 1, 14, 28 and 42), and antimicrobial resistance cluster membership in a trial evaluating single-dose florfenicol treatment of steers.

	Frequency	Percentage of E. coli isolates by source, sampling, and cluster membership									
Cluster A B C D E F G H I J K L M		Day 1*		Day	14**	Day	28**	Day 42**			
		А	В	А	В	А	В	А	В		
А	2518	97	76	73	64	85	81	88	87		
В	263	3	6	2	5	6	13	6	10		
С	23	0	3	0	0	2	0	0	0		
D	5	0	0	1	0	0	0	0	0		
Е	54	0	6	0	3	0	2	1	1		
F	31	0	5	0	2	0	2	0	0		
G	71	0	0	3	7	4	2	2	1		
Н	12	0	3	0	0	0	0	0	1		
Ι	12	0	0	0	3	0	0	0	0		
J	8	0	0	0	2	2	0	0	0		
Κ	31	0	0	17	5	0	0	3	0		
L	107	0	0	1	7	0	0	0	0		
М	10	0	0	2	1	0	0	0	0		

* Source group B had significantly higher proportion of isolates belonging to more resistant clusters than group A (Wilcoxon-Mann-Whitney test, *p*-value < 0.0001).

** No significant difference in resistance between group A and B at day 14, 28 and 42, Wilcoxon-Mann-Whitney test p-value = 0.16, 0.23 and 0.46 respectively.

isolates belonging to clusters with chloramphenicol resistance. Apart from the 1% of isolates belonging to chloramphenicol resistant cluster D found at Day 14, there was no indication of chloramphenicol resistance in control steers. The faecal *E. coli* isolates obtained from non-treated steers housed in pens with treated steers compared to nontreated steers housed in pens with only nontreated steers revealed no significant difference in cluster distribution (Wilcoxon-Mann-Whitney test *p*-value = 0.46)

The multinomial logistic regression model showed that treatment group, sampling time and animal source, as well as interactions between these variables, were important predictors of the odds of *E. coli* belonging to a more resistant cluster (Tab. V). There was no association with nutrition treatment and pen mates to treated steers did not show increasing levels of resistance compared to steers in pens without treatment.

The main effect of treatment indicates a clear shift toward carriage of more resistant *E. coli* in calves treated with florfenicol. The statistical interactions indicate an effect-measure modification. In order to determine the probability of a steer belonging to a higher order cluster, the log-odds of all variables, main effects and interactions, must be calculated. As a result, the GEE estimate of log odds ratio associated with single main effect variables cannot be simply interpreted as the magnitude of effect. The model is therefore discussed in terms of trends for different variables. Abtx × time interaction formally indicated what was

	Frequency -	Percentage of <i>E. coli</i> isolates by treatment group, sampling time, and cluster membership									
Cluster		Day 1		Day 14*		Day 28*		Day 42*			
		control	abtx	control	abtx	control	abtx	control	abtx		
А	2518	88	87	91	0	87	72	88	83		
В	263	5	5	5	0	10	8	9	5		
С	23	2	0	0	0	2	0	0	0		
D**	5	0	0	1	0	0	0	0	0		
Е	54	3	3	2	0	1	2	2	1		
F	31	3	0	1	0	1	0	0	0		
G**	71	0	0	0	20	0	12	0	6		
Н	12	0	4	0	0	0	0	1	0		
I**	12	0	0	0	7	0	0	0	0		
J**	8	0	0	0	4	0	0	0	0		
K**	31	0	0	0	17	0	1	0	0		
L**	107	0	0	0	47	0	6	0	6		
M**	10	0	0	0	6	0	0	0	0		

Table IV. Percentage faecal *E. coli* isolates (total number = 3145) belonging to antimicrobial resistance clusters A–M from steers in a clinical trial having received a single-dose florfenicol treatment on study day 11 (abtx) group and control group by sampling occasion (day 1, 14, 28 and 42).

* Steers receiving florfenicol treatment on day 11 had a significantly greater proportion of isolates belonging to clusters with chloramphenicol resistance than non-treated steers (Chi-square *p*-value < 0.05).

** Clusters containing chloramphenicol resistance.

evident in the contingency tables, that *E. coli* isolates of treated calves tended to return to less resistant cluster groupings as time from treatment increased.

Calves from source B tended to carry a greater proportion of resistant *E. coli* than calves from source A. Furthermore, calves from source B that were weaned one month prior to shipment carried a higher proportion of resistant *E. coli* compared to calves gathered up directly from pasture.

Over time, *E. coli* antimicrobial resistance between sources moved toward a common profile. Though source A steers displayed a greater proportion of susceptible *E. coli* initially, time \times source interaction indicated that source A tended to display more resistance over time, relative to source B steers. Furthermore, abtx \times source interaction indicates source A calves tended to carry a greater proportion of resistant *E. coli* due to treatment than calves from source B.

4. DISCUSSION

Florfenicol is a fluorinated analogue of chloramphenicol that was approved in 1996 in the USA for bovine respiratory disease pathogens. It is routinely used in the USA feedlot industry and has been shown to be clinically successful in treating respiratory disease and fever [4]. It has broad spectrum activity, including respiratory pathogens and gram-negative bacterial flora, and has an extended duration of action. It is found in high concentration in urine and bile, and would be expected to impact faecal *E. coli* flora [2]. Chloramphenicol, the parent

Table V. Cumulative logistic model: GEE Estimates of log odds faecal *E. coli* isolates belonging to clusters with higher level antimicrobial resistance in a trial evaluating single-dose florfenicol treatment of steers on antimicrobial resistance in faecal *E. coli*.

		GEE	Standard		Lower	Upper	Chi-		
		estimate	error	Df	C.I.	C.I.	square	P-value	
Treatment (abtx)									
No		Reference							
Yes		6.34	0.66	1	5.05	7.63	9.64	< 0.0001	
Sampling time	e (time)								
1		Reference							
2		-0.80	0.39	1	-1.57	-0.03	-2.03	0.0427	
3		-0.52	0.40	1	-1.31	0.27	-1.29	0.1962	
4		-0.76	0.39	1	-1.53	0.01	-1.93	0.0542	
Source of stee	ers (source)								
В		Reference							
А		-2.07	0.41	1	-2.87	-1.28	-5.1	< 0.0001	
Weaned one n	nonth prior s	hipment (we	ean)						
Yes		Reference							
No		-0.59	0.29	1	-1.16	-0.02	-2.04	0.0416	
Interaction tin	ne and sourc	e							
Time $= 4$	Source A	1.88	0.53	1	0.83	2.92	3.52	0.0004	
Time $= 4$	Source B	Reference							
Time $= 3$	Source A	1.69	0.57	1	0.57	2.80	2.96	0.003	
Time $= 3$	Source B	Reference							
Time $= 2$	Source A	1.30	0.62	1	0.09	2.51	2.1	0.0357	
Time $= 2$	Source B	Reference							
Time $= 1$	Source A	Reference							
Time $= 1$	Source B	Reference							
Interaction ab	tx and source	e							
abtx = 1	Source A	1.31	0.48	1	0.36	2.25	2.72	0.0065	
abtx = 1	Source B	Reference							
abtx = 0	Source A	Reference							
abtx = 0	Source B	Reference							
Interaction ab	tx and time								
abtx = yes	Time $= 4$	-6.51	0.85	1	-8.17	-4.84	-7.66	< 0.0001	
abtx = yes	Time = 3	-5.89	0.75	1	-7.36	-4.43	-7.9	< 0.0001	
abtx = yes	Time = 2	Reference							
abtx = no	Time $= 4$	Reference							
abtx = no	Time $= 3$	Reference							
abtx = no	Time $= 2$	Reference							
abtx = no	Time $= 1$	Reference							

compound of florfenicol, was prohibited from use in food animals in 1986.

This study has demonstrated a very clear but transitory shift in antimicrobial resistance patterns of faecal *E. coli* of feedlot steers in response to systemic florfenicol treatment. Results here indicate that florfenicol selects for exclusively resistant faecal flora three days post treatment, during the period when the antimicrobial would be expected to be present in the feces. This suggests that post-therapy, treated animals shed a larger proportion of resistant organisms in their faeces.

Treated steers had a significantly higher proportion of isolates belonging to clusters with chloramphenicol resistance, compared to non-treated steers. We hypothesize that this is due to selection of E. coli with a plasmid or chromosomally located flo gene conferring resistance to florfenicol and chloramphenicol [27, 33]. Additionally, florfenicol treatment selected for E. coli isolates that exhibited simultaneous resistance to several other antimicrobial classes. We hypothesize that this resistance co-selection is due to linkage of antimicrobial resistance genes. E. coli have been shown to harbour a large array of plasmids with different antimicrobial resistance genes. Florfenicol treatment likely selected for E. coli with plasmids containing multiple antimicrobial resistance genes. Several plasmid types carrying the flo gene in E. coli isolates of bovine origin have been identified, and multiple resistance was a common feature [7].

Whether multi-resistant *E. coli* persist in the environment, or establish in non-treated animals following casual contact is not known. The 21 untreated pen mates of treated cattle showed no tendency to shed faecal *E. coli* with higher levels of resistance. This finding suggested that the environmental spread of resistant bacteria from treated to untreated steers was limited in this study.

Due to limited resources, we did not confirm all isolates as *E. coli*, so a small proportion of non-*E. coli* organisms almost certainly were included in the analysis. Assuming they occurred sporadically and at random, they would not be expected to affect the analysis. Berge et al. demonstrated no difference in antimicrobial resistance pattern distribution in non-*E. coli* isolated from MacConkey agar compared to isolates biochemically confirmed as *E. coli* [5].

Our study has shown that the antimicrobial resistance associated with florfenicol treatment decreased dramatically over 4 weeks post-treatment. Similar to other range beef populations, this population of steers had generally susceptible flora at sample day 1. It appears that antimicrobial treatment as described here did not have a large long-term effect on E. coli resistance pattern of treated cattle. By study end, some resistant organisms were evident in treated steers, compared to controls. The significance of this small proportion, about 6% of E. coli, is unknown. It is unknown what effect another antimicrobial treatment. florfenicol or other, would have in treated steers

Use of cluster analysis allowed descriptive assessment of antimicrobial susceptibility to multiple antimicrobials. As a result of florfenicol treatment, resistance to chloramphenicol was observed, as well as resistance to other antimicrobials. Higher order clusters were associated with higher orders of resistance, and florfenicol treatment was associated with selection of *E. coli* possessing chloramphenicol, tetracycline, sulfisoxazole, streptomycin, and some beta-lactam resistance. It is unknown if the resistance observed was transferable, so it is not possible to fully assess the direct implication of this observation.

As indicated by the stratified analysis and multinomial model, antibiotic treatment had a strong effect on *E. coli* antimicrobial resistance patterns. However, the model and stratified analysis both indicated that *E. coli* antimicrobial resistance patterns in treated steers moved back toward baseline levels over time. Resistant bacteria may incur a fitness cost to retain resistance genes in the absence of further antibiotic selective pressure [21]. At the end of our study a higher proportion of faecal *E. coli* were resistant than at the time of arrival. It is unknown whether this proportion would further decrease over time, or stabilize at this level.

Though of lesser magnitude, environmental factors influenced level of antimicrobial resistance following treatment. Source differences are intriguing, in that they suggest some attribute or dynamic that existed at arrival and impacted subsequent antimicrobial resistance cluster membership. Though individual antimicrobial use records were not available from ranch sources, antimicrobials were not included in feed prior to feedlot entry. Additionally, ranch personnel indicated that use of antimicrobials for individual animal treatment was very infrequent. The difference between source A and source B was not solely attributed to the fact that some calves were weaned and rounded up one month prior shipment, but that other environmental factors were also involved. It is unknown how these effects would appear after a longer time in the feedlot.

Though E. coli of source A steers appeared more susceptible initially, they paradoxically tended to shift to higher levels of resistance with time, compared to source B. In a similar manner, swine not exposed to antimicrobials as piglets, then exposed subsequently showed a similar response, compared to swine exposed to antimicrobials early in life [10]. Initial antimicrobial susceptibility differences observed between sources decreased over time in the feedlot, as proportions of isolates in source A and source B calves appeared to become similar. Changes observed over time between sources reflect a natural selection toward a uniform flora. This flora is apparently influenced by factors apart from antimicrobial use, possibly including feedstuffs, feed management, and pen environment [9, 16, 20]. This observation has ramifications in future studies, and suggests that source may

impact faecal *E. coli* resistance patterns, but that these patterns will tend to migrate toward the patterns dictated by feedlot environment. This suggests that antimicrobial resistance of feedlot cattle may be partially manipulated by management interventions apart from controlling antimicrobial use in the feedlot, and may suggest that antimicrobial resistance of enteric bacteria in animals not exposed to antimicrobials may still exhibit changes over time [22].

ACKNOWLEDGEMENTS

Supported by the Ethel Austin Martin Program for Human Nutrition (South Dakota State University, Brookings, SD, 57007, USA). Special thanks to C.A. Gates for technical assistance.

REFERENCES

- Anonymous, Performance standards for antimicrobial disk and dilution susceptibility tests for bacteria isolated from animals; Approved standard, 2nd ed., National Committee for Clinical Laboratory Standards, Wayne, Pennsylvania, 2002.
- [2] Adams P.E., Varma K.J., Powers T.E., Lamendola J.F., Tissue concentrations and pharmacokinetics of florfenicol in male veal calves given repeated doses, Am. J. Vet. Res. 48 (1987) 1725–1732.
- [3] Agresti A., Categorical data analysis, John Wiley & Sons, Inc., New York, 2002.
- [4] Bauer A.W., Kirby W.M., Sherris J.C., Turck M., Antibiotic susceptibility testing by a standardized single disk method, Am. J. Clin. Pathol. 45 (1966) 493–496.
- [5] Berge A.C.B., Atwill E.R., Sischo W.M., Assessing antibiotic resistance in fecal *Escherichia coli* in young calves using cluster analysis techniques, Prev. Vet. Med. 61 (2003) 91–102.
- [6] Blake D.P., Hillman K., Fenlon D.R., Low J.C., Transfer of antibiotic resistance between commensal and pathogenic members of the Enterobacteriaceae under ileal conditions, J. Appl. Microbiol. 95 (2003) 428–436.

- [7] Cloeckaert A., Baucheron S., Chaslus-Dancla E., Nonenzymatic chloramphenicol resistance mediated by IncC plasmid R55 is encoded by a *flo*R gene variant, Antimicrob. Agents Chemother. 45 (2001) 2381–2382.
- [8] Courvalin P., Transfer of antibiotic resistance genes between gram-positive and gram-negative bacteria, Antimicrob. Agents Chemother. 38 (1994) 1447–1451.
- [9] Dargatz D.A., Fedorka-Cray P.J., Ladely S.R., Ferris K.E., Green A.L., Headrick M.L., Antimicrobial susceptibility patterns of *Sal-monella* isolates from cattle in feedlots, J. Am. Vet. Med. Assoc. 221 (2002) 268–272.
- [10] Dawson K.A., Langlois B.E., Stahly T.S., Cromwell G.L., Antibiotic resistance in anaerobic and coliform bacteria from the intestinal tract of swine fed therapeutic and subtherapeutic concentrations of chlortetracycline, J. Anim. Sci. 58 (1984) 123–131.
- [11] DeFrancesco K.A., Cobbold R.N., Rice D.H., Besser T.E., Hancock D.D., Antimicrobial resistance of commensal *Escherichia coli* from dairy cattle associated with recent multiresistant salmonellosis outbreaks, Vet. Microbiol. 98 (2004) 55–61.
- [12] Everitt B.S., Cluster analysis, Halsted Press, John Wiley & Sons Inc., New York, 1993.
- [13] Food and Drug Administration, Nuflor[®] (florfenicol), FOI summary, NADA 141-063 (original), Laurel, Md, 1996.
- [14] Hagedorn C., Robinson S.L., Filtz J.R., Grubbs S.M., Angier T.A., Reneau R.J., Determining sources of fecal pollution in a rural Virginia watershed with antibiotic resistance patterns in fecal streptococci, Appl. Environ. Microbiol. 65 (1999) 5522–5531.
- [15] Hanley J.A., Statistical analysis of Correlated Data using GEE, An orientation, Am. J. Epidemiol. 157 (2003) 364–375.
- [16] Hinton M., Rixson P.D., Allen V., Linton A.H., The persistence of drug resistant *Escherichia coli* strains in the majority faecal flora of calves, J. Hyg. (Lond.) 93 (1984) 547– 557.
- [17] Hoyle D.V., Knight H.I., Shaw D.J., Hillman K., Pearce M.C., Low J.C., Gunn G.J., Woolhouse M.E., Acquisition and epidemiology of antibiotic-resistant *Escherichia coli* in a cohort of newborn calves, J. Antimicrob. Chemother. 53 (2004) 867–871.
- [18] Khachatryan A.R., Hancock D.D., Besser T.E., Call D.R., Role of calf-adapted *Escherichia coli* in maintenance of antimicro-

bial drug resistance in dairy calves, Appl. Environ. Microbiol. 70 (2004) 752–757.

- [19] Kleinbaum D.G., Kupper L.L., Muller K.E., Nizam A., Applied regression analysis and other multivariable methods, Duxbury Press, Pacific Grove, CA, 1998, pp. 1–718.
- [20] Langlois B.E., Dawson K.A., Leak I., Aaron D.K., Antimicrobial resistance of fecal coliforms from pigs in a herd not exposed to antimicrobial agents for 126 months, Vet. Microbiol. 18 (1988) 147–153.
- [21] Lenski R.E., The cost of antibiotic resistance – from the perspective of a bacterium, Ciba Found. Symp. 207 (1997) 131–140.
- [22] Marshall B., Petrowski D., Levy S.B., Interand intraspecies spread of *Escherichia coli* in a farm environment in the absence of antibiotic usage, Proc. Natl. Acad. Sci. USA 87 (1990) 6609–6613.
- [23] Mathew A.G., Saxton A.M., Upchurch W.G., Chattin S.E., Multiple antibiotic resistance patterns of *Escherichia coli* isolates from swine farms, Appl. Environ. Microbiol. 65 (1999) 2770–2772.
- [24] McEwen S.A., Fedorka-Cray P.J., Antimicrobial use and resistance in animals, Clin. Infect. Dis. 34 Suppl. 3 (2002) S93–S106.
- [25] Phillips I., Casewell M., Cox T., De Groot B., Friis C., Jones R., Nightingale C., Preston R., Waddell J., Does the use of antibiotics in food animals pose a risk to human health? A critical review of published data, J. Antimicrob. Chemother. 53 (2004) 28–52.
- [26] Saenz Y., Zarazaga M., Lantero M., Gastanares M.J., Baquero F., Torres C., Antibiotic resistance in *Campylobacter* strains isolated from animals, foods, and humans in Spain in 1997-1998, Antimicrob. Agents Chemother. 44 (2000) 267–271.
- [27] Singer R.S., Patterson S.K., Meier A.E., Gibson J.K., Lee H.L., Maddox C.W., Relationship between phenotypic and genotypic florfenicol resistance in *Escherichia coli*, Antimicrob. Agents Chemother. 48 (2004) 4047–4049.
- [28] Smith D.L., Harris A.D., Johnson J.A., Silbergeld E.K., Morris J.J., Animal antibiotic use has an early but important impact on the emergence of antibiotic resistance in human commensal bacteria, Proc. Natl. Acad. Sci. USA 99 (2002) 6434–6439.
- [29] Stabler S.L., Fagerberg D.J., Quarles C.L., Effects of oral and injectable tetracyclines on

bacterial drug resistance in feedlot cattle, Am. J. Vet. Res. 43 (1982) 1763–1766.

- [30] van den Bogaard A.E., Stobberingh E.E., Epidemiology of resistance to antibiotics. Links between animals and humans, Int. J. Antimicrob. Agents 14 (2000) 327–335.
- [31] Wagner B.A., Dargatz D.A., Salman M.D., Morley P.S., Wittum T.E., Keefe T.J., Comparison of sampling techniques for measuring the antimicrobial susceptibility of enteric

Escherichia coli recovered from feedlot cattle, Am. J. Vet. Res. 63 (2002) 1662–1670.

- [32] Ward J.H., Hierarchical grouping to optimize an objective function, J. Am. Stat. Assoc. 58 (1963) 236–244.
- [33] White D.G., Hudson C., Maurer J.J., Ayers S., Zhao S., Lee M.D., Bolton L., Foley T., Sherwood J., Characterization of chloramphenicol and florfenicol resistance in *Escherichia coli* associated with bovine diarrhea, J. Clin. Microbiol. 38 (2000) 4593–4598.

To access this journal online: www.edpsciences.org