



HAL
open science

Draft Genome Sequences of *Mycoplasma auris* and *Mycoplasma yeatsii*, Two Species of the Ear Canal of Caprinae

Emilie Dordet-Frisoni, Eric Baranowski, Aurélien Barré, Alain Blanchard, Marc Breton, Carole Couture, Virginie Dupuy, Patrice Gaurivaud, Daniel Jacob, Claire Lemaitre, et al.

► To cite this version:

Emilie Dordet-Frisoni, Eric Baranowski, Aurélien Barré, Alain Blanchard, Marc Breton, et al.. Draft Genome Sequences of *Mycoplasma auris* and *Mycoplasma yeatsii*, Two Species of the Ear Canal of Caprinae. *Genome Announcements*, 2013, 1 (3), pp.e00280-13. 10.1128/genomeA.00280-13. hal-00907449

HAL Id: hal-00907449

<https://hal.science/hal-00907449>

Submitted on 4 Aug 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

Draft Genome Sequences of *Mycoplasma auris* and *Mycoplasma yeatsii*, Two Species of the Ear Canal of *Caprinae*

Emilie Dordet-Frisoni,^{a,b} Eric Baranowski,^{a,b} Aurélien Barré,^c Alain Blanchard,^{d,e} Marc Breton,^{d,e} Carole Couture,^{d,e} Virginie Dupuy,^{f,g} Patrice Gaurivaud,^{h,i} Daniel Jacob,^{c,*} Claire Lemaître,^{c,*} Lucía Manso-Silván,^{f,g} Macha Nikolski,^{c,j} Laurent-Xavier Nouvel,^{a,b} François Poumarat,^{h,i} Pascal Sirand-Pugnet,^{d,e} Patricia Thébault,^{c,j} Sébastien Theil,^{f,g,*} François Thiaucourt,^{f,g} Christine Citti,^{a,b} Florence Tardy^{h,i}

INRA, UMR 1225, IHAP, Toulouse, France^a; Université de Toulouse, INP, ENVT, UMR 1225, IHAP, Toulouse, France^b; Université de Bordeaux, Centre de Bioinformatique et Génomique Fonctionnelle, Bordeaux, CBiB, Bordeaux, France^c; INRA, UMR 1332 de Biologie du Fruit et Pathologie, Villenave d'Ornon, France^d; Université de Bordeaux, UMR 1332 de Biologie du Fruit et Pathologie, Villenave d'Ornon, France^e; CIRAD, UMR CMAEE, Montpellier, France^f; INRA, UMR 1309 CMAEE, Montpellier, France^g; Anses, Laboratoire de Lyon, UMR Mycoplasmoses des Ruminants, Lyon, France^h; Université de Lyon, VetAgro Sup, UMR Mycoplasmoses des Ruminants, Marcy L'Etoile, Franceⁱ; Université Bordeaux, LaBRI, UMR 5800, Talence, France^j

* Present address: Daniel Jacob and Sébastien Theil, INRA, UMR 1332 de Biologie du Fruit et Pathologie, Villenave d'Omon, France, and Université de Bordeaux, UMR 1332 de Biologie du Fruit et Pathologie, Villenave d'Omon, France; Claire Lemaître, INRIA Rennes - Bretagne Atlantique/IRISA, EPI GenScale, Rennes, France.

We report here the draft genome sequences of *Mycoplasma auris* and *Mycoplasma yeatsii*, two species commonly isolated from the external ear canal of *Caprinae*.

Received 16 April 2013 Accepted 1 May 2013 Published 13 June 2013

Citation Dordet-Frisoni E, Baranowski E, Barré A, Blanchard A, Breton M, Couture C, Dupuy V, Gaurivaud P, Jacob D, Lemaître C, Manso-Silván L, Nikolski M, Nouvel L-X, Poumarat F, Sirand-Pugnet P, Thébault P, Theil S, Thiaucourt F, Citti C, Tardy F. 2013. Draft genome sequences of *Mycoplasma auris* and *Mycoplasma yeatsii*, two species of the ear canal of *Caprinae*. *Genome Announc*. 1(3):e00280-13. doi:10.1128/genomeA.00280-13.

Copyright © 2013 Dordet-Frisoni et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 3.0 Unported license](https://creativecommons.org/licenses/by/3.0/).

Address correspondence to Florence Tardy, florence.tardy@anses.fr, or Christine Citti, c.citti@envt.fr.

Mycoplasma auris and *Mycoplasma yeatsii*, of the hominis and spiroplasma phylogenetic groups, respectively, are two recently described species of the *Mycoplasma* genus (1, 2). Both species appear to have little clinical relevance, yet they are commonly isolated from the external ear canal of *Caprinae* (3–8), often in association with *Mycoplasma agalactiae* (hominis group) and/or *Mycoplasma mycoides* subsp. *capri* (spiroplasma group), two pathogens of small ruminants that are responsible for contagious agalactia (9). Ruminant *Mycoplasma* species of these two remote groups were shown to have exchanged a significant amount of genetic information by horizontal gene transfer (HGT) (10). Thus, the ear canal may offer an ecological niche in which HGT can occur between pathogenic and nonpathogenic *Mycoplasma* species.

Genome sequences are available for *M. agalactiae* and for *M. mycoides* subsp. *capri*, but not for *M. auris* and for *M. yeatsii*. To fill this gap, we report here the complete genome sequence of *M. yeatsii* strain 13926 and *M. auris* strain 15026, isolated in France from ear swabs of a domestic goat (within a flock with a clinical history of mycoplasmosis [7]) and an Alpine ibex (during an abnormal mortality episode with recurrent isolations of *M. agalactiae* [11]), in 2003 and 2008, respectively. Whole-genome sequences were obtained using a combination of Illumina (single reads) and 454 (mate paired with 8-kb insert size), resulting in a 50× (*M. auris*) and 69× (*M. yeatsii*) median coverages. Assembly was performed using Newbler 2.3, and the annotation was conducted using a customized version of the CAAT-Box platform (12) with an automatic preannotation for coding sequences (CDSs) followed by expert validation, as detailed previously (11). Genome analysis and comparisons were mainly conducted using the MolliGen 3.0 platform (13).

Sequence data showed that the estimated genome sizes (767,867 bp and 896,612 bp for *M. auris* and *M. yeatsii*, respectively), the G+C% (27.20 and 25.76% for *M. auris* and *M. yeatsii*, respectively), the gene density (89.13 and 85.24% for *M. auris* and *M. yeatsii*, respectively), and the highest similarity scores of CDSs were consistent with the phylogenetic grouping of these species (2). *M. auris* displayed 26 CDSs (out of 616) with no homolog in ruminant *Mycoplasma* species sequenced so far, one of which has similarities with an *Mycoplasma hominis* CDS annotated as the virulence-associated protein D. Of the 32 *M. yeatsii*-specific CDSs (out of 687), 16 correspond to lipoproteins containing a DUF285 domain, also found in a large family of surface proteins shared by other ruminant *Mycoplasma* species (14). Finally, both genomes displayed sequences related to an integrative conjugative element, with only *M. auris* having a complete element. Altogether, these data indicate that *M. auris* and *M. yeatsii* may be equipped with or able to acquire virulence genes from other *Mycoplasma* species sharing their habitat.

Nucleotide sequence accession numbers. These Whole-Genome Shotgun projects have been deposited at DDBJ/EMBL/GenBank under the accession no. [AORI00000000](https://www.ncbi.nlm.nih.gov/nuclink/AORI00000000) for *M. auris* 15026 and [AORK00000000](https://www.ncbi.nlm.nih.gov/nuclink/AORK00000000) for *M. yeatsii* 13926. The versions described in this paper are the first versions, accession no. [AORI01000000](https://www.ncbi.nlm.nih.gov/nuclink/AORI01000000) and [AORK01000000](https://www.ncbi.nlm.nih.gov/nuclink/AORK01000000), respectively.

ACKNOWLEDGMENTS

Financial support was provided by the EVOLMYCO project (ANR-07-GMGE-001) from ANR to A.B. (principal investigator [PI]), F. Thiaucourt (Co-PI), F.P. (Co-PI), and C.C. (Co-PI).

REFERENCES

1. DaMassa AJD, Tully JG, Rose DL, Pitcher D, Leach RH, Cottew GS. 1994. *Mycoplasma auris* sp. nov., *Mycoplasma cottewii* sp. nov., and *Mycoplasma yeatsii* sp. nov., new sterol-requiring mollicutes from the external ear canals of goats. *Int. J. Syst. Bacteriol.* 44:479–484.
2. Heldtander M, Pettersson B, Tully JG, Johansson KE. 1998. Sequences of the 16S rRNA genes and phylogeny of the goat mycoplasmas *Mycoplasma adleri*, *Mycoplasma auris*, *Mycoplasma cottewii* and *Mycoplasma yeatsii*. *Int. J. Syst. Bacteriol.* 48(Pt 1):263–268.
3. Chazel M, Tardy F, Le Grand D, Calavas D, Poumarat F. 2010. Mycoplasmoses of ruminants in France: recent data from the national surveillance network. *BMC Vet. Res.* 6:32.
4. Cottew GS, Yeats FR. 1982. Mycoplasmas and mites in the ears of clinically normal goats. *Aust. Vet. J.* 59:77–81.
5. DaMassa AJ, Brooks DL. 1991. The external ear canal of goats and other animals as a mycoplasma habitat. *Small Rumin. Res.* 4:85–93.
6. Jimena ON, Laura JM, Elena MM, Alonso NH, Teresa QM. 2009. Association of *Raillietia caprae* with the presence of mycoplasmas in the external ear canal of goats. *Prev. Vet. Med.* 92:150–153.
7. Tardy F, Mercier P, Solsona M, Saras S, Poumarat F. 2007. *Mycoplasma mycoides* subsp. *mycoides* biotype large colony isolates from healthy and diseased goats: prevalence and typing. *Vet. Microbiol.* 121:268–277.
8. Cottew GS, Yeats FR. 1981. Occurrence of mycoplasmas in clinically normal goats. *Aust. Vet. J.* 57:52–53.
9. Corrales JC, Esnal A, De la Fe C, Sánchez A, Assunção P, Poveda JB, Contreras A. 2007. Contagious agalactia in small ruminants. *Small Rumin. Res.* 68:154–166.
10. Sirand-Pugnet P, Lartigues C, Marena M, Jacob D, Barré A, Barbe V, Schenowitz C, Mangelot S, Couloux A, Segurens B, de Daruvar A, Blanchard A, Citti C. 2007. Being pathogenic, plastic, and sexual while living with a nearly minimal bacterial genome. *PLoS Genet.* 3:e75.
11. Tardy F, Baranowski E, Nouvel LX, Mick V, Manso-Silvàn L, Thiaucourt F, Thébault P, Breton M, Sirand-Pugnet P, Blanchard A, Garnier A, Gibert P, Game Y, Poumarat F, Citti C. 2012. Emergence of atypical *Mycoplasma agalactiae* strains harboring a new prophage and associated with an alpine wild ungulate mortality episode. *Appl. Environ. Microbiol.* 78:4659–4668.
12. Frangeul L, Glaser P, Rusniok C, Buchrieser C, Duchaud E, Dehoux P, Kunst F. 2004. CAAT-Box, contigs-assembly and annotation Tool-box for genome sequencing projects. *Bioinformatics* 20:790–797.
13. Barré A, de Daruvar A, Blanchard A. 2004. MolliGen, a database dedicated to the comparative genomics of Mollicutes. *Nucleic Acids Res.* 32:D307–D310.
14. Nouvel LX, Sirand-Pugnet P, Marena MS, Sagné E, Barbe V, Mangelot S, Schenowitz C, Jacob D, Barré A, Claverol S, Blanchard A, Citti C. 2010. Comparative genomic and proteomic analyses of two *Mycoplasma agalactiae* strains: clues to the macro- and micro-events that are shaping mycoplasma diversity. *BMC Genomics* 11:86.