

Semi-classical measures on Quantum Graphs and the Gauss map of the determinant manifold

Yves Colin de Verdière

▶ To cite this version:

Yves Colin de Verdière. Semi-classical measures on Quantum Graphs and the Gauss map of the determinant manifold. 2013. hal-00907569v1

HAL Id: hal-00907569 https://hal.science/hal-00907569v1

Preprint submitted on 21 Nov 2013 (v1), last revised 15 Feb 2014 (v2)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Semi-classical measures on Quantum graphs and the Gauß map of the determinant manifold

Yves Colin de Verdière^{*}

November 21, 2013

Introduction

The so-called Quantum Ergodic Theorem (QE), due mainly to Alexander Shnirelman [Shn74, Shn93, Zel87, CdV85], asserts that the eigenfunctions of the Laplace operator on a smooth closed Riemannian manifold whose geodesic flow is ergodic are equi-distributed on the manifold in the limit of large eigenvalues provided one removes a sub-sequence of density 0. The paper [JSS13] gives QE Theorems in the case of metrics with discontinuities on smooth manifolds: the Assumptions are ergodicity of the geodesic flow which in this context is a Markov process and the fact that recombining geodesics are exceptional. The results apply also to piecewise smooth metrics on simplicial complexes, in particular to metric graphs (also called "quantum graphs"). The geodesic flow is then ergodic, but they are many recombining geodesics. Hence the second assumption is not satisfied for quantum graphs. It is currently believed that QE does not hold in general for a FIXED quantum graph. This is proved for star graphs in [BKW04] and [KMW02] (put $f(x) = x^2$ in Equation (51)). In the litterature, QE for graphs means some property of the eigenfunctions in the limit of a sequence of graphs whose number of vertices is going to infinity (see for example [GKF10]).

In the present paper, our goal is to describe all weak limits of measures $|\phi_j|^2 |dx|_{\vec{l}}$, where (ϕ_j) is an orthonormal basis of eigenfunctions for the Laplacian $\Delta_{\vec{l}}$ on a metric graph (G, \vec{l}) and $|dx|_{\vec{l}}$ is the Riemannian measure, and to get upper bounds on the densities of the associated sub-sequences of eigenfunctions. In particular we show that QE does not hold for a generic metric \vec{l} on a fixed graph, except if the graph is homeomorphic to an interval or to a circle. In order to avoid confusion, we will denote this last property by QEF (Quantum Ergodicity for Fixed graph).

^{*}Université de Grenoble, Institut Fourier, Unité mixte de recherche CNRS-UJF 5582, BP 74, 38402-Saint Martin d'Hères Cedex (France); yves.colin-de-verdiere@ujf-grenoble.fr

If G = (V, E) is a finite graph, we introduce, following ideas in [BG00, BG01], an algebraic sub-manifold Z_G , which we call the "determinant manifold", of the torus $(\mathbb{R}/2\pi\mathbb{Z})^E$ which allows to compute the eigenvalues thanks to the so-called "secular equation". The Gauß map Γ associates to any point of the smooth part Z_G^{reg} of Z_G the half-line of the cone $\mathcal{P}_E := [0, +\infty[^E$ which is orthogonal to Z_G . Let us denote by \mathcal{M}_G the semi-algebraic set which is the closure of the image of Γ . Let us state the main result of this note:

Theorem 0.1 Let us fix G. For a generic metric \vec{l} , the set of non-normalized semi-classical measures is the set of all $\sum_{e \in E} m_e |dt_e|$ where $\vec{m} \in \mathcal{M}_G$ and where $|dt_e|$ is the Riemannian measure on the edge e, i.e. t_e is an arc-length parametrization of the edge e. Moreover the densities of the corresponding eigenfunctions sub-sequences are bounded in terms of the image by Γ of a measure $\mu_{\mathrm{BG},\vec{l}}$ on Z_G^{reg} introduced in [BG00, BG01].

Another related result is the description of all semi-classical measures with *minimal support*, which are analogs of the so-called "scars" in the case of manifolds: the supports of these scars are simple paths joining two vertices of degree one and simple cycles (this is related to the paper [SK03]). From this, we deduce that QEF does not hold for a generic \vec{l} , except for graphs homeomorphic to intervals or circles.

A good introduction to Quantum graphs is the recent book [BK13].

1 The determinant manifold of a quantum graph

In this section, we recall the way to compute the spectra of the Laplacians $\Delta_{\vec{l}}$ on a graph G = (V, E) from an algebraic hyper-surface Z_G of a torus $\mathcal{T}_E = (\mathbb{R}/2\pi\mathbb{Z})^E$ which depends only of the combinatorics of the graph and not of the lengths \vec{l} . This approach is closely related to what people do in several papers like [BG00, BKW04, BW08, Ba12, BB13].

Let us consider a finite connected graph G = (V, E) where loops and multiple edges are allowed. We choose an orientation of the edges. We associate, in the usual way, to G a 1D singular manifold |G| by gluing together #E intervals using the combinatorics of G. A (Riemannian) metric on |G| is given by the lengths of the edges. We will denote by $\vec{l} \in]0, +\infty[^E$ the collection of the lengths and the pair (G, \vec{l}) is called a *Quantum Graph*.

Definition 1.1 We will say that \vec{l} is irrational if the components of \vec{l} are independent over the rational numbers.

To any metric \vec{l} is associated the Riemannian measure on |G| denoted by $|dx|_{\vec{l}} = \sum_{e \in E} |dt_e|$ where $t_e \in [-l_e/2, l_e/2]$ is the arc-length parametrization of the edge e following the given orientation and with the origin at the center of the edge e. We

have $\int_{|G|} |dx|_{\vec{l}} = L$ where $L = \sum_{e \in E} l_e$ is the total length of (G, \vec{l}) . The Laplacian $\Delta_{\vec{l}}$ is the self-adjoint operator on $L^2(|G|, |dx|_{\vec{l}})$ with domain the functions ϕ : $|G| \to \mathbb{C}$ which are in the Sobolev space $\bigoplus_{e \in E} H^2(e)$, are continuous on |G| and satisfy the Kirchhoff condition that the sums of outgoing derivatives of ϕ at each vertex vanish. Then $\Delta_{\vec{l}}\phi$ is given by $-d^2\phi/dt_e^2$ on each edge. The operator $\Delta_{\vec{l}}$ has a discrete spectrum given by $\lambda_j = k_j^2$ with $k_1 = 0 < k_2 \leq \cdots$. We denote by $\phi_j : |G| \to \mathbb{C}$ an associated orthonormal basis of eigenfunctions. Let us remark that we can assume that G has no vertex of degree 2 if we allow multiple edges and loops.

Let us try to compute the non-zero part of the spectrum of $\Delta_{\vec{l}}$: let us look for an eigenfunction ϕ with eigenvalue k^2 with k > 0. On the edge e, we have

$$\phi(t_e) = a_e \cos kt_e + b_e \sin kt_e . \tag{1}$$

Let us define $\xi_e = \cos k l_e/2$ and $\eta_e = \sin k l_e/2$. From the continuity conditions and the Kirchhoff conditions, we get, for the function $\phi = \phi_{a,b}$, a linear system

$$L_{\xi,\eta}(a,b) = 0$$

of 2#E equations with the same number of unknowns, because the values of the function ϕ and its derivatives divided by k at the vertices are (linear) functions of ξ_e and η_e . Let us remark that if we change any (ξ_e, η_e) into $(-\xi_e, -\eta_e)$, the solutions of the new system are changed by $(a_e, b_e) \rightarrow (-a_e, -b_e)$.

The determinant of the system $L_{\xi,\eta}(a, b) = 0$ is an homogeneous polynomial of degree 2 with respect to each $\zeta_e = (\xi_e, \eta_e)$. Hence, it can be expressed as a polynomial of total degree #E in the variables $z_e = (x_e, y_e)$ with $x_e = \cos k l_e$, $y_e =$ $\sin k l_e$ which is of degree 1 w.r. to each pair (x_e, y_e) . Let θ_e be the corresponding angles so that $x_e = \cos \theta_e$, $y_e = \sin \theta_e$. Let us denote this trigonometric polynomial $\delta_G(\theta)$ viewed as a function on the torus $\mathcal{T}_E = (\mathbb{R}/2\pi\mathbb{Z})^E$ whose coordinates are denoted by $\theta = (\theta_e)_{e \in E}$. The following fact is clear from what we said, the number k^2 , k > 0 is an eigenvalue of $\Delta_{\vec{l}}$ if and only if $\delta_G([k\vec{l}]) = 0$ where [] means that we take the components modulo 2π . In other words, $[k\vec{l}]$ belongs to the algebraic hyper-surface Z_G of equation $\delta_G = 0$ of \mathcal{T}_E . Let us remark that, by an elementary check, $\theta = 0$ belongs too to Z_G .

Let us discuss a few elementary property of Z_G :

Theorem 1.1 If $k^2 > 0$ is a non degenerate eigenvalue of $\Delta_{\vec{l}}$, the point [kl] is a smooth point of Z_G and the tangent space to Z_G at that point is given by $\sum_e m_e d\theta_e = 0$ where $m_e = a_e^2 + b_e^2$ for an eigenfunction given by Equation (1). The manifold Z_G is singular in a set of co-dimension at least 1.

We will denote by Z_G^{reg} the open dense subset of Z_G consisting of points where the corresponding eigenspace is of dimension 1.

Questions: as we will see from the examples in Sections 6.2 and 6.3, Z_G can be reducible. For which graphs is Z_G irreducible? Is the set of singular point of co-dimension ≥ 2 . *Proof.*–

Using the homogeneity of the spectrum, i.e. $\forall c > 0, k_j(c\vec{l}) = c^{-1}k_j(\vec{l})$, We can assume that k = 1. In other words Z_G is identified with the $[\vec{l}]$'s so that 1 is an eigenvalue of $\Delta_{\vec{l}}$. Let $z_0 = [\vec{l}] \in Z_G$. We assume that the eigenvalue 1 is non-degenerate for $\Delta_{\vec{l}}$. Close to z_0 , the set Z_G is defined by the equation $k([\vec{l}]) = 1$ meaning that 1 is an eigenvalue of $\Delta_{\vec{l}}$, so that Z_G is smooth if $dk \neq 0$ and the tangent space is the kernel of dk. From the calculation of dk in Appendix A, we get the result.

Using a result of L. Friedlander on the genericity of non degeneracy of the eigenvalues of $\Delta_{\vec{l}}$ [Fr05], we get the fact that the set Z_G^{reg} is dense in Z_G and hence, because Z_G is an algebraic manifold, the set of singular points Z_G^{sing} has co-dimension larger than one.

The singularities of Σ are associated to degenerate eigenvalues. A description of these singularities would be of interest, already for general star graphs or complete graphs. The case of diabolical singularities, corresponding to eigenvalues of multiplicity two, is used in [BG01] in order to study the behavior as $s \to 0$ of the distribution P(s)ds of the level spacing $\lambda_{j+1} - \lambda_j$ defined by

$$P(s)ds = \lim_{\lambda \to \infty} \frac{\#\{j | s \le \lambda_{j+1} - \lambda_j \le s + ds\}}{\#\{j | \lambda_j \le \lambda\}}$$

2 The Gauß map and semi-classical measures

Let us denote by \mathcal{P}_E the cone $[0, +\infty[^E \setminus \{0\}]$. For any $\vec{m} = (m_e)_{e \in E} \in \mathcal{P}_E$ and any \vec{l} , we identify \vec{m} with the measure $\sum_{e \in E} m_e |dt_e|$ on |G|. In other words, if \vec{l} is given, we consider \mathcal{P}_E as a sub-cone of the cone of positive Radon measures on |G|. Our goal is to characterize the measures which are semi-classical limits. We will need a

Definition 2.1 If ϕ is an eigenfunction of some $\Delta_{\vec{l}}$ with eigenvalue $k^2 > 0$, the restriction of ϕ to the edge e writes $\phi(t_e) = a_e \cos kt_e + b_e \sin kt_e$; we denote by μ_{ϕ} the measure on |G| defined by $\mu_{\phi} = \sum_e (a_e^2 + b_e^2) |dt_e|$ which belongs to \mathcal{P}_E .

Definition 2.2 A semi-classical limit for $\Delta_{\vec{l}}$ is a weak limit of a sequence of measures $\sum_{e \in E} |\phi_{k_j}(t_e)|^2 |dt_e|$ on |G| where ϕ_{k_j} are non-zero eigenfunctions of $\Delta_{\vec{l}}$ with eigenvalues $k_j^2 \to +\infty$.

Remark 2.1 We do not use a lift of the measures to the cotangent space as in the Shnirelman Theorem.

Remark 2.2 We do not ask the eigenfunctions to be L^2 -normalized. Our semiclassical limits are in general not probability measures.

We can summarize the results as follows:

- **Theorem 2.1** 1. Let $\mu = \sum_{e \in E} m_e |dt|_e$ be a semi-classical limit for $\Delta_{\vec{l}_0}$, then there exists $\vec{l_{\infty}}$ and an eigenfunction ϕ_{∞} of $\Delta_{\vec{l_{\infty}}}$ associated to the eigenvalue 1 so that $\mu = \mu_{\phi_{\infty}}$.
 - 2. Let us assume that ϕ is an eigenfunction of some $\Delta_{\vec{l_0}}$ with a simple eigenvalue k_0^2 , then μ_{ϕ} is a semi-classical measure for $\Delta_{\vec{l_0}}$ and for any $\Delta_{\vec{l}}$ with \bar{l} irrational.
 - 3. Let \mathcal{G}_G be the set of the \vec{l} so that \vec{l} is irrational and the line $\{[k\vec{l}] \mid k > 0\}$ does not meet Z_G^{sing} . The set \mathcal{G}_G is Baire generic and, if $\vec{l}_0 \in \mathcal{G}_G$, the set of semi-classical measures for $\Delta_{\vec{l}_0}$ is the closure of the image of Z_G^{reg} by the Gauß map.

Proof.-

- 1. Let us consider a sequence of eigenfunctions of $\Delta_{\vec{l}}$ defined by $\phi_{k_j}(t_e) = a_{e,j} \cos k_j t_e + b_{e,j} \sin k_j t_e$. Then for large k_j 's, the measures $\sum_e \phi_{k_j}(t_e)^2 |dt_e|$ have a weak limit if and only if the limits $m_e = \lim_{j \to \infty} (a_{e,j}^2 + b_{e,j}^2)$ exist and this limit measure is given by $\sum_{e \in E} m_e |dt|_e$. By compactness of Z_G , we can take assume that $[k_j l_0]$ converges to $[\vec{l_{\infty}}]$ and we can assume that all components of $\vec{l_{\infty}}]$ are > 0. Then we can again extract a sub-sequence so that the numbers $a_{e,j}$ and $b_{e,j}$ converge to some limit which will be associated, by the closeness of the eigenfunctions equations, to an eigenfunction of $\Delta_{\vec{l_{\infty}}}$ with eigenvalue 1.
- 2. From the assumption and Theorem 1.1, we get that the set Z_G is smooth near $[k_0 \vec{l_0}]$ and transverse to any curve $t \to [t\vec{l}]$ if the components of \vec{l} are all > 0. From this and the Poincaré recurrence Theorem, we can get a sequence $k_j \to +\infty$ so that $[k_j \vec{l_0}] \to [k_0 \vec{l_0}]$ as $j \to \infty$ and we have that $\mu_{\phi_{k_j}} \to \mu_{\phi}$. Similarly, if \vec{l} is irrational, we can find $k_j \to \infty$ so that $[k_j \vec{l}] \in Z_G$ and $[k_j \vec{l}] \to [k_0 \vec{l_0}]$.

3. The genericity comes from the fact that Z_G^{sing} is of co-dimension at least two in \mathcal{T}_E . From the previous part, we know already that each measure in the image of the Gauß map is a semi-classical measure and the same is true for a measure in the closure of this image, because the set of semi-classical measures is closed. On the other hand, we know that each semi-classical measure is the limit of a sequence of $\mu_{\phi_{k_j}}$; because of the assumption on \vec{l} this implies that each semi-classical measure lies in the closure of the image of the Gauß map.

3 The Barra-Gaspard measure, Weyl formula and the densities of semi-classical limits

In [BG00], the authors introduce a \vec{l} -dependent measure on Z_G^{reg} , which we will denote by $\mu_{\text{BG},\vec{l}}$. The manifold Z_G^{reg} is transverse to any vector with > 0 coordinates and hence oriented. The measure $\mu_{\text{BG},\vec{l}}$ is defined by

$$\mu_{\mathrm{BG},\vec{l}} = (2\pi)^{-|E|} |\iota(\vec{l}) \wedge_{e \in E} d\theta_e|$$

 $(\iota(\vec{V})\omega$ is the inner product), which is a Radon measure with > 0 density everywhere on Z_G^{reg} . We will need the following:

Lemma 3.1 If \vec{l} is irrational and $D \subset Z_G^{\text{reg}}$ is a compact domain with piecewise smooth boundary, we have

$$\lim_{K \to \infty} \frac{1}{K} \# \{ k_j \mid 0 < k_j \le K, \ [k_j \vec{l}] \in D \} = \int_D \mu_{\mathrm{BG}, \vec{l}} \, .$$

The same result holds if D is the interior of a piecewise smooth compact sub-set of Z_G^{reg} or by integrating a smooth function $f: Z_G \to \mathbb{R}$ with compact support in Z_G^{reg} :

$$\lim_{K \to \infty} \frac{1}{K} \sum_{0 < k_j \le K, \ [k_j \vec{l}] \in Z_G} f([k_j \vec{l}]) = \int_{Z_G} f\mu_{\mathrm{BG}, \vec{l}} \, dk_{\mathrm{BG}, \vec{l}} \, dk_{\mathrm{BG}$$

Proof.-

Let us choose $\varepsilon > 0$ small enough so that the map $F : [-\varepsilon, \varepsilon] \times D \to \mathcal{T}_E$, defined by $F(t, \theta) = \theta + t\vec{l}$, is a smooth embedding of image D_{ε} . From the unique ergodicity of the Kronecker flows on the tori, we get

$$\lim_{K \to \infty} \frac{1}{K} |\{k \mid 0 < k \le K, [k\vec{l}] \in D_{\varepsilon}\}| = (2\pi)^{-|E|} \operatorname{vol}(D_{\varepsilon})$$
(2)

where |X| denote the Lebesgue measure of a set $X \subset \mathbb{R}$. We observe that $[k\vec{l}]$ belongs to D_{ε} if and only if there exists j so that $[k_j\vec{l}] \in D$ and $k\vec{l} = [k_j\vec{l}] + t\vec{l}$ with $|t| \leq \varepsilon$. The lefthandside of Equation (2) is equal to

$$\lim_{K \to \infty} \frac{2\varepsilon}{K} \#\{k_j \mid 0 < k_j \le K, \ [k_j \vec{l}] \in D\}$$

while, using the definition of $\mu_{\text{BG},\vec{l}}$, the righthandside of Equation (2) is equal to $2\varepsilon\mu_{\text{BG},\vec{l}}(D)$.

We have also

Lemma 3.2 If \vec{l} belongs to \mathcal{G}_G , we have

$$\lim_{K \to \infty} \frac{1}{K} \# \{ k_j \mid 0 < k_j \le K, \ [k_j \vec{l}] \in Z_G \} = \int_{Z_G} \mu_{\mathrm{BG}, \vec{l}} \, .$$

A similar result holds for the integration of a continuous function f on Z_G :

$$\lim_{K \to \infty} \frac{1}{K} \sum_{0 < k_j \le K, \ [k_j \vec{l}] \in Z_G} f([k_j \vec{l}]) = \int_{Z_G} f\mu_{\mathrm{BG}, \vec{l}}$$

Proof.-

The proof is an adaptation of the previous one. We first split $Z_G = D \cup R$ with R a small neighborhood of Z_G^{sing} . The asymptotics of the set of k's with $[kl] \in D$ is given by Lemma 3.1, while an upper bound is given by remarking that the pre-images of the map $F : R \times [-\varepsilon, \varepsilon] \rightarrow \mathcal{T}_E$ given by $F(\theta, t) = \theta + tl$ have a bounded number of elements: this comes from the boundedness of the remainder term in Weyl's formula for $\Delta_{\vec{l}}$ (see [BK13] p. 95).

Let us consider, for $e \in E$, the canonical projection $\pi_e : Z_G \to (\mathbb{R}/2\pi\mathbb{Z})^{E\setminus e}$. The map π_e is orientation preserving and hence we get

$$\int_{Z_G} \mu_{\mathrm{BG},\vec{l}} = \sum_{e \in E} l_e \int_{(\mathbb{R}/2\pi\mathbb{Z})^{E \setminus e}} \deg_e(\theta) |\wedge \widehat{d\theta_e}|$$

where $\deg_e(\theta)$ is the cardinal of $\pi_e^{-1}(\theta)$. The fact that δ_G is of degree 1 w.r. to each $(\cos \theta_e, \sin theta_e)$ implies that $\deg_e \leq 2$.

In fact $\deg_e \equiv 2$ as follows from Lemma 3.2 and the Weyl asymptotic formula (see [BK13] p. 95)

Theorem 3.1 We have (Weyl's law)

$$\lim_{K \to \infty} \frac{1}{K} \# \{ 0 \le k \le K | [k\vec{l}] \in Z_G \} = \frac{L}{\pi} .$$

The Barra-Gaspard measure is related to the densities of sequences of eigenvalues giving a given semi-classical limit. This density vanishes in general, so we have to say it in the following way:

Theorem 3.2 Let $D \subset \mathcal{P}_E$ be a compact domain with smooth boundary and consider a sequence of eigenfunctions ϕ_{k_j} of some $\Delta_{\vec{l}}$ with \vec{l} generic. Let us assume that all semi-classical limits of the $\mu_{\phi_{k_j}}$ lie in D, then the density of the sequence k_j is bounded from above in terms of $\mu_{BG,\vec{l}}$ as follows:

$$\limsup_{K \to \infty} \frac{\#\{k_j \le K\}}{K} \le \mu_{\mathrm{BG},\vec{l}} \left(\Gamma^{-1}(D)\right) .$$

The converse is true; for any generic \vec{l} and for any compact domain $D \subset \mathcal{P}_E$, there exists a sub-sequence ϕ_{k_j} of density $\mu_{\mathrm{BG},\vec{l}}(\Gamma^{-1}(D))$ so that the semi-classical limits associated to sub-sequences of ϕ_{k_j} belongs to D.

Proof.-

Let $D' \subset \mathcal{P}_E$ a neighborhood of D. Then for j large enough, $\Gamma_{\vec{l}}([k_j \vec{l}])$ belongs to D'. So that, from a slight extension of Lemma 3.2 to domains in Z_G , we get the upper bound $\mu_{\mathrm{BG},\vec{l}}(\Gamma_{\vec{l}}^{-1}(D'))$.

For, the converse, it is enough to take the sequence of k_j 's so that $[k_i \vec{l}]$ belongs to $\Gamma^{-1}(D)$.

We get the following

Corollary 3.1 If \vec{l} belongs to Z_G^{reg} and \vec{l} is generic, let ϕ be an eigenfunction of $\Delta_{\vec{l}}$ with eigenvalue 1. Then, for all neighborhoods U of μ_{ϕ} , there exists sub-sequences of positive densities of the eigenfunctions of $\Delta_{\vec{l}}$ so that the semi-classical limits of sub-sequences belongs to $\Gamma(U)$.

Moreover, we have the following link with the Liouville measure:

Theorem 3.3 Let $\Gamma_{\vec{l}}$ be the map Γ normalized so that the sums $\sum m_e |dt_e|$ are probabilities, then

$$\int \Gamma_{\vec{l}}(\theta) d\mu_{\mathrm{BG},\vec{l}} = \frac{1}{\pi} \left(\sum_{e} |dt_{e}| \right) \; .$$

This can be reformulated as "The average of the Gauß map with respect to $\mu_{\mathrm{BG},\vec{l}}$ is the Liouville measure".

Proof.-

Le us define

$$A_e(K) = \frac{l_e}{2K} \sum_{0 \le k_j \le K} (a_{j,e}^2 + b_{j,e}^2)$$

where the normalized eigenfunction ϕ_j writes, on the edge e, $\phi_j(t_e) = a_{j,e} \cos kt_e + b_{j,e} \sin kt_e$. From the local Weyl Theorem and the fact that

$$\int_{e} |\phi_j(t_e)|^2 dt_e = \frac{l_e}{2} (a_{j,e}^2 + b_{j,e}^2) + O(1/k) ,$$

we get that

$$\lim_{K \to \infty} A_e(K) = l_e/\pi \; .$$

On the other hand, we have

$$\Gamma_{\vec{l}}([k_j\vec{l}]) = \frac{1}{2}(a_{j,e}^2 + b_{j,e}^2) + O(1/k) \; .$$

Using Lemma 3.2, we get

$$\lim_{K \to \infty} A_e(K) = l_e \int_{Z_G} \left(\Gamma_{\vec{l}}(\theta) \right)_e d\mu_{\mathrm{BG},\vec{l}} \,.$$

From this, we deduce:

Corollary 3.2 If \vec{l} is generic and there exists an eigenfunction ϕ of $\Delta_{\vec{l}}$ so that μ_{ϕ} is not the Liouville measure, then QEF does not holds for (G, \vec{l}) .

4 Scars

Let us look at semi-classical measures with small supports; we have the:

Theorem 4.1 Let G be given. The minimal supports of the semi-classical measures for a generic \vec{l} are the simple cycles and the simple paths between two vertices of degree 1 of G. These measures are extremal points of the convex hull of all semi-classical measures.

We start with the

Lemma 4.1 Let $K \subset |G|$ be the support of a semi-classical measure μ . Then K is an union of edges of G and every vertex of K is of degree ≥ 2 in K or is of degree 1 in G.

Proof of the Lemma.-

From the first Assumption of Theorem 2.1, we deduce that $\mu = \mu_{\phi}$ where ϕ is an eigenfunction of some $\Delta_{l_{\infty}}$ on G. The Lemma follows then from the Kirchoff conditions applied to ϕ .

Proof of the Theorem.-

• Simple paths joigning to vertices of degree 1 are minimal supports: Let γ be a simple oriented path whose N vertices are $(1, 2, \dots, N)$ and choose a vector \vec{l} so that the lengths of the edges of the path are $\vec{l_0} = (1/2, 1, 1, \dots, 1, 1/2)$. If we parametrize this path by $0 \leq t \leq N-2$, the function ϕ_1 defined by $\phi_1 = \cos t$ on γ and 0 outside is an eigenfunction of $\Delta_{\vec{l}}$ with eigenvalue π^2 . I claim that we can choose such a vector \vec{l} extending $\vec{l_0}$ so that the eigenvalue π^2 of $\Delta_{\vec{l}}$ is simple: if it is not, let $X := \ker (\Delta_{\vec{l}} - \pi^2)$, and l_t defined by $(l_t)_e = (1 - t)l_e$ for $e \notin \gamma$ and $(l_t)_e = l_e$ for $e \in \gamma$. Then the quadratic form on X defined by $f \to \langle \Delta f | f \rangle$ vanishes on ϕ_1 and is > 0 in any other direction (because as we have seen the support of ϕ_1 is minimal). It implies that π^2 is a simple eigenvalue for $\Delta_{\vec{l}}$ for small non-zero t. It follows that the uniform measure on γ is a semi-classical measure for this choice of \vec{l} and hence for all generic \vec{l} 's.

The minimality follows from Lemma 4.1.

• Simple cycles are minimal supports: Let γ be a simple oriented cycle whose N vertices are $(1, 2, \dots, N - 1, N = 1)$ and choose a vector \vec{l} so that the lengths of the edges of γ are 1. By an argument similar to the case of a simple path, we get that the uniform measure on γ is a semi-classical measure for this choice of \vec{l} and all irrational \vec{l} 's.

Again the minimality follows from lemma 4.1.

• If K is the support of a semi-classical measure, K contains a simple path joigning 2 vertices of degree 1 of G or a simple cycle of G: if K contains no simple cycles, K is a forest and every vertex of degree 1 of K is also of degree 1 in G. So that every sub-tree of K contains simple paths joigning 2 vertices of degree 1 in G.

As a Corollary, we get

Theorem 4.2 For a generic \vec{l} , QEF holds for $\Delta_{\vec{l}}$ if and only if |G| is homeomorphic to a circle or to an interval.

Any connected graph which is not homeomorphic to a circle or an interval contains as strict sub-graphs a simple cycle or a path whose end are of degree one. The Theorem 3.3 implies that the image of Z_G^{reg} by the Gauß map contains at least one line distinct from the Liouville measure. The proof is completed by applying Corollary 3.2.

5 Ergodicity of the geodesic flow on graphs

In the case of a connected quantum graph, the classical dynamics is ergodic as shown from the study of the associated Perron-Frobenius operator done in [BG01]. The phase space Z of a Quantum graph $G = (V, E, \vec{l})$ (the unit cotangent bundle) can be identified with the set of oriented edges: to a point x of an oriented edge, we associate the unit co-vector pointing in the direction given from the orientation. Using the probabilities given from the transition coefficients in Appendix B, we get the geodesic flow on Z as a Markov process. We have the

Theorem 5.1 If G is connected and is not homeomorphic to a circle, the geodesic flow on (V, E, \vec{l}) is ergodic.

Proof.-

It is enough to prove that the only functions invariant by the Perron-Frobenius semi-group $(T_t)_{t>0}$ are the constant functions. We see first that such a function f has to be constant on each oriented edge by using at a point x the invariance by T_t for t small. Moreover if Gis connected and is not homeomorphic to a circle, the geodesic flow is transitive: for any pair $x, y \in Z$ there exists a geodesic from xto y. Let us choose f constant on the oriented edges and invariant by the semi-group $(T_t)_{t>0}$. Let us choose an edge e_0 so that $\forall x \in$ Z, $f(x) \leq f(e_0)$ and $x_0 \in e_0$. We have $T_t f(x_0) = f(x_0)$ and $T_t f(x_0) =$ $\sum_{\gamma:[0,t]\to Z} w(\gamma) f(\gamma(t))$. Using the fact that $\sum_{\gamma:[0,t]\to Z} w(\gamma) = 1$, we get that $f(y) = f(x_0)$ if there is a geodesic γ so that $\gamma(0) = x_0$ and $\gamma(t) = y$. This holds for all $y \in Z$ if the geodesic flow is transitive.

From the previous study, we see that the QEF Theorem is not valid for Quantum Graphs even if the geodesic flow is ergodic, which is the case for all connected graphs which are not homeomorphic to a circle, This is due to the fact that there are a lot of recombining geodesics: for example in a star graph, they are many geodesics starting from x in an edge e_1 an coming back to x in the opposite direction following a geodesic containing k_1 times the edge e_1 and k_2 times the edge e_2 with $k_1, k_2 \ge 1$ and with different orders.

6 Examples



Figure 1: The 3 examples: the black circles are the vertices and the white circles are the middle of the edges

6.1 The star graph with three edges

Let us consider for simplicity the case of a star graph with three edges G = (V, E)where l_1, l_2, l_3 are the lengths of three edges. The function δ_G can be computed as follows: we assume that, we parametrize the *j*th edge by $0 \le t_j \le l_j$ where 0 corresponds to the vertex of degree 1. Then an eigenfunction of eigenvalue k^2 will be of the form $\phi(t_j) = a_j \cos kt_j$. Putting $x_j = \cos kt_j$ and $y_j = \sin kt_j$, we get the three equations

$$a_1x_1 = a_2x_2 = a_3x_3, a_1y_1 + a_2y_2 + a_3y_3 = 0 .$$

Hence

$$\delta_G(x,y) = x_1 x_2 y_3 + x_2 x_3 y_1 + x_3 x_1 y_2 \, .$$

Singularities of Σ : The surface $\Sigma_G \subset \mathcal{T}_E$ is invariant by the eighth translations whose vectors have coordinates 0 or π . We will restrict ourselves to the intersection of Σ_G with the cube $C = [-\pi/2, \pi/2]^3$. The seven other cubes are deduced by translations. The surface Σ_G is also invariant by the central symmetry $\sigma: (\theta_j) \to (-\theta_j)$.

We can cover Σ by four types of charts and use adapted coordinates taken from the set $X_j = x_j/y_j, Y_j = y_j/x_j$:

- 1. In the domain $x_1x_2x_3 \neq 0$, we get $Y_1 + Y_2 + Y_3 = 0$ which is smooth.
- 2. In the domain $y_1y_2y_3 \neq 0$, we get $X_1X_2+X_2X_3+X_3X_1=0$ which is singular at the eight points where all x_j vanish. They are "diabolical" points located at the vertices of the cubes.
- 3. In the domains, $x_1x_2y_3 \neq 0$ and $x_1y_2y_3 \neq 0$, Σ_G is smooth.

The boundary of the closure of the intersection $\Sigma_0 = \Sigma \cap [-\pi/2, \pi/2]^3$ is the union of the six edges of C which have no vertex $(-\pi/2, -\pi/2, -\pi/2)$ or $(\pi/2, \pi/2, \pi/2)$. We call this closed path the *equator* of C. The surface Σ_0 is a smooth 2-disk whose boundary is the equator of C.



Figure 2: the surface Z_{star} and the equator of the cube; the map Γ from Z_{star} to the semi-classical measures.

The set \mathcal{M}_G is the cone $\sum_{i=1}^{3} m_i^2 \leq 2 \sum_{1 \leq i < j \leq 3} m_i m_j$. In particular, this closed set is not equal to \mathcal{P}_E .

6.2 The eight figure

Let us denote by G_8 the graph with one vertex and two edges (two loops). We have

$$\delta_{G_8}(\theta_1, \theta_2) = \sin \frac{\theta_1}{2} \sin \frac{\theta_2}{2} \sin \frac{\theta_1 + \theta_2}{2} \, .$$

The set Z_{G_8} is the union of the circles $\theta_1 = 0$, $\theta_2 = 0$ and $\theta_1 + \theta_2 = 0$. The image by the Gauß map is the union of the three lines $m_2 = 0$, $m_1 = 0$ and $m_1 = m_2$. The associated densities are up to global normalization l_1 , l_2 , $l_1 + l_2$. This implies that there exactly three semi-classical limits for an irrational length vector. The Liouville measure $|dt_1| + |dt_2|$ has relative density $\frac{1}{2}$. On this example we see that the set of semi-classical measure is not even connected and that the scars give measures which are extremal point of their convex hull.

6.3 The cherry

Let G_C be the cherry graph consisting of one loop with an attached edge. Let us denote by l_1 the length of the loop and by l_2 the length of the edge. We get for Z_{G_C} the equation:

$$\sin \theta_1 \sin \theta_2 + 2(1 - \cos \theta_1) \cos \theta_2 = 0.$$

The set Z_{G_C} is the union of the three circles

$$\{\theta_1 = 0\} \cup \{\tan \theta_2 + 2\tan \theta_1/2 = 0\}$$

The circle $\{\theta_1 = 0\}$ correspond to the scars localized on the loop ; the other part corresponds to a continuum of semi-classical measures $m_1|dt_1| + m_2|dt_2|$ with $m_1l_1 + m_2l_2 = 1, 1 \le m_2/m_1 \le 4.$

7 Limits of graphs

From the characterization of the spectrum of $\Delta_{\vec{l}}$ as the set of k^2 so that $[k\vec{l}]$ belongs to Z_G , on can still define the spectrum of $\Delta_{\vec{l}}$ for \vec{l} a real vector. In particular, one can take some lengths to be zero and the others > 0. What is then the interpretation of this "spectrum"?

Let G be a graph as before. Given a set of edges $X \subset E$, we introduce a new graph G_X obtained by contracting all edges $e \in E$. From the point of view of Riemannian, metrics, the graph G_X can be interpreted as the graph G where the lengths of the edges in X vanish. We have the

Lemma 7.1 For any connected graph G = (V, E) not homeomorphic to the circle or the interval, there exists a set $X \subset E$ so that G_X is homeomorphic to the star graph with three edges, to G_8 or to G_C .

The first case occur if G is a tree not reduced to an interval, the second one if $b_1(X) \ge 2$ and the third one if $b_1 = 1$ and G is not a circle.

Let $(Z_G)_X$ be the intersection of Z_G with the torus $\mathcal{T}_X = \{\theta_e = 0, \forall e \in X\}$. We have clearly an identification of $(Z_G)_X$ with Z_{G_X} .

Any non singular point of Z_{G_X} is in this way associated to an non singular point of Z_G .

A The derivatives of the eigenvalues w.r. to the edge lengths

This Lemma is contained in [Fr05]:

Lemma A.1 If λ is a non degenerate eigenvalue of $\Delta_{\vec{l}}$ with a normalized eigenfunction $\phi(t_e) = a_e \cos kt_e + b_e \sin kt_e$ on the edge e, the derivative of λ w.r. to l_e is given by

$$\frac{\partial \lambda}{\partial l_e} = -\lambda \left(a_e^2 + b_e^2 \right) \; .$$

Proof.-

Let us put $t = t_e$, $I =] - l_e/2$, $l_e/2[$ and choose a function $\psi \in C_o^{\infty}(I, \mathbb{R})$ and the metric $g_u = \exp(4u\psi(t))|dt|^2$ on the edge e and independent of u on the other edges. Then we have $dl_e/du = 2\int_I \psi(t)|dt|$,

$$\|\phi\|_{u}^{2} = \|\phi\|'_{0}^{2} + \int_{I} e^{2u\psi(t)}\phi^{2}(t)|dt| ,$$

and the Dirichlet integral

$$q_u(\phi) = q'_0(\phi) + \int_I e^{-2u\psi(t)} {\phi'}^2(t) |dt| ,$$

where $_{0}'$ denotes the integrals on the other edges which are independent of u. Let now take the u-derivative of the Rayleigh quotient $q_{u}(\phi)/\|\phi\|_{u}^{2}$ at u = 0, we get

$$d\lambda/du = -2\int_{I}\psi(t)(\phi'(t)^{2} + \lambda\phi(t)^{2})|dt$$

from which the result follows.

B Appendix: the transition probabilities for a quantum graph

We want to describe the way a wave arriving at a vertex of a quantum graph splits into several waves. Let us denote by O the vertex of degree d and by e_j , $j = 1, \dots, d$ the d-edges arriving at O. Let us denote the arc-length coordinate x_j along e_j starting from 0 at O. Let us consider a function f_1 compactly supported in $]0, +\infty[$ and a wave u(x,t) defined on e_1 by $u(x_1,t) = f_1(x_1+t)+g_1(x_1-t)$ and, on e_j for $2 \le j \le d$, by $u(x_j,t) = g_2(x_j-t)$. From the Kirchhoff conditions, we get $f_1(t) + g_1(-t) = g_2(-t)$ and $f'_1(t) + g'_1(-t) + (d-1)g'_2(-t) = 0$. Integrating the second equation, we get $f_1(t) = g_1(-t) + (d-1)g_2(-t)$ (we see that the integration constant vanishes by putting t = 0). So that we get

$$g_1(t) = \frac{2-d}{d} f_1(-t), \ g_2(t) = \frac{2}{d} f_1(-t) \ .$$

The transition probabilities are the numbers

$$p_{j,j} = \left(\frac{d-2}{d}\right)^2$$

and for $i \neq j$,

$$p_{i,j} = \left(\frac{2}{d}\right)^2 \; .$$

The quantum graph is said *classically ergodic* if the following Markov process defined on the unit tangent bundle of the metric graph is ergodic: follow the edges with unit speed and arriving at a vertex use the transition probabilities defined before. We can assume that the graph has no vertex of degree two.

C Appendix: the case $\vec{l} = (1, 1, \cdots, 1)$

In this case the spectrum is given by the k_j^2 , $k_j \ge 0$ so that $[k] \in Z_G$. Using Weyl asymptotics, the number of k_j 's with multiplicity is 2#E. On the other hand, we find from [Ni84, Ni87, Catt97], that the eigenvalues of $\Delta_{\vec{1}}$ are given in terms of the eigenvalues μ_l , $l = 1, \dots, \#V$, of the weighted adjacency operator defined

$$A_G f(i) = \frac{1}{d_i} \sum_{j \sim i} f(j) \; .$$

The result is the following one for the k_j :

- For each $|\mu_l| < 1$, 2 values of k defined by $\cos k = \mu_l$.
- The value k = 0 with multiplicity $1 + b_1$.
- The value $k = \pi$ with multiplicity $b_1 + 1$ if G is bipartite and $b_1 1$ if-not.

At the end, we have, in the bipartite case $2\#E = 2(\#V - 2) + 2b_1 + 2$, and in the non-bipartite case $2\#E = 2(\#V - 1) + 2b_1$. Both formulas give the Euler formula for G:

$$\#V - \#E = 1 - b_1$$
.

References

- [Ba12] Ram Band. The Nodal Count $\{0, 1, 2, 3, ...\}$ Implies The Graph is a Tree. ArXiv 1212.6710 (2012).
- [BB13] Ram Band & Gregory Berkolaiko. Universality of the momentum band density of periodic networks. ArXiv 1304.6028 (2013).
- [BG00] Felipe Barra & Pierre Gaspard, On the Level Spacing Distribution in Quantum Graphs, *Journal of Statistical Physics* **101**:283–319 (2000).
- [BG01] Felipe Barra & Pierre Gaspard, Classical dynamics on graphs, *Physical Review E* 63:066215 (2001).

- [BK13] Gregory Berkolaiko & Peter Kuchment, Introduction to quantum graphs, Mathematical Surveys and Monographs (AMS), 186 (2013).
- [BKW04] Gregory Berkolaiko, Jonathan Keating & Brian Winn, No Quantum Ergodicity for Star Graphs. Commun. Math. Phys. 250:259–285 (2004).
- [BW08] Gregory Berkolaiko & Brian Winn, Relationship between scattering matrix and spectrum of quantum graphs. ArXiv 0801.4104 (2008).
- [Catt97] Carla Cattaneo. The Spectrum of the Continuous Laplacian on a Graph. Monatshefte für Mathematik 124:215–235 (1997).
- [CdV85] Yves Colin de Verdière, Ergodicité et fonctions propres du laplacien. Commun. Math. Phys. 102:497–502, (1985).
- [Fr05] Leonid Friedlander, Genericity of simple eigenvalues for a metric graph, Israel J. Math., 146:149–156 (2005).
- [GKF10] Sven Gnutzman, Jon P. Keating & Fabien Piotet. Eigenfunction statistics on quantum graphs, Ann. Phys. 325:2595-2640 (2010).
- [JSS13] Dmitry Jakobson, Yuri Safarov & Alexander Strohmaier, The semiclassical theory of discontinuous systems and ray-splitting billiards (with an Appendix of Yves Colin de Verdière). *American Journal of Maths (to appear)* and *ArXiv* 1301.6783v1 (2013).
- [KMW02] Jon P. Keating, Jens Marklof & Brian Winn. Value Distribution of the Eigenfunctions and Spectral Determinants of Quantum Star Graphs. *Commun. Math. Phys.* 241:421–452 (2003).
- [Ni84] Serge Nicaise, Some results on spectral theory over networks, applied to nerve impulse transmission, Orthogonal polynomials and applications (Barle-Duc, 1984), Lecture Notes in Math. (Springer), 1171:532–541 (1985).
- [Ni87] Serge Nicaise, Approche spectrale des problèmes de diffusion sur les réseaux, Séminaire de Théorie du Potentiel, Paris, No. 8, Lecture Notes in Math. (Springer), 1235:120–140 (1987).
- [SK03] Holger Schanz & Tsampikos Kottos. Scars on Quantum Networks Ignore the Lyapunov Exponent. *Phys. Rev. Lett.* **90**:234101 (2003).
- [Shn74] Alexander I. Shnirelman, Ergodic properties of eigenfunctions. Uspehi Mat. Nauk 29:181–182, (1974).
- [Shn93] Alexander I. Shnirelman, On the asymptotic properties of eigenfunctions in the regions of chaotic motion. In V. Lazutkin KAM theory and semiclassical approximations to eigenfunctions. Ergebnisse der Mathematik und ihrer Grenzgebiete (3), 24. Springer-Verlag, Berlin, (1993].

[Zel87] Steve Zelditch, Uniform distribution of eigenfunctions on compact hyperbolic surfaces. Duke Math. J. 55:919–941 (1987).