



HAL
open science

Crystal structure of lanthanum triiodate iodic acid $\text{La}[\text{IO}_3]_3[\text{HIO}_3]$

Mohamed Benabdallah Taouti, Aissa Gacemi, Djamal Benbortal, Isabelle
Gautier-Luneau

► **To cite this version:**

Mohamed Benabdallah Taouti, Aissa Gacemi, Djamal Benbortal, Isabelle Gautier-Luneau. Crystal structure of lanthanum triiodate iodic acid $\text{La}[\text{IO}_3]_3[\text{HIO}_3]$. *Zeitschrift für Kristallographie - New Crystal Structures*, 2008, 223, pp.179-180. 10.1524/ncrs.2008.0074 . hal-00908370

HAL Id: hal-00908370

<https://hal.science/hal-00908370>

Submitted on 24 Nov 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Crystal structure of lanthanum triiodate iodic acid, $\text{La}(\text{IO}_3)_3(\text{HIO}_3)$


Mohamed Benabdallah Taouti^{I,II,III}, Aissa Gacemi^{II}, Djamal Benbertal^{II} and Isabelle Gautier-Luneau^{*I}

^I Institut Néel CNRS, associated to Université Joseph Fourier, 25 avenue des Martyrs, Bât. F BP166, 38042 Grenoble Cedex 9, France

^{II} Université de Laghouat, Laboratoire des Sciences Fondamentales, BP 37G Laghouat 03000, Algeria

^{III} Université Hadj Lakhdar, 5 Avenue Chahid Boukhlof, Batna 05000, Algeria

Received January 31, 2008, accepted and available on-line July 17, 2008; CSD no. 409943



Abstract

$\text{HI}_4\text{LaO}_{12}$, monoclinic, $P12_1/c1$ (no. 14), $a = 10.685(1) \text{ \AA}$, $b = 7.626(1) \text{ \AA}$, $c = 14.314(1) \text{ \AA}$, $\beta = 110.23(1)^\circ$, $V = 1094.4 \text{ \AA}^3$, $Z = 4$, $R_{\text{gt}}(F) = 0.019$, $wR_{\text{ref}}(F^2) = 0.040$, $T = 293 \text{ K}$.

* Correspondence author (e-mail: isabelle.gautier-luneau@grenoble.cnrs.fr)

Source of material

0.5 mmol of anhydrous lanthanum chloride and 2.5 mmol of lithium iodate were dissolved in 40 ml of 7 M nitric acid. The solution was evaporated slowly at 323 K. After six days, transparent and colourless prismatic crystals suitable for X-ray crystal structure analysis were obtained. They were filtered, washed with deionised water and dried at room temperature (yield 0.33 g, 78 %).

Experimental details

Hydrogen atom from iodic acid molecule could not be located by difference Fourier analysis.

Discussion

Until now, $\text{La}(\text{IO}_3)_3(\text{HIO}_3)$ was only characterized by powder XRD (JCPDS file no 51-0075). The asymmetric unit of the title compound is made up of one lanthanum(III) cation, three iodate anions and one iodic acid ligand. The coordination number of the lanthanum cation is nine according to the bond valence sum concept with a valence value of 3.054 (for the expected value of 3) [1,2]. The La—O bond lengths are in the range of 2.428(3) to 2.674(3) Å and the O—La—O angles vary from 65.00(1)° to 143.98(1)°. The nine oxygen atoms come from seven iodate anions (two IIO_3 , two I_2O_3 , three I_3O_3 groups) and two iodic acid molecules (HI_4O_3) (figure, top). The polyhedron can be described as a monocapped square antiprism with the $\text{O}42^{\text{v}}$ (symmetry code v: $2-x, 2-y, 2-z$) capping atom corresponding to the longest La—O bond length. The mean deviations from calculated mean planes of the two square faces ($\text{O}11-\text{O}21-\text{O}31-\text{O}41$) and ($\text{O}12^{\text{ii}}-\text{O}22^{\text{iv}}-\text{O}32^{\text{i}}-\text{O}33^{\text{iii}}$) are equal to 0.480(3) Å and 0.122(3) Å, respectively (symmetry code i: $x, y+1, z$; ii: $-x, y+1/2, 1/2-z$; iii: $1-x, -y, 1-z$; iv: $x, 1/2-y, z+1/2$). The dihedral angle between these faces is equal to 3.1(4)°. The environment of iodine is formed by three I—O strong bonds (mean bond length is 1.810(3) Å, except for the $\text{I}4-\text{O}43$ longest bond length of the iodic acid for which $\text{O}43$ is protonated with $d(\text{I}4-\text{O}43) = 1.896(3) \text{ \AA}$ corresponding to an AX_3E configuration of the iodate anion and three additional weak bonds (bond lengths in the range 2.468(3) Å to 3.215(3) Å). This coordination is described as an octahedral with the iodine atom displaced off centre along the ternary axis [3]. Iodate groups shows two different coordination scheme towards lanthanum atom: IIO_3 , I_2O_3 and I_4O_3 groups bridge two cations in a bis-monodentate way via $\text{O}11$, $\text{O}12$, $\text{O}21$, $\text{O}22$, $\text{O}41$ and $\text{O}42$ respectively, and I_3O_3 group bridge three cations in a tris-monodentate way via its three oxygens. The crystal structure reveals a three-dimensional packing in which each lanthanum atom is connected to eight other lanthanum atoms through iodate bridges (figure, bottom). One lanthanum neighbour is linked via two I_3O_3 and two I_4O_3 groups, through an inversion centre, leading to eight-membered rings and to the shortest La...La distance equal to

5.739 Å. An other lanthanum neighbour is linked *via* two IO₃ groups leading also to eight-membered ring and to a La...La distance equal to 6.045 Å. The six other lanthanum neighbours are connected by simple iodate bridges. The distances La...La are equivalent to two by two and they vary from 7.161 to 7.626 Å.

Table 1. Data collection and handling.

Crystal:	colorless prism, size 0.10 × 0.11 × 0.12 mm
Wavelength:	Ag K α radiation (0.56087 Å)
μ :	80.66 cm ⁻¹
Diffractometer, scan mode:	Nonius Kappa CCD, φ/ω
$2\theta_{\max}$:	42.82°
$N(hkl)_{\text{measured}}$, $N(hkl)_{\text{unique}}$:	20421, 2509
Criterion for I_{obs} , $N(hkl)_{\text{gt}}$:	$I_{\text{obs}} > 2 \sigma(I_{\text{obs}})$, 2348
$N(\text{param})_{\text{refined}}$:	154
Programs:	SIR92 [5], SHELXL-97 [6]

Table 2. Atomic coordinates and displacement parameters (in Å²).

Atom	Site	<i>x</i>	<i>y</i>	<i>z</i>	<i>U</i> ₁₁	<i>U</i> ₂₂	<i>U</i> ₃₃	<i>U</i> ₁₂	<i>U</i> ₁₃	<i>U</i> ₂₃
La	4e	0.31892(2)	0.26551(3)	0.36111(2)	0.0057(1)	0.0069(1)	0.0060(1)	0.00031(8)	0.00166(9)	-0.00025(9)
I(1)	4e	-0.04958(3)	0.13711(4)	0.22986(2)	0.0080(1)	0.0091(1)	0.0103(1)	-0.0013(1)	0.0026(1)	0.0005(1)
I(2)	4e	0.14970(3)	0.33711(4)	0.06055(2)	0.0098(1)	0.0082(1)	0.0073(1)	0.0004(1)	0.0033(1)	0.0002(1)
I(3)	4e	0.46513(2)	-0.23389(4)	0.36078(2)	0.0062(1)	0.0074(1)	0.0081(1)	0.00030(9)	0.0029(1)	0.0002(1)
I(4)	4e	0.70401(3)	0.31737(4)	0.44496(2)	0.0088(1)	0.0075(1)	0.0086(1)	0.00114(9)	0.0018(1)	-0.0001(1)
O(11)	4e	0.1215(3)	0.0701(4)	0.2887(3)	0.006(1)	0.012(2)	0.027(2)	-0.002(1)	-0.003(1)	0.003(1)
O(12)	4e	-0.1045(3)	-0.0864(4)	0.1922(3)	0.009(1)	0.011(2)	0.022(2)	-0.002(1)	0.003(1)	-0.001(1)
O(13)	4e	-0.0558(3)	0.2143(5)	0.1089(2)	0.018(2)	0.023(2)	0.013(2)	-0.002(1)	0.004(1)	0.007(1)
O(21)	4e	0.2673(3)	0.2699(5)	0.1800(2)	0.018(2)	0.025(2)	0.008(2)	0.008(1)	0.005(1)	0.005(1)
O(22)	4e	0.2738(3)	0.3996(4)	0.0056(2)	0.019(2)	0.014(2)	0.015(2)	0.001(1)	0.011(1)	0.002(1)
O(23)	4e	0.1071(3)	0.5509(4)	0.0924(2)	0.023(2)	0.009(2)	0.018(2)	0.005(1)	0.007(1)	-0.000(1)
O(31)	4e	0.3724(3)	-0.0356(4)	0.3175(2)	0.023(2)	0.011(2)	0.017(2)	0.009(1)	0.003(1)	0.001(1)
O(32)	4e	0.3362(3)	-0.3978(4)	0.3197(2)	0.015(2)	0.014(2)	0.016(2)	-0.007(1)	0.003(1)	-0.001(1)
O(33)	4e	0.4703(3)	-0.2195(4)	0.4900(2)	0.010(1)	0.020(2)	0.008(2)	-0.000(1)	0.003(1)	-0.001(1)
O(41)	4e	0.5539(3)	0.2930(5)	0.3362(2)	0.009(1)	0.032(2)	0.009(2)	-0.001(1)	-0.001(1)	0.002(1)
O(42)	4e	0.6721(3)	0.5311(4)	0.4845(3)	0.026(2)	0.007(2)	0.018(2)	0.003(1)	0.011(1)	-0.002(1)
O(43)	4e	0.8180(3)	0.4000(5)	0.3783(3)	0.016(2)	0.020(2)	0.022(2)	-0.002(1)	0.012(1)	0.005(1)

Acknowledgments. Part of this work and scientific stays for two of us (MBT and AG) was funded by the Comité Mixte d'Evaluation et de Prospective de coopération interuniversitaire franco - algérien (CMEP, project N° 04 MDU 618).

References

- Zocchi, F.: Accurate bond valence parameters for M—O bonds (M = C, N, La, Mo, V). *Chem. Phys. Lett.* **421** (2006) 277- 280.
- Zocchi, F.: Accurate bond valence parameters for lanthanide-oxygen bonds. *J. Mol. Struct.: THEOCHEM* **805** (2007) 73 -78.
- Brown, I. D.: Bond valence as an aid to understanding the stereochemistry of O and F complexes of Sn(II), Sb(III), Te(IV), I(V) and Xe(VI). *J. Solid State Chem.* **11** (1974) 214-233.
- Farrugia, L. J.: WinGX suite for small-molecule single-crystal crystallography. *J. Appl. Crystallogr.* **32** (1999) 837-838.
- Altomare, A.; Casciarano, M.; Giacovazzo, C.; Guagliardi, A.: Completion and refinement of crystal structures with SIR92. *J. Appl. Crystallogr.* **27** (1994) 435.
- Sheldrick, G. M.: SHELXL-97. Program for the Refinement of Crystal Structures. University of Göttingen, Germany 1997.