



**HAL**  
open science

# Charge renormalisation in a mean-field approximation of QED

Jérémy Sok

► **To cite this version:**

| Jérémy Sok. Charge renormalisation in a mean-field approximation of QED. 2014. hal-00909168v2

**HAL Id: hal-00909168**

**<https://hal.science/hal-00909168v2>**

Preprint submitted on 15 May 2014

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# Charge renormalisation in a mean-field approximation of QED

Sok Jérémy

Ceremade, UMR 7534, Université Paris-Dauphine,  
Place du Maréchal de Lattre de Tassigny,  
75775 Paris Cedex 16, France.

May 15, 2014

## Abstract

We study the Bogoliubov-Dirac-Fock (BDF) model, a no-photon, mean-field approximation of quantum electrodynamics that allows to study relativistic electrons interacting with the vacuum. It is a variational model in which states are represented by Hilbert-Schmidt operators. We prove a charge renormalisation formula that holds close to the non-relativistic limit: the density of a ground state is shown to be integrable although such a state is known not to be trace-class. We prove that we can take the non-relativistic limit by keeping track of the vacuum polarisation. We get an altered Hartree-Fock model due to the screening effect.

## 1 Introduction

The relativistic quantum theory of electrons is based on the Dirac operator [24]:  $mc^2\beta - \sum_{j=1}^3 i\hbar c\alpha_j \cdot \partial_j$ . Here  $c$  is the speed of light,  $m$  the mass of electron,  $\hbar$  the Planck's constant,

$$\beta := \begin{pmatrix} \text{id}_{\mathbb{C}^2} & 0 \\ 0 & -\text{id}_{\mathbb{C}^2} \end{pmatrix}, \quad \alpha_j := \begin{pmatrix} 0 & \sigma_j \\ \sigma_j & 0 \end{pmatrix} \in \text{End}(\mathbb{C}^4),$$

where the  $\sigma_j$ 's are the Pauli matrices:

$$\sigma_1 = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \quad \sigma_2 = \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix}, \quad \sigma_3 = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}. \quad (1)$$

The Dirac operator is a self-adjoint operator acting on  $\mathfrak{H} := L^2(\mathbb{R}^3, \mathbb{C}^4)$  and whose domain is  $H^1(\mathbb{R}^3, \mathbb{C}^4)$ . In the one-particle theory, the energy of a free particle  $\psi \in L^2(\mathbb{R}^3, \mathbb{C}^4)$  is given by  $\langle D_0\psi, \psi \rangle$ , while the spectrum of  $D_0$  is  $(-\infty, -mc^2] \cup [mc^2, +\infty)$ . According to Dirac's interpretation, all the negative energy states are already occupied by "virtual" electrons, the so-called Dirac sea. By the Pauli principle a real electron can only have positive energy.

In this paper we study the Bogoliubov-Dirac-Fock (BDF) model which is a mean-field approximation of Quantum Electrodynamics (QED). This model, introduced by Chaix and Iracane in [2], enables us to consider a system of relativistic electrons interacting with the vacuum in the presence of an electrostatic field. This paper is a continuation of previous works by Hainzl, Gravejat, Lewin, Séré, Siedentop, Solovej [12, 8, 9, 11, 10, 7] and Sok (unpublished work [23]). In this paper we will extend some results of [7] and of [10].

We use relativistic units  $\hbar = c = 4\pi\epsilon_0 = 1$  and set the bare particle mass equal to 1. The fine structure constant is written  $\alpha$ . The free Dirac operator is written  $D^0 =$

$-i\boldsymbol{\alpha} \cdot \nabla + \beta$ , furthermore we write  $\mathfrak{H} := L^2(\mathbb{R}^3, \mathbb{C}^4)$  and define  $P_-^0$  (resp.  $P_+^0$ ) as the negative (resp. positive) spectral projector of  $D_0$ .

We will not recall here how the BDF energy is derived from QED but refer the reader to [2] or [8, Appendix]. Let us just say that the starting point is the Hamiltonian of QED  $\mathbb{H}_{\text{QED}}$ , defined on the electronic Fock space  $\mathcal{F}_{el}$ . The mean-field approximation consists in restricting the Hamiltonian of QED  $\mathbb{H}_{\text{QED}}$  to "Hartree-Fock" states, the so-called BDF states.

These BDF states are fully characterized by their one-body density matrix (1pdm)  $P$ , an orthogonal projector of  $L^2(\mathbb{R}^3, \mathbb{C}^4)$ . For instance, the projector  $P_-^0$  is the 1pdm of the free vacuum  $\Omega_0$  of the Fock space  $\mathcal{F}_{el}$ . Taking  $P_-^0$  as a reference state, we consider the reduced 1pdm  $Q := P - P_-^0$ . Not all projectors are admissible: a projector  $P$  defines a BDF state if and only if the difference  $P - P_-^0$  is Hilbert-Schmidt.

*Remark 1.* We recall that a Hilbert-Schmidt operator is a compact operator  $Q$  whose integral kernel  $Q(x, y)$  is square-integrable, or equivalently whose singular values form a sequence in  $\ell^2$ . If this sequence is in  $\ell^1$ , then the corresponding operator is trace-class.

Let  $\Omega_P$  be a BDF state with 1pdm  $P$ . The *formal* difference of the energy  $\langle \Omega_P | \mathbb{H} | \Omega_P \rangle$  of the state  $\Omega_P$  and that of  $\Omega_0$  gives a function of  $Q$ , the so-called BDF energy.

We assume the presence of an external density of charge  $\nu$  (real-valued) of finite Coulomb norm:

$$D(\nu, \nu) = \|\nu\|_{\mathcal{C}}^2 := 4\pi \int \frac{|\widehat{\nu}(k)|^2}{|k|^2} dk = \iint \frac{\nu(x)\nu(y)^*}{|x-y|} dx dy. \quad (2)$$

The last equality holds for suitable  $\nu$  (for instance  $\nu \in \mathcal{C} \cap L^{6/5}(\mathbb{R}^3)$ ).

Formally the BDF energy of a state with reduced 1pdm  $Q$  is:

$$\begin{cases} \text{Tr}_{P_-^0} (D_0 Q) - \alpha D(\rho_Q, \nu) + \frac{\alpha}{2} (D(\rho_Q, \rho_Q) - \text{Ex}[Q]), \\ \text{Tr}_{P_-^0} (D_0 Q) := \text{Tr} \{ P_-^0 (D_0 Q) P_-^0 + P_+^0 (D_0 Q) P_+^0 \}, \\ \text{Ex}[Q] := \iint \frac{|Q(x, y)|^2}{|x-y|} dx dy. \end{cases} \quad (3)$$

Here,  $\alpha > 0$  is the coupling constant,  $Q(x, y)$  the integral kernel of the operator  $Q$  and  $\rho_Q$  is its density:  $\rho_Q(x) = \text{Tr}_{\mathbb{C}^4} (Q(x, x))$ . We recognize the kinetic energy, the interaction energy with  $\nu$ , the direct term and the exchange term as in Hartree-Fock theory.

This expression is not always well defined, in particular the formula for the density  $\rho_Q$  makes sense *a priori* only if  $Q$  is (locally) trace-class.

An ultraviolet cut-off  $\Lambda > 0$  is needed: many choices are possible. In [8, 9, 11, 10], Hainzl *et al.* have considered a "sharp" cut-off in which  $L^2(\mathbb{R}^3, \mathbb{C}^4)$  is replaced by its subspace  $\mathfrak{H}_\Lambda$  made of functions whose Fourier transforms vanish outside a ball  $B(0, \Lambda)$ .

In [11], Hainzl *et al.* proposed another BDF energy based on an altered Dirac operator  $\mathcal{D}^0$  and on its spectral projectors

$$\mathcal{P}_\pm^0 := \chi_{\mathbb{R}_\pm^*}(\mathcal{D}^0) \quad (4)$$

In fact Hainzl *et al.* studied the periodized Hamiltonian  $\mathbb{H}_L$  in a finite box  $[-\frac{L}{2}, \frac{L}{2}]$  (with periodic boundary conditions). Setting an ultraviolet cut-off, the problem becomes finite dimensional: for  $L$  large enough they prove there exists a unique ground state which tends to  $\mathcal{P}_-^0$  as  $L$  tends to  $+\infty$ . Thus the BDF energy with respect to this minimizer ("substracting  $\langle \Omega_{\mathcal{P}_-^0} | \mathbb{H} | \Omega_{\mathcal{P}_-^0} \rangle$ ") gives a more relevant model.

The operator  $\mathcal{D}^0$  has the same structure as the Dirac operator:  $\mathcal{D}^0 := \boldsymbol{\alpha} \cdot \mathbf{g}_1(-i\nabla) + \beta g_0(-i\nabla)$  and it satisfies the following equation:

$$\mathcal{D}^0 = D_0 + \frac{\alpha}{2} \frac{\text{sgn}(\mathcal{D}^0)(x, y)}{|x-y|}. \quad (5)$$

Here  $g_0$  and  $\mathbf{g}_1$  are smooth functions of  $B(0, \Lambda)$ .

In this paper the energy functional  $\mathcal{E}_{\text{BDF}}^\nu$  is defined on a subspace  $\mathcal{K}$  of  $\mathfrak{S}_2(\mathfrak{H}_\Lambda)$ , made of convex combinations of reduced 1pdm's of form  $P - \mathcal{P}_-^0$ . The set  $\mathcal{K}$  is properly defined

in the next section and  $\mathcal{E}_{\text{BDF}}^\nu$  is defined as in (3) except that we replace the  $P_-^0$ -trace by a  $\mathcal{P}_-^0$ -trace:

$$\text{Tr}_0(\mathcal{D}^0 Q) := \text{Tr}\{\mathcal{P}_-^0(\mathcal{D}^0 Q)\mathcal{P}_-^0 + \mathcal{P}_+^0(\mathcal{D}^0 Q)\mathcal{P}_+^0\}, \quad (6)$$

A global minimizer of  $\mathcal{E}_{\text{BDF}}^\nu$  is interpreted as the polarized vacuum in the presence of  $\nu$ .

The charge of a state  $Q \in \mathcal{K}$  is given by  $\text{Tr}_0(Q)$ . Thus the ground state of a system with  $M$  electrons is given by a minimizer of  $\mathcal{E}_{\text{BDF}}^\nu$  over the corresponding charge sector.

Furthermore, we define then the energy functional for  $q \in \mathbb{R}$ :

$$\begin{cases} E_{\text{BDF}}^\nu(q) & := \inf \{\mathcal{E}_{\text{BDF}}^\nu(Q), Q \in \mathcal{Q}(q)\}, \\ \mathcal{Q}(q) & := \{Q \in \mathcal{K}, \text{Tr}_0(Q) = q\}. \end{cases}$$

The question becomes: does there exist a minimizer for  $E_{\text{BDF}}^\nu(q)$ ?

In [10], Hainzl *et al.* proved that a sufficient condition for the existence is the validity of binding inequalities at level  $q$ :

$$\forall q' \in \mathbb{R} \setminus \{0, q\}, \quad E_{\text{BDF}}^\nu(q) < E_{\text{BDF}}^\nu(q - q') + E_{\text{BDF}}^0(q'). \quad (7)$$

A much more difficult task is to check that these inequalities hold.

In [10], the authors showed the following.

Let a density  $\nu \in L^1(\mathbb{R}^3, \mathbb{R}_+) \cap \mathcal{C}$ , an integer  $0 \leq M < \int \nu + 1$  and a cut-off level  $\Lambda_0 > 0$  be given, then there exists minimizer for  $E_{\text{BDF}}^\nu(M)$  provided  $\alpha \leq \varepsilon_0(\nu, \Lambda_0)$  for some number  $\varepsilon_0(\nu, \Lambda_0) > 0$ .

In [23] we proved that  $E_{\text{BDF}}^0(1)$  admits a minimizer provided that  $\alpha, \Lambda^{-1}$  and  $L := \alpha \log(\Lambda)$  are small enough. In other words, surprisingly an electron can bind alone in the Dirac sea without any external density, due to the vacuum polarisation.

In both cases the results hold in the non-relativistic regime  $\alpha \ll 1$ .

Let  $M \in \mathbb{Z}$ : a minimizer for  $E_{\text{BDF}}^\nu(M)$  satisfies a self-consistent equation of the form [10]

$$Q + \mathcal{P}_-^0 = \chi_{(-\infty, \mu]} \left( \mathcal{D}^0 + \alpha \left( (\rho_Q - \nu) * \frac{1}{|\cdot|} - \frac{Q(x, y)}{|x - y|} \right) \right) =: \chi_{(-\infty, \mu]}(D_Q). \quad (8)$$

Here,  $\mu$  is a Lagrange multiplier due to the charge constraint  $M$ , interpreted as a chemical potential. For  $M > 0$ , it is positive, the projector  $\chi_{(-\infty, 0)}(D_Q)$  is interpreted as the 1pdm of the polarized vacuum while  $\chi_{[0, \mu]}(D_Q)$  is the 1pdm of the "real" electrons. For  $\alpha$  sufficiently small, the last projector is indeed of rank  $M$ . Furthermore in the limit  $\alpha \rightarrow 0$ ,  $\Lambda_0 > 0$  fixed, its scaling by  $\alpha^{-1}$  tends (up to extraction) to a minimizer of the Hartree-Fock energy  $\mathcal{E}_{\text{HF}}^Z$  for  $M$  electrons and  $Z := \int \nu$ , restricted to  $L^2(\mathbb{R}^3, \mathbb{C}^2 \oplus 0)$ .

In [23], a similar result is obtained with a minimizer for  $\mathcal{E}_{\text{BDF}}^0(1)$  in the non-relativistic limit  $\alpha \rightarrow 0$ ,  $\alpha \log(\Lambda) := L_0$  fixed, the limit is then the Choquard-Pekar model [15].

In this paper we show that, assuming  $L = \alpha \log(\Lambda) \leq L_0$ , there exists a minimizer for  $E_{\text{BDF}}^\nu(M)$  as soon as  $M < \int \nu + 1$  and  $\alpha \leq \alpha_1(\nu, L)$ . The nonrelativistic limit is an altered Hartree-Fock model: writing  $Z = \int \nu$  and  $a = (\frac{2}{3\pi}L)/(1 + \frac{2}{3\pi}L) < 1$  the energy is

$$\forall \Gamma \in \mathfrak{S}_1(L^2(\mathbb{R}^3, \mathbb{C}^4)), \quad 0 \leq \Gamma \leq 1, \quad \text{Tr}(\Gamma) = M :$$

$$\mathcal{E}_{\text{nr}}^Z(\Gamma) := \frac{1}{2} \text{Tr}(-\Delta \Gamma) - Z(1 - a) \text{Tr}\left(\frac{1}{|\cdot|} \Gamma\right) + \frac{1}{2} \left\{ \|\rho_\Gamma\|_{\mathcal{C}}^2 - \text{Ex}[\Gamma] \right\} - \frac{a}{2} \|\rho_\Gamma\|_{\mathcal{C}}^2.$$

The vacuum polarizes due to the presence of  $\nu$  and the electrons: the positive charge  $\nu$  attracts a cloud of negative charge which makes it appear smaller (hence the term  $Z(1 - a)$ ) while the electrons repelled them resulting to an attractive well created by the distortion (hence the term  $-\frac{a}{2} \|\rho_\Gamma\|_{\mathcal{C}}^2$  like in a polaron model). This result gives a wider range of existence of ground state in the space of parameters  $(\alpha, \Lambda)$  compared to that of [10], where the quantity  $\alpha \log(\Lambda_0)$  is neglected and considered as  $\frac{\alpha}{\alpha \rightarrow 0}(1)$ .

To prove it, it is necessary to have a good understanding of a minimizer  $Q_0$  and of its density  $\rho_{Q_0}$ . In [7] the authors proved that, in the simplified model without the exchange term, the density of a minimizer is integrable. This is a natural result: the distortion of

the vacuum due to a finite number of charged particles with finite Coulomb energy should also be finite.

Mathematically speaking however this is a non-trivial fact because a minimizer for  $E_{\text{BDF}}^\nu(M)$  is *not* trace-class. As in [7] we prove that, assuming that  $L$  is small enough and  $M, \|\nu\|_{\mathcal{C}}^2 \lesssim \log(\Lambda)$ , then the density  $\rho_Q$  of a minimizer  $Q$  is in  $L^1 \cap \mathcal{C}$ . Moreover, the following *charge renormalisation* formula holds:

$$\int (\rho_Q - \nu) =: Z_3(M - Z) \simeq \frac{M - Z}{1 + \frac{2}{3\pi}L}, \quad (9)$$

where  $Z_3$  is interpreted as the *renormalization constant* [6]. This means that the total observed charge  $\int (\rho_Q - \nu)$  is different from the real charge  $M - Z$  of the system.

The quantity  $L = \alpha \log(\Lambda)$  is related to  $Z_3$ . In the reduced BDF model where the exchange term is neglected, Gravejat *et al.* showed in [7] that the density  $\rho_Q$  of a minimizer of the reduced energy  $E_{\text{rBDF}}^\nu(M)$  is radial as soon as  $\nu$  is radial and that, in this case, away from the origin, the electrostatic potential of the system is

$$\alpha(\rho_Q - \nu) * \frac{1}{|\cdot|}(x) \underset{x \rightarrow +\infty}{\sim} \frac{\alpha Z_3(M - Z)}{|x|}.$$

In the full model we were unable to prove such behaviour at infinity but we think this is true. Taking  $L$  small corresponds then to considering  $Z_3$  close to 1.

The main contribution of this paper is the integrability result stating that the density of a minimizer is in  $L^1$  together with the charge renormalisation formula (9). It cannot be easily obtained from [7], the presence of the exchange term complicates the study. In our results, we were unable to remove the technical conditions  $M, \|\nu\|_{\mathcal{C}}^2 \lesssim \log(\Lambda)$ . We emphasize here that we can prove the same results with another choice of cut-off considered in [7], the one consisting in replacing  $\mathcal{D}^0$  by  $D_0(1 - \frac{\Delta}{\Lambda^2})$  in  $L^2(\mathbb{R}^3, \mathbb{C}^4)$ .

The paper is organized as follows: in the next section we properly define the variational problem  $\mathcal{E}_{\text{BDF}}^\nu$  and states the main results.

In Section 3, we derive two fixed point schemes from the equation satisfied by a minimizer, using the Cauchy expansion. Moreover *a priori* estimates are proved in Subsection 3.2.

In Section 4 we prove important estimates on a term of the Cauchy expansion (" $Q_{1,0}$ ") and prove Theorem 1.

Section 5 is devoted to prove estimates for the fixed point method and apply it to prove that the density of a minimizer is in  $L^1$  (under some assumptions).

We prove the formula of charge renormalization (Theorem 2) and the existence of minimizers close to the nonrelativistic limit (Theorem 3) in Section 6.

The nonrelativistic energy is studied in Appendix B. The very technical Appendix C is devoted to prove Proposition 1. We prove Lemma 8 which is used for Sections 4 and 5 in Appendix A.

*Remark 2* (Fourier transform). Throughout this paper, the Fourier transform  $\mathcal{F}$  is defined as the extension of

$$\forall f \in L^2(\mathbb{R}^3) \cap L^1(\mathbb{R}^3) : \widehat{f}(p) := \frac{1}{(2\pi)^{3/2}} \int_{\mathbb{R}^3} f(x) e^{-ip \cdot x} dx.$$

*Remark 3* (Form of  $\mathcal{D}^0$ ). The operator  $\mathcal{D}^0$  was first studied by Lieb and Siedentop in [18] in another context. We know  $\mathbf{g}_1(-i\nabla) = \frac{-i\nabla}{|-i\nabla|} g_1(-i\nabla)$  and  $g_0, g_1$  are radial functions satisfying

$$\forall p \in B(0, \Lambda), |p| \leq g_1(p) \leq g_0(p)|p| \text{ and } 1 \leq g_0(p) \leq 1 + \text{Cst} \times \alpha \log(\Lambda). \quad (10)$$

We define

$$m := \inf \sigma(|\mathcal{D}^0|). \quad (11)$$

For  $\alpha \log(\Lambda)$  and  $\alpha$  sufficiently small,  $m$  is equal to  $g_0(0)$ .

Useful estimates on  $g_0, \mathbf{g}_1$  are proved in [23].

## 2 Description of the model and main results

**BDF Energy** We assume there is an external density of charge  $\nu$  (real-valued) of finite Coulomb norm ( $\|\nu\|_{\mathcal{C}} < +\infty$ ).

Let us recall our choice of cut-off: following [10], we replace  $D_0$  by  $\mathcal{D}^0$  and work in  $\mathfrak{H}_\Lambda$ , defined by

$$\mathfrak{H}_\Lambda := \{\psi \in L^2(\mathbb{R}^3, \mathbb{C}^4), \text{supp } \widehat{\psi} \subset B_{\mathbb{R}^3}(0, \Lambda)\}, \quad \Lambda > 0.$$

We write  $\mathfrak{S}_p(\mathfrak{H}_\Lambda)$  the Schatten class of compact operators  $A$  in  $\mathfrak{H}_\Lambda$  such that  $\text{Tr}(|A|^p) < +\infty$  [22]. The set of  $\mathcal{P}_-^0$ -trace operators is [10]:

$$\mathfrak{S}_1^{\mathcal{P}_-^0} := \{Q \in \mathfrak{S}_2(\mathfrak{H}_\Lambda), Q^{++}, Q^{--} \in \mathfrak{S}_1(\mathfrak{H}_\Lambda)\} \quad (12)$$

where  $Q^{\varepsilon_1 \varepsilon_2} := \mathcal{P}_{\varepsilon_1}^0 Q \mathcal{P}_{\varepsilon_2}^0$ . This set is a Banach space with

$$\|Q\|_{\mathfrak{S}_1^{\mathcal{P}_-^0}} := \|Q^{+-}\|_{\mathfrak{S}_2} + \|Q^{-+}\|_{\mathfrak{S}_2} + \|Q^{--}\|_{\mathfrak{S}_1} + \|Q^{++}\|_{\mathfrak{S}_1}. \quad (13)$$

We recall that  $\text{Tr}_0(|\mathcal{D}^0|(Q^{++} - Q^{--}))$  is the kinetic energy functional.

We work in a subset of this space, namely

$$\mathcal{K} := \{Q, -\mathcal{P}_-^0 \leq Q \leq \mathcal{P}_+^0\} \cap \mathfrak{S}_1^{\mathcal{P}_-^0} \subset \{Q, Q^* = Q\} \cap \mathfrak{S}_1^{\mathcal{P}_-^0}. \quad (14)$$

It is the closed convex hull of the  $P - \mathcal{P}_-^0 \in \mathfrak{S}_1^{\mathcal{P}_-^0}$ , where  $P$  is an orthogonal projection.

The density  $\rho_Q$  must be defined consistently with the usual formula when  $Q$  is (locally) trace-class and it must also be of finite Coulomb energy.

Let  $Q$  be in  $\mathfrak{S}_1^{\mathcal{P}_-^0}$ , then  $\rho_Q$  is defined by duality:

$$\forall V \in \mathcal{C}', QV \in \mathfrak{S}_1^{\mathcal{P}_-^0} \text{ and } \text{Tr}_0(QV) = \mathcal{C}'\langle V, \rho_Q \rangle_{\mathcal{C}}. \quad (15)$$

The map  $Q \in \mathfrak{S}_1^{\mathcal{P}_-^0} \mapsto \rho_Q \in \mathcal{C}$  is continuous [7, Proposition 2].

The exchange term is well defined: thanks to Kato's inequality [1, 11, 8]

$$\begin{aligned} \frac{2}{\pi} \iint \frac{|Q(x, y)|^2}{|x - y|} dx dy &\leq \text{Tr}(|\nabla|Q^2) \leq \text{Tr}(|D_0|Q^2) = \text{Tr}\{|D_0|^{1/2}Q^2|D_0|^{1/2}\} \\ \text{and for } Q \in \mathcal{K} : &\leq \text{Tr}\{|D_0|^{1/2}(Q^{++} - Q^{--})|D_0|^{1/2}\} \leq \text{Tr}_{\mathcal{P}_-^0}(\mathcal{D}^0 Q), \end{aligned} \quad (16)$$

The BDF energy is defined as follows:

$$\mathcal{E}_{\text{BDF}}^\nu(Q) := \text{Tr}_{\mathcal{P}_-^0}(\mathcal{D}^0 Q) - \alpha D(\nu, \rho_Q) + \frac{\alpha}{2} \left( D(\rho_Q, \rho_Q) - \iint \frac{|Q(x, y)|^2}{|x - y|} dx dy \right), \quad Q \in \mathcal{K}. \quad (17)$$

As said in the introduction we define the energy functional  $E_{\text{BD}}^\nu(q)$  by the infimum over  $\mathcal{Q}(q) = \{Q \in \mathcal{K}, \text{Tr}_{\mathcal{P}_-^0}(Q) = q\}$ .

For  $M \in \mathbb{N}^*$ , let us say that the problem  $E_{\text{BDF}}^\nu(M)$  has a minimizer: as pointed out in [10, 7] such a minimizer  $\gamma' = \gamma + N$  must be of the following form:

$$\begin{cases} \gamma + \mathcal{P}_-^0 = \chi_{(-\infty, 0)} \{ \mathcal{D}^0 + \alpha((\rho[\gamma'] - \nu) * \frac{1}{|\cdot|} - R[\gamma']) \} =: \chi_{(-\infty, 0)}(D_{\gamma'}), \\ N = \chi_{(0, \mu]} \{ \mathcal{D}^0 + \alpha((\rho_{\gamma'} - \nu) * \frac{1}{|\cdot|} - (R_{\gamma'})) \} = \sum_{j=1}^{M_0} |\psi_j\rangle \langle \psi_j|, \\ \text{so } D_{\gamma'} \psi_j = \mu_j \psi_j \text{ and we write: } n := \rho_N = \sum_j |\psi_j|^2. \end{cases} \quad (18)$$

We choose  $0 \leq \mu_1 \leq \mu_2 \leq \dots \leq \mu_{M_0} = \mu < m$ . *A priori*  $M_0 \neq M$  but in our regime they are equal (Lemma 3). Indeed in the spirit of [8] the equation of the dressed vacuum  $\gamma$  enables us to say that  $(\gamma', \rho_{\gamma'} - \nu)$  is the only fixed point of some function  $F^{(1)}$  defined in (a ball of) the Banach space  $\mathcal{X}_1 = \mathbf{Q}_1 \times \mathcal{C}$  where

$$\|Q\|_{\mathbf{Q}_1}^2 = \|Q\|_{\mathbb{T}}^2 := \iint (\tilde{E}(p) + \tilde{E}(q)) |\widehat{Q}(p, q)|^2 dp dq.$$

*Notation 4.* For a density  $\rho \in \mathcal{C}$  we write:  $v_\rho = v[\rho] := \rho * \frac{1}{|\cdot|}$ .

For an operator  $Q \in \mathfrak{S}_1^{P_0}$  with integral kernel  $Q(x, y)$  we define the operator  $R_Q = R[Q]$  by the formula:

$$R_Q(x, y) := \frac{Q(x, y)}{|x - y|}.$$

We remark that  $\text{Ex}[Q] = \text{Tr}(R_Q^* Q) =: \|Q\|_{\text{Ex}}^2$ .  
Moreover we write

$$B_Q := v[\rho_Q - \nu] - R_Q \text{ and } D_Q := \mathcal{D}^0 + \alpha B_Q. \quad (19)$$

**The Cauchy expansion** Let  $\gamma' = \gamma + N$  be a minimizer for  $E_{\text{BDF}}^\nu(M)$ , the decomposition being that of (18).

*Notation 5.* Throughout this paper  $n := \rho_N$ , moreover we write  $\rho_\gamma'$  for  $\rho_{\gamma'}$  and the double prime means  $-\nu$  is added:

$$\rho_\gamma'' := \rho_\gamma + n - \nu, \quad n'' = n - \nu.$$

We also write  $B_\gamma' = B_{\gamma'} := \rho_\gamma'' * \frac{1}{|\cdot|} - R[\gamma']$ .

By functional calculus, we expand  $\chi_{(-\infty, 0)}(D_Q) - \mathcal{P}_-^0$  in power of  $\alpha$ : this is the Cauchy expansion [8]

$$\begin{cases} \gamma + N &= N - \frac{1}{2\pi} \int_{-\infty}^{+\infty} d\eta \left( \frac{1}{D_{\gamma'} + i\eta} - \frac{1}{\mathcal{D}^0 + i\eta} \right) = \sum_{j=1}^{+\infty} \alpha^j Q_j(\gamma', \rho_\gamma''), \\ Q_j(\gamma', \rho_\gamma'') &:= -\frac{1}{2\pi} \int_{-\infty}^{+\infty} d\eta \frac{1}{\mathcal{D}^0 + i\eta} \left( B_{\gamma'} \frac{1}{\mathcal{D}^0 + i\eta} \right)^j. \end{cases} \quad (20)$$

We define  $Q_{k,l}$  as the part of  $Q_{k+l}(Q, \rho)$  which is a homogeneous polynomial of degree  $k$  in  $R_Q$  and of degree  $l$  in  $\rho$ ;  $\rho_{k,l}(Q, \rho)$  denotes its density. For  $\ell \geq 1$  and  $(Q, \rho) \in \mathfrak{S}_2(H^{1/2}) \in \mathcal{C}$ ,  $\tilde{Q}_\ell[Q, \rho]$  is the operator:

$$\tilde{Q}_\ell[Q, \rho] := \sum_{j=\ell}^{+\infty} \alpha^{j-\ell} Q_j[Q, \rho].$$

As shown in [8, 7] we have

$$\rho_{0,1}[\rho] = -\mathcal{F}^{-1}(B_\Lambda) * \rho \quad (21)$$

where  $\mathcal{F}^{-1}(B_\Lambda)$  is a radial  $L^1$  function.

In the following Lemma, we refer to the Banach spaces  $\mathbf{Q}_w$  and  $\mathfrak{C}_w$ : they are defined below (26). This Lemma is proved in Section 4.

**Lemma 1.**  $F_{1,0} : Q \mapsto Q_{1,0}(Q)$  is a bounded linear map of  $\mathfrak{S}_p$  for  $p = 1$  and  $p = 2$  with respective norms  $\mathcal{O}(\log(\Lambda))$  and  $\mathcal{O}(\sqrt{\log(\Lambda)})$ . By interpolation  $F_{1,0}$  is in  $L(\mathfrak{S}_p)$  for  $1 < p = 1 + \varepsilon < 2$  with norm  $\mathcal{O}((\log(\Lambda))^{1-\frac{\varepsilon}{2}})$ .

Moreover it is also a bounded operator in  $L(\mathbf{Q}_w)$  with norm  $\mathcal{O}(1)$ , and the function

$$\rho F_{1,0} : Q \in \mathbf{Q}_w \mapsto \rho(F_{1,0}[Q]) \in \mathfrak{C}_w$$

is bounded with norm  $\mathcal{O}(\sqrt{\log(\Lambda)})$ . Provided that  $\alpha \log(\Lambda)$  is sufficiently small, the operator  $(\text{Id} - \alpha F_{1,0})$  is invertible with inverse  $\mathbf{T}$  in all those Banach spaces with norm  $\mathcal{O}(1)$ . The function  $\mathbf{t} : Q \in \mathbf{Q}_w \mapsto \rho(\mathbf{T}[Q] - Q) \in \mathfrak{C}_w$  is bounded and

$$\|\mathbf{t}_Q\|_{\mathfrak{C}_w} \lesssim \sqrt{L\alpha} \|Q\|_{\mathbf{Q}_w}.$$

We write

$$\mathfrak{T} := \mathbf{T} - \text{Id}, \quad \tau_Q := \rho_{\mathbf{T}(Q)}, \tau_{j,k} := \rho_{\mathbf{T}(Q_{j,k})} \text{ and } \mathfrak{t}_Q := \rho_{\mathfrak{T}(Q)}. \quad (22)$$

If  $Q \in \mathbf{Q}_{w=1} \cap \mathfrak{S}_1^{P^0}$  then  $\tau_Q \in \mathcal{C}$  and if  $(Q, \rho_Q) \in \mathbf{Q}_w \times \mathfrak{C}_w$  then  $\tau_Q \in \mathfrak{C}_w$ .  
The self-consistent equation (18) is rewritten as follows:

$$(\text{Id} - \alpha F_{1,0})(\gamma') = N + \alpha Q_{0,1}(\rho''_\gamma) + \sum_{j=2}^{+\infty} Q_j(\gamma', \rho''_\gamma).$$

Taking the inverse  $\mathbf{T}$ , we get:

$$\gamma' = \mathbf{T} \left\{ N + \alpha Q_{0,1}(\rho''_\gamma) + \sum_{j=2}^{+\infty} Q_j(\gamma', \rho''_\gamma) \right\}. \quad (23)$$

The important proposition holds:

**Proposition 1.** *For  $\rho \in \mathcal{C}$  we have  $\alpha \tau_{0,1}(\rho) = -\check{f}_\Lambda * \rho$  where  $\check{f}_\Lambda$  is a radial  $L^1$  function whose  $L^1$ -norm is  $\mathcal{O}(\alpha \log(\Lambda))$ .*

Its technical proof is in Appendix C.

There holds a Theorem *à la Furry* [5, 8]:

**Theorem 1.** *There exists  $K > 0$  such that for any  $\rho_0, \rho_1$  (say in  $\mathcal{C}$ ) and  $\alpha \sqrt{\log(\Lambda)} \leq K$  there holds:*

$$\rho \{ \mathbf{T}(Q_{0,2}(\rho_0)) \} = \rho \{ \mathbf{T}(Q_{1,1}(\mathbf{T}Q_{0,1}(\rho_1), \rho_0)) \} = 0. \quad (24)$$

*Remark 6.*  $\mathbf{T}(Q_{0,2}(\rho_0))$  and  $\mathbf{T}(Q_{1,1}(\mathbf{T}(Q_{0,1}(\rho_1)), \rho_0)$  may not vanish but their density do due to the fact that the trace  $\text{Tr}_{\mathbb{C}^4}$  is taken. The smallness of  $\alpha \sqrt{\log(\Lambda)}$  is to ensure the  $\mathbf{T}$  operator is well defined on  $\mathbf{Q}_1$ .

## Main Theorems

**Theorem 2** (Computation of  $\int_{\mathbb{R}} \rho_\gamma(x) dx$ ). *Let  $M$  be in  $\mathbb{N}$  and  $\gamma' = \gamma + N$  be a minimizer of  $E_{BDF}^\nu(M)$  and assume  $M, \|\nu\|_{\mathcal{C}}^2 \lesssim \log(\Lambda)$  and (28), the decomposition of  $\gamma'$  is that of (18). Then  $\rho_\gamma \in L^1$  and*

$$\int \rho_\gamma(x) dx = -\frac{\alpha f_\Lambda(0)}{1 + \alpha f_\Lambda(0)} (M - Z) \quad (25)$$

**Theorem 3** (Existence of minimizers). *There exists  $K_0 > 0$  satisfying the following result: for any non-negative function  $\nu \in \mathcal{C} \cap L^1$  with  $Z = \int \nu$  and  $0 < L \leq 1/(MK_0)$ , there exists  $\alpha_1 = \alpha_1(\nu, L) > 0$  such that if  $\alpha \leq \alpha_1$  then for any integer  $0 \leq M < Z + 1$  the problem  $E_{BDF}^\nu(M)$  admits a minimizer.*

*Let  $\gamma' = \chi_{(0,\mu]}(D_{\gamma'})$  be a minimizer, decomposed as in (18) and let  $U_\alpha$  be defined as follows:*

$$U_\alpha : \begin{array}{ccc} L^2(\mathbb{R}^3, \mathbb{C}^4) & \rightarrow & L^2(\mathbb{R}^3, \mathbb{C}^4) \\ \phi(x) & \mapsto & \alpha^{-3/2} \phi(\frac{x}{\alpha}) \end{array} .$$

*We write  $\frac{f_\Lambda(0)}{1+f_\Lambda(0)} = a$ , then as  $\alpha$  tends to 0,  $U_\alpha^* \chi_{(0,\mu]}(D_{\gamma'}) U_\alpha$  tends to a minimizer of*

$$\mathcal{E}_{nr}^Z(\Gamma) := \frac{1}{2} \text{Tr}(-\Delta \Gamma) - Z(1-a) \text{Tr}\left(\frac{1}{|\cdot|} \Gamma\right) + \frac{1}{2} (D(\rho_\Gamma, \rho_\Gamma) - \text{Ex}[\Gamma]) - \frac{a}{2} D(\rho_\Gamma, \rho_\Gamma),$$

for  $0 \leq \Gamma \leq 1$ ,  $\text{Tr}(\Gamma) = M$  and  $\frac{1+\beta}{2} \Gamma = 0$ .

*Remark 7.* Thanks to Section C and [7] we have

$$\frac{f_\Lambda(0)}{1+f_\Lambda(0)} = \frac{\frac{2}{3\pi} \alpha \log(\Lambda)}{1 + \frac{2}{3\pi} \alpha \log(\Lambda)} + \mathcal{O}(\alpha + (\alpha \log(\Lambda))^2).$$



**Banach spaces** We use several Banach spaces. For  $p \in [1, +\infty]$ ,  $s \geq 0$ ,  $\|\cdot\|_{L^p}$  (resp.  $\|\cdot\|_{H^s}$ ) is the norm of the usual  $L^p$  (resp. Sobolev) space. We write  $\|\cdot\|_{\mathfrak{S}_p}$  for the norm of Schatten class operators  $\mathfrak{S}_p$  [22]. The norm of bounded linear operator in  $\mathfrak{H}$  is written  $\|\cdot\|_{\mathfrak{B}}$ . We recall  $\|\cdot\|_{\text{Ex}}$  and  $\|\cdot\|_{\mathcal{C}}$  have already been defined in Sections 1 and 2 and  $\|\cdot\|_{\mathbf{Q}_w}, \|\cdot\|_{\mathfrak{C}_w}$  are defined in Remark 8.

*Notation 8.* From now on, for any  $w : \mathbb{R}^3 \rightarrow [1, +\infty)$  satisfying the condition

$$\exists K_{(w)} > 0 \mid \forall p, q, p_1 \in \mathbb{R}^3, w(p - q) \leq K_{(w)}(w(p - p_1) + w(p_1 - q)),$$

we define two Hilbert spaces:

$$\begin{aligned} \mathbf{Q}_w &:= \{Q \in \mathfrak{S}_2, \iint (\sqrt{1 + |p|^2} + \sqrt{1 + |q|^2}) w(p - q) |\widehat{Q}(p, q)|^2 dp dq < +\infty\}, \\ \mathfrak{C}_w &:= \{\rho \in \mathcal{S}'(\mathbb{R}^3), \int \frac{w(k)}{|k|^2} |\widehat{\rho}(k)|^2 dk < +\infty\}. \end{aligned} \quad (26)$$

The letter  $w$  always refers to a function of this kind. The case  $w \equiv 1$  gives the space  $\mathbf{Q}_1$  of operators  $Q$  with  $\text{Tr}(|\mathcal{D}^0| |Q|^2 + Q^* |\mathcal{D}^0| Q) < +\infty$  and  $\mathfrak{C}_1 = \mathcal{C}$ . Typically, we consider  $w(p - q) := E(p - q)^a$  for  $a > 1$ .

By the fixed point method we may estimate together

- $\|F_Q(Q, \rho)\|_{\mathfrak{T}}$  and  $\|F_\rho(Q, \rho)\|_{\mathcal{C}}$ ,
- In general  $\|F_Q(Q, \rho)\|_{\mathbf{Q}_w}$  and  $\|F_\rho(Q, \rho)\|_{\mathfrak{C}_w}$ . We define  $\mathcal{X}_g := \mathbf{Q}_w \times \mathfrak{C}_w$

## Notations

*Notation 9* (On  $D_0$  and  $\mathcal{D}^0$ ). The operator  $\text{sign}(\mathcal{D}^0)$  is a Fourier multiplier that we write  $s_p = \frac{\widehat{\mathcal{D}^0(p)}}{\sqrt{g_0(p)^2 + g_1(p)^2}}$ . We also write

$$E(p) := \sqrt{1 + |p|^2} \text{ and } \widetilde{E}(p) := \sqrt{g_0(p)^2 + g_1(p)^2}. \quad (27)$$

*Remark 10* (Regime). We will work in the regime

$$\alpha \leq \alpha_0 \ll 1 \text{ and } L := \alpha \log(\Lambda) \leq L_0 \ll 1. \quad (28)$$

We consider systems with  $M$  electrons and an external charge density  $\nu \geq 0$  with  $\|\nu\|_{\mathcal{C}}, Z := \|\nu\|_{L^1} < +\infty$ . We will often consider  $M = \mathcal{O}(Z)$  and  $\|\nu\|_{\mathcal{C}}^2 + M = \mathcal{O}(\log(\Lambda))$ .

Throughout this paper the letter  $K$  denotes a constant independent of the parameters  $\alpha, \Lambda, M, \nu$ .  $K(M, \nu)$  is a constant depending on  $M, \nu$  and so on. The inequality  $a \lesssim b$  means that  $a \leq Kb$  for  $a, b > 0$ . When  $a > 1$  is some integer, then as in [8] we write

$$K_a := \frac{1}{2\pi} \int_{-\infty}^{+\infty} \frac{d\eta}{E(\eta)^a} = \mathcal{O}_{a \rightarrow +\infty}(a^{-1/2}). \quad (29)$$

*Notation 11* (On  $\widetilde{Q}_{k,\ell}$ ). For  $(\varepsilon_1, \dots, \varepsilon_{J+1}) \in \{+, -\}^{J+1}$  we define  $Q_J^{\varepsilon_1 \dots \varepsilon_{J+1}}$  with the same formula as in (20) except that we replace the  $J + 1$  operators  $(\mathcal{D}^0 + i\eta)^{-1}$ 's by  $P_{\varepsilon_j}^0 / (\mathcal{D}^0 + i\eta)$ . We define  $Q_{k,\ell}^{\varepsilon_1 \dots \varepsilon_{J+1}}$  analogously.

We write  $Q_{k,\ell}^{\varepsilon_1 a_1 \varepsilon_2 \dots \varepsilon_J a_J \varepsilon_{J+1}}$  with  $a_j \in \{v, R\}$  for the operator

$$-\frac{1}{2\pi} \int_{-\infty}^{+\infty} d\eta \frac{P_{\varepsilon_1}^0}{\mathcal{D}^0 + i\eta} A_1 \frac{P_{\varepsilon_2}^0}{\mathcal{D}^0 + i\eta} \dots A_J \frac{P_{\varepsilon_{J+1}}^0}{\mathcal{D}^0 + i\eta},$$

where  $A_j = v = \rho_\gamma'' * \frac{1}{|\cdot|}$  if  $a_j = v$  or  $A_j = -R(\gamma')$  if  $a_j = R$ .

*Notation 12* (On  $f_\Lambda$ ). We introduce the function  $F_\Lambda := \frac{f_\Lambda}{1 + f_\Lambda}$ , studied in Appendix C. We prove in particular that  $F_\Lambda \in L^1$  and that  $\|F_\Lambda\|_{L^1} \lesssim L$ .

### 3 Description of minimizers

#### 3.1 Minimizers and fixed point schemes

Let  $\gamma' = \gamma + N$  be a minimizer for  $E_{\text{BDF}}^\nu(M)$ . From Eq. (20) and (21), we define a fixed-point scheme:

$$F^{(1)} = F_Q^{(1)} \times F_\rho^{(1)} : \mathcal{X}_1 \rightarrow \mathcal{X}_1,$$

$$F_Q^{(1)}(Q', \rho'') = N + \sum_{\ell=1}^{\infty} \alpha^\ell Q_\ell(Q', \rho''), \quad (30a)$$

$$\mathcal{F}(F_\rho^{(1)}(Q', \rho''); k) = \frac{1}{1 + \alpha B_\Lambda(k)} \widehat{n}''(k) + \frac{1}{1 + \alpha B_\Lambda(k)} \left( \alpha \widehat{\rho}_{1,0}(Q'; k) + \sum_{\ell=2}^{\infty} \alpha^\ell \widehat{\rho}_\ell(Q', \rho''); k \right) \quad (30b)$$

To prove  $F^{(1)}$  is well-defined we use the following Lemma proved in Section 5.

**Lemma 2.** *Let  $w$  be some function satisfying (8), with constant  $K_{(w)} > 0$ . There exists  $C_0 > 0$  such that for any  $J \geq 2$ , the linear operator:*

$$(Q, \rho) \in \mathbf{Q}_w \times \mathbf{C}_w \mapsto (Q_J(Q, \rho), \rho_J(Q, \rho)) \in \mathbf{Q}_w \times \mathbf{C}_w$$

is bounded with norm lesser than  $2K_{(w)}^J C_0^J J^{1/2}$ .

We apply the Banach-Picard Theorem.

**Lemma 3.** *Let  $\gamma' = \gamma + N$  be a minimizer for  $E_{\text{BDF}}^\nu(M)$ . In the regime of Remark 10 the following holds:*

1.  $F^{(1)} : B_{\mathcal{X}_1}(0, R_0) \rightarrow B_{\mathcal{X}_1}(0, R_0)$  is well-defined for some  $R_0 > 0$  and this restriction is a Lipschitz function with constant lesser than 1.
2.  $(\gamma', \rho_\gamma'')$  is in the previous ball and so is the unique fixed point of  $F^{(1)}$ , moreover:

$$\|F^{(1)}(\gamma', \rho_\gamma'') - (N, n'')\|_{\mathcal{X}_1} = o(1).$$

3. As a consequence  $N = \chi_{(0, \mu]}(D_Q)$  has rank  $M_0 = M$ .

**Proof of part 3.** If we assume the first two points, the last one is clear. Indeed on the one hand we have:  $|\text{Tr}_0(\gamma)| \lesssim \|\gamma\|_{\mathfrak{S}_2}^2 = o(1)$ , on the other hand, as  $\gamma$  is a difference of an orthogonal projector and  $\mathcal{P}_-^0$ , it must be an integer [8, Lemma 2]. Thus  $\text{Tr}_0(\gamma) = 0$  and

$$\text{Tr}(N) = \text{Tr}_0(N) = \text{Tr}_0(\gamma') - \text{Tr}_0(\gamma) = M.$$

□

To prove that  $\rho_\gamma$  is integrable we need another fixed point scheme.

We see  $\rho_\gamma''$  as the fixed point of a function  $F^{(2)}$  defined in (a ball of)  $\mathcal{C}$  and also in (a ball of)  $\mathcal{C} \cap L^1$ . We write:

$$\begin{cases} h_2 &= \alpha^2 \tau_{1,1} \left\{ \mathbf{T}[N] + \alpha^2 \left\{ \alpha \mathbf{T} \widetilde{Q}_3(\gamma', \rho_\gamma'') + \mathbf{T} Q_{2,0}(\gamma', \rho_\gamma'') \right\}, \rho_\gamma'' \right\} + \alpha^2 \tau_{2,0}(\gamma') \\ F_2^{(2)}(\rho'') &= \alpha^2 (\tau_{1,1} \left\{ \alpha^2 [\mathbf{T} Q_{1,1}(\gamma', \rho'') + \mathbf{T} Q_{0,2}(\rho'')], \rho'' \right\}) \\ h_3 &= \alpha^4 \tau(\widetilde{Q}_4(\gamma', \rho_\gamma'')) + \alpha^3 \left\{ \tau_{3,0}(\rho_\gamma'') + \tau_{2,1}(\gamma', \rho_\gamma'') \right\} \\ F_3^{(2)}(\rho'') &= \alpha^3 \tau_{0,3}(\rho'') + \alpha^3 \tau_{1,2}(\gamma', \rho'') \end{cases} \quad (31)$$

$$\mathcal{F}\{F^{(2)}(\rho'')\} = \frac{1}{1 + f_\Lambda(\cdot)} \widehat{n}'' + \frac{1}{1 + f_\Lambda(\cdot)} \left\{ \widehat{h}_2 + \mathcal{F}\{F_2^{(2)}\} + \widehat{h}_3 + \mathcal{F}\{F_3^{(2)}\} \right\}(\rho'') \quad (32)$$

*Remark 13.* The definition of  $F^{(2)}$  may appear complicated. It is built on the following self-consistent equation:

$$\rho_\gamma' = \tau \left\{ N + \alpha Q_{0,1}(\rho_\gamma'') + \alpha^2 (\widetilde{Q}_2(\gamma', \rho_\gamma'') - Q_{1,1}(\gamma', \rho_\gamma'')) \right\} + \alpha^2 \tau [Q_{1,1}(F_Q^{(1)}(\gamma', \rho_\gamma''), \rho_\gamma'')].$$

**Lemma 4.** Let  $\gamma' = \gamma + N$  be a minimizer for  $E_{BDF}^\nu(M)$  and  $F^{(2)}$  the function (31). In the regime of Remark 10, there exists  $R_0 > 0$  such that  $F^{(2)}$  is well-defined in  $B_C(0, R_0)$  and in  $B_{C \cap L^1}(0, R_0)$ .

Furthermore these balls are  $F^{(2)}$ -invariant and  $F^{(2)}$  is a contraction on them;  $\rho_\gamma'$  is the only fixed point in both Banach spaces. In particular  $\rho_\gamma \in L^1$ .

*Remark 14.* The linear response of the vacuum to the presence of electrons  $N$  and the external potential  $\nu$  is:

$$\begin{cases} \gamma &= \alpha \mathbf{T}[Q_{0,1}((\delta_0 - \check{F}_\Lambda) * (n - \nu + \mathbf{t}_N))] + \mathfrak{T}_N + \dots \\ \rho_\gamma &= -\check{F}_\Lambda * (n - \nu) + (\delta_0 - \check{F}_\Lambda) * \mathbf{t}_N + \dots \end{cases}$$

### 3.2 A priori estimates

**Lemma 5** (Estimates on the energy). Let  $M \in \mathbb{N}$  and  $Q$  a test function for  $E_{BDF}^\nu(M)$ . We assume:  $\mathcal{E}_{BDF}^\nu(Q) \leq E_{BDF}^\nu(M) + \varepsilon$  where  $0 < \varepsilon = o(\alpha \|\nu\|_C^2)$ .

Then we have  $\|Q\|_{\mathfrak{S}_2}^2 \lesssim M + \alpha \|\nu\|_C^2$  and

$$\begin{aligned} \mathrm{Tr}(|\nabla|Q^2) &\lesssim \alpha \|\nu\|_C^2 + \alpha^{1/2} M + \sqrt{\alpha M} \|\nu\|_C, \\ \alpha \|\rho_Q - \nu\|_C^2 &\lesssim \alpha \|\nu\|_C^2 + \alpha^{3/2} M + \sqrt{\alpha M} \alpha \|\nu\|_C. \end{aligned}$$

As a corollary we get the following.

**Lemma 6** (Estimates on the mean-field operator). In the regime of Remark 10 and for  $Q$  as in Lemma 5 we have in the sense of self-adjoint operator:

$$(1 - o(1))|\mathcal{D}^0| \leq |\mathcal{D}^0 + \alpha B_Q| \leq (1 + o(1))|\mathcal{D}^0|. \quad (33)$$

Both  $o(1)$  are  $\mathcal{O}(\alpha \|\nu\|_C + \alpha^{5/4} M^{1/2} + (\alpha M)^{1/4} \alpha \|\nu\|_C^{1/2})$ .

**Lemma 7** (A priori estimates of a minimizer). Let  $\gamma' = \gamma + N$  be a minimizer for  $E_{BDF}^\nu(M)$ , decomposed as in (18). Then we have in the regime (28)

$$\begin{aligned} \mathrm{Tr}(|\mathcal{D}^0|N) &\lesssim \log(\Lambda), & \left| \|\gamma\|_{\mathbf{T}} \right| &\lesssim L, \\ \|n''\|_C &\lesssim \sqrt{\log(\Lambda)}, & \|\rho_\gamma\|_C &\lesssim L \sqrt{\log(\Lambda)}. \end{aligned}$$

**Proof of Lemma 5:** It is known that  $E_{BDF}^\nu(M) \leq M$  [10]. There holds:

$$\begin{aligned} M + \varepsilon + \frac{\alpha}{2} \|\nu\|_C^2 &\geq \mathcal{E}_{BDF}^\nu(Q) + \frac{\alpha}{2} \|\nu\|_C^2 \geq (1 - \alpha \frac{\pi}{4}) \mathrm{Tr}_0(\mathcal{D}^0 Q) + \frac{\alpha}{2} \|\rho_Q - \nu\|_C^2 \\ &\geq (1 - \alpha \frac{\pi}{4}) \|Q\|_{\mathfrak{S}_2}^2 + \frac{\alpha}{2} \|\rho_Q - \nu\|_C^2. \end{aligned}$$

Furthermore:

$$\begin{aligned} \mathrm{Tr}_0(\mathcal{D}^0 Q) - M &= \mathrm{Tr}(|\mathcal{D}^0|^{1/2} (Q^{++} - Q^{--}) |\mathcal{D}^0|^{1/2}) - \mathrm{Tr}_0(Q) \\ &\geq \mathrm{Tr}(|\mathcal{D}^0|^{1/2} Q^2 |\mathcal{D}^0|^{1/2}) - \mathrm{Tr}(Q^2) \\ &\geq \frac{1}{(2\pi)^3} \iint (\tilde{E}(p) - 1) |\hat{Q}(p, q)|^2 dp dq, \end{aligned} \quad (34)$$

and  $\tilde{E}(p) - 1 \geq \frac{1}{2} \frac{p^2}{E(p)}$ . Then thanks to Kato's inequality (61):

$\mathrm{Tr}(QR_Q) \leq \frac{\pi}{2} \mathrm{Tr}(|\nabla|Q^2)$  which leads to:

$$\frac{1}{2} \mathrm{Tr} \left( \frac{-\Delta}{|D_0|} Q^2 \right) + \frac{\alpha}{2} \|\rho_Q - \nu\|_C^2 \leq \varepsilon + \alpha \left( \frac{\|\nu\|_C^2}{2} + \frac{\pi}{4} \mathrm{Tr}(|\nabla|Q^2) \right).$$

Splitting at level  $r_0 = \frac{\alpha\pi}{\sqrt{1-(\alpha\pi)^2}}$  (to get  $\alpha \frac{|p|\pi}{4} \leq \frac{1}{4} \frac{|p|^2}{E(p)}$  for  $|p| \geq r_0$ ) we obtain:

$$\mathrm{Tr} \left( \frac{-\Delta}{|D_0|} Q^2 \right) \lesssim \alpha (\|\nu\|_C^2 + M), \quad (35)$$

thus by the Cauchy-Schwartz inequality:  $\mathrm{Tr}(|\nabla|Q^2) \lesssim \alpha \|\nu\|_C^2 + \sqrt{\alpha} M + \sqrt{\alpha M} \|\nu\|_C$ .  $\square$

**Proof of Lemma 6:**

For all  $f \in \mathfrak{H}_\Lambda$  we have:

$$\langle |\mathcal{D}^0|^2 f, f \rangle (1 - \alpha \|\mathcal{D}^0|^{-1} B\|_{\mathcal{B}})^2 \leq \langle \mathcal{D}^0 + \alpha B|^2 f, f \rangle \leq \langle |\mathcal{D}^0|^2 f, f \rangle (1 + \alpha \|\mathcal{D}^0|^{-1} B\|_{\mathcal{B}})^2. \quad (36)$$

However thanks to Ineq.(58) and the second point of Lemma 8:

$$\|R_Q |\nabla|^{-1/2}\|_{\mathcal{B}} \lesssim \sqrt{\text{Tr}(QR_Q)} \text{ and } \|(\rho_Q - \nu) * \frac{1}{|\cdot|} |\nabla|^{-1/2}\|_{\mathcal{B}} \lesssim \|(\rho_Q - \nu) * \frac{1}{|\cdot|}\|_{L^6} \lesssim \|\rho_Q - \nu\|_c.$$

As the square root is monotone, there holds

$$(1 - \alpha \|\mathcal{D}^0|^{-1} B_Q\|_{\mathcal{B}}) |\mathcal{D}^0| \leq |\mathcal{D}^0 + \alpha B_Q| \leq (1 + \alpha \|\mathcal{D}^0|^{-1} B_Q\|_{\mathcal{B}}) |\mathcal{D}^0|, \quad (37)$$

and in the regime of Remark 10, this gives  $(1 - o(1)) |\mathcal{D}^0| \leq |\mathcal{D}^0 + \alpha B_Q| \leq (1 + o(1)) |\mathcal{D}^0|$ . This  $o(1)$  is of order  $\mathcal{O}(\alpha(\|\rho_Q - \nu\|_c + \|\nabla|^{1/2} Q\|_{\mathfrak{S}_2}))$ , that is of order

$$\mathcal{O}(\alpha \|\nu\|_c + \alpha^{5/4} M^{1/2} + (\alpha M)^{1/4} \alpha \|\nu\|_c^{1/2}). \quad \square$$

**Proof of Lemma 7:** For  $E_{\text{BDF}}^{\nu}(M)$  with  $M, \|\nu\|_c^2 \lesssim \log(\Lambda)$ , we have thanks to Lemma 5:

$$\alpha(\|\rho_\gamma''\|_c + \sqrt{\text{Tr}(|\nabla|\gamma')}) \lesssim \sqrt{\alpha}(\alpha^{1/2} \|\nu\|_c + \alpha^{3/4} M^{1/2} + (\alpha M)^{1/4} \alpha^{1/2} \|\nu\|_c^{1/2}) =: \alpha^{1/2} \ell.$$

We have  $\ell = \mathcal{O}(\sqrt{L})$ . Using Eq. (23) and assuming Lemma 2 and Proposition 1 above we get that:

$$\|\rho_\gamma\|_c \leq \|\check{F}_\Lambda * n''\|_c + \|(\delta_0 - \check{F}_\Lambda) * (\mathfrak{t}_N + \sum_{j \geq 2} \alpha^j \tau_j)\|_c \lesssim L \|n''\|_c + \sqrt{L\alpha} \|N\|_{\mathfrak{T}} + \mathcal{O}(L\alpha).$$

As  $\|n''\|_c \leq \|\rho_\gamma''\|_c + \|\rho_\gamma\|_c$  we get

$$\|n''\|_c \lesssim \|\nu\|_c + (\alpha M)^{1/4} (M^{1/4} + \sqrt{\|\nu\|_c}) + \sqrt{L\alpha M} + \mathcal{O}(\alpha \ell^2) \lesssim \sqrt{\log(\Lambda)}.$$

Thanks to the equations  $\mathcal{D}^0 \psi_j = \mu_j \psi_j - B \psi_j$ , there holds:

$$\text{Tr}(|\mathcal{D}^0| N) \lesssim M(1 + \mathcal{O}(\sqrt{\alpha} \ell)) \lesssim \log(\Lambda).$$

Finally we have

$$\begin{aligned} \|\gamma\|_{\mathfrak{T}} &\lesssim \sqrt{L\alpha} \|n''\|_c + \alpha \sqrt{\text{Tr}(|\nabla|Q^2)} + \mathcal{O}(L\alpha) \lesssim L + \mathcal{O}(L\alpha) \lesssim L \\ \|\rho_\gamma\|_c &\lesssim L \|n''\|_c + \sqrt{L\alpha M} + \mathcal{O}(L\alpha) \lesssim L \sqrt{\log(\Lambda)}. \end{aligned} \quad (38)$$

□

## 4 The operator $F_{1,0}^1$

*Remark 15.* • If  $Q$  is a nonnegative operator then so is  $R_Q$  when it is well defined. Moreover if  $Q$  is self-adjoint then so is  $R_Q$ .

- The  $R$ . operator commutes with Fourier multiplier of the form  $g(p - q)$ , indeed we have

$$\widehat{R_Q}(p, q) = \frac{1}{2\pi^2} \int \frac{\widehat{Q}(p - l, q - l)}{|l|^2}.$$

In particular there holds:

$$[\partial_j, R_Q] = R([\partial_j, Q]). \quad (39)$$

**Lemma 8.** *Let  $Q$  be in  $\mathcal{S}(\mathbb{R}^3 \times \mathbb{R}^3)$  (Schwartz class).*

1. *We have:*

$$\|\nabla|^{-1/2} R_Q\|_{\mathfrak{S}_2} \lesssim \sqrt{\text{Tr}(R_Q^* Q)}.$$

*In particular for any  $w \geq 1$  there holds:*

$$\iint \frac{w(p - q)}{|p|} |\widehat{R_Q}(p, q)|^2 dp dq \lesssim \iint |p + q| w(p - q) |\widehat{Q}(p, q)|^2 dp dq.$$

2. There exists  $K > 0$  such that for all  $0 < \epsilon \leq 1$

$$\begin{aligned} \||D_0|^{-\frac{1+\epsilon}{2}} R_Q |D_0|^{-\frac{1+\epsilon}{2}}\|_{\mathfrak{S}_1} &\leq \frac{K}{\epsilon} \|Q\|_{\mathfrak{S}_1}, \\ \||D_0|^{-(1+\epsilon)} R_Q\|_{\mathfrak{S}_2} &\leq \frac{K}{\sqrt{\epsilon}} \|Q\|_{\mathfrak{S}_2}. \end{aligned}$$

For  $Q \in \mathfrak{S}_2(\mathfrak{H}_\Lambda)$ , we can replace  $|D_0|^{-(1+\epsilon)/2}$  by  $|D^0|^{-1/2}$ , provided that  $\epsilon^{-1}$  is replaced by  $\log(\Lambda)$ .

By density, these inequalities hold for  $Q$  in the Banach spaces corresponding to the norms in the r.h.s.

We prove this Lemma in Appendix A.

## 4.1 Proof of Lemma 1

**In the Schatten norms** We recall  $F_{1,0}$  is defined as

$$F_{1,0} : Q \mapsto Q_{1,0}(Q) := -\frac{1}{2\pi} \int_{-\infty}^{+\infty} d\eta \frac{1}{\mathcal{D}^0 + i\eta} R_Q \frac{1}{\mathcal{D}^0 + i\eta}. \quad (40)$$

The integral kernel of its Fourier transform is [8]:

$$\widehat{Q}_{1,0}(p, q) = \frac{1}{2} \frac{1}{\widetilde{E}(p) + \widetilde{E}(q)} \left( \widehat{R}(p, q) - \mathbf{s}_p \widehat{R}(p, q) \mathbf{s}_q \right). \quad (41)$$

It corresponds to a difference of two operators which are in  $\mathfrak{S}_p$  if  $Q$  is in  $\mathfrak{S}_p$  for both cases  $p = 1$  and  $p = 2$  (see below). By interpolation, for  $p \in [1, 2]$ , if  $Q \in \mathfrak{S}_p$  then so is  $F_{1,0}(Q)$ . Let us show the  $\mathfrak{S}_1$ -norm is  $\mathcal{O}(\log(\Lambda))$  while the  $\mathfrak{S}_2$ -norm is  $\mathcal{O}(\sqrt{\log(\Lambda)})$ . Indeed

$$\frac{1}{f(p) + f(q)} = \int_{s=0}^{+\infty} e^{-sf(p) - sf(q)} ds,$$

therefore if  $Q$  is nonnegative, then so is

$$\int_{s=0}^{+\infty} \frac{\mathcal{D}^0}{|\mathcal{D}^0|} \mathcal{F}^{-1}(e^{-s\widetilde{E}(\cdot)}) R_Q \mathcal{F}^{-1}(e^{-s\widetilde{E}(\cdot)}) \frac{\mathcal{D}^0}{|\mathcal{D}^0|} ds.$$

Writing  $Q = \frac{Q+Q^*}{2} + \frac{Q-Q^*}{2}$  and splitting each self-adjoint operator into nonnegative and nonpositive part, we may assume that  $Q \geq 0$ . Then from Eq. (41), we get:

$$\|F_{1,0}(Q)\|_{\mathfrak{S}_1} \leq K \log(\Lambda) \|Q\|_{\mathfrak{S}_1}.$$

As  $(\widetilde{E}(p) + \widetilde{E}(q))^{-1} \leq \widetilde{E}(p)^{-1/2} \widetilde{E}(q)^{-1/2}$ , it follows that

$$\begin{aligned} \||\mathcal{D}^0|^{-\frac{1}{2}} R(\mathcal{F}^{-1}(|\widehat{Q}(p, q)|)) |\mathcal{D}^0|^{-\frac{1}{2}}\|_{\mathfrak{S}_2} &\leq K \sqrt{\log(\Lambda)} \|\mathcal{F}^{-1}(|\widehat{Q}(p, q)|)\|_{\mathfrak{S}_2} \\ &= K \sqrt{\log(\Lambda)} \|\widehat{Q}\|_{\mathfrak{S}_2} = K \sqrt{\log(\Lambda)} \|Q\|_{\mathfrak{S}_2}. \end{aligned}$$

By interpolation ( $1 < p = 1 - \epsilon + 2\epsilon < 2$ ), there exists  $K_{(1,0)}^{\mathfrak{S}} > 0$

$$\|Q_{1,0}(Q)\|_{\mathfrak{S}_p} \leq K_{(1,0)}^{\mathfrak{S}} (\log(\Lambda))^{1-\frac{\epsilon}{2}} \|Q\|_{\mathfrak{S}_p}, \quad (42)$$

*Remark 16.* The operators  $Q_{1,0}(Q_0)$  (and  $Q_{0,1}(\rho_0)$ ) can be rewritten as

$$\mathcal{J}_t(x - y) := \mathcal{F}^{-1}(\exp(-t\widetilde{E}(p)))(x - y) \quad (43a)$$

$$\begin{cases} Q_{1,0}(Q_0) &= \frac{1}{2} \int_{t=0}^{+\infty} (\mathcal{J}_t R_{Q_0} \mathcal{J}_t - \mathcal{J}_t \frac{\mathcal{D}^0}{|\mathcal{D}^0|} R_{Q_0} \frac{\mathcal{D}^0}{|\mathcal{D}^0|} \mathcal{J}_t) dt \\ Q_{0,1}(\rho_0) &= -\frac{1}{2} \int_{t=0}^{+\infty} (\mathcal{J}_t v_{\rho_0} \mathcal{J}_t - \mathcal{J}_t \frac{\mathcal{D}^0}{|\mathcal{D}^0|} v_{\rho_0} \frac{\mathcal{D}^0}{|\mathcal{D}^0|} \mathcal{J}_t) dt \end{cases} \quad (43b)$$

$\rho[Q_{1,0}(\cdot)]$  We show here inequalities needed to estimate  $\mathbf{T}(Q_\ell(Q, \rho))$  and  $\tau_\ell(Q, \rho)$  in norms  $\|\cdot\|_{\mathbf{Q}_w}, \|\cdot\|_{\mathfrak{E}_w}$ . There exists a constant  $C_R$  (defined in [8]) such that for any function  $w \geq 0$

$$\iint (\tilde{E}(p) + \tilde{E}(q))w(p-q)|\widehat{Q}_{1,0}(Q, p, q)|^2 dpdq \leq C_R^2 \iint w(p-q)E(p+q)|\widehat{Q}(p, q)|^2 dpdq. \quad (44)$$

By Cauchy-Schwartz inequality (cf [8] and inequality (98)):

$$|\widehat{\rho}_{1,0}(Q, k)|^2 \lesssim |k|^2 \int_{B(0, \Lambda)} \frac{|\widehat{R}(u + \frac{k}{2}, u - \frac{k}{2})|^2}{1 + \tilde{E}(u, k/2)} du \int_{B(0, \Lambda)} \frac{du}{1 + \tilde{E}(u, k/2)} \frac{1}{1 + |u|^2 + |k|^2/4}, \quad (45)$$

where  $\tilde{E}(u, k/2) := \max(\tilde{E}(u + k/2), \tilde{E}(u - k/2))$ . Thus we have:

$$|\widehat{\rho}_{1,0}(Q, k)|^2 \leq C_{(1,0)} \int E(2u)|\widehat{Q}(u + \frac{k}{2}, u - \frac{k}{2})|^2 du, \quad (46)$$

where  $0 < C_{(1,0)} = C_{(1,0)}(\Lambda)$  satisfies  $C_{(1,0)} \lesssim \log(\Lambda)$ .

**Well-definedness of  $\mathbf{T}$  and  $\tau$**  Thanks to (42) we can prove Lemma 1: for  $\alpha \log(\Lambda)$  sufficiently small the function  $\mathbf{T}$  is a linear bounded operator in  $L(\mathfrak{S}_p)$  for  $1 \leq p = 1 + \varepsilon \leq 2$  with norm lesser than

$$C_{\mathbf{T}, \mathfrak{S}}^{(p)} := \sum_{\ell=0}^{+\infty} (\alpha K_{(1,0)}^{\mathfrak{S}} (\log(\Lambda))^{1-\frac{\varepsilon}{2}})^\ell = \frac{1}{1 - \alpha (\log(\Lambda))^{1-\frac{\varepsilon}{2}} K_{(1,0)}^{\mathfrak{S}}}$$

which is finite as soon as  $\alpha \log(\Lambda)$  is sufficiently small. We write  $C_{\mathbf{T}, \mathfrak{S}} := C_{\mathbf{T}, \mathfrak{S}}^{(1)}$ .

As  $\mathbf{T} = (\text{Id} - \alpha F_{1,0})^{-1} = \sum_{\ell=0}^{+\infty} \alpha^\ell F_{1,0}^{\circ(\ell)}$ , let us show that  $\alpha F_{1,0}$  is a bounded operator in  $L(\mathbf{Q}_w)$  with norm lesser than 1. Thanks to inequality (44),  $\alpha F_{1,0}$  is bounded with norm lesser than  $\alpha C_R$ . Thus  $\mathbf{T}$  is a bounded linear operator with norm lesser than

$$C_{\mathbf{T}, \mathbf{Q}_w} := \frac{1}{1 - \alpha C_R}. \quad (47)$$

Then thanks to Ineq. (44) and (46), for  $\ell \geq 1$  we have:

$$|\widehat{\rho}(F_{1,0}^{\circ(\ell)}(Q); k)|^2 \leq \alpha^{2\ell} C_{(1,0)}^\ell |k|^2 \int E(2u)|\widehat{Q}(u + \frac{k}{2}, u - \frac{k}{2})|^2 du$$

Therefore:

$$\int \frac{g(k)}{|k|^2} |\widehat{\rho}(F_{1,0}^{\circ(\ell)}; k)|^2 \leq \alpha^{2\ell} C_{(1,0)}^\ell \iint g(p-q)E(p+q)|\widehat{Q}(p, q)|^2 dpdq \quad (48)$$

and  $\mathfrak{t}$  is a bounded linear operator in  $L(\mathfrak{E}_w)$  with norm lesser than

$$C_{\mathfrak{t}, \mathfrak{E}} := \sum_{\ell=1}^{+\infty} (\alpha \sqrt{C_{(1,0)}})^\ell = \mathcal{O}(\alpha \sqrt{\log(\Lambda)}) \quad (49)$$

for  $\alpha \sqrt{\log(\Lambda)}$  sufficiently small.

*Notation 17.* Let us define for  $1 \leq p = 1 + \varepsilon \leq 2$ :

$$Y_{\alpha, \Lambda}(p) = Y(p) \lesssim C_{\mathbf{T}, \mathfrak{S}}^{(p)}, \quad (50)$$

which is an upper bound of the  $L(\mathfrak{S}_p)$ -norm of  $Q \mapsto |D_0|^{-7/12} R(\mathbf{T}[Q]) |D_0|^{-7/12}$ : cf Lemma 8 in Appendix A.1.

We have thus proved:

$$\begin{cases} \|\mathbf{T}(Q)\|_{\mathbf{Q}_w} & \leq C_{\mathbf{T}, \mathbf{Q}_w} \|Q\|_{\mathbf{Q}_w} = \frac{\|Q\|_{\mathbf{Q}_w}}{1 - \alpha C_R}, \\ \|\tau_Q\|_{\mathfrak{E}_w} & \leq C_{\mathfrak{t}, \mathfrak{E}} \|Q\|_{\mathbf{Q}_w}. \end{cases} \quad (51)$$

## 4.2 Proof of Theorem 1

First we recursively define the function  $A_J^{(\ell_j)_{j=1}^J}$  as follows:

$$\begin{cases} A_1^{\ell_1} \widehat{Q}(p, q) & := \widehat{Q}(p - \ell_1, q - \ell_1) - \mathbf{s}_p \widehat{Q}(p - \ell_1, q - \ell_1) \mathbf{s}_q, \\ A_J^{(\ell_1, \mathbf{L})} \widehat{Q}(p, q) & := A_1^{\ell_1} (A_{J-1}^{\mathbf{L}} \widehat{Q})(p, q) \text{ with } J \in \mathbb{N}^*, \ell_j \in \mathbb{R}^3. \end{cases} \quad (52)$$

These functions appear in the Fourier transform of  $Q_{1,0}^{\circ J}[Q]$  (see Appendix C).

**Proof:** It is based on the following fact:

**Lemma 9.** *The trace  $\text{Tr}_{\mathbb{C}^4}$  of the product of an odd number of Dirac matrices (that is  $\alpha_1, \alpha_2, \alpha_3, \beta$ ) vanishes.*

Writing  $\langle a_1, \dots, a_M \rangle$  the algebra spanned by the  $a_j$ 's, we define:

$$\begin{cases} \mathcal{A}_D & := \langle \alpha_1, \alpha_2, \alpha_3, \beta \rangle, \\ \mathcal{A}_D^+ & := \langle \text{Id}, (1 - \delta_{jk}) \alpha_j \alpha_k, \beta \alpha_j \rangle \\ \mathcal{A}_D^- & := \alpha_1 \mathcal{A}_D^+ + \alpha_2 \mathcal{A}_D^+ + \alpha_3 \mathcal{A}_D^+ + \beta \mathcal{A}_D^+ \end{cases} \quad (53)$$

It is clear that  $\mathcal{A}_D = \mathcal{A}_D^+ + \mathcal{A}_D^-$  and Lemma 9 just says that

$$\forall M \in \mathcal{A}_D^- : \text{Tr}_{\mathbb{C}^4}(M) = 0.$$

Remark 24 and Appendix C implies that for almost all  $(p, q) \in \mathbb{R}^3 \times \mathbb{R}^3$ :

- $\widehat{F_{1,0}^{\circ J}}(Q_{0,1}(\rho); p, q) \in \mathcal{A}_D^+$ ,
- if  $\widehat{Q}(p, q) \in \mathcal{A}_D^-$  then so is  $\widehat{F_{1,0}^{\circ J}}(Q; p, q)$ .

Now let us study  $Q_{0,2}(\rho)$ :

$$Q_{0,2} = -\frac{1}{2\pi} \int_{\eta=-\infty}^{+\infty} \frac{d\eta}{\mathcal{D}^0 + i\eta} v_\rho \frac{1}{\mathcal{D}^0 + i\eta} v_\rho \frac{1}{\mathcal{D}^0 + i\eta}.$$

where  $Q_{0,2}^{\varepsilon_1 \varepsilon_2 \varepsilon_3}$  is defined in Notation 11 (as  $Q_{1,2}^{\varepsilon_1, R, \varepsilon_2, v, \varepsilon_3}$  and so on). By the residuum formula in the case  $\varepsilon_1 = \varepsilon_2 = \varepsilon_3$  the term vanishes. We deal with  $Q_{0,2}^{+-}$  and  $Q_{0,2}^{-++}$  together, like  $Q_{0,2}^{++}$  and  $Q_{0,2}^{--}$ ,  $Q_{0,2}^{-+}$  and  $Q_{0,2}^{+-}$ . We compute the first couple with  $A = Q_{0,2}^{+-}$  and  $B = Q_{0,2}^{-++}$ :

$$\begin{aligned} A &= \frac{1}{2\pi} \int_{-\infty}^{+\infty} d\eta \int_{p_1} dp_1 \frac{P_+^0(p)}{\widetilde{E}(p) + i\eta} \widehat{v}(p - p_1) \frac{P_-^0(p_1)}{-\widetilde{E}(p_1) + i\eta} \widehat{v}(p_1 - q) \frac{P_-^0(q)}{-\widetilde{E}(q) + i\eta} \\ &= \int_{p_1} \frac{dp_1}{8} \frac{1}{\widetilde{E}(p) + \widetilde{E}(p_1)} \frac{1}{\widetilde{E}(p) + \widetilde{E}(q)} (1 + \mathbf{s}_p) \widehat{v}(p - p_1) (1 - \mathbf{s}_{p_1}) \widehat{v}(p_1 - q) (1 - \mathbf{s}_q), \\ B &= \frac{1}{2\pi} \int_{-\infty}^{+\infty} d\eta \int_{p_1} dp_1 \frac{P_-^0(p)}{-\widetilde{E}(p) + i\eta} \widehat{v}(p - p_1) \frac{P_+^0(p_1)}{\widetilde{E}(p_1) + i\eta} \widehat{v}(p_1 - q) \frac{P_+^0(q)}{\widetilde{E}(q) + i\eta} \\ &= - \int_{p_1} \frac{dp_1}{8} \frac{1}{\widetilde{E}(p) + \widetilde{E}(p_1)} \frac{1}{\widetilde{E}(p) + \widetilde{E}(q)} (1 - \mathbf{s}_p) \widehat{v}(p - p_1) (1 + \mathbf{s}_{p_1}) \widehat{v}(p_1 - q) (1 + \mathbf{s}_q), \end{aligned}$$

However

$$\begin{aligned} &\frac{1}{2} ((1 + \mathbf{s}_p) \widehat{v}(p - p_1) (1 - \mathbf{s}_{p_1}) \widehat{v}(p_1 - q) (1 - \mathbf{s}_q) - (1 - \mathbf{s}_p) \widehat{v}(p - p_1) (1 + \mathbf{s}_{p_1}) \widehat{v}(p_1 - q) (1 + \mathbf{s}_q)) \\ &= \mathbf{s}_p \widehat{v}(p - p_1) \mathbf{s}_{p_1} \widehat{v}(p_1 - q) \mathbf{s}_q + \mathbf{s}_p \widehat{v}(p - p_1) \widehat{v}(p_1 - q) - \widehat{v}(p - p_1) \widehat{v}(p_1 - q) \mathbf{s}_q - \widehat{v}(p - p_1) \mathbf{s}_{p_1} \widehat{v}(p_1 - q). \end{aligned} \quad (54)$$

In (54) there only remains matrices in  $\mathcal{A}_D^-$ . Symmetrically, the other two couples give:

$$\begin{aligned} &\bullet \frac{1}{2} ((1 + \mathbf{s}_p) \widehat{v}(p - p_1) (1 - \mathbf{s}_{p_1}) \widehat{v}(p_1 - q) (1 + \mathbf{s}_q) - (1 - \mathbf{s}_p) \widehat{v}(p - p_1) (1 + \mathbf{s}_{p_1}) \widehat{v}(p_1 - q) (1 - \mathbf{s}_q)) \\ &= -\mathbf{s}_p \widehat{v}(p - p_1) \mathbf{s}_{p_1} \widehat{v}(p_1 - q) \mathbf{s}_q + \mathbf{s}_p \widehat{v}(p - p_1) \widehat{v}(p_1 - q) + \widehat{v}(p - p_1) \widehat{v}(p_1 - q) \mathbf{s}_q - \widehat{v}(p - p_1) \mathbf{s}_{p_1} \widehat{v}(p_1 - q), \\ &\bullet \frac{1}{2} ((1 - \mathbf{s}_p) \widehat{v}(p - p_1) (1 - \mathbf{s}_{p_1}) \widehat{v}(p_1 - q) (1 + \mathbf{s}_q) - (1 + \mathbf{s}_p) \widehat{v}(p - p_1) (1 + \mathbf{s}_{p_1}) \widehat{v}(p_1 - q) (1 - \mathbf{s}_q)) \\ &= \mathbf{s}_p \widehat{v}(p - p_1) \mathbf{s}_{p_1} \widehat{v}(p_1 - q) \mathbf{s}_q - \mathbf{s}_p \widehat{v}(p - p_1) \widehat{v}(p_1 - q) + \widehat{v}(p - p_1) \widehat{v}(p_1 - q) \mathbf{s}_q - \widehat{v}(p - p_1) \mathbf{s}_{p_1} \widehat{v}(p_1 - q). \end{aligned} \quad (55)$$

Therefore for almost all  $(p, q)$ :  $\widehat{Q}_{0,2}(\rho; p, q) \in \mathcal{A}_D^-$ : its trace  $\text{Tr}_{\mathbb{C}^4}$  vanishes. Furthermore for all  $J \geq 1$ :

$$\widehat{\rho}(F_{1,0}^{\circ J}(Q_{0,2}(\rho)); k) = \text{Cst} \iint_{u, \ell_1} \cdots \int \frac{dud\ell}{\prod_{\ell_j, 1 \leq j \leq J} |\ell_j|^2} \text{Tr}_{\mathbb{C}^4} \frac{A_J^{(\ell_j)_{j=1}^J} \widehat{Q}_{0,2}(\rho)(u + \frac{k}{2}, u - \frac{k}{2})}{\prod_{0 \leq j \leq J} (\widetilde{E}(u + k/2 - L_j) + \widetilde{E}(u - k/2 - L_j))} \quad (56)$$

where for almost all  $(p, q, \ell_j)$ :  $\text{Tr}_{\mathbb{C}^4} \left\{ A_J^{(\ell_j)_{j=1}^J} \widehat{Q}_{0,2}(\rho; p, q) \right\} = 0$  because these matrices are in  $\mathcal{A}_D^-$ . Thus  $\widehat{\rho}(F_{1,0}^{\circ J}(Q_{0,2}(\rho)); k) = 0$  for almost all  $k \in \mathbb{R}^3$  and so  $\widehat{\tau}_{0,2}(\rho; k) = 0$  for almost all  $k \in \mathbb{R}^3$ . In other words  $\widehat{\tau}_{0,2}(\rho) = 0$ .

There remains to prove that  $\tau_{1,1}(\alpha \mathbf{T}(Q_{0,1}(\rho_0)), \rho_1) = 0$ : it suffices to show that for all  $J, J' \geq 0$ :  $\rho \left\{ F_{1,0}^{\circ J} [Q_{1,1}(\alpha F_{1,0}^{\circ J'} [Q_{0,1}(\rho_0)], \rho_1)] \right\}$  vanishes. As before we treat together

- $Q_{1,1}^{+R-v-}(F_{1,0}^{\circ J'}(Q_{0,1}(\rho_0)), \rho_1)$  and  $Q_{1,1}^{-R+v+}(F_{1,0}^{\circ J'}(Q_{0,1}(\rho_0)), \rho_1)$ ,
- then  $Q_{1,1}^{+v-R-}(F_{1,0}^{\circ J'}(Q_{0,1}(\rho_0)), \rho_1)$  and  $Q_{1,1}^{-v+R+}(F_{1,0}^{\circ J'}(Q_{0,1}(\rho_0)), \rho_1)$ , and so on.

As  $F_{1,0}^{\circ J'}(Q_{0,1}(\rho_0); p, q) \in \mathcal{A}_D^+$  for almost all  $p, q$ , then  $\widehat{Q}_{1,1}^{+R-v-}(F_{1,0}^{\circ J'}(Q_{0,1}(\rho_0); p, q), \rho_1) + \widehat{Q}_{1,1}^{-R+v+}(F_{1,0}^{\circ J'}(Q_{0,1}(\rho_0)), \rho_1; p, q) \in \mathcal{A}_D^-$  for almost all  $p, q$  thanks to (54) and (55). So its trace  $\text{Tr}_{\mathbb{C}^4}$  vanishes. The same result holds for the other cases:  $Q_{1,1}^{+v-R-} + Q_{1,1}^{-v+R+}$ ,  $Q_{1,1}^{+-+} + Q_{1,1}^{-+-}$  and  $Q_{1,1}^{-+-} + Q_{1,1}^{+-+}$ . Finally as in (56) we have:

$$\widehat{\rho}(F_{1,0}^{\circ J}(Q_{1,1}(F_{1,0}^{\circ J'}(\rho_0), \rho_1)); k) = 0 \text{ for almost all } k.$$

□

## 5 The fixed point method

We prove here Lemmas 2, 3 and 4 and start with some inequalities.

### 5.1 Tools

- We recall the following Sobolev inequalities in  $\mathbb{R}^3$ : for suitable  $f$  –say  $H^1$ – we have

$$\|f\|_{L^6} \lesssim \|\nabla f\|_{L^2}, \quad \|f\|_{L^4} \lesssim \| |\nabla|^{3/4} f \|_{L^2}, \quad \|f\|_{L^3} \lesssim \| |\nabla|^{1/2} f \|_{L^2}. \quad (57)$$

We use them to prove the following inequalities: for  $\rho \in \mathcal{C}$ ,  $v_\rho := \rho * \frac{1}{|\cdot|}$  and  $\phi \in H^{1/2}$ :

$$\|v_\rho \phi\|_{L^2} \lesssim \|v_\rho\|_{L^6} \|\phi\|_{L^3} \lesssim \|\rho\|_{\mathcal{C}} \| |\nabla|^{1/2} \phi \|_{L^2}. \quad (58)$$

$$\|\rho * \frac{1}{|\cdot|}\|_{L^4} \lesssim \| |\nabla|^{3/4} \rho * \frac{1}{|\cdot|} \|_{L^2} \lesssim \sqrt{\int \frac{|\widehat{\rho}(k)|^2}{|k|^{5/2}} dk} \lesssim \left( \inf_{\varepsilon > 0} \left\{ 2\pi \varepsilon^{1/2} \|\widehat{\rho}\|_{L^\infty}^2 + \varepsilon^{-1/2} \|\rho\|_{\mathcal{C}}^2 \right\} \right)^{1/2}. \quad (59)$$

With  $v_\rho := \rho * \frac{1}{|\cdot|}$  Eq. (59) is used in:

$$\| \frac{1}{\mathcal{D}^{0+i\eta}} v_\rho \|_{\mathfrak{S}_4}, \| \frac{1}{|\mathcal{D}^{0+i\eta}|^{1/2}} v_\rho \frac{1}{|\mathcal{D}^{0+i\eta}|^{1/2}} \|_{\mathfrak{S}_4} \leq \frac{K_2^{1/4}}{E(\eta)^{1/4}} \|\rho * \frac{1}{|\cdot|}\|_{L^4} \quad (60)$$

- We recall Kato's and Hardy's inequalities for  $\phi \in L^2(\mathbb{R}^3)$ :

$$\begin{cases} \int_{\mathbb{R}^3} \frac{|\varphi(x)|^2}{|x|} dx & \leq \frac{\pi}{2} \langle |\nabla| \varphi, \varphi \rangle, \\ \int_{\mathbb{R}^3} \frac{|\varphi(x)|^2}{|x|^2} dx & \leq 4 \langle (-\Delta) \varphi, \varphi \rangle, \end{cases} \quad (61)$$



and the Kato-Seiler-Simon's inequality (KSS) for compact operators in  $\mathcal{B}(L^2(\mathbb{R}^3))$ :

$$\forall 2 \leq p \leq +\infty : \|f(-i\nabla)g(x)\|_{\mathfrak{S}_p} \leq (2\pi)^{-3/p} \|f\|_{L^p} \|g\|_{L^p}. \quad (62)$$

- We recall that for any  $p, q \in B(0, \Lambda)$  we have (see [23].)

$$|\widehat{\mathcal{P}}_-^0(p) - \widehat{\mathcal{P}}_-^0(q)| = |\widehat{\mathcal{P}}_+^0(p) - \widehat{\mathcal{P}}_+^0(q)| \lesssim \frac{|p - q|}{\max(\widetilde{E}(p), \widetilde{E}(q))}. \quad (63)$$

By Ineq. (63) we get the following.

**Lemma 10.** *Let  $\rho \in \mathcal{C}$ , then there exists  $K > 0$  such that for any  $a > 1/2$  and  $\varepsilon \in \{+, -\}$  we have:*

$$\|\mathcal{P}_\varepsilon^0 v_\rho \mathcal{P}_{-\varepsilon}^0 |D_0|^{-a}\|_{\mathfrak{S}_2} \leq \frac{K}{\sqrt{2a-1}} \|\rho\|_{\mathcal{C}}.$$

**Proof:** It is obvious once we have seen that the norm of the integral kernel of its Fourier transform is lesser than:

$$K \frac{|\widehat{\rho}(p - q)|}{|p - q|} \frac{1}{E(q)^a \max(E(q), E(p))}.$$

□

## 5.2 Estimate on $Q_{0,1}$

We estimate  $\|Q_{0,1}\|_{\mathfrak{Q}_w}$  as in [8]. We have

$$\begin{aligned} \int_{B(0,\Lambda)} \frac{du}{E(u + \varepsilon k/2)^2} \frac{\widetilde{E}(u + k/2) + \widetilde{E}(u - k/2)}{(\widetilde{E}(u + k/2) + \widetilde{E}(u - k/2))^2} &\leq 4\pi \int_0^\Lambda \frac{du}{\sqrt{1+r^2}} \\ &\leq 4\pi(1 + \log(\Lambda)) \lesssim \log(\Lambda), \end{aligned} \quad (64)$$

leading to:

$$\iint w(p - q) (\widetilde{E}(p) + \widetilde{E}(q)) |\widehat{Q}_{0,1}(\rho; p, q)|^2 dp dq \lesssim (1 + \log(\Lambda)) \|\rho\|_{\mathfrak{C}_w}^2, \quad (65)$$

where we have used (63).

## 5.3 Proof of Lemma 2

We recall that for  $J \geq 1$ :

$$Q_J(Q, \rho) := \frac{1}{2\pi} \int_{-\infty}^{+\infty} \frac{d\eta}{\mathcal{D}^0 + i\eta} \prod_{1 \leq j \leq J} \left( (v_\rho - R_Q) \frac{1}{\mathcal{D}^0 + i\eta} \right)$$

We write

$$\mathfrak{a}(Q) := \mathcal{F}^{-1}(|\widehat{Q}|) \text{ and } \mathfrak{a}(\rho) := \mathcal{F}^{-1}(|\widehat{\rho}|).$$

It is clear that  $|\widehat{Q}_{k,\ell}(p, q)|$  is lesser than the integral kernel of the Fourier transform of

$$\mathfrak{a}(Q_{k,\ell}) := \frac{1}{2\pi} \int_{-\infty}^{+\infty} \frac{d\eta}{\sqrt{|\mathcal{D}^0|^2 + \eta^2}} \left( \mathfrak{a}(\rho) * \frac{1}{|\cdot|} + R[\mathfrak{a}(Q)] \right)^J.$$

We write  $\mathfrak{a}(v_\rho) = v_{\mathfrak{a}(\rho)}$  and  $\mathfrak{a}(R_Q) := R_{\mathfrak{a}(Q)}$  and  $d_\eta := \sqrt{|\mathcal{D}^0|^2 + \eta^2}$ . We have:

$$\left| \begin{array}{lll} \|\mathfrak{a}(v_\rho)\|_{L^6} & \lesssim & \|\nabla \mathfrak{a} v_\rho\|_{L^2} \lesssim \|\mathfrak{a}(\rho)\|_{\mathcal{C}} = \|\rho\|_{\mathcal{C}}, \\ \|\mathfrak{a}(v_\rho)\|_{L^4} & \lesssim & \|\nabla|^{3/2} \mathfrak{a}(v_\rho)\|_{L^2} \lesssim \|\widehat{\mathfrak{a}(\rho)}\|_{L^\infty} + \|\mathfrak{a}(\rho)\|_{\mathcal{C}} = \|\widehat{\rho}\|_{L^\infty} + \|\rho\|_{\mathcal{C}}, \\ \|\frac{1}{|\cdot|^{1/2}} \mathfrak{a}(R_Q)\|_{\mathfrak{S}_2} & \lesssim & \|\mathfrak{a}(R_Q)\|_{\text{Ex}} \lesssim \|\mathfrak{a}(Q)\|_{\text{T}} = \|Q\|_{\text{T}}. \end{array} \right.$$

By the KSS inequality, there exist  $C_6, C_4 > 0$  such that:

$$\begin{aligned} \|d_\eta^{-1/2} v_\rho d_\eta^{-1/2}\|_{\mathfrak{S}_6} &\leq C_6 E(\eta)^{-1/2} \|\rho\|_c, \\ \|d_\eta^{-5/12} v_\rho d_\eta^{-7/12}\|_{\mathfrak{S}_4} &\leq C_4 E(\eta)^{-1/4} \|v_\rho\|_{L^4}. \end{aligned} \quad (66)$$

As  $w$  satisfies (8), we have:

$$w(p-q)\widehat{\mathfrak{a}}(Q_J(Q, \rho); p, q) \leq JK_{(w)}^J \widehat{\mathfrak{a}}\left(Q_J[\mathcal{F}^{-1}(w(p'-q')\widehat{Q}(p', q')), \mathcal{F}^{-1}(\rho)]; p, q\right).$$

It suffices to check that for  $p_0 = p, p_{J+1} = q$  and  $p_1, \dots, p_J \in \mathbb{R}^3$  we have:

$$w(p-q) \leq \sum_{j=1}^{J+1} K_{(w)}^j w(p_{j-1} - p_j) \leq JK_{(w)}^J \prod_{j=1}^{J+1} w(p_{j-1} - p_j).$$

In the definition of  $\|\cdot\|_{\mathbf{Q}_w}$ , there remains to multiply by  $\widetilde{E}(p)^{1/2} + \widetilde{E}(q)^{1/2}$ . We use the first or the last  $d_\eta^{-1}$  to get:

$$\frac{\widetilde{E}(r)^{1/2}}{\sqrt{\widetilde{E}(r)^2 + \eta^2}} \leq \frac{1}{(\widetilde{E}(r)^2 + \eta^2)^{1/4}} \text{ with } r \in \{p, q\}.$$

For the terms  $Q_J(Q, \rho)$  with  $J \geq 3$  we get that:

$$\|\mathfrak{a}Q_J(Q, \rho)\|_{\mathbf{Q}_w} \leq \frac{JK_{(w)}^J}{2\pi} \left( \|\frac{1}{|\cdot|^{1/2}} R[\mathfrak{a}(Q)]\|_{\mathfrak{S}_2} + C_6 \|\rho\|_c \right)^J \int_{-\infty}^{+\infty} \frac{d\eta}{\widetilde{E}(\eta)^{(J+1)/2}}.$$

For  $J = 2$ , we treat  $Q_{0,2}(\rho)$  in another way because the product of two operators in  $\mathfrak{S}_6$  is not necessarily Hilbert-Schmidt. By the Cauchy expansion we have [8]

$$Q_J^{+\dots+} = Q_J^{-\dots-} = 0.$$

So it suffices to treat  $Q_{0,2}^{\varepsilon_1, \varepsilon_2, \varepsilon_3}$  with  $(\varepsilon_1, \varepsilon_2, \varepsilon_3) \neq (++++), (----)$ . In particular there is a change of sign  $+-$  or  $-+$ . By Hölder inequality and Lemma 10 we have for  $\varepsilon \in \{+, -\}$ :

$$\|d_\eta^{-1/2} v_\rho^{\varepsilon, -\varepsilon} d_\eta^{-1/4}\|_{\mathfrak{S}_2} \leq \|\rho\|_c \left\{ \int \frac{dq}{E(q)^{7/2}} \right\}^{1/2} \leq \|\rho\|_c.$$

Hence using the above inequality and (66) we get:

$$\|Q_{0,2}(\rho)\|_{\mathbf{Q}_w} \leq \|\rho\|_c^2 \int_{-\infty}^{+\infty} \frac{d\eta}{E(\eta)^{1+4^{-1}}}.$$

By (29), there exists  $K > 0$  such that

$$\|Q_J(Q, \rho)\|_{\mathbf{Q}_w} \leq J^{1/2} (K \times K_{(w)} (\|Q\|_{\mathbf{Q}_w} + \|\rho\|_{\mathfrak{C}_w}))^J.$$

To deal with  $\rho_J$ , we use the same method as in [8] and estimate  $\|\rho_J\|_c$  by duality. We take a Schwartz function  $\zeta \in \mathcal{S}(\mathbf{R}^3)$  and prove that for any  $k, \ell \geq 0$  with  $k + \ell \geq 2$  we have:

$$|\mathrm{Tr}(Q_{k,\ell}\zeta)| \leq K(Q, \rho, k, \ell) \sqrt{\int \frac{|p|^2 |\widehat{\zeta}(p)|^2}{g(p)^2} dp} = K(Q, \rho, k, \ell) \|\zeta\|_{\mathfrak{C}_w'}.$$

We emphasize that by Furry's Theorem [5, 8] we have  $\rho_{0,2J} = 0$  for any  $J \in \mathbb{N}^*$ . First we must prove that  $Q_{k,\ell}\zeta$  is trace-class. We use the same method as in [8]:

$$\|Q_{k,\ell}\zeta\|_{\mathfrak{S}_1} \leq \|Q_{k,\ell} \mathcal{D}^0\|_{\mathfrak{S}_2} \|\frac{1}{|\mathcal{D}^0|^2} \zeta\|_{\mathfrak{S}_2} \leq E(\Lambda)^2 \|Q_{k,\ell}\|_{\mathfrak{S}_2} \|\zeta\|_{L^2}.$$

It is clear that  $|\widehat{Q_{k,\ell}\zeta}(p, p)| \leq |\mathfrak{a}(\widehat{Q_{k,\ell}})\zeta|$ .

Writing  $d_\eta(p) := \sqrt{\widehat{E}(p)^2 + \eta^2}$ ,  $p_0 = p$  and  $\mathbf{m} = (m_1, \dots, m_J) \in \{v_\rho, R_Q\}^J$  we have:

$$2\pi |\mathfrak{a}(\widehat{Q_{k,\ell}^{\mathbf{m}}})\zeta(p, p)| \leq \int_{\mathbb{R}} d\eta \int_{(B(0,\Lambda))^J} \frac{d\mathbf{p}}{d_\eta(p)} \prod_{j=1}^J |\widehat{m}_j(p_j, p_{j-1})| d_\eta(p_j)^{-1} |\widehat{\zeta}(p_J - p)|$$

We replace  $|\widehat{\zeta}(p_J - p)|$  by:

$$|\widehat{\zeta}(p_J - p)| \times \frac{w(p_J - p)}{w(p_J - p)} \leq JK_{(w)}^J \frac{|\widehat{\zeta}(p_J - p)|}{w(p_J - p)} \prod_{j=1}^J w(p_j - p_{j-1}) =: JK_{(w)}^J |\widehat{\zeta}'(p_J - p)| \prod_{j=1}^J w(p_j - p_{j-1}). \quad (67)$$

We write  $R' := R[\mathcal{F}^{-1}(w(p-q)|\widehat{Q}(p,q))]$  and  $V' := v[\mathcal{F}^{-1}(w(p)|\widehat{\rho}(p))]$ .

For  $(k, \ell)$  different from  $(0, 3), (1, 1), (0, 2J)$  we have:

$$\begin{aligned} |\mathrm{Tr}(Q_{k,\ell}\zeta)| &\leq \frac{(k+\ell)K_{(w)}^{k+\ell}}{2\pi} \binom{k+\ell}{k} \int_{\mathbb{R}} d\eta \|d_\eta^{-1/2}\zeta' d_\eta^{-1/2}\|_{\mathfrak{S}_6} \|d_\eta^{-1/2}R' d_\eta^{-1/2}\|_{\mathfrak{S}_2}^k \|d_\eta^{-1/2}V' d_\eta^{-1/2}\|_{\mathfrak{S}_6}^\ell \\ &\leq \frac{(k+\ell)K_{(w)}^{k+\ell}}{2\pi} \binom{k+\ell}{k} K^{k+\ell} \int_{\mathbb{R}} \frac{d\eta}{E(\eta)^{(1+j+\ell)/2}} \|Q\|_{\mathbf{Q}_w}^k \|\rho\|_{\mathfrak{C}_w}^\ell. \end{aligned}$$

To deal with  $\rho_{1,1}, \rho_{0,3}$  we use the same method as the one used for  $\|Q_{0,2}\|_{\mathbf{Q}_w}$ . We treat the case of  $\rho[Q_{1,1}^{+R-v-}]$  as an example and the other cases are similar and left to the reader.

$$|\mathrm{Tr}_{\mathfrak{C}^4}(\widehat{Q_{1,1}^{+R-v-}}(p_0, p_2)\widehat{\zeta}(p_2 - p_0))| \leq \int_{\mathbb{R}} \int_{(B(0,\Lambda))^3} \frac{d\eta dp_1 dp_2 |\widehat{R}_Q(p_0, p_1)| |\widehat{v}(p_1 - p_2)|}{d_\eta(p_0) d_\eta(p_1) d_\eta(p_2)} |\widehat{\zeta}^-(p_2 - p_0)|.$$

Using Lemma 10 and (67) we get that:

$$|\mathrm{Tr}(Q_{1,1}^{+R-v-}\zeta)| \leq \|Q\|_{\mathbf{Q}_w} \|\rho\|_{\mathfrak{C}_w} K_{5/4} \|\zeta\|_{\mathfrak{C}'_w}.$$

□

## 5.4 Estimates for $F^{(2)}$

We consider  $\gamma' = \gamma + N$  a minimizer of  $E_{\mathrm{BDF}}^v(M)$  and define the function  $F^{(2)}$  (32). Two Banach spaces will be considered: first  $\mathcal{C}$  and then  $\mathcal{C} \cap L^1$ . We recall that for  $\eta \in \mathbb{R}$  we write  $d_\eta = \sqrt{|\mathcal{D}^0|^2 + \eta^2}$ .

### 5.4.1 Estimates on the $\mathcal{C}$ -norm

Thanks to previous estimates (Lemmas 5, 6, *a priori* estimates (38) and estimates in the  $\|\cdot\|_{\mathfrak{C}_w}$ -norm), in the regime  $M, \|\nu\|_{\mathcal{C}} \leq \log(\Lambda)$  there hold the following *non-sharp* estimates:

$$\begin{cases} \|h_2\|_{\mathcal{C}} &\leq \alpha^2 \left\{ \|\rho''\|_{\mathcal{C}} [\|N\|_{\mathcal{T}} + \alpha^2 (\|\gamma'\|_{\mathcal{T}} + \|\rho''\|_{\mathcal{C}})^2] + \|\gamma'\|_{\mathcal{T}}^2 \right\} \\ &\leq \alpha^2 \times \log(\Lambda) = L\alpha \\ \|h_3\|_{\mathcal{C}} &\leq \alpha^3 (\|\gamma'\|_{\mathcal{T}} + \|\rho''\|_{\mathcal{C}})^3 \leq (L\alpha)^{3/2}. \end{cases} \quad (68)$$

Then  $F_2^{(2)}(\rho'')$  and  $F_3^{(2)}(\rho'')$  are at most cubic in  $\rho''$ :

$$\begin{cases} \|F_2^{(2)}(\rho')\|_{\mathcal{C}} &\leq \alpha^4 (\|\gamma'\|_{\mathcal{T}} + \|\rho''\|_{\mathcal{C}}) \|\rho''\|_{\mathcal{C}}^2 \\ \|F_3^{(2)}(\rho')\|_{\mathcal{C}} &\leq \alpha^3 (\|\rho''\|_{\mathcal{C}} + \|\gamma'\|_{\mathcal{T}}) \|\rho''\|_{\mathcal{C}}^2 \\ \|dF_2^{(2)}(\rho')\|_{\mathcal{L}(\mathcal{C})} &\leq \alpha^4 (\|\gamma'\|_{\mathcal{T}} \|\rho''\|_{\mathcal{C}} + \|\rho''\|_{\mathcal{C}}^2) \\ \|dF_3^{(2)}(\rho')\|_{\mathcal{L}(\mathcal{C})} &\leq \alpha^3 (\|\gamma'\|_{\mathcal{T}} \|\rho''\|_{\mathcal{C}} + \|\rho''\|_{\mathcal{C}}^2). \end{cases} \quad (69)$$

### 5.4.2 Estimates on the $L^1$ -norm

Our aim in this part is to prove Lemma 11 below which states that  $F^{(2)}$  is a well-defined  $\mathcal{C}^1$  function of  $\mathcal{C} \cap L^1$  (differentiable with a continuous differential).

• We first prove that  $h_2, h_3 \in L^1$  (we recall they are defined in (31)). In fact they are densities of trace-class operators: to see this we use the methods of the proof of Lemma 2.

1.  $N = \sum_j |\psi_j\rangle\langle\psi_j| \in \mathfrak{S}_1$  so  $\mathbf{T}[N] \in \mathfrak{S}_1$  and

$$\|\tau_N\|_{L^1} \leq \|\mathbf{T}[N]\|_{\mathfrak{S}_1} \leq C_{\mathbf{T}, \mathfrak{S}} \|N\|_{\mathfrak{S}_1}. \quad (70)$$

2.  $Q_{2,0}(\gamma') \in \mathfrak{S}_1$  : We have:

$$\|Q_{2,0}(\gamma')\|_{\mathfrak{S}_1} \lesssim \|\gamma'\|_{\mathbb{E}_x}^2 K_2. \quad (71)$$

3.  $Q_{0,\ell}(\rho''_\gamma)$  with  $\ell \geq 4$ . As  $Q_{0,\ell}^{+\dots+} = Q_{0,\ell}^{-\dots-} = 0$  there is at least one change of sign  $+-$  or  $-+$ . Then with the help of Lemma 10 and Estimates (66) we have

$$\|Q_{0,\ell}(\rho''_\gamma)\|_{\mathfrak{S}_1} \lesssim \|\rho''_\gamma\|_{\mathcal{C}}^\ell K_{\frac{\ell+1}{2} + \frac{1}{4}},$$

the product of  $\ell - 1$  operators in  $\mathfrak{S}_6$  and one in  $\mathfrak{S}_2$  is trace-class.

4. Similarly  $Q_{k,\ell}(\gamma', \rho''_\gamma) \in \mathfrak{S}_1$  with  $k \geq 2$  or  $k \geq 1$  and  $\ell \geq 3$  :

$$\|Q_{k,\ell}(\gamma', \rho''_\gamma)\|_{\mathfrak{S}_1} \lesssim \binom{k+\ell}{k} (K\|\gamma'\|_{\mathbf{T}})^k (K\|\rho''_\gamma\|_{\mathcal{C}})^\ell K_{1+(k+\ell)/2}. \quad (72)$$

5. Thanks to Furry's Theorem and Theorem 1:

$$\tau\{Q_{0,2}(\rho''_\gamma)\} = \tau_{1,1}\{\mathbf{T}[Q_{0,1}(\rho''_\gamma)], \rho''_\gamma\} = 0. \quad (73)$$

6. By the same methods as before we have  $Q_{0,3}(\rho''_\gamma), Q_{1,2}(\gamma', \rho''_\gamma) \in \mathfrak{S}_{6/5}$  with:

$$\|Q_{0,3}(\rho''_\gamma)\|_{\mathfrak{S}_{6/5}} \lesssim \|\rho''_\gamma\|_{\mathcal{C}}^3 K_{2+1/4} \text{ and } \|Q_{1,2}(\gamma', \rho''_\gamma)\|_{\mathfrak{S}_{6/5}} \lesssim \|\gamma'\|_{\mathbf{T}} \|\rho''_\gamma\|_{\mathcal{C}}^2 K_{1+3/2}.$$

Furthermore the following inequalities hold (we recall that  $Y$  is defined in (50)):

$$\|d_\eta^{-3/8} v_\rho d_\eta^{-5/8}\|_{\mathfrak{S}_6} \lesssim E(\eta)^{-1/2} \|\rho\|_{\mathcal{C}} \text{ and } \|d_\eta^{-5/8} R(\mathbf{T}[Q]) d_\eta^{-5/8}\|_{\mathfrak{S}_{6/5}} \lesssim Y(\frac{6}{5}) \|Q\|_{\mathfrak{S}_{6/5}}.$$

Thus

$$\begin{cases} \|\mathbf{T}_{1,1}\{\mathbf{T}Q_{0,3}(\rho''_\gamma), \rho''_\gamma\}\|_{\mathfrak{S}_1} & \lesssim 2C_{\mathbf{T}, \mathfrak{S}} K_{5/4} \|\rho''_\gamma\|_{\mathcal{C}} (Y(\frac{6}{5}) \|\rho''_\gamma\|_{\mathcal{C}}^3 K_{2+1/4}), \\ \|\mathbf{T}_{1,1}\{\mathbf{T}Q_{1,2}(\gamma', \rho''_\gamma), \rho''_\gamma\}\|_{\mathfrak{S}_1} & \lesssim 2C_{\mathbf{T}, \mathfrak{S}} K_{5/4} \|\rho''_\gamma\|_{\mathcal{C}} (3Y(\frac{6}{5}) \|\gamma'\|_{\mathbf{T}} \|\rho''_\gamma\|_{\mathcal{C}}^2 K_{1+3/2}) \\ \|\mathbf{T}_{1,1}\{\mathbf{T}N, \rho''_\gamma\}\|_{\mathfrak{S}_1} & \lesssim 2C_{\mathbf{T}, \mathfrak{S}} K_{5/4} \|\rho''_\gamma\|_{\mathcal{C}} Y(\frac{6}{5}) M. \end{cases} \quad (74)$$

7. We apply  $\mathbf{T}$ ,  $h_2$  (resp.  $h_3$ ) is the density of  $Q(h_2)$  (resp.  $Q(h_3)$ ) with

$$\begin{cases} Q(h_2) & = \alpha^2 \left\{ \mathbf{T}Q_{1,1}[\mathbf{T}N + \alpha^2 \mathbf{T}[Q_{2,0}(\gamma') + \tilde{Q}_3(\gamma', \rho''_\gamma)]; \rho''_\gamma \right\} + \mathbf{T}Q_{2,0}(\gamma') \\ Q(h_3) & = \alpha^3 \left\{ \mathbf{T}Q_{3,0}(\gamma') + \mathbf{T}Q_{2,1}(\gamma', \rho''_\gamma) + \alpha \tilde{Q}_4(\gamma', \rho''_\gamma) \right\} \end{cases}$$

The previous estimates lead to a sequence of numbers  $(b_\ell)_{\ell \geq 2}$  with the following asymptotic behaviour:

$$b_\ell = \mathcal{O}_{\ell \rightarrow +\infty}(\ell^{1/2}) \quad (75)$$

and a constant  $C_0 > 0$  such that:

$$\begin{aligned} & \left\| \alpha^2 Q_{2,0}(\gamma') + \alpha^3 [Q_{3,0} + Q_{2,1}](\gamma', \rho''_\gamma) + \alpha^4 \tilde{Q}_4(\gamma', \rho''_\gamma) \right\|_{\mathfrak{S}_1} \\ & + \alpha^3 \left\| Q_{0,3}(\rho''_\gamma) + Q_{1,2}(\gamma', \rho''_\gamma) \right\|_{\mathfrak{S}_{6/5}} \leq \sum_{\ell=2}^{+\infty} b_\ell (\alpha C_0)^\ell (\|\rho''_\gamma\|_{\mathcal{C}} + \|\gamma'\|_{\mathbf{T}})^\ell =: A_{h, \mathfrak{S}}. \end{aligned} \quad (76)$$

We have:

$$\|Q(h_2)\|_{\mathfrak{S}_1} \lesssim \alpha^2 C_{\mathbf{T},\mathfrak{S}} (2K_{5/4} Y(\frac{6}{5})(M + A_{h,\mathfrak{S}}) + \|\gamma'\|_{\mathbf{Q}_w}^2) \quad (77)$$

and write  $B_{h_2,\mathfrak{S}}$  this upper bound. Similarly:

$$\|Q(h_3)\|_{\mathfrak{S}_1} \leq C_{\mathbf{T},\mathfrak{S}} \sum_{\ell=3}^{+\infty} b_\ell (\alpha C_0)^\ell (\|\rho''\|_c + \|\gamma'\|_{\mathbf{T}})^\ell =: B_{h_3,\mathfrak{S}_1}. \quad (78)$$

*Remark 18.* The introduced numbers  $A_{h,\mathfrak{S}}, B_{h_2,\mathfrak{S}_1}, B_{h_3,\mathfrak{S}}$  are *not* constants: they all depend on  $\alpha$  and the minimizer  $\gamma'$ . As *a priori* estimates hold (Lemma 5), these upper bounds are small provided that we are in the regime of Remark 10. Indeed we have

$$(1 - \frac{\alpha\pi}{4}) \|\gamma'\|_{\mathbf{T}}^2 + \frac{\alpha}{2} \|\rho''\|_c^2 \leq \frac{\alpha}{2} \|\nu\|_c^2 + M,$$

so  $\alpha(\|\gamma'\|_{\mathbf{T}} + \|\rho''\|_c) \lesssim \alpha\|\nu\|_c + \sqrt{\alpha M} = \mathcal{O}((L\alpha)^{1/4})$ . In particular those upper bounds are  $o(1)$ .

• Let us estimate the  $L^1$ -norm of  $F_2^{(2)}(\rho'')$  and  $F_3^{(2)}(\rho'')$  with  $\rho'' \in \mathcal{C} \cap L^1$ . To this end we use (60) and (59) at level  $\varepsilon = 1$  for instance: there exists  $K_{L^4}^{(v)} > 0$  such that:

$$\|v_{\rho''}\|_{L^4} \leq K_{L^4}^{(v)} \{\|\rho''\|_{L^1} + \|\rho''\|_c\}. \quad (79)$$

We use the second inequality of (66) and Lemma 10 with  $a = 7/12$ . Using the method of the proof of Lemma 2, we obtain the following.

**Lemma 11.** *Let  $\rho''$  be in  $\mathcal{C} \cap L^1$  and  $\gamma'$  a minimizer for  $E_{BDF}^\nu(M)$  with density  $\rho'_j$ . We have:*

$$\begin{aligned} \|\mathbf{T}Q_{0,3}(\rho'')\|_{\mathfrak{S}_1} &\lesssim 6K_{13/12} C_{\mathbf{T},\mathfrak{S}} \{\|\rho''\|_{L^1} + \|\rho''\|_c\}^2 \|\rho''\|_c \\ \|\mathbf{T}Q_{1,2}(\gamma', \rho'')\|_{\mathfrak{S}_1} &\lesssim \binom{3}{1} K_2 C_{\mathbf{T},\mathfrak{S}} \|\gamma'\|_{\mathbf{T}} \{\|\rho''\|_{L^1} + \|\rho''\|_c\}^2 \\ \|Q_{0,2}(\rho'')\|_{\mathfrak{S}_{4/3}} &\lesssim 4K_{7/3} \|\rho''\|_c \{\|\rho''\|_{L^1} + \|\rho''\|_c\} \\ \|Q_{1,1}(\gamma', \rho'')\|_{\mathfrak{S}_{4/3}} &\lesssim 2K_{7/4} \|\gamma'\|_{\mathbf{T}} \{\|\rho''\|_{L^1} + \|\rho''\|_c\} \\ \|\mathbf{T}Q_{1,1}\{\mathbf{T}Q_{0,2}(\rho''), \rho''\}\|_{\mathfrak{S}_1} &\lesssim 2K_{13/12} Y(\frac{4}{3}) C_{\mathbf{T},\mathfrak{S}} \|Q_{0,2}(\rho'')\|_{\mathfrak{S}_{4/3}} \{\|\rho''\|_{L^1} + \|\rho''\|_c\} \\ \|\mathbf{T}Q_{1,1}\{\mathbf{T}Q_{1,1}(\gamma', \rho''), \rho''\}\|_{\mathfrak{S}_1} &\lesssim 2K_{13/12} Y(\frac{4}{3}) C_{\mathbf{T},\mathfrak{S}} \|Q_{1,1}(\gamma', \rho'')\|_{\mathfrak{S}_{4/3}} K_{L^4}^{(v)} \{\|\rho''\|_{L^1} + \|\rho''\|_c\} \end{aligned} \quad (80)$$

Similarly we can estimate  $\|dF_j^{(2)}\|_{L(\mathcal{C} \cap L^1)}$ . As  $\|\gamma'\|_{\mathbf{T}} \lesssim \sqrt{\log(\Lambda)}$  we have:

$$\begin{cases} \|F_2^{(2)}(\rho'')\|_{\mathcal{C} \cap L^1} &\lesssim \alpha^4 \|\rho''\|_{\mathcal{C} \cap L^1}^2 \{\sqrt{\log(\Lambda)} + \|\rho''\|_{\mathcal{C} \cap L^1}\} \\ \|F_3^{(2)}(\rho'')\|_{\mathcal{C} \cap L^1} &\lesssim \alpha^3 \|\rho''\|_{\mathcal{C} \cap L^1}^2 \{\sqrt{\log(\Lambda)} + \|\rho''\|_{\mathcal{C} \cap L^1}\}, \\ \|dF_2^{(2)}(\rho'')\|_{L(\mathcal{C} \cap L^1)} &\lesssim \alpha^4 \|\rho''\|_{\mathcal{C} \cap L^1}^2 \{\sqrt{\log(\Lambda)} + \|\rho''\|_{\mathcal{C} \cap L^1}\}, \\ \|dF_3^{(2)}(\rho'')\|_{L(\mathcal{C} \cap L^1)} &\lesssim \alpha^3 \|\rho''\|_{\mathcal{C} \cap L^1}^2 \{\sqrt{\log(\Lambda)} + \|\rho''\|_{\mathcal{C} \cap L^1}\}. \end{cases} \quad (81)$$

## 5.5 Application of the Banach fixed point theorem

### 5.5.1 $F^{(1)}$

With *exactly* the same method of [8] let us apply the Banach fixed point theorem to  $F^{(1)}$  with the help of estimates of the previous subsections. We recall the different steps. We define (where  $K_{(w)} > 0$  is defined in (8) and  $C_0 > 0$  is the constant of Lemma 2)

$$\mathcal{X}_w := \mathbf{Q}_w \times \mathfrak{C}_w, \quad \text{with } \|(Q, \rho)\|_{\mathcal{X}_w} := K_{(w)} C_0 (\|Q\|_{\mathbf{Q}} + \|\rho\|_{\mathfrak{C}_w}). \quad (82)$$

Thanks to the previous estimates we can say that the function  $F^{(1)}$  is well defined in a ball  $B_{\mathcal{X}_g}(0, \bar{R})$  with  $\bar{R} = \mathcal{O}(\sqrt{\log(\Lambda)})$ , say  $\bar{R} = K_0 \sqrt{\log(\Lambda)}$ . Indeed:

$$\|F^{(1)}(Q', \rho'')\|_{\mathcal{X}_w} \leq \|(N, n'')\|_{\mathcal{X}_w} + \alpha \kappa_1(\Lambda) \|(Q', \rho'')\|_{\mathcal{X}_w} + \sum_{\ell=2}^{+\infty} \alpha^\ell \kappa_\ell \|(Q', \rho'')\|_{\mathcal{X}_w}^\ell, \quad (83)$$

where

$$\begin{cases} \kappa_1(\Lambda) &= \mathcal{O}_{\Lambda \rightarrow +\infty}(\sqrt{\log(\Lambda)}) \\ \kappa_\ell &= \mathcal{O}_{\ell \rightarrow +\infty}(\ell^{1/2}). \end{cases} \quad (84)$$

In particular the radius of convergence of the power series  $f(x) = \sum_{\ell=2}^{+\infty} \kappa_\ell x^\ell$  is 1 and:

$$\|dF^{(1)}(Q', \rho'')\|_{L(\mathcal{X}_g)} \leq \alpha \kappa_1(\Lambda) + \alpha f'(\alpha \| (Q', \rho'') \|_{\mathcal{X}_w}). \quad (85)$$

For  $\|(N, n'')\|_{\mathcal{X}_w} \neq (0, 0)$  it is clear that  $F^{(1)}(0, 0) = (N, \mathcal{F}^{-1}(-\frac{1}{1+\alpha B_\Lambda(\cdot)} \hat{n}'')) \neq 0$ . So

$$\sup_{(Q', \rho'') \in B_{\mathcal{X}_g}(0, \bar{R})} \|dF^{(1)}(Q', \rho'')\|_{L(\mathcal{X}_g)} \leq \alpha \kappa_1(\Lambda) + \alpha f'(\alpha \bar{R}) =: \nu(\bar{R}). \quad (86)$$

For  $(Q', \rho'') \in B_{\mathcal{X}_g}(0, \bar{R})$  we have

$$\begin{aligned} \|F^{(1)}(Q', \rho'')\|_{\mathcal{X}_w} &\leq \|F^{(1)}(Q', \rho'') - F^{(1)}(0, 0)\|_{\mathcal{X}_w} + \|F^{(1)}(0, 0)\|_{\mathcal{X}_w} \\ &\leq \nu(\bar{R}) \|(Q', \rho'')\|_{\mathcal{X}_w} + \|F^{(1)}(0, 0)\|_{\mathcal{X}_w}. \end{aligned}$$

Thus  $B_{\mathcal{X}_g}(0, \bar{R})$  is invariant under  $F^{(1)}$  provided that:

$$\|F^{(1)}(0, 0)\|_{\mathcal{X}_w} \leq (1 - \nu(\bar{R}))\bar{R}. \quad (87)$$

As  $F^{(1)}(0, 0) \neq 0$  this gives  $\nu(\bar{R}) < 1$ .

Let us say that  $\|(N, n'')\|_{\mathcal{X}_w} = \varepsilon_0 \bar{R} = \varepsilon_0 K_0 \sqrt{\log(\Lambda)}$ ,  $\varepsilon_0 < 1$ . We have:

$$\|F^{(1)}(0, 0)\|_{\mathcal{X}_w} \leq \varepsilon_0 \bar{R}, \quad (88)$$

it suffices to take  $\alpha > 0$  such that  $\sqrt{L\alpha}K_0 \ll 1$  and then take  $\bar{R}$  accordingly. The constant  $K_0$  depends on the constants in the conditions  $M, \|\nu\|_{\mathcal{C}} \lesssim \sqrt{\log(\Lambda)}$ : we get  $\bar{R} = K_0 \sqrt{\log(\Lambda)}$  and for sufficiently small  $\alpha$  the Theorem can be applied on that ball.

### 5.5.2 $F^{(2)}$

We work with  $(\mathcal{C}, \|\cdot\|_{\mathcal{C}})$  and  $(\mathcal{C} \cap L^1, \max(\|\cdot\|_{\mathcal{C}}, \|\cdot\|_{L^1}))$ . In Appendix C it is proved that  $\|f_\Lambda\|_{L^1} \leq K\alpha B_\Lambda(0)$  where we can choose  $K = 2$  for  $\alpha \log(\Lambda)$  sufficiently small. Thus:

$$\mathcal{F}^{-1}(F_\Lambda) = \mathcal{F}^{-1}\left\{\frac{f_\Lambda}{1+f_\Lambda}\right\} = \sum_{\ell=1}^{+\infty} (-1)^{\ell+1} \check{f}_\Lambda^{*\ell} \in L^1$$

and its  $L^1$ -norm is lesser than  $\frac{2\alpha B_\Lambda(0)}{1-2\alpha B_\Lambda(0)} \leq 4\alpha B_\Lambda(0)$  as soon as  $\alpha B_\Lambda(0) \leq 4^{-1}$ . Moreover we can write

$$\frac{1}{1+f_\Lambda} = 1 - \frac{f_\Lambda}{1+f_\Lambda};$$

therefore if  $\rho \in L^1$  then  $\mathcal{F}^{-1}\{\frac{1}{1+f_\Lambda} \hat{\rho}\}^{-1} \in L^1$  and its  $L^1$ -norm is lesser than

$$(1 + 4\alpha B_\Lambda(0))\|\rho\|_{L^1} \leq 2\|\rho\|_{L^1}.$$

In particular:

$$\|\mathcal{F}^{-1}(\frac{1}{1+f_\Lambda} \hat{n}'')\|_{L^1} \leq 2(M + Z).$$

So we have:

$$\begin{cases} \|F^{(2)}(\rho'')\|_{\mathcal{C} \cap L^1} &\leq 2(M + Z) + \|h_2 + h_3\|_{\mathcal{C} \cap L^1} + K\alpha^3(\sqrt{\log(\Lambda)} + \|\rho''\|_{\mathcal{C} \cap L^1})\|\rho''\|_{\mathcal{C} \cap L^1}^2 \\ \|dF^{(2)}(\rho'')\|_{L(\mathcal{C} \cap L^1)} &\leq K\alpha^3\|\rho''\|_{\mathcal{C} \cap L^1}(2\sqrt{\log(\Lambda)} + 3\|\rho''\|_{\mathcal{C} \cap L^1}). \end{cases} \quad (89)$$

where the constants  $K$  can be chosen independently of  $\alpha \leq \alpha_0$  and  $\alpha \log(\Lambda) \leq L_0$  for  $\alpha_0, L_0$  sufficiently small. The term  $\sqrt{\log(\Lambda)}$  is due to  $\|\gamma'\|_{\mathcal{T}} \lesssim \sqrt{\log(\Lambda)}$  (see Lemma 5 and the regime of Remark 10). We get similar estimates for  $F^{(2)}$  defined in  $\mathcal{C}$ . So it suffices to take  $\bar{R} > 2$  sufficiently large so that  $B_{\mathcal{C} \cap L^1}(0, \bar{R})$  is invariant under  $F^{(2)}$ . This function is a contraction and we can apply the fixed point theorem. To end the proof we remark:

- There is only one fixed point of  $F^{(2)}$  in  $B_C(0, \overline{R})$  by the Banach-Picard Theorem and  $\rho_\gamma + n - \nu$  is a fixed point. Indeed by Section 3.2,  $(\gamma + N, \rho_\gamma + n - \nu)$  has norm  $\mathbf{Q}_1 \times \mathcal{C}$  bounded by  $K\sqrt{\log(\Lambda)}$  in the regime of Remark 10 and is a fixed point of  $F^{(1)}$ . So it is a fixed point of  $F^{(2)}$ .
- There is only one fixed point of  $F^{(2)}$  in  $B_{C \cap L^1}(0, \overline{R})$  by the same theorem. In particular it is also a fixed point of  $F^{(2)}$  in  $B_C(0, \overline{R})$  as  $B_{C \cap L^1}(0, \overline{R}) \subset B_C(0, \overline{R})$ . By unicity  $\rho_\gamma \in L^1$ .

## 6 Proofs of Theorems 2 and 3

### 6.1 Proof of Theorem 2

**Proof:** The fact that  $\rho_\gamma \in L^1$  is a result of Section 5.5. We recall that if  $Q \in \mathfrak{S}_1$ , then  $\int \rho_Q = \text{Tr}(Q) = \text{Tr}_{P_-^0}(Q)$ . Writing

$$\begin{aligned} A &:= \alpha \mathbf{T}[Q_{0,1}(\rho_\gamma'')] & C &:= \alpha^3 \mathbf{T}\{Q_{1,1}[\mathbf{T}[Q_{0,1}(\rho_\gamma'')], \rho_\gamma''\} \\ B &:= \alpha^2 \mathbf{T}(Q_{0,2}(\rho_\gamma'')) & S &:= \gamma - (A + B + C) \end{aligned} \quad (90)$$

it has been shown in Section 5 that  $S \in \mathfrak{S}_1$ . Theorem 1 says  $\rho_B = \rho_C = 0$ .

Let us show that  $B^{++}, B^{--}, C^{++}, C^{--}$  are trace-class. First for any  $Q$  in  $\mathfrak{S}_2$ , we have

$$\mathcal{P}_-^0 Q_{1,0}(Q) \mathcal{P}_-^0 = \mathcal{P}_+^0 Q_{1,0}(Q) \mathcal{P}_+^0 = 0.$$

It follows that  $B^{\pm\pm} = \alpha^2 Q_{0,2}(\rho_\gamma'')^{\pm\pm}$  and  $C^{\pm\pm} = \alpha^3 Q_{1,1}(\mathbf{T}Q_{0,1}(\rho_\gamma''), \rho_\gamma'')^{\pm\pm}$ . And as

$$Q_{0,2}^{+++} = Q_{0,2}^{---} = Q_{1,1}^{+++} = Q_{1,1}^{---} = 0$$

there only remain  $Q_{0,2}^{+-+}, Q_{0,2}^{-+-}, Q_{1,1}^{+--}, Q_{1,1}^{-+-}$ . Using Lemma 10 with  $a = \frac{3}{4}$  and Cauchy-Schwartz inequality we have

$$\left\| \frac{1}{|D_0|^{3/8}} \mathcal{P}_\pm^0 v_\rho \mathcal{P}_\mp^0 \frac{1}{|D_0|^{3/8}} \right\|_{\mathfrak{S}_2} \lesssim \|\rho\|_C \lesssim \|\rho\|_C \quad (91)$$

We recall that  $\left\| \frac{1}{|\nabla|^{1/2}} R_Q \right\|_{\mathfrak{S}_2} \lesssim \|Q\|_{\text{Ex}}$ : these two estimates enables us to prove the following:

$$\begin{aligned} \|Q_{0,2}^{\pm\mp\pm}(\rho_\gamma'')\|_{\mathfrak{S}_1} &\lesssim K_{3/2} \|\rho_\gamma''\|_C^2, \\ \|Q_{1,1}^{\pm\mp\pm}(\gamma', \rho_\gamma'')\|_{\mathfrak{S}_1} &\lesssim K_{7/4} \|\gamma'\|_{\text{Ex}} \|\rho_\gamma''\|_C. \end{aligned}$$

As shown in Sections 5 and C we have  $Q_{0,1}^{++} = Q_{0,1}^{--} = 0$  and  $\rho_A = -\check{f}_\Lambda * (\rho_\gamma') \in L^1$ .

$$\begin{aligned} \int \rho_\gamma &= \int (\rho_{\gamma^{++}} + \rho_{\gamma^{--}}) + \int \{\rho_{A^{+-}} + \rho_{A^{-+}} + \rho_{B^{+-}} + \rho_{B^{-+}} + \rho_{C^{+-}} + \rho_{C^{-+}}\} \\ &= \text{Tr}_{P_-^0}(\gamma) - \alpha f_\Lambda(0) \int \{\rho_\gamma + n - \nu\} - \int \{\rho_{B^{++}} + \rho_{B^{--}} + \rho_{C^{--}} + \rho_{C^{++}}\} \\ &= 0 - \alpha f_\Lambda(0) \left\{ \int \rho_\gamma + M - Z \right\} - \text{Tr}_{P_-^0}(B) - \text{Tr}_{P_-^0}(C). \end{aligned}$$

To end the proof we have to show that  $\text{Tr}(B^{++} + B^{--}) = \text{Tr}(C^{++} + C^{--}) = 0$ : this is straightforward when written in Fourier space (see [8] for formulae).  $\square$

### 6.2 Proof of Theorem 3

We follow the method of [10]. We apply a Lemma of Borwein and Preiss [10, Theorem 4] and consider an approximate minimizer  $\gamma'_0 = \gamma_0 + N_0$  of  $E^\nu(M)$ .

Indeed, we can extend  $\mathcal{E}_{BDF}^\nu$  to  $\mathfrak{K} = \cap \{Q \in \mathfrak{S}_2 : Q^* = Q, 0 \leq Q + P_-^0 \leq 1\}$  by setting  $\mathcal{E}_{BDF}^\nu(Q) := +\infty$  whenever  $Q \notin \mathfrak{K}$ . This extension is lower semi-continuous and bounded from below in the  $\mathfrak{S}_2$ -topology and the set

$$\mathcal{M} := \{Q \in \mathfrak{K}, (Q + P_-^0)^2 = Q + P_-^0, \text{Tr}_0(Q) = M\}$$

is closed in the same topology. Its convex closure in  $\mathfrak{S}_2$  is

$$\mathfrak{K}(M) := \{Q \in \mathfrak{K}, \text{Tr}_0(Q) = M\}.$$

Applying the lemma, for each  $\varepsilon > 0$  there exists a projector  $P$  and  $A \in \mathfrak{K}(M)$  such that  $\gamma'_0 := P - P_-^0$  minimizes the functional  $\mathcal{E}_{\text{BDF}}^\nu + \varepsilon \text{Tr}((A - \cdot)^2)$  on  $\mathcal{M}$  and

$$\mathcal{E}_{\text{BDF}}^\nu(\gamma'_0) \leq E_{\text{BDF}}^\nu(M) + \varepsilon^2, \quad \|\gamma'_0 - A\|_{\mathfrak{S}_2} \leq \sqrt{\varepsilon}.$$

As in [10],  $\gamma'_0$  satisfies the self-consistent equation

$$\begin{aligned} \gamma'_0 + P_-^0 &= \chi_{(-\infty, \mu_0]}(D_{\gamma'_0} + 2\varepsilon(\text{sgn}(D_0) - A)) \\ &= \chi_{(-\infty, \mu_0]}(\tilde{D} + \alpha B_{\gamma'_0} - 2\varepsilon A) \end{aligned} \quad (92)$$

where  $\mu_0 \in \mathbb{R}$  and  $\tilde{D} := \mathcal{D}^0 + D_0 \frac{2\varepsilon}{|D_0|}$ . We choose  $\varepsilon = \lambda^{-1}$  small *e.g.*  $\varepsilon = \Gamma(\frac{\Delta}{\alpha})^{-1}$ . Using the proof of Lemma 5 we show that the following *a priori* estimate holds for  $\gamma'_0$ :

$$\text{Tr}(|\nabla|(\gamma'_0)^2) + \alpha \|\rho''_{\gamma'_0}\|_{\mathcal{C}}^2 \lesssim \alpha \|\nu\|_{\mathcal{C}}^2 + \sqrt{\alpha} M + \sqrt{\alpha M} \|\nu\|_{\mathcal{C}}.$$

Using the Cauchy expansion, we can write

$$\gamma_0 = \sum_{j=0}^{+\infty} \alpha^j \mathcal{O}_j(\rho''_{\gamma'_0}, \gamma'_0) + \frac{2}{\lambda} W_\lambda(A, \alpha B(\gamma'_0)),$$

where the  $\mathcal{O}_j$ 's are defined as the  $Q_j$ 's with  $\tilde{D}$  replacing  $\mathcal{D}^0$  (see (20)). By the same method as in Section 5 we have:

$$\|\mathcal{D}^0\|^{1/2} W_\lambda\|_{\mathfrak{S}_2} + \|\rho[W_\lambda]\|_{\mathcal{C}} \lesssim \|A\|_{\mathfrak{S}_2} (1 + \alpha[\|\rho''_{\gamma'_0}\|_{\mathcal{C}} + \|\nabla\|^{1/2} \gamma'_0\|_{\mathfrak{S}_2}]).$$

Indeed it suffices to replace one  $R[\gamma'_0]$  in the  $\mathcal{O}_j$ 's by  $A$  and remark that  $A \in \mathfrak{S}_2$ . Replacing  $\mathcal{D}^0$  by  $\tilde{D}$  is harmless; as before, by defining some function  $\tilde{F}^{(1)}$  we can show that  $\text{Tr}_0(\gamma_0) = 0$  (but with an alternative  $B_\Lambda$  cf Section C).

In particular we can write

$$\rho_{\gamma_0} := -\mathcal{F}^{-1}(\tilde{F}_\Lambda) * n''_0 + (\delta_0 - \mathcal{F}^{-1}(\tilde{F}_\Lambda)) * \tau_{\text{rem}} \in \mathcal{C}$$

where  $\|\tau_{\text{rem}}\|_{\mathcal{C}} \lesssim \|\mathfrak{t}[N_0]\|_{\mathcal{C}} + \alpha^2 \|\tilde{\tau}_2\|_{\mathcal{C}} + \|A\|_{\mathfrak{S}_2}/\lambda$  and  $\tilde{F}_\Lambda$  is defined in Section C. We write  $\mathfrak{f}_\Lambda := \mathcal{F}^{-1}(\tilde{F}_\Lambda)$  for short. As in Section 5 we get:

$$\begin{cases} \|\gamma_0\|_{\mathfrak{S}_2} \lesssim \alpha(\|\rho''_{\gamma'_0}\|_{\mathcal{C}} + \|\gamma'_0\|_{\mathfrak{T}}) \\ \|\rho_{\gamma_0} + \mathfrak{f}_\Lambda * n''_0 - (\delta_0 - \mathfrak{f}_\Lambda) * \mathfrak{t}[N_0]\|_{\mathcal{C}} \lesssim \alpha^2(\|\gamma'_0\|_{\mathfrak{T}} + \|\rho''_{\gamma'_0}\|_{\mathcal{C}})^2. \\ \|\mathfrak{f}_\Lambda * n''_0 + (\delta_0 - \mathfrak{f}_\Lambda) * \mathfrak{t}[N_0]\|_{L^1} \lesssim L(Z + M). \end{cases} \quad (93)$$

Let  $(\psi_j)_{1 \leq j \leq M}$  be an orthonormal family of eigenvectors of  $\tilde{D} + \alpha B_{\gamma'_0} + 2/\varepsilon(1 - P_-^0 - A)$  spanning  $\text{Ran}(N_0)$  (with eigenvalues  $(\mu_j)$ ).

We then scale  $\gamma'_0$  by  $\alpha^{-1}$  (this procedure is emphasized by an underline) as in [10] we get:

$$\left[ \left( \frac{g_0(-i\alpha\nabla)\beta}{\alpha^2} - \frac{ig_1(-i\alpha\nabla)}{\alpha^2} \alpha \cdot \nabla \right) + \rho[\underline{\gamma}'_0] * \frac{1}{|\cdot|} - R[\underline{\gamma}'_0] + \frac{2}{\alpha^2 \lambda} \left( \frac{1}{2} - \underline{P}_-^0 - \underline{A} \right) \right] \underline{\psi}_j = \frac{\mu_j}{\alpha^2} \underline{\psi}_j. \quad (94)$$

*Remark 19.* We have  $U_\alpha \underline{\psi}(x) = \alpha^{\frac{3}{2}} \underline{\psi}(\alpha x) = \psi(x)$  and for an operator  $S$  we define:

$$\underline{S} := U_\alpha^* S U_\alpha.$$



This mean-field operator  $H_{\alpha^{-1}}$  is decomposed as follows:  $H_{\alpha^{-1}} = H_{\alpha^{-1}}^{(1)} + h_{\text{rem}}$  where

$$H_{\alpha^{-1}}^{(1)} := \frac{\mathcal{D}^0}{\alpha^2} + (\delta_0 - \underline{f}_\Lambda) * \underline{n}_0'' - R[N_0], \quad \underline{n}_0''(x) = \alpha^{-3} n''(x/\alpha), \quad \tilde{F}_\Lambda(k) = \tilde{F}_\Lambda(\alpha k).$$

As in the Lemma 13 and 14 of [10] we can show that there exists  $\varepsilon > 0$  such that  $\limsup_{\alpha \rightarrow 0} (\alpha^{-2}(\mu_j - 1)) < -\varepsilon < 0$  for all  $1 \leq j \leq M$  and that  $(\underline{\psi}_j)_j$  is bounded in  $H^1(\mathbb{R}^3, \mathbb{C}^4)^M$  (as  $\alpha$  tends to 0). Lemma 13 is based on a min-max description of eigenvalues in the gap of the mean-field operator  $H_{\alpha^{-1}}$ . We refer to this paper for the proofs. The only difference lies in the presence of  $-\underline{f}_\Lambda * (\underline{n}_0'') * \frac{1}{|\cdot|}$  and  $(\delta_0 - \underline{f}_\Lambda) * \underline{t}_{N_0}$ : we deal with these terms in the following lemma, proved below.

**Lemma 12.** *Let  $\chi$  be a Schwartz function and for  $R > 0$ :  $\chi_R(x) := R^{-3/2} \chi(x/R)$ . Then there holds:*

$$\begin{aligned} \left| \langle \underline{f}_\Lambda * (\underline{n}_0'') * \frac{1}{|\cdot|} \chi_R - Z \tilde{F}_\Lambda(0) \frac{\chi_R}{|\cdot|}, \chi_R \rangle \right| &\lesssim \frac{ZL}{R^2} \|\nabla \chi\|_{L^2}^2 \\ &+ \frac{1}{R} \|\nabla \chi\|_{L^2} \|\chi\|_{L^2} \left( L \int_{|y| > \frac{1}{\alpha}} \nu(y) dy + Z \int_{|y| > \frac{1}{\alpha}} |\underline{f}_\Lambda(y)| dy \right), \end{aligned}$$

and  $\int_{|y| > \frac{1}{\alpha}} |\underline{f}_\Lambda(y)| dy \leq L\alpha^{1/2}$ . Moreover for  $r_0 > 0$  we have

$$\begin{aligned} R \left| \langle (\delta_0 - \underline{f}_\Lambda) * \underline{t}_{N_0} * \frac{1}{|\cdot|} \chi_R, \chi_R \rangle \right| &\lesssim \frac{\alpha \|\underline{t}_{N_0}\|_{L^1}}{R} \int \frac{|\mathcal{F}(|\chi|^2; k)|}{|k|} dk + \int_{|y| \geq r_0} |\underline{t}_{N_0}(y)| dy \int \frac{|\chi(x)|^2}{|x|} dx \\ &+ \|\underline{t}_{N_0}\|_{L^1} \int_{|x| \leq \frac{r_0}{R}} \frac{|\chi(x)|^2}{|x|} dx. \end{aligned}$$

*Remark 20.* This is because of the last term that the bound on  $L$  depends on  $M$ . If we could prove that  $\int_{|x| \geq r_0} |\underline{t}_{N_0}(y)| dy$  tends to 0 as  $r_0 \rightarrow +\infty$  uniformly in  $\varepsilon$  (the parameter of Borwein and Preiss's Lemma), then we could take  $L \leq L_0$  instead of  $L \leq 1/(K_0 M)$  in Theorem 3.

To prove  $(\underline{\psi}_j)_j$  is  $H^1$ -bounded we show that:

$$\frac{M}{\alpha^4} + \frac{\text{Tr}(-\Delta N_0)}{\alpha^2} \leq \frac{\text{Tr}(\mathcal{D}^{02} N_0)}{\alpha^4} \leq \frac{M}{\alpha^4} + K(M, \nu) \left\{ \text{Tr}(-\Delta N_0) + \frac{\|\nabla N_0\|_{\mathfrak{S}_2}}{\alpha^2} \right\}. \quad (95)$$

The lower bound is clear and the upper bound follows from Eq. (94), Lemma 5 and Proposition 5 (for estimations of  $g_*(\alpha p)^2$ ,  $\star \in \{0, 1\}$ ). We get:

$$\begin{aligned} \left| \left| \rho[\underline{\gamma}_0] + \rho[\underline{f}_\Lambda * \underline{n}_0'' - (\delta_0 - \underline{f}_\Lambda) * \underline{t}_{N_0}] \underline{\psi}_j \right| \right|_{L^2} &\lesssim \alpha^{3/2} (\|\rho'_{\gamma_0}\|_c + \|\gamma'_0\|_{\mathbb{T}})^2 \|\nabla|^{1/2} \underline{\psi}_j\|_{L^2} \\ &\lesssim K(M, \nu) \|\nabla|^{1/2} \underline{\psi}_j\|_{L^2}. \end{aligned}$$

Moreover:

$$\begin{cases} \|R[\underline{\gamma}_0] \underline{\psi}_j\|_{L^2} \leq \|\gamma_0\|_{\mathfrak{S}_2} \|\nabla \underline{\psi}_j\|_{L^2} \lesssim \alpha^{3/4} K(M, \nu) \|\nabla \underline{\psi}_j\|_{L^2} \\ \|v[\underline{f}_\Lambda * \underline{n}_0'' - (\delta_0 - \underline{f}_\Lambda) * \underline{t}_{N_0}] \underline{\psi}_j\|_{L^2}^2 \leq 4 \|\nabla \underline{\psi}_j\|_{L^2}^2 \|\rho[\underline{f}_\Lambda * \underline{n}_0'' - (\delta_0 - \underline{f}_\Lambda) * \underline{t}_{N_0}]\|_{L^1}^2 \\ \leq L^2 (Z + M)^2 \|\nabla \underline{\psi}_j\|_{L^2}^2 \\ |\langle v[\underline{f}_\Lambda * \underline{n}_0'' - (\delta_0 - \underline{f}_\Lambda) * \underline{t}_{N_0}] \underline{\psi}_j, \underline{\psi}_j \rangle| \leq |D(\rho[\underline{f}_\Lambda * \underline{n}_0'' - (\delta_0 - \underline{f}_\Lambda) * \underline{t}_{N_0}], |\underline{\psi}_j|^2)| \\ \leq L(Z + M) \langle |\nabla| \underline{\psi}_j, \underline{\psi}_j \rangle. \end{cases} \quad (96)$$

Summing over  $1 \leq j \leq M$  the inequalities (96) we get (95) because

$$\sum_{j=1}^M \|\nabla \underline{\psi}_j\|_{L^2}^2, \text{Tr}(|\nabla| \overline{N_0}) \leq \sqrt{M} \sqrt{\text{Tr}(-\Delta N_0)}.$$

We conclude as in [10] (the proof uses [20]) provided that there hold binding inequalities for the non-relativistic limit: this is the result of Proposition 2 in Appendix B.

In particular there holds

$$\lim_{\alpha \rightarrow 0} \alpha^{-2} (E_{BDF}^\nu(M) - M + \frac{\alpha}{2} D(\tilde{F}_\Lambda * \nu, \nu)) = E_{nr}(M), \quad (97)$$

where  $E_{nr}$  is the non-relativistic energy cf Appendix B.

**Proof of Lemma 12** With  $f(x) = |\chi_R|^2 * \mathcal{F}^{-1}(\tilde{F}_\Lambda)$ , we first estimate  $|\iint f(x)\nu(y)(1/|x - \alpha y| - 1/|x|)dxdy|$ : it is lesser than

$$\iint |f(x)|\nu(y)|\alpha y| \frac{dxdy}{|x||x - \alpha y|}.$$

Splitting at level  $\alpha^{-1}$  for  $y$ , we use Hardy's and Kato's inequalities:

$$\left\{ \begin{array}{l} \int_{|y| \leq \frac{1}{\alpha}} \nu(y)dy \int \frac{|f(x)|dx}{|x||x - \alpha y|} \leq (4Z\|\tilde{F}_\Lambda\|_{L^1}) \frac{\|\nabla \chi\|_{L^2}^2}{R^2} \\ \int_{|y| > \frac{1}{\alpha}} \nu(y)dy \int |\alpha y| \frac{dx}{|x||x - \alpha y|} |f(x)| \leq \frac{2\pi}{2} \int_{|y| > \frac{1}{\alpha}} \nu(y)dy \|\tilde{F}_\Lambda\|_{L^1} \frac{\|\nabla \chi\|_{L^2} \|\chi\|_{L^2}}{R}. \end{array} \right.$$

We estimate  $Z|\iint |\chi_R(x)|^2 \tilde{f}_\Lambda(y)(1/|x - \alpha y| - 1/|x|)dxdy|$  analogously, with the help of Lemma 15. To treat the terms with  $\underline{t}_{N_0}$  we use the fact that:

$$\|\underline{t}_{N_0}\|_{L^1} = \|\underline{t}_{N_0}\|_{L^1} \lesssim LM \text{ and } \int \underline{t}_{N_0} = \int t_{N_0} = 0.$$

The first term in the upper bound corresponds to the error term that we get when we replace  $\mathcal{F}^{-1}(\tilde{F}_\Lambda) * \frac{1}{|\cdot|}$  by  $\tilde{F}_\Lambda(0)$ . To see this, we write  $a := \widehat{t_{N_0}}$  and  $b := \widehat{|\chi|^2}$ : we have

$$\begin{aligned} \int \frac{a^*(k)b(Rk)}{|k|^2} (\tilde{F}_\Lambda(\alpha k) - \tilde{F}_\Lambda(0)) &= \int \frac{a^*(\frac{k}{R})b(k)}{|k|^2} (\tilde{F}_\Lambda(\frac{\alpha k}{R}) - \tilde{F}_\Lambda(0)), \\ \left| \int \frac{a^*(k)b(Rk)}{|k|^2} (\tilde{F}_\Lambda(\alpha k) - \tilde{F}_\Lambda(0)) \right| &\lesssim \frac{\alpha \|\underline{t_{N_0}}\|_{L^1}}{R} \int \frac{|b(k)|}{|k|} dk. \end{aligned}$$

Let  $\varrho$  be in  $L^1$ . Thanks to Newton's Theorem (for radial functions) we have

$$\begin{aligned} R \times D(|\chi_R|^2, \varrho) &= \int \varrho(y) \left( \frac{R}{|y|} \int_{|x| \leq \frac{|y|}{R}} |\chi(x)|^2 dx + \int_{|x| \geq \frac{|y|}{R}} \frac{|\chi(x)|^2}{|x|} dx \right) \\ &= \int_{|y| > r_0} \varrho(y) \int_{|x| \leq \frac{|y|}{R}} |\chi(x)|^2 \left( \frac{R}{|y|} - \frac{1}{|x|} \right) dx + \int_{|y| \leq r_0} \varrho(y) \left( \int_{|x| \leq \frac{|y|}{R}} |\chi(x)|^2 \left( \frac{R}{|y|} - \frac{1}{|x|} \right) dx \right) \\ &\lesssim \|\nabla^{1/2} \chi\|_{L^2} \int_{|y| > r_0} |\varrho(y)| dy + \|\varrho\|_{L^1} \int_{|x| \leq \frac{r_0}{R}} \frac{|\chi(x)|^2}{|x|}. \end{aligned}$$

□

## A Estimates and inequalities

*Notation 21.* In Section A and C,  $\mathbf{e}$  refers to any unitary vector in  $\mathbb{R}^3$  and for  $p \in \mathbb{R}^3$ , we write  $\omega_p := \frac{p}{|p|}$ .

We recall that  $\mathbf{s}_p = \mathcal{F}(\text{sign}(\mathcal{D}^0); p)$ . There exists  $C_s > 0$  such that:

$$\left| \begin{array}{l} \text{Id} - \mathbf{s}_p \mathbf{s}_q = \mathbf{s}_p (\mathbf{s}_p - \mathbf{s}_q) = (\mathbf{s}_p - \mathbf{s}_q) \mathbf{s}_q \\ |\text{Id} - \mathbf{s}_p \mathbf{s}_q| \leq |\mathbf{s}_p - \mathbf{s}_q| = \left| \widehat{\mathcal{P}}_-^0(p) - \widehat{\mathcal{P}}_-^0(q) \right| \leq C_s \frac{|p-q|}{\max(\widehat{E}(p), \widehat{E}(q))}. \end{array} \right. \quad (98)$$

## A.1 Proof of Lemma 8

We have [16]  $\frac{1}{|\nabla|}(x-y) = \text{Cst}/|x-y|^2$ . By Cauchy-Schwartz inequality there holds:

$$\begin{aligned} \text{Tr}(R_Q^* |\nabla|^{-1} R_Q) &= \iiint \text{Tr}_{\mathbb{C}^4} \frac{{}^t \overline{Q}(x, y)}{|x-y|} \frac{\text{Cst}}{|y-z|^2 |z-x|} Q(z, x) dx dy dz \\ &\leq \iiint |Q(x, y)|^2 \frac{1}{|y-z|^2 |z-x|^2} dx dy dz \\ &\lesssim \iint \frac{|Q(x, y)|^2}{|x-y|} = \text{Tr}(R_Q^* Q). \end{aligned}$$

We write  $m(|p+q|)$  the multiplication in Fourier space by  $|p+q|$ : the operators  $R$ . and  $\frac{1}{|\nabla|^{1/2}}$  commute with the multiplication in Fourier space by  $w(p-q)$  (written  $m(w)$ ). By Kato's inequality we have

$$\|m(w) \cdot \frac{1}{|\nabla|^{1/2}} R[Q]\|_{\mathfrak{S}_2} = \|\frac{1}{|\nabla|^{1/2}} R[m(w) \cdot Q]\|_{\mathfrak{S}_2} \lesssim \|m(|p+q|)m(w) \cdot Q\|_{\mathfrak{S}_2}.$$

Similarly for  $a > 0$  the operator  $|D_0|^{-a}$  is a convolution operator associated to a *positive* function  $\phi_a$ . Indeed there holds [16]:  $\frac{1}{\omega^{2-\Delta}}(x-y) = \frac{e^{-\omega|x-y|}}{4\pi|x-y|}$ ,  $\omega \geq 0$  and for any  $0 < \varepsilon < 1$  (see [17, footnote p. 87]):

$$\frac{1}{|D_0|^{2\varepsilon}} = \frac{\sin(\varepsilon\pi)}{\pi} \int_0^{+\infty} t^{\varepsilon-1} \frac{1-\Delta}{t+1-\Delta} dt.$$

Thus for  $a = 1 + \varepsilon > 1$  we have by Cauchy-Schwarz inequality:

$$\begin{aligned} \text{Tr}(R_Q^* \frac{1}{|D_0|^{2a}} R_Q) &\leq \iint |Q(x, y)|^2 \frac{1}{|x-y|} * \phi_{2a}(x-y) dx dy, \\ &\leq \iint |Q(x, y)|^2 \|\frac{1}{|\cdot|} * \phi_{2a}\|_{L^\infty} \\ &\leq \iint |Q(x, y)|^2 dx dy \int \frac{dp}{|p|E(p)^{2a}} \leq \frac{\|Q\|_{\mathfrak{S}_2}^2}{2(a-1)}. \end{aligned}$$

Let us consider a finite rank operator  $Q(x, y)$ . As  $Q = \frac{Q+Q^*}{2} + \frac{Q-Q^*}{2}$  one may suppose it is self-adjoint, writing  $Q = Q_+ - Q_-$  one may suppose it is nonnegative: then so is  $R_Q$  and  $|D_0|^{-a/2} R_Q |D_0|^{-a/2}$ . We have

$$\begin{aligned} \int \frac{\text{Tr}_{\mathbb{C}^4}(\widehat{R}(p, p))}{E(p)^{2a}} dp &= \frac{1}{2\pi^2} \iint \frac{d\ell}{|\ell|^2} \text{Tr}(\widehat{Q}(p-\ell, p-\ell)) \frac{dp}{E(p)^{2a}} \\ &= \frac{1}{2\pi^2} \int dp \text{Tr}(\widehat{Q}(p, p)) \int \frac{d\ell}{|\ell|^2} \frac{1}{E(p+\ell)^{2a}} \\ &\lesssim \frac{\|\widehat{Q}\|_{\mathfrak{S}_1}}{2a-2}. \end{aligned}$$

In Fourier space we have:  $\mathcal{F}(|\mathcal{D}^0|^{-1/2}) : f(p) \mapsto \chi_{|p| < \Lambda} \frac{f(p)}{E(p)^{1/2}}$ . Thus writing  $\Pi_A$  the projection onto  $\{f \in L^2, \text{supp } \widehat{f} \subset B(0, A)\}$  we get

$$\| |\mathcal{D}^0|^{1/2} R_Q \|_{\mathfrak{S}_2} \leq \| |\mathcal{D}^0|^{1/2} \Pi_{2\Lambda} R_Q \Pi_{3\Lambda} \|_{\mathfrak{S}_2}.$$

As  $|\mathcal{D}^0|^{-1/2} \Pi_{2\Lambda} \leq e |D_0|^{-\frac{1}{2} - \frac{1}{2 \log(\Lambda)}}$  for  $\Lambda \geq e$  we finally have:

$$\text{Tr}(\Pi_{3\Lambda} R_Q^* \frac{\Pi_{2\Lambda}}{|D_0|} R_Q \Pi_{3\Lambda}) \leq \text{Tr}(R_Q^* \frac{e}{|D_0|^{\frac{1}{2} + \frac{1}{2 \log(\Lambda)}}} R_Q) \lesssim \log(\Lambda) \|Q\|_{\text{Ex}}^2.$$

## B The non relativistic limit

We fix the value  $F_\Lambda(0) = a$ . For any trace-class operator  $0 \leq \Gamma \leq 1$  with density  $\rho_\Gamma$  the non-relativistic energy is

$$\begin{aligned} \mathcal{E}_{nr}^Z(\Gamma) &:= \frac{1}{2}\text{Tr}(-\Delta\Gamma) - Z(1-a)\text{Tr}\left(\frac{1}{|\cdot|}\Gamma\right) \\ &\quad + \frac{1}{2}(D(\rho_\Gamma, \rho_\Gamma) - \text{Ex}[\Gamma]) - \frac{a}{2}D(\rho_\Gamma, \rho_\Gamma). \end{aligned} \quad (99)$$

If we drop the last term, this is exactly the Hartree-Fock energy  $\mathcal{E}_{\text{HF}}$  with a nucleus of charge  $Z_0 := Z(1-a)$  and if we drop  $\text{Tr}\left(\frac{1}{|\cdot|}\Gamma\right)$  we get the Pekar-Tomasevitch energy  $\mathcal{E}_{nr}^0 = \mathcal{E}_{\text{PT}}\left[\frac{a}{2}, U = \frac{1}{2}\right]$  (cf [4]).

*Remark 22.* We can easily show stability of matter of the second kind for  $a \leq a_0$  by splitting the energy in two: a Hartree-Fock one and a Pekar-Tomasevitch one,

$$\begin{aligned} \mathcal{E}_{nr}^Z(\Gamma) &= \frac{x^2}{2}\text{Tr}(-\Delta\Gamma) + \frac{y^2}{2}(D(\rho_\Gamma, \rho_\Gamma) - \text{Ex}[\Gamma]) - Z(1-a)\text{Tr}\left(\frac{1}{|\cdot|}\Gamma\right) \\ &\quad + \frac{1-x^2}{2}\text{Tr}(-\Delta\Gamma) + \frac{1-y^2}{2}(D(\rho_\Gamma, \rho_\Gamma) - \text{Ex}[\Gamma]) - \frac{a}{2}D(\rho_\Gamma, \rho_\Gamma) \text{ with } 0 < x, y < 1. \end{aligned}$$

Optimizing in  $x$  and  $y$  we get a lower bound  $\mathcal{O}(K(a)M)$  for  $M \geq 2Z_0 + 1$ .

We define

$$\mathcal{G}(x) = \{\Gamma \in \mathfrak{S}_1 : \Gamma^* = \Gamma, 0 \leq \Gamma \leq 1, \sqrt{-\Delta}\Gamma \in \mathfrak{S}_2 \text{ and } \text{Tr}(\Gamma) = x\} \text{ with } x \in \mathbb{R}_+^*.$$

$E_{nr}(M)$  corresponds to the infimum over  $\mathcal{G}(M)$ . We want to prove:

**Proposition 2.** *For any  $M < Z+1$ , the variational problem  $E_{nr}^Z(M)$  admits a minimizer.*

By Lieb's method in [15], it is easy to see that there is a minimizer for  $E_{nr}^Z(1)$ . To prove binding for  $2 \leq M \leq Z(1-a)$  we can follow Lieb's and Simon's method [19, 20]. We will however prove it with the method of concentration-compactness. We prove the problem  $E_{nr}^Z(M)$  admits a minimizer by induction over  $M$  by using:

**Proposition 3.** *For each  $\ell > 0$  the following assertions are equivalent*

- $\forall 0 < k < \ell : E_{nr}^Z(\ell) < E_{nr}^Z(\ell - k) + E_{nr}^0(k)$ .
- *Each minimizing sequence for  $E_{nr}^Z(\ell)$  is precompact in  $H^1(\mathbb{R}^3 \times \mathbb{R}^3)$ .*

*In the case  $\ell \in \mathbb{N}^*$ , it suffices to prove binding inequalities for  $K \in (0, \ell) \cap \mathbb{N}$ .*

This proposition is standard and we will not give the proof here but refer to [14, 13, 20]. In [4] Frank *et al.* prove that  $E_{nr}^0(M_0) = M_0 E_{nr}^0(1)$  for  $M_0 \in \mathbb{N}^*$  provided that  $a$  is sufficiently small. Thus we just have to show

$$E_{nr}^Z(M) < E_{nr}^Z(M-1) + E_{nr}^0(1).$$

To this end, we exhibit a test function  $Q$  whose energy is lesser than  $E_{nr}^Z(M-1) + E_{nr}^0(1)$ .

Lieb's variational principle still holds (cf [10, Proposition 3]). In fact for any orthonormal family  $(\phi_1, \phi_2)$ , with  $P_\phi := |\phi\rangle\langle\phi|$  and  $0 < t < 1$ , we have

$$\begin{aligned} \mathcal{E}_{nr}^Z(\Gamma + t(P_{\phi_1} - P_{\phi_2})) - \mathcal{E}_{nr}^Z(\Gamma) &= \frac{t}{2}(\|\nabla\phi_1\|_{L^2}^2 - \|\nabla\phi_2\|_{L^2}^2 + 2(1-a)D(\rho_\Gamma, |\phi_1|^2 - |\phi_2|^2)) \\ &\quad - t\Re[\text{Tr}(\Gamma R[P_{\phi_1} - P_{\phi_2}])] - t^2\{D(|\phi_1|^2, |\phi_2|^2) - D(\phi_1^*\phi_2, \phi_1^*\phi_2) + \frac{a}{2}\||\phi_1|^2 - |\phi_2|^2\|_{\mathcal{C}}^2\}. \end{aligned} \quad (100)$$

This shows that  $E_{nr}^Z(m)$  is also the infimum of  $\mathcal{E}_{nr}^Z$  over

$$\{\Gamma \in \mathcal{G}(m) : \Gamma = P + (m - [m])|\phi\rangle\langle\phi|, P^2 = P = P^*, \phi \in \text{Ker}(P)\}.$$

Taking  $\phi_2 = 0$  in (100) shows that  $E_{nr}^Z(\cdot)$  is concave in  $[M_0, M_0 + 1]$  with  $M_0 \in \mathbb{N}$ . It is also clear that  $E_{nr}^Z$  is decreasing since large binding inequalities hold.

We consider a minimizer of  $E_{nr}^Z(M-1)$  of the form  $\Gamma = \sum_{1 \leq j \leq M-1} |\psi_j\rangle\langle\psi_j|$ , each  $\psi_j$  satisfying

$$-\frac{\Delta}{2}\psi_j - \frac{Z_0}{|\cdot|}\psi_j + (1-a)\rho[\Gamma] * \frac{1}{|\cdot|}\psi_j - R[\Gamma]\psi_j + \varepsilon_j\psi_j = 0, \text{ with } \varepsilon_j > 0.$$

In particular we can easily show the  $\psi_j$ 's are in  $H^2(\mathbb{R}^3)$  and fast decaying.

We also consider a minimizer for  $E_{nr}^0(1)$ : this is a minimizer  $\phi_{CP}$  of  $E_{PT}(1)$  scaled by  $a$ :  $\phi_0(x) = a^{3/2}\phi_{CP}(ax)$ , we chose it to be radial [15]. Following [13], we take a Schwartz function  $0 \leq \chi \leq 1$  that satisfies  $\chi(x) = 1$  for  $|x| \leq 1$  and  $\chi(x) = 0$  for  $|x| \geq 2$  and  $\chi_R(x) = \chi(x/R)$  with  $R > 0$  to be chosen.

We define the trial state as follows: for some  $\mathbf{e} \in \mathbb{S}^2$  we write

$$\Gamma' := \chi_R \Gamma \chi_R + \tau_{-5R\mathbf{e}} |\chi_R \phi_0\rangle \langle \chi_R \phi_0 | \tau_{5R\mathbf{e}}$$

where  $\tau_{x_0} \psi(x) := \psi(x - x_0)$ . We have  $0 \leq \Gamma' \leq 1$  and  $\text{Tr}(\Gamma') \leq M$ , so  $\mathcal{E}_{nr}(\Gamma') \geq E_{nr}(M)$ . As the wave functions  $(\psi_j)$ 's and  $\phi_0$  are fast decaying, the following holds:

$$\begin{aligned} \mathcal{E}_{nr}^Z(\Gamma') &= \mathcal{E}_{nr}^Z(\Gamma) + \mathcal{E}_{nr}^0(\phi_0) + \int (\rho[\Gamma] * \frac{1}{|\cdot|}(x) - \frac{Z_0}{|x|}) |\tau_{5R\mathbf{e}} \phi_0(x)|^2 dx \\ &\quad - aD(\rho[\Gamma], |\tau_{5R\mathbf{e}} \phi_0|^2) + o(R^{-1}). \end{aligned}$$

As  $R$  tends to infinity we get:

$$\mathcal{E}_{nr}^Z(\Gamma') \leq E_{nr}^Z(M-1) + E_{nr}^0(1) + \frac{(M-1)(1-a) - Z_0}{5R} + o(R^{-1}) < E_{nr}(M-1) + E_{nr}^0(1).$$

## C Proof of Proposition 1

*Notation 23.* We write:

$$\begin{aligned} E(u, k/2) &:= \max(E(u+k/2), E(u-k/2)) \geq \sqrt{1 + |u|^2 + \frac{|k|^2}{4}}, \\ \tilde{E}(u, k/2) &:= \max(\tilde{E}(u+k/2), \tilde{E}(u-k/2)) \geq E(u, k/2). \end{aligned}$$

Our aim is to prove Proposition 4 below.

**Proposition 4.** *Let  $\rho_0 \in \mathcal{C}$ . Then we have:*

$$\alpha \rho(\mathbf{T}[Q_{0,1}(\rho_0)]) = -\check{f}_\Lambda * \rho_0$$

where  $\check{f}_\Lambda \in L^1$  is a radial function. Moreover

$$f_\Lambda = \sum_{J=0}^{+\infty} \alpha^J f_{\Lambda, J}, \quad f_{\Lambda, 0} = \alpha B_\Lambda \quad \text{and} \quad g_\Lambda := \sum_{J=1}^{+\infty} \alpha^J f_{\Lambda, J},$$

with

$$\|\check{f}_\Lambda\|_{L^1} \lesssim L \quad \text{and} \quad \|\check{g}_\Lambda\|_{L^1} \lesssim L\alpha.$$

In particular  $\check{F}_\Lambda := \mathcal{F}^{-1}\left(\frac{f_\Lambda}{1+f_\Lambda}\right) \in L^1$ .

We also study an alternative function  $F_\Lambda$ , needed for the proof of Theorem 3, at the end of this section.

We need the following proposition.

**Proposition 5.** *The function  $\widehat{\mathcal{D}}^0 : \overline{B(0, \Lambda)} \rightarrow \mathbb{R}^3$  is infinitely differentiable. In particular so is  $\tilde{E}(\cdot)$  and there exists  $L_0 \geq 0$  such that if  $L := \alpha \log(\Lambda) \leq L_0$  then for any  $J \geq 1$  there exists  $C_J > 0$  such that:*

$$\|d^J g_0\|_{L^\infty} \leq \alpha C_J \quad \text{and} \quad \|d^J \mathbf{g}_1\|_{L^\infty} \leq \chi_{J=1} + LC_J.$$

**Proof:** In the spirit of [23], we can prove it by induction over  $J$ : in [11] Hainzl *et al.* proved that  $\widehat{\mathcal{D}}^0$  is infinitely differentiable. Thus the function

$$|\widehat{\mathcal{D}}^0(p)| = \sqrt{g_0(p)^2 + \mathbf{g}_1(p) \cdot \mathbf{g}_1(p)},$$

is infinitely differentiable and does not vanish on  $\overline{B(0, \Lambda)}$ . Thanks to the self-consistent equation one has:

$$d^J \widehat{\mathcal{D}^0}(p) = d^J \widehat{D}_0(p) + \frac{\alpha}{4\pi^2} \frac{1}{|\cdot|^2} * d^J \left( \frac{\mathcal{D}^0}{|\mathcal{D}^0|} \right)(p).$$

□

**Proof of Proposition 4:** Throughout this proof we write  $k := r\mathbf{e}$ .

1. Let us first prove the following:

$$\widehat{\tau}_{1,0}(\rho) = -f_\Lambda(\cdot)\widehat{\rho}(\cdot),$$

We recall that for any  $Q \in \mathfrak{S}_2(\mathfrak{H}_\Lambda)$  we have (41):

$$\widehat{Q}_{1,0}(Q, p, q) = \frac{1}{4\pi^2} \frac{1}{\widetilde{E}(p) + \widetilde{E}(q)} \int_{\ell} \frac{d\ell}{|\ell|^2} (\widehat{Q}(p - \ell, q - \ell) - \mathbf{s}_p \widehat{Q}(p - \ell, q - \ell) \mathbf{s}_q),$$

and (cf [8])

$$\widehat{Q}_{0,1}(\rho; p, q) = \frac{4\pi}{2^{5/2}\pi^{3/2}} \frac{\widehat{\rho}(p - q)}{|p - q|^2} \frac{1}{\widetilde{E}(p) + \widetilde{E}(q)} (\mathbf{s}_p \mathbf{s}_q - 1). \quad (101)$$

The functions  $A_J^{(\ell_j)_{j=1}^J}$  are defined recursively in (52). We have for instance:

$$\begin{aligned} A_2^{(\ell_1, \ell_2)} \widehat{Q}(p, q) &= A_1^{\ell_1} (\widehat{Q}(\cdot_p - \ell_2, \cdot_q - \ell_2) - \mathbf{s}_p \widehat{Q}(\cdot_p - \ell_2, \cdot_q - \ell_2) \mathbf{s}_q)(p, q) \\ &= \left\{ \widehat{Q}(p - \ell_1 - \ell_2, q - \ell_1 - \ell_2) - \mathbf{s}_{p - \ell_1} \widehat{Q}(p - \ell_1 - \ell_2, q - \ell_1 - \ell_2) \mathbf{s}_{q - \ell_1} \right\} \\ &\quad - \mathbf{s}_p \left\{ \widehat{Q}(p - \ell_1 - \ell_2, q - \ell_1 - \ell_2) - \mathbf{s}_{p - \ell_1} \widehat{Q}(p - \ell_1 - \ell_2, q - \ell_1 - \ell_2) \mathbf{s}_{q - \ell_1} \right\} \mathbf{s}_q. \end{aligned}$$

Writing  $L_J := \sum_{j=1}^J \ell_j$  with  $L_0 := 0 \in \mathbb{R}^3$  we have:

$$\widehat{F_{1,0}^{\circ J}}(Q; p, q) = \frac{\alpha^J}{(4\pi^2)^J} \int_{\ell_1} \cdots \int_{\ell_J} \frac{d\ell}{\prod_{1 \leq j \leq J} |\ell_j|^2} \frac{A_J^{(\ell_j)_{j=1}^J} \widehat{Q}(p, q)}{\prod_{0 \leq j \leq J-1} (\widetilde{E}(p - L_j) + \widetilde{E}(q - L_j))}. \quad (102)$$

In particular the Fourier transform of the density  $\rho(F_{1,0}^{\circ J}(Q))$  is

$$\begin{aligned} \widehat{\rho}(F_{1,0}^{\circ J}(Q); k) &= \frac{1}{(2\pi)^{3/2}} \int_u \mathrm{Tr}_{\mathbb{C}^4} \widehat{F_{1,0}^{\circ J}}(Q; u + \frac{k}{2}, u - \frac{k}{2}) du \\ &= \frac{\alpha^J}{(2\pi)^{3/2} (4\pi^2)^J} \iint_{u, \ell_1} \cdots \int_{\ell_J} \mathrm{Tr}_{\mathbb{C}^4} \frac{dud\ell}{\prod_{1 \leq j \leq J} |\ell_j|^2} \frac{A_J^{(\ell_j)_{j=1}^J} \widehat{Q}(u + \frac{k}{2}, u - \frac{k}{2})}{\prod_{0 \leq j \leq J-1} (\widetilde{E}(u + \frac{k}{2} - L_j) + \widetilde{E}(u - \frac{k}{2} - L_j))} \\ &= \frac{\alpha^J}{(2\pi)^{3/2} (4\pi^2)^J} \iint_{u, \ell_1} \cdots \int_{\ell_J} \frac{dud\ell}{\prod_{1 \leq j \leq J} |\ell_j|^2} \frac{\mathrm{Tr}_{\mathbb{C}^4} \left\{ (1 - \mathbf{s}_{u - \frac{k}{2}} \mathbf{s}_{u + \frac{k}{2}}) A_J^{(\ell_j)_{j=2}^{J-1}} \widehat{Q}(u + \frac{k}{2}, u - \frac{k}{2}) \right\}}{\prod_{0 \leq j \leq J-1} (\widetilde{E}(u + \frac{k}{2} - L_j) + \widetilde{E}(u - \frac{k}{2} - L_j))}. \end{aligned} \quad (103)$$

Above the domain of  $\ell_j$  is:

$$\widetilde{B}_j(r) := \{\ell_j, |u - L_j \pm \frac{r}{2} \mathbf{e}| < \Lambda\},$$

and the domain of  $u$  is  $\widetilde{B}_0(r) := \{u, |u \pm \frac{r}{2} \mathbf{e}| < \Lambda\}$ . In particular

$$\mathrm{supp} \widehat{\rho}(F_{1,0}^{\circ J}(Q)) \subset B(0, 2\Lambda).$$

*Remark 24.* We would like to apply (103) to the operator  $Q_{0,1}(\rho)$ . From (101) we realize that  $\widehat{Q}_{0,1}(p, q)$  is not a scalar matrix because of the term  $\mathbf{s}_p \mathbf{s}_q - \mathrm{Id}$ . Yet it is in the algebra spanned by the Dirac matrices  $\alpha_1, \alpha_2, \alpha_3, \beta$  as a sum of *even* products of Dirac matrices. The form of  $\widehat{Q}_{1,0}(Q)$  is similar to  $\widehat{Q}_{0,1}$ : it only adds an *even* number of Dirac matrices to  $\widehat{Q}$ . This is an important remark to be done to prove Theorem 1.

For any  $J \geq 1$  and  $\rho \in \mathcal{C}$ , the density  $\widehat{\rho}(F_{1,0}^{\circ J}(Q_{0,1}[\rho]); k)$  is equal to:

$$\begin{aligned} & \frac{4\pi\alpha^J \widehat{\rho}(k)}{(2^5\pi^3)^{\frac{1}{2}}(2\pi)^{3/2}(4\pi^2)^J} \int_{\prod_{0 \leq j \leq J} \widetilde{B}_j(r)} \frac{dud\ell}{|k|^2 \prod_{1 \leq j \leq J} |\ell_j|^2} \frac{\text{Tr}_{\mathbb{C}^4} \left\{ (1 - \mathbf{s}_{u-\frac{k}{2}} \mathbf{s}_{u+\frac{k}{2}}) A_{J-1}^{(\ell_j)_{j=2}^J} (\mathbf{s}_{u-\frac{k}{2}} \mathbf{s}_{u+\frac{k}{2}} - 1) \right\}}{\prod_{0 \leq j \leq J} (\widetilde{E}(u + \frac{k}{2} - L_j) + \widetilde{E}(u - \frac{k}{2} - L_j))} \\ &= \widehat{\rho}(k) \int_{\prod_{0 \leq j \leq J} \widetilde{B}_j(r)} dud\ell S_J(u - L_j \pm \frac{k}{2}) T_J(u - L_j \pm \frac{k}{2}) \end{aligned} \quad (104)$$

where  $S_J(u - L_j \pm \frac{k}{2})$  is a scalar which is a function of  $|u - L_j \pm \frac{k}{2}|$  while  $T_J(u - L_j \pm \frac{k}{2})$  is the trace  $\text{Tr}_{\mathbb{C}^4}$  of a sum of products of  $\mathbf{s}_{u-L_j-\frac{k}{2}}$ .

We have to deal with  $\frac{1}{|k|^2}$  and we must show that this integral is well defined. The first problem is easy, the quantity

$$\frac{1}{|k|^2} (\mathbf{s}_{u-L_j+k/2} \mathbf{s}_{u-L_j-k/2} - 1) (1 - \mathbf{s}_{u-k/2} \mathbf{s}_{u+k/2}) = \frac{(\mathbf{s}_{u-L_j+k/2} \mathbf{s}_{u-L_j-k/2} - 1)}{|k|} \frac{(1 - \mathbf{s}_{u-k/2} \mathbf{s}_{u+k/2})}{|k|}$$

defines a smooth function by Taylor's formula (for  $|k|$  or for  $k$  in  $\mathbb{R}^3 \setminus \{0\}$ ). Moreover from (98), we get the estimates:

$$\begin{aligned} \left| \frac{\mathbf{s}_{u-L_j+k/2} \mathbf{s}_{u-L_j-k/2} - 1}{|k|} \frac{1 - \mathbf{s}_{u-k/2} \mathbf{s}_{u+k/2}}{|k|} \right| &\leq \frac{4C_s^2}{E(u - L_j, k/2)} \\ &\leq \frac{4C_s^2}{|u - L_j| E(u, k/2)}. \end{aligned}$$

For any  $U$ , we have:

$$\begin{aligned} \int_{\ell} \frac{d\ell}{|\ell|^2} \frac{1}{|U - \ell| \widetilde{E}(U - \ell, k/2)} &\leq \int_{\ell} \frac{d\ell}{|\ell|^2} \frac{1}{|U - \ell|^2}, \\ &\leq \frac{1}{|U|} \int \frac{d\ell}{|\ell|^2 |\mathbf{e} - \ell|^2}. \end{aligned}$$

Integrating over the  $\ell_{j+1}$ 's one after the other from  $\ell = J - 1$  down to  $j = 1$  as above with  $U = U_j = u - L_j$ , there remains but the integral over  $u$ :

$$\begin{aligned} \int_{u \in \widetilde{B}_0(k)} \frac{2C_s^2 du}{\widetilde{E}(u, k/2)} \frac{1}{|u| E(u, k/2)} &\times \left\{ 2 \int \frac{d\ell}{|\ell|^2 |\mathbf{e} - \ell|^2} \right\}^J \\ &\leq \left\{ 2 \int \frac{d\ell}{|\ell|^2 |\mathbf{e} - \ell|^2} \right\}^J \int_{u \in \widetilde{B}_0(r)} \frac{2C_s^2 du}{|u|^2 \widetilde{E}(u, k/2)} \\ &= (K \log(\Lambda)) \times (C'_{1,0})^J. \end{aligned}$$

At last we have:

$$\begin{aligned} \alpha |\widehat{\rho}(F_{1,0}^{\circ J}(Q_{0,1}(\rho)); k)| &\leq \frac{\alpha^{J+1}}{(2\pi)^{3/2} (4\pi^2)^J} 2^{J+1} C_s^2 \left\{ \int \frac{d\ell}{|\ell|^2 |\mathbf{e} - \ell|^2} \right\}^J \int_{u \in \widetilde{B}_0(r)} \frac{du}{|u|^2 E(u)} |\widehat{\rho}(k)| \\ &\lesssim C_{1,0} (\alpha C'_{1,0})^J \alpha \log(\Lambda) |\widehat{\rho}(k)|. \end{aligned} \quad (105)$$

As a consequence there holds  $\alpha \widehat{\rho}(F_{1,0}^{\circ J}(Q_{0,1}(\rho)); k) = -g_{\Lambda, J}(k) \widehat{\rho}(k)$ , and  $\sum_{J=0}^{\infty} f_{\Lambda, J}$  is well defined (at least in  $L^\infty \cap L^2$ ) as soon as  $\alpha$  is sufficiently small. We have

$$\alpha \widehat{\tau_{0,1}}(\rho, k) = - \left( \alpha B_\Lambda(k) + \sum_{J=1}^{+\infty} g_{\Lambda, J}(k) \right) \widehat{\rho}(k) =: -f_\Lambda(k) \widehat{\rho}(k), \quad (106)$$

with

$$|f_\Lambda(k)| \leq \alpha B_\Lambda(k) + \alpha^2 \log(\Lambda) K = \mathcal{O}(\alpha \log(\Lambda)). \quad (107)$$

2. Let us prove this function is radial. Let  $\mathbf{e}_1$  and  $\mathbf{e}_2$  in  $\mathbb{S}^2$  and  $r > 0$ . We must show that  $f_\Lambda(r\mathbf{e}_1) = f_\Lambda(r\mathbf{e}_2)$ . There exists  $\mathcal{R} \in \text{SO}_3(\mathbb{R})$  such that  $\mathbf{e}_2 = \mathcal{R}\mathbf{e}_1$ . In (104) for  $k = r\mathbf{e}_2$ , we change variables in the integrals:  $v = \mathcal{R}^{-1}u$  and  $m_j = \mathcal{R}^{-1}l_j$ . Writing  $M_j = m_1 + \dots + m_j$ , we get:  $S_J(\mathcal{R}(v - M_j \pm \frac{r}{2}\mathbf{e}_1)) = S_J(v - M_j \pm \frac{r}{2}\mathbf{e}_1)$ . We must show the same holds for  $T_J$ . Let  $\mathbf{b} = (b_1, b_2, b_3)$  be the canonical base of  $\mathbb{R}^3$ . We define

$$\alpha'_j := \boldsymbol{\alpha} \cdot \mathcal{R}b_j.$$

These new matrices satisfy the same relation as the  $\alpha$ 's:

$$\{\alpha'_j, \alpha'_k\} = 2\delta_{jk} \text{ and } \{\alpha'_j, \beta\} = 0.$$

Thus we have  $T_J(\mathcal{R}(v - M_j \pm \frac{r}{2}\mathbf{e}_1)) = T_J(v - M_j \pm \frac{r}{2}\mathbf{e}_1)$  and  $f_\Lambda$  is radial.

From now on we change variables:

$$\begin{cases} u_0 := u \text{ and for } 1 \leq j \leq J, u_j := u - L_j, l_j = u_j - u_{j-1}, \\ u_j \in B(|k|) := \{v \in B(0, \Lambda), |v \pm \frac{|k|}{2}\mathbf{e}| < \Lambda\}. \end{cases} \quad (108)$$

3. Our purpose is to show that  $f_\Lambda$  is in  $\mathcal{F}(L^1)$  with a (rather) precise bound on  $\|\check{f}_\Lambda\|_{L^1}$ . We already know:  $f_\Lambda(k) = \alpha B_\Lambda(k) + \mathcal{O}_{L^\infty}(\alpha^2 \log(\Lambda)) = \mathcal{O}(\alpha \log(\Lambda))$ .

As  $f_\Lambda$  is radial we take a fixed vector  $\mathbf{e} \in \mathbb{S}^2$  and study  $f_\Lambda(k) = f_\Lambda(|k|)$  with the help of the integral formulae where  $k$  is replaced by  $|k|\mathbf{e}$ .

The strategy is to differentiate  $f_\Lambda$  and prove that its Sobolev norms  $\|-\Delta f_\Lambda\|_{L^2}$  and  $\|-\Delta f_\Lambda\|_{L^p}$  are "small" where  $p < 2$  is some constant to be chosen later. By Cauchy-Schwartz inequality in Direct space, we obtain an upper bound of  $\|\check{f}_\Lambda\|_{L^1}$ . We will use the *co-area formula* [3].

We show that  $f_\Lambda \in L^1$  with  $L^1$ -norm lesser than 1 in order to give a meaning to  $\sum_{\ell=1}^{\infty} (-1)^\ell \{f_\Lambda\}^{*\ell}$ .

*Remark 25.* 1. As  $f_\Lambda$  is radial we have:

$$(-\Delta)f_\Lambda = (-\Delta_r)f_\Lambda = -(\partial_r^2 + \frac{2}{r}\partial_r)f_\Lambda. \quad (109)$$

2. For any  $u \in \mathbb{R}^3$  and  $r \geq 0$  Taylor's formula gives:

$$\begin{cases} (1 - \mathbf{s}_{u+2^{-1}r\mathbf{e}}\mathbf{s}_{u-2^{-1}r\mathbf{e}}) = r\{\mathbf{s}_u m_1(-\frac{r}{2}) - m_1(\frac{r}{2})\mathbf{s}_u\} \\ \text{with } m_1(\frac{x}{2}) := \int_{t=0}^1 d\mathbf{s}_{u+tx\mathbf{e}/2} \cdot \left(\frac{\mathbf{e}}{2}\right) dt. \end{cases} \quad (110)$$

We write  $\mathbf{g}(p) := \begin{pmatrix} g_0(p) \\ \mathbf{g}_1(p) \end{pmatrix} \in \mathbb{R}^4$  and  $\sigma(p) := \frac{\mathbf{g}(p)}{E(p)}$ .

As we have  $\langle \sigma(u), d\sigma(u) \rangle = 0$ , Taylor's Formula at order 2 gives

$$\begin{cases} \frac{1 - \langle \sigma(u + r\frac{\mathbf{e}}{2}), \sigma(u - r\frac{\mathbf{e}}{2}) \rangle}{r^2} := \langle a(u), a(u) \rangle + \langle \sigma(u), m_2(r) + m_2(-r) \rangle \\ \quad + r\langle a(u), m_2(r) - m_2(-r) \rangle + r^2\langle m_2(r), m_2(-r) \rangle, \\ a(u) := d\sigma(u) \cdot \frac{\mathbf{e}}{2} \text{ and } m_2(\frac{x}{2}) := \iint_{[0,1]^2} d^2\sigma_u + stx\mathbf{e}/2 \cdot \left(\frac{\mathbf{e}}{2}, \frac{\mathbf{e}}{2}\right) tdsdt. \end{cases} \quad (111)$$

3. For any  $-\frac{1}{2} \leq x \leq \frac{1}{2}$ :

$$\tilde{E}(u + x\mathbf{e}) \geq E(u + x\mathbf{e}) \geq \frac{E(u)}{2}. \quad (112)$$

In particular if one takes the modulus of the derivative over  $r$  in (110) or (111) for  $0 \leq r \leq 1$ , we get the following upper bounds:

- (a)  $K/\tilde{E}(u)$  for the first derivative,
- (b)  $K/\tilde{E}(u)^2$  for the second.



**Lemma 13.** *The functions  $\partial_r f_\Lambda$  and  $\partial_r^2 f_\Lambda$  are well defined in  $\mathbb{R}^3$  with support in  $\overline{B}(0, 2\Lambda)$ . Furthermore for  $J \in \mathbb{N}$  we have:*

$$\left\{ \begin{array}{l} |\partial_r f_{\Lambda, J}(p)| \lesssim J \frac{\alpha^{J+1} \log(\Lambda) K^{J+1}}{E(p)} \chi_{|p| < 2\Lambda} \quad \left| \partial_r f_\Lambda(p) \right| \lesssim \frac{L}{E(p)} \chi_{|p| < 2\Lambda}, \\ |\partial_r^2 f_{\Lambda, J}(p)| \lesssim J \frac{\alpha^{J+1} \log(\Lambda) K^{J+1}}{E(p)^2} \chi_{|p| < 2\Lambda} \quad \left| \partial_r^2 g_\Lambda \right| \lesssim \frac{\alpha^2 \log(\Lambda)}{E(r)^2} \chi_{r < 2\Lambda}. \end{array} \right. \quad (113)$$

As a consequence:

**Lemma 14.** *For  $\alpha$  sufficiently small,  $\check{g}_\Lambda \in L^1$  and*

$$\|\check{g}_\Lambda\|_{L^1} \lesssim (\alpha \log(\Lambda))^2. \quad (114)$$

*Remark 26.* At the very end of the proof of Lemma 13, we refer the reader to the thesis of the author for a (last) technical assumption: proving that  $\lim_{|x| \rightarrow 2\Lambda^-} \partial_r^2 f_\Lambda(x) = 0$ .

**Proof of Lemma 14**

We assume Lemma 13. As  $(-\Delta_r) = -(\partial_r^2 + \frac{2}{r}\partial_r)$  we have  $f_\Lambda \in H^2(\mathbb{R}^3)$  with:

$$|\Delta f_\Lambda(p)| \lesssim \frac{L}{|p|E(p)}. \quad (115)$$

**Proof of  $\int_{B(0,1)} |\check{f}_\Lambda(x)| dx \lesssim L$ :** The function  $-\Delta f_\Lambda$  has a singularity at  $r = 0$  due to the term  $\frac{2\partial_r f_\Lambda(r)}{r}$ . We split  $-\Delta f_\Lambda$  w.r.t.  $\chi_{|x| \leq 1} + \chi_{|x| > 1}$ . We have

$$I_\Lambda^{(2)} := -\Delta f_\Lambda \chi_{|x| \leq 1} \in L^{p_1} \cap L_w^3, \text{ and } E_\Lambda^{(2)} := -\Delta f_\Lambda \chi_{|x| > 1} \in L_w^{3/2} \cap L^{p_2}, \quad |p_2| > \frac{3}{2}, 1 \leq p_1 < 3. \quad (116)$$

The corresponding norms are respectively  $\mathcal{O}(LK(p_1))$  and  $\mathcal{O}(LK(p_2))$ . We use the Hausdorff-Young inequality and the generalized Young inequality [21, Vol. II]. The decomposition (116) implies the decomposition  $f_\Lambda = I_\Lambda^{(0)} + E_\Lambda^{(0)}$  by multiplication by  $\frac{1}{\Delta}$ .

For  $p = 1, a = \frac{1}{2}, q = 2$  and  $q' = \frac{q}{q-1}$  we have

$$\begin{aligned} \int_{|x| \leq 1} |\check{I}_\Lambda^{(0)}(x)| dx &\leq \left( \int |x|^{aq'} |\check{I}_\Lambda^{(0)}(x)|^{q'} dx \right)^{1-1/q} \left( \int_{|x| \leq 1} \frac{dx}{|x|^{aq}} \right)^{1/q}, \\ &\lesssim \|\nabla\|^a I_\Lambda^{(0)}\|_{L^q} \lesssim \left\| \frac{1}{|\cdot|^{1+a}} * I_\Lambda^{(2)} \right\|_{L^q}, \\ &\lesssim \|I_\Lambda^{(0)}\|_{L^p} \left\| \frac{1}{|\cdot|^{1+a}} \right\|_{L_w^{\frac{3}{1+a}}} \lesssim L. \end{aligned}$$

Similarly let  $0 < \varepsilon < 1$  to be chosen: we have  $|\nabla|^{2-\varepsilon} E_\Lambda^{(0)} = \frac{K(\varepsilon)}{|\cdot|^{3-\varepsilon}} * E_\Lambda^{(2)}$ . This last function is in  $L^2$  provided that  $E_\Lambda^{(2)} \in L^{\frac{6}{3-2\varepsilon}}$ . We choose  $\varepsilon = 3/4$  for instance: this gives

$$\int_{B(0,1)} |E_\Lambda^{(0)}(x)| dx \leq \sqrt{\int |x|^{5/2} |E_\Lambda^{(0)}(x)|^2 dx} \int_{B(0,1)} \frac{dx}{|x|^{5/2}} \lesssim \|E_\Lambda^{(2)}\|_{L^4} \left\| \frac{1}{|\cdot|^{9/4}} \right\|_{L_w^{4/3}} \lesssim L.$$

**Proof of  $\int_{|x| \geq 1} |\check{f}_\Lambda(x)| dx \lesssim L$ :** Then it is clear that

$$\int_{|x| \geq 1} |\check{f}_\Lambda(x)| \lesssim \|-\Delta f_\Lambda\|_{L^2} \sqrt{\int_{|x| \geq 1} \frac{dx}{|x|^4}} \lesssim L.$$

□

**Proof of Lemma 13** The idea of the proof is that each time we differentiate with respect with the radius  $r > 0$ , it leads to an additional term  $\frac{1}{E(u)}$  in the integrand or a change of the domains and so a better upper bound of the integral.

We will often use the following inequality:

$$\int_{B(0,\Lambda)} \frac{dv}{|u-v|^2(\tilde{E}(v+\frac{k}{2})+\tilde{E}(v-\frac{k}{2}))|u+\varepsilon\frac{k}{2}|} \leq \frac{1}{|u+\varepsilon\frac{k}{2}|} \int \frac{dv}{|v|^2|v-\mathbf{e}|^2}, \quad (117)$$

and for convenience we write:

$$u^\varepsilon := u + \varepsilon\frac{k}{2}, \quad \varepsilon \in \{1, -1\}. \quad (118)$$

That the function (and its derivatives) has an extension in 0 is clear from (110) and (112): differentiating under the integral sign of the Taylor's formula, we get:

$$\left| d^{J+1} \mathbf{s}_{u+tr\mathbf{e}/2} \cdot ((t\mathbf{e}/2)^J, \frac{\mathbf{e}}{2}) \right| \leq K^J \frac{1}{E(u)^{J+1}}, \quad 0 < r, t < 1, \quad (119)$$

thus the problem of singularity at  $r = 0$  drops thanks to (112).

More generally the variable  $r$  appears

1. either in the domains  $B(r)^{J+1}$ ,
2. or in a function of  $v_j \pm r\frac{\mathbf{e}}{2}$ .

One may write:

$$f_{\Lambda,J}(r) =: \int_{B(r)^{J+1}} G_J(u_0 \pm r\frac{\mathbf{e}}{2}, \dots, u_J \pm r\frac{\mathbf{e}}{2}) d\mathbf{u}, \quad (120)$$

$$G_J =: G_J^0(u_0 \pm r\frac{\mathbf{e}}{2}, \dots, u_J \pm r\frac{\mathbf{e}}{2}) \prod_{1 \leq j \leq J} \frac{1}{|u_j - u_{j-1}|^2}.$$

It is easy to see that  $G_J^0 : (\mathbb{R}^3)^{2J+2} \rightarrow \mathbb{R}$  is a differentiable function and that each time we take  $\partial_{u_j+r\frac{\mathbf{e}}{2}} - \partial_{u_j-r\frac{\mathbf{e}}{2}}$  we get a term  $K(r^{-1} + \tilde{E}(u \pm \frac{k}{2})^{-1})$  for  $r > 1$  or  $K\tilde{E}(u)^{-1}$  for  $r \leq 1$  (see Remark 25). This enables us to get upper bounds of the terms of  $\partial_r^j f_{\Lambda,J}$  corresponding to derivatives of  $G_J^0$ . Indeed for the first derivative: for  $\varepsilon, \varepsilon' \in \{+, -\}$  one has for  $1 < |k| < 2\Lambda$ :

$$\left\{ \begin{array}{l} \int \frac{du_j}{|u_j - u_{j-1}|^2 \tilde{E}(u_j + \varepsilon k/2)^2} \leq \frac{1}{|u_{j-1} + \varepsilon k/2|} \int_{\mathbb{R}^3} \frac{du_j}{|u_j|^2 |u_j - \mathbf{e}|^2}, \\ \int \frac{du_i}{|u_i - u_{i-1}|^2 |u_i + \varepsilon k/2| \tilde{E}(u_i, k/2) \tilde{E}(u_i + \varepsilon' k/2)} \leq \frac{1}{|k|} \left( \frac{1}{|u_{i-1} + k/2|} + \frac{1}{|u_i - k/2|} \right) \\ \quad \times \int_{(\mathbb{R}^3)} \frac{du_i}{|u_i - \mathbf{e}|^2 |u_i|^2}. \end{array} \right. \quad (121)$$

For the term  $(\partial_{u_0+k/2} - \partial_{u_0-k/2})G_0$  we have:

$$\int_{B(r)} \frac{du_0}{|u_0 - \varepsilon k/2| \tilde{E}(u + \varepsilon k/2)} \left( \frac{1}{\tilde{E}(u + \varepsilon k/2)^2} + \frac{1}{\tilde{E}(u - \varepsilon k/2)^2} \right) \lesssim \frac{2}{|k|} \int_{B(0,2\Lambda)} \frac{du_0}{\tilde{E}(u_0)^2 |u_0|} \lesssim \frac{\log(\Lambda)}{|k|}. \quad (122)$$

If  $r \leq 1$ , Remark 25 enables us to say that

$$\int_{B(r)^{J+1}} \frac{|\partial_r G_J^0(u_j \pm r\mathbf{e}/2)|}{\prod_{1 \leq j \leq J} |u_j - u_{j-1}|^2} \lesssim \alpha^{J+1} J \left( K \int \frac{du}{|u|^2 |u - \mathbf{e}|^2} \right)^J \log(\Lambda).$$




Figure 1: Level sets of the  $z$  function

– In the case of the terms corresponding to  $\partial_{v_1}\partial_{v_2}G_J^0$  with  $v_a = u_{i(a)} + \varepsilon(a)\frac{k}{2}$ , the above upper bounds enable us to say that if  $i(1) \neq i(2)$  then it suffices to apply twice (121),(122) and we get an upper bound of the form:

$$KJ^2(\chi_{|k|\leq 1} + \frac{\chi_{1<|k|<2\Lambda}}{|k|^2})\alpha^{J+1}\left(K\int\frac{du}{|u|^2|u-\mathbf{e}|^2}\right)^J\log(\Lambda),$$

If  $i(1) = i(2)$ , then as:

$$\int\frac{du}{|u-v|^2|u|\tilde{E}(u,\frac{k}{2})}\left(\frac{1}{\tilde{E}(u+k/2)^2} + \frac{1}{\tilde{E}(u-k/2)^2}\right)\lesssim\frac{1}{|k|^2|u|}, \quad (123)$$

we obtain an upper bound of the form

$$KJ(\chi_{|k|\leq 1} + \frac{\chi_{1<|k|<2\Lambda}}{|k|^2})\alpha^{J+1}\left(K\int\frac{du}{|u|^2|u-\mathbf{e}|^2}\right)^J\log(\Lambda).$$

If  $i(1) = i(2) = 0$ , we integrate first over  $u_0$ , then over  $u_1, u_2, \dots, u_J$  and use (123) with  $u = u_0, v = u_1$ : this gives

$$\text{for } 1 < r < 2\Lambda, \left|\partial_r^2\frac{1-\mathbf{s}_{u_0+\frac{k}{2}}\mathbf{s}_{u_0-\frac{k}{2}}}{r(\tilde{E}(u_0+\frac{k}{2})+\tilde{E}(u_0-\frac{k}{2}))}\right|\lesssim\frac{r^{-2}+\tilde{E}(u_0+\frac{k}{2})^{-2}+\tilde{E}(u_0-\frac{k}{2})^{-2}}{|u|\tilde{E}(u,\frac{k}{2})}.$$

If  $r \leq 1$  we use Remark 25 as before.

– There remains to deal with the terms corresponding to differentiation over  $r$  in the domain  $B(r)^{J+1}$ . We rewrite (120) using the *co-area formula*. Indeed, let us write for  $\varepsilon \in \{1, -1\}$  and  $r \in [0, 2\Lambda]$ :

$$B_\varepsilon(r) := \{p, |p + \frac{\varepsilon r}{2}\mathbf{e}| < \Lambda, \langle p, \varepsilon\mathbf{e} \rangle > 0\} \text{ and } B(r) := B_1(r) \cup B_{-1}(r) \subset B(0, \Lambda).$$

In particular  $B(\Lambda) = \{p \in B(0, \Lambda), \langle p, \mathbf{e} \rangle \neq 0\}$ . We define the level function:

$$\begin{aligned} B(\Lambda) &\rightarrow [0, 2\Lambda] \\ z: p \in B_\varepsilon(\Lambda) &\mapsto r \text{ such that } \left|u + \frac{r\varepsilon\mathbf{e}}{2}\right| = \Lambda. \end{aligned}$$

We apply the co-area formula with respect to  $z$ . If  $p \in B_{\varepsilon_0}$ , we write  $\varepsilon(p) := \varepsilon_0$  and

$$n(p) := \frac{p + \varepsilon(p)z(p)\frac{\mathbf{e}}{2}}{|p + \varepsilon(p)z(p)\frac{\mathbf{e}}{2}|} = \Lambda^{-1}(p + \varepsilon(p)z(p)\frac{\mathbf{e}}{2}).$$

For  $0 \leq r < 2\Lambda$  we write  $S(r) := \{p \in B, z(p) = r\}$  and  $S_\varepsilon(r) := S \cap B_\varepsilon$ ; each  $S_\varepsilon(r)$  is a spherical cap of  $S(-\frac{r\varepsilon\mathbf{e}}{2}, \Lambda)$ . The measure of  $B(0, \Lambda) \setminus B(\Lambda)$  is zero and the function  $z$  is differentiable with

$$\nabla z(p) = \frac{-2\varepsilon(p)}{\langle n(p), \mathbf{e} \rangle} n(p).$$

Thus for any integrable function  $F : B(0, \Lambda) \rightarrow \mathbb{R}$  and  $0 \leq r < 2\Lambda$  we have:

$$\int_{B(r)} F(p) |\nabla z(p)| dp = \int_{t=r}^{2\Lambda} dt \int_{S(t)} F(p) d\mathcal{H}_2(p), \quad (124)$$

where  $d\mathcal{H}_2(p)$  is the Hausdorff measure on  $S(r)$ . If we take spherical coordinates with axis  $\mathbb{R}e$  in  $S_\varepsilon(r)$  there holds  $d\mathcal{H}_2(p) = \Lambda^2 \sin(\theta) d\theta d\phi$  in the domain:

$$M_\pm(r) = \{(\theta, \phi) \in (\frac{\pi}{2}, \frac{\pi}{2} \mp \frac{\pi}{2}) \times (-\pi, \pi), \cos(\theta) \geq \frac{r}{2\Lambda}\}.$$

*Notation 27.* For convenience we write  $du$  for both  $d\mathcal{H}_2(u)$  (integration over a spherical cap) or  $d\mathcal{H}_1(u)$  (integration over a curve).

– For each  $u_j$  we rewrite the integration over  $u_i \in B(r)$ . For each  $0 \leq j \leq J$  we need to estimate

$$\int_{B(r)^{j-1} \times S(r) \times B(r)^{J-j}} \frac{du_0 \cdots \widehat{du_j} \cdots du_J d\mathcal{H}_2(u_j)}{\prod_{1 \leq j \leq J} |u_j - u_{j-1}|^2} |G_J^0(u_i \pm \frac{k}{2})|.$$

In  $S_\varepsilon(r)$  we take spherical coordinates and write  $v = u_{j-1} + \frac{\varepsilon r}{2} \mathbf{e}$ , if  $j = 0$  we replace  $u_{j-1}$  by  $u_2$  and integrate over  $u_1, u_2, \dots, u_J$ . Using (121) we have:

$$\begin{aligned} \int_{M_\varepsilon(r)} \frac{\Lambda^2 \sin(\theta) d\theta d\phi}{|v - \Lambda n|^2} \frac{1}{|\Lambda n|(\widetilde{E}(\Lambda n) + \widetilde{E}(\Lambda n - k))} &\leq \int_{\mathbb{S}^2} \frac{\sin(\theta) d\theta d\phi}{|v - \Lambda n|^2} \\ &\leq \frac{2\pi}{\Lambda|v|} \log\left(\frac{\Lambda+|v|}{\Lambda-|v|}\right). \end{aligned}$$

Then writing  $v := u_{i-1} + \varepsilon \frac{k}{2}$  we have:

$$\begin{aligned} \int_{B(r)} \frac{du_i}{|u_i - u_{i-1}|^2 |u_i + \varepsilon \frac{k}{2}|} \log\left(\frac{\Lambda+|u_i + \varepsilon \frac{k}{2}|}{\Lambda-|u_i + \varepsilon \frac{k}{2}|}\right) \frac{1}{\widetilde{E}(u_i, k/2)} &= \int_{B(r) + \frac{\varepsilon k}{2}} \frac{du_i}{|u_i - v|^2 |u_i| \widetilde{E}(u_i)} \log\left(\frac{\Lambda+|u_i|}{\Lambda-|u_i|}\right) \\ &\leq \int_{B(0,1)} \frac{du}{|u - v\Lambda^{-1}|^2 |u| \widetilde{E}(\Lambda u)} \log\left(\frac{1+|u|}{1-|u|}\right) \leq 2\pi \int_{r=0}^1 \frac{\Lambda dr}{|v| \widetilde{E}(\Lambda r)} \log\left(\frac{1+r}{1-r}\right) \log\left|\frac{\Lambda^{-1}|v|+r}{\Lambda^{-1}|v|-r}\right| \\ &\leq 2\pi \int_0^1 \frac{\Lambda dr}{|v| \widetilde{E}(\Lambda r)} \left(\log^2\left(\frac{1+r}{1-r}\right) + \log^2\left|\frac{\Lambda^{-1}|v|+r}{\Lambda^{-1}|v|-r}\right|\right) \lesssim |v|^{-1}. \end{aligned}$$

Finally for sufficiently small  $\alpha$ , we have

$$|\partial_r f_{\Lambda, J}(r)| \leq KL(\alpha K)^J (\chi_{r \leq 1} + \frac{\chi_{1 < r < 2\Lambda}}{r}).$$

So by dominated convergence, as  $r$  tends to  $(2\Lambda)^-$ ,  $\partial_r f_{\Lambda, J}$  tends to 0 and  $g_\Lambda \in H^1(\mathbb{R}^3)$ .

– Let us deal with the second derivative. There remains the three cases:

1. One derivative in  $B(r)$  and one derivative in the integrand.
2. Two derivatives in two different  $B(r)$ .
3. Two derivatives in the same  $B(r)$ .

In fact, we have to deal with the last two cases together because each term alone is not well defined but the sum gives a finite term. Seeing the second derivative as the coefficient of the second term in the Taylor series of  $g_{\Lambda, J}(r + \delta r)$ , each term is  $\mathcal{O}_{\delta r \rightarrow 0}(-\delta r \log(\delta r))$  but the sum is  $\mathcal{O}(\delta r)$  due to some cancellation.

1.

1.1. One derivative in  $u_{i_1} \pm \frac{\varepsilon}{2} \mathbf{e}$  and one in the domain of  $u_{i_2}$  with  $i_1 \neq i_2$ . Up to integrating over  $u_j$  from  $j = 0$  to  $j = J$ , we can suppose that  $i_1 < i_2$ . We split  $S(r)$  between  $S_+(r)$  and  $S_-(r)$ . In  $S_\varepsilon(r)$ , we use (117) (and (98) at the beginning), this gives:

$$\begin{aligned} \int_{S_\varepsilon(r)} \frac{du_{i_2}}{|u_{i_2-1} - u_{i_2}|^2 \widetilde{E}(u_{i_2} + \varepsilon \frac{k}{2}) |u_{i_2} + \varepsilon \frac{k}{2}|} &\leq \int_{\mathbb{S}^2} \frac{du_{i_2}}{\Lambda^2 \left|\frac{|u_{i_2-1}^\varepsilon|}{\Lambda} - u_{i_2}\right|^2} \\ &\lesssim \frac{1}{\Lambda |u_{i_2-1}^\varepsilon|} \log\left(\frac{1 + \frac{|u_{i_2-1}^\varepsilon|}{\Lambda}}{\Lambda - |u_{i_2-1}^\varepsilon|}\right). \end{aligned}$$

We take spherical coordinates with respect to  $-\varepsilon \frac{k}{2}$ : for any  $v \in B = B_+ \cup B_-$  we have

$$\begin{aligned} \int_{B(0,\Lambda)} \frac{du}{|u_{i_2-1} - v^\varepsilon|^2 |u_{i_2-1}| \tilde{E}(u_{i_2-1})} \log \left( \frac{1 + \frac{|u_{i_2-1}|}{\Lambda}}{1 - \frac{|u_{i_2-1}|}{\Lambda}} \right) &\lesssim \frac{1}{|v|} \int_0^1 dz \log \left( \frac{1+z}{1-z} \right) \log \left( \frac{\frac{|v^\varepsilon|}{\Lambda} + z}{\frac{|v^\varepsilon|}{\Lambda} - z} \right) \\ &\lesssim \frac{1}{|v^\varepsilon|} \int_0^2 \log \left( \frac{1+z}{1-z} \right)^2, \\ \int_{B(0,\Lambda)} \frac{du}{|u_{i_2-1}| \tilde{E}(u_{i_2-1}^2)} \log \left( \frac{1 + \frac{|u_{i_2-1}|}{\Lambda}}{1 - \frac{|u_{i_2-1}|}{\Lambda}} \right) &\lesssim \int_0^\Lambda \frac{dz}{E(z)} \log \left( \frac{1 + \frac{z}{\Lambda}}{1 - \frac{z}{\Lambda}} \right) \\ &\lesssim 1 + \Lambda^{-1}. \end{aligned}$$

Then we use the same method as for the first derivative: when integrating over  $u_{i_1}$ , we use  $(\tilde{E}(u_{i_1} + \frac{k}{2}) + \tilde{E}(u_{i_1} - \frac{k}{2}))^{-1} \leq \tilde{E}(\frac{k}{2})^{-1}$ . In this first subcase, we get an upper bound of the form:

$$\frac{J^2(K\alpha)^{J+1} \log(\Lambda)}{\Lambda E(k)}.$$

1.2. One derivative in  $u_i \pm \frac{r}{2} \mathbf{e}$  and one in the domain of  $u_i$ . Splitting the integration over  $S_+(r)$  and  $S_-(r)$ , and using (117), we have to estimate

$$\int_{S_\varepsilon(r)} \frac{du_i}{|u_i - v|^2 |u_i + \varepsilon \frac{k}{2}| \tilde{E}(u_i, k/2) \tilde{E}(u_i + \varepsilon' \frac{k}{2})} \leq \int_{S_\varepsilon(r)} \frac{du_i}{|u_i - v|^2 |u_i + \varepsilon \frac{k}{2}| \tilde{E}(u_i, k/2) \tilde{E}(u_i - \varepsilon \frac{k}{2})}. \quad (125)$$

Above  $v$  is either  $u_{i+1}$  or  $u_{i-1}$  depending on the order of integration (from  $u_J$  to  $u_0$  or from  $u_0$  to  $u_J$  if the derivatives act on  $u_0 + \frac{k}{2}$ ). Moreover  $\varepsilon, \varepsilon' \in \{1, -1\}$  and the term with  $\varepsilon'$  comes from the derivative in the integrand. By using (117) several times (starting with (98)) we get the term  $|u_i + \varepsilon \frac{k}{2}| = |u_i^\varepsilon|$  in (125).

In (125), we use spherical coordinates to obtain the following upper bound:

$$\int_{\mathbb{S}^2} \frac{\Lambda^2 du}{\Lambda^2} \frac{1}{|\Lambda u - v^\varepsilon|^2 \tilde{E}(\Lambda u - r\mathbf{e})} \leq 2 \int_{\mathbb{S}^2} \frac{du}{|\Lambda u - |v^\varepsilon| \mathbf{e}|^2 E(u - r\mathbf{e})}. \quad (126)$$

Let us assume for the moment that this integral is lesser than:

$$\frac{K}{\Lambda^2 |v^\varepsilon|} \left( 1 - \chi_{|v^\varepsilon| > (2-\sqrt{3})\Lambda} \log \left( 1 - \frac{|v^\varepsilon|}{\Lambda} \right) \right). \quad (127)$$

In the process of integrating over the  $u_i$ 's, we have to integrate over  $v$  with this upper bound. Taking spherical coordinates with respect to  $-\frac{\varepsilon r}{2} \mathbf{e}$ , we have:

$$\left\{ \begin{array}{l} \int_{B(0,\Lambda)} \frac{dv}{|v' - v|^2 \tilde{E}(v) |v|} \lesssim \frac{1}{|v'|} \int \frac{dv}{|v|^2 |v - \mathbf{e}|^2} \\ \int_{B(0,\Lambda)} \frac{dv}{|v| \tilde{E}(v)^2} \lesssim \log(\Lambda). \end{array} \right.$$

Moreover, writing  $A_\Lambda := A(0, (2 - \sqrt{3})\Lambda, \Lambda)$  the annulus, we have:

$$\left\{ \begin{array}{l} \int_{A_\Lambda} \frac{\log(1 - \frac{|v|}{\Lambda}) dv}{|v' - v|^2 |v| \tilde{E}(v) (\Lambda^2 + |v|^2)} \lesssim \frac{1}{\Lambda^2 |v'|} \int_{2-\sqrt{3}}^1 \frac{-\log(1-z) dz}{z(1+z^2)} \log \left| \frac{\frac{|v'|}{\Lambda} + z}{\frac{|v'|}{\Lambda} - z} \right| \\ \int_{A_\Lambda} \frac{\log(1 - \frac{|v|}{\Lambda}) dv}{|v| \tilde{E}(v)^2 (\Lambda^2 + |v|^2)} \lesssim \frac{1}{\Lambda^2 |v'|}, \\ \int_{A_\Lambda} \frac{\log(1 - \frac{|v|}{\Lambda}) dv}{|v| \tilde{E}(v)^2 (\Lambda^2 + |v|^2)} \lesssim \frac{1}{\Lambda^2} \int_{2-\sqrt{3}}^1 \frac{-\log(1-z) dz}{z^3}. \end{array} \right.$$

*Proof of (126) ≤ (127)* We write

$$x := |v^\varepsilon|, \quad A := \Lambda^2 + x^2, \quad B := \sqrt{1 + \Lambda^2 + r^2}, \quad a := \frac{2x\Lambda}{x^2 + \Lambda^2} \quad \text{and} \quad b := \frac{2\Lambda r}{1 + \Lambda^2 + r^2}.$$

The upper bound (126) is equal to

$$\frac{4\pi}{AB} \int_{-1}^1 \frac{dy}{(1-ay)\sqrt{1-by}} = \frac{4\pi}{ABa} \int_0^{\frac{2b}{\sqrt{1+b}+\sqrt{1-b}}} \frac{dz}{z^2 + 2z\sqrt{1-b} + b(\frac{1}{a}-1)}. \quad (128)$$

If  $a \leq \frac{1}{2}$ , then this integral is  $\mathcal{O}\left(\frac{1}{AB} \int_{-1}^1 \frac{dy}{\sqrt{1-by}}\right) = \mathcal{O}\left(\frac{1}{\Lambda^2 E(v^\varepsilon)}\right)$ .

Similarly, if  $b \leq \frac{1}{2}$ , we get:  $\mathcal{O}\left(\frac{1}{AB} \int_{-1}^1 \frac{dy}{1-ay}\right) = \mathcal{O}\left(\frac{1}{\Lambda^2 E(v^\varepsilon)}\right)$ .

If  $\frac{1}{2} < a, b \leq 1$ , we consider formula (128).

We have  $z^2 \geq 2z\sqrt{1-b}$  for  $z \geq 2\sqrt{1-b}$  and  $2\sqrt{1-b} < \frac{2b}{\sqrt{1+b}+\sqrt{1-b}}$  iff  $b > \frac{4}{5}$ .

For  $\frac{1}{2} < b \leq \frac{4}{5}, a > \frac{1}{2}$  we get the upper bound:

$$\frac{20\pi}{AB} \int_{-1}^1 \frac{dy}{1-ay} \lesssim \frac{1}{\Lambda^2 |v^\varepsilon|}.$$

For  $b > \frac{4}{5}, a > \frac{1}{2}$ , we have the upper bound

$$\frac{4\pi}{AaB} \left( \int_0^{2\sqrt{1-b}} \frac{dz}{2z\sqrt{1-b} + b(\frac{1}{a}-1)} + \int_{2\sqrt{1-b}}^{\frac{2b}{\sqrt{1+b}+\sqrt{1-b}}} \frac{dz}{z^2 + b(\frac{1}{a}-1)} \right) \quad (129)$$

The first integral of (129) gives (without  $4\pi/(AB)$ )

$$\frac{1}{2a\sqrt{1-b}} \log \left( 1 + \frac{4(1-b)}{b(\frac{1}{a}-1)} \right) \leq \frac{1}{\sqrt{1-b}} \log \left( 1 + 5 \frac{1-b}{1-a} \right).$$

If  $1-b \leq 1-a$ , then this gives  $\mathcal{O}((1-b)^{-1/2})$ , else this gives  $\mathcal{O}(\frac{\log(1-a)}{\sqrt{1-b}})$ .

The second integral of (129) gives (without  $4\pi/(AaB)$  and writing  $X := (a^{-1}-1)^{-1}$ ):

$$\begin{aligned} \sqrt{\frac{X}{b}} \int_{2\sqrt{\frac{1-b}{b}X}}^{\frac{2\sqrt{bX}}{\sqrt{1-b}+\sqrt{1+b}}} \frac{dz}{z^2+1} &\lesssim \int_{2\sqrt{1-b}}^2 \frac{Xz^2}{1+Xz^2} \frac{dz}{z^2} \\ &\lesssim \frac{1}{\sqrt{1-b}} = \frac{\sqrt{1+\Lambda^2+r^2}}{\sqrt{1+(\Lambda-r)^2}}. \end{aligned}$$

We have:  $\frac{\log(1-a)}{AB\sqrt{1-b}} = 2 \frac{\log \left| \frac{\sqrt{\Lambda^2+x^2}}{\Lambda-x} \right|}{(\Lambda^2+x^2)\sqrt{1+(\Lambda-x)^2}} \lesssim \frac{1 + \log(1 - \frac{|v^\varepsilon|}{\Lambda})}{(\Lambda^2 + |v^\varepsilon|^2)\sqrt{1+(\Lambda-|v^\varepsilon|)^2}}$ .

Let us emphasize that the condition  $a > 2^{-1}$  is equivalent to  $\frac{|v^\varepsilon|}{\Lambda} \geq 2 - \sqrt{3}$ .

Bringing all those estimates together ends the proof of (126)  $\leq$  (127).

2.

2.1. One derivative in the domain of  $u_j$  and one in the domain of  $u_i$  with  $i-j \geq 2$ . We integrate over  $u_{j'}$  from  $j'=0$  to  $j'=j$  and from  $j'=J$  to  $j'=i$  using the method for the first derivative. The integration over  $u$  with  $u$  either  $u_j$  or  $u_i$  (resp. with  $v$  either  $u_{j+1}$  or  $u_{i-1}$ ) gives:

$$\begin{aligned} \sum_{\varepsilon \in \{1, -1\}} \int_{S_\varepsilon(r)} \frac{du}{|u-v|^2} \frac{1}{|u + \varepsilon \frac{k}{2}| \tilde{E} \left( u + \varepsilon \frac{k}{2} \right)} &\lesssim \frac{1}{\Lambda^2} \sum_{\varepsilon} \int_{\frac{r}{2\Lambda}}^1 \frac{dy}{\Lambda^2 + |v^\varepsilon|^2 - 2\Lambda|v^\varepsilon|y} \\ &\lesssim \sum_{\varepsilon} \frac{1}{\Lambda|v^\varepsilon|} \log \left( \frac{\Lambda + |v^\varepsilon|}{\Lambda - |v^\varepsilon|} \right). \end{aligned} \quad (130)$$

If  $j + 2 \leq i$ , then by integrating over  $u_{j+1}$  we have:

$$\begin{aligned} & \int_{B(r)} \frac{du_{j+1}}{|u_{j+1} - u_{j+2}|^2} \frac{1}{|u_{j+1}^\varepsilon| |(\tilde{E}(u_{j+1}^+) + \tilde{E}(u_{j+1}^-))|} \log \left( \frac{\Lambda + |u_{j+1}^\varepsilon|}{\Lambda - |u_{j+1}^\varepsilon|} \right) \\ & \leq \frac{1}{\Lambda} \int_{B(0,1)} \frac{du}{|u|^2 |u - \Lambda^{-1} u_{j+2}^\varepsilon|^2} \log \left( \frac{1 + |u|}{1 - |u|} \right) \\ & \leq \int_0^1 \frac{dr}{|u_{j+2}^\varepsilon|} \log \left( \frac{1+r}{1-r} \right) \log \left( \frac{r + \frac{|u_{j+2}^\varepsilon|}{\Lambda}}{r - \frac{|u_{j+2}^\varepsilon|}{\Lambda}} \right) \lesssim \frac{1}{|u_{j+2}^\varepsilon|}, \end{aligned}$$

and we conclude as before. Else  $j + 1 = i$  and we have:

$$\begin{aligned} \int_{B(r)} \frac{du_{j+1}}{|u_{j+1}^\varepsilon|^2} \frac{1}{\tilde{E}(u_{j+1}^+) + \tilde{E}(u_{j+1}^-)} \log \left( \frac{\Lambda + |u_{j+1}^\varepsilon|}{\Lambda - |u_{j+1}^\varepsilon|} \right)^2 & \lesssim \frac{1}{\Lambda} \int_{z=0}^\Lambda \frac{dz}{\tilde{E}(z)} \log \left( \frac{1 + \frac{z}{\Lambda}}{1 - \frac{z}{\Lambda}} \right)^2 (1 - \frac{r}{2\Lambda}) \\ & \lesssim (1 - \frac{r}{2\Lambda})(\log(\Lambda) + 1). \end{aligned}$$

2.2. One derivative in the domain of  $u_j$  and one in the domain of  $u_{j+1}$ . We only look at the corresponding coefficient in the Taylor series of  $g_{\Lambda, J}(r + \delta r)$  with  $r' = r + \delta r$ . Indeed, let us treat for instance

$$\begin{aligned} & \iint_{(u_j, u_{j+1}) \in B(r') \times S(r)} \frac{du_j du_{j+1}}{|u_j - u_{j+1}|^2} \frac{|\langle n(u_{j+1}), \mathbf{e} \rangle|}{2} \int_{B(r)^{J-1}} \frac{G_J^0(u_\ell \pm \frac{k}{2})}{\prod_{a \neq j+1} |u_a - u_{a+1}|^2} \\ =: & \iint_{(u_j, u_{j+1}) \in B(r') \times S(r)} \frac{du_j du_{j+1}}{|u_j - u_{j+1}|^2} \frac{|\langle n(u_{j+1}), \mathbf{e} \rangle|}{2} G_{J, j}(u_j, u_{j+1}). \end{aligned}$$

We subtract the integral of the same function but over  $(u_j, u_{j+1}, \mathbf{u}')$  in  $B(r) \times S(r) \times B(r)^{J-1}$  where  $\mathbf{u}' = (u_0, \dots, \hat{u}_j, \hat{u}_{j+1}, \dots)$  and use the co-area formula. This gives

$$\int_{r+\delta r}^r dt \int_{S(t)S(r)} \frac{du_j du_{j+1}}{|u_j - u_{j+1}|^2} \frac{|\langle n(u_j), \mathbf{e} \rangle|}{2} \frac{|\langle n(u_{j+1}), \mathbf{e} \rangle|}{2} G_{J, j}(u_j, u_{j+1}). \quad (131)$$

We deal with  $G_{J, j}(u_j, u_{j+1})$  as in the case 2.1. Let us say for instance  $0 < \delta r \ll 1$ , then for any  $(u_{j+1}, t) \in S(r) \times (r, r')$  we have:

$$\text{dist}(u_{j+1}, S(t)) \geq \Lambda \left| \sqrt{1 + \frac{\langle n_{u_{j+1}}, \mathbf{e} \rangle}{\Lambda} \delta r + \left( \frac{t-r}{2} \right)^2} - 1 \right| = \mathcal{O}_{\delta r \rightarrow 0}(\Lambda |(t-r) \langle n_{u_{j+1}}, \mathbf{e} \rangle|).$$

By the Theorem of projection onto a closed convex  $\mathbb{R}^3$ , we have

$$|u_{j+1} - u_j|^2 \geq |u_{j+1} - \Pi_{S(t)} u_{j+1}|^2 + |\Pi_{S(t)} u_{j+1} - u_j|^2.$$

If  $r' < r$ , then we consider instead the projection of  $u_j \in S(r)$  onto  $B(t)$ . Anyway the quantity in (131) is

$$\mathcal{O}_{\delta r \rightarrow 0} \left( \frac{(\alpha K)^J}{\Lambda^2} \int_r^{r+\delta r} dt \int_{\mathbb{S}^2} da |\langle a, \mathbf{e} \rangle| \log \left( 1 + \frac{1}{|t-r|^2 |\langle a, \mathbf{e} \rangle|^2} \right) = \frac{(\alpha K)^J}{\Lambda^2} \delta r (1 - \log(\delta r)) \right).$$

The corresponding term is not Lipschitz because of the term  $-\delta r \log(\delta r)$ .

3. Let us write the expansion of

$$\int_{B(r')} \frac{|\langle n_{u_j}, \mathbf{e} \rangle| du_j}{2} \int_{B(r)^J} du_0 \cdots \widehat{du_j} \cdots du_J G_J(u_\ell \pm \frac{k}{2}) =: \int_{B(r')} \tilde{G}_{J, j}(u_j) du_j. \quad (132)$$

We subtract  $\int_{B(r)} \tilde{G}_{J,j}(u_j) du_j$  in (132) and get

$$\int_{r+\delta r}^r dt \int_{S(t)} du_j \tilde{G}_{J,j}(u_j). \quad (133)$$

We split (133) between integration over  $S_+(t)$  and  $S_-(t)$ . For any  $t \in (r, r']$ , we write  $s := t - r$  and:

$$\begin{aligned} \Phi_t : \quad S(t) &\rightarrow S(r) \\ u \in S_\varepsilon(t) &\mapsto v(u) := u + z_t(u)n_u \in S_\varepsilon(r) \text{ where } |z(u)| = \mathcal{O}_{\delta r \rightarrow 0}(\delta r). \end{aligned} \quad (134)$$

From now on we assume  $v \in S(r)$  and  $u \in S(t)$  and write  $\bar{n}_u$  instead of  $n_u$  to emphasize this is a function of  $u \in S(t)$  and not of  $v \in S(r)$ . The function  $z_t : S(t) \rightarrow \mathbb{R}$  satisfies the equation

$$\left| u + z_t(u)\bar{n}_u + \varepsilon \frac{r}{2} \mathbf{e} \right|^2 = \Lambda^2 \text{ that is } z_t \left( 1 + \frac{z_t}{2\Lambda} - \frac{\varepsilon s \langle \bar{n}_u, \mathbf{e} \rangle}{2\Lambda} \right) = \frac{\varepsilon s \langle \bar{n}_u, \mathbf{e} \rangle}{2} - \frac{s^2}{8\Lambda}. \quad (135)$$

Changing variables in the integration over  $S(t)$  we have:

$$\int_{S(t)} du_j \tilde{G}_{J,j}(u_j) = \int_{\Phi_t(S(t))} dv \tilde{G}_{J,j}(\Phi_t^{-1}(v)) J(\Phi_t; \Phi_t^{-1}(v))^{-1} dv.$$

– We need to compute  $\Phi_t^{-1}(v)$  and  $J(\Phi_t; \Phi_t^{-1}(v))$ . First we have:

$$\bar{n}_u = \frac{v - z_t \bar{n}_u + \varepsilon \frac{r+s}{2} \mathbf{e}}{\Lambda} = n_v + \varepsilon \frac{s}{2\Lambda} \mathbf{e} - \frac{z_t}{\Lambda} \bar{n}_u,$$

thus

$$\bar{n}_u = \frac{1}{1 + \frac{z_t}{\Lambda}} (n_v + \varepsilon \frac{s}{2\Lambda} \mathbf{e}), \quad (136)$$

and

$$n_v = \left( 1 + \frac{z_t}{\Lambda} \right) \bar{n}_u - \frac{\varepsilon s}{2\Lambda} \mathbf{e}. \quad (137)$$

Using the formula (136) in (135), we obtain the following equation satisfied by  $z_t$ :

$$z_t \left( 1 + \frac{z_t}{\Lambda} - \frac{\varepsilon s}{2\Lambda \left( 1 + \frac{z_t}{\Lambda} \right)} \langle n_v, \mathbf{e} \rangle + \frac{\varepsilon s}{2\Lambda} \right) = \frac{\varepsilon s}{2 \left( 1 + \frac{z_t}{\Lambda} \right)} \left( \langle n_v, \mathbf{e} \rangle + \frac{\varepsilon s}{2\Lambda} \right) - \frac{s^2}{8\Lambda}. \quad (138)$$

In particular there holds:

$$z_t(u) = \frac{\varepsilon s}{2\Lambda} \langle n_v, \mathbf{e} \rangle + \mathcal{O}_{\delta r \rightarrow 0}((\delta r)^2). \quad (139)$$

We differentiate  $z_t$  in (135) and get:

$$\begin{aligned} T_u S_\varepsilon(t) &\rightarrow \mathbb{R} \\ dz_t(u) : \quad h &\mapsto \frac{\varepsilon s}{2\Lambda} \frac{\langle h, \mathbf{e} \rangle \left( 1 + \frac{z_t}{\Lambda} \right)}{1 + \frac{z_t}{\Lambda} - \frac{\varepsilon s}{2\Lambda} \langle \bar{n}_u, \mathbf{e} \rangle}. \end{aligned} \quad (140)$$

Thus differentiating in (134) and using (136) in (140) give

$$\begin{aligned} T_u S_\varepsilon(t) &\rightarrow T_v S_\varepsilon(r) \\ d\Phi_t(u) : \quad h &\mapsto \left( 1 + \frac{z_t}{\Lambda} \right) h + \frac{\varepsilon s}{2\Lambda} \frac{\langle h, \mathbf{e} \rangle \left( 1 + \frac{z_t}{\Lambda} \right)}{1 + \frac{z_t}{\Lambda} - \frac{\varepsilon s}{2\Lambda} \langle \bar{n}_u, \mathbf{e} \rangle} \frac{n_v + \frac{s}{2\Lambda} \mathbf{e}}{1 + \frac{z_t}{\Lambda}} \end{aligned} \quad (141)$$



Let  $(a, b)$  be an orthonormal basis of  $T_u S_\varepsilon(t)$  with  $b \times \bar{n}_u = a$ , then we have:

$$\begin{aligned}
J(\Phi_t; u) &= \langle d\Phi_t(u)a \times d\Phi_t(u)b, n_v \rangle \\
&= \langle ([1 + \frac{z_t}{\Lambda}]a + \bar{n}_u dz_t(u)a) \times ([1 + \frac{z_t}{\Lambda}]b + \bar{n}_u dz_t(u)b) \times, n_v \rangle \\
&= (1 + \frac{z_t}{\Lambda})^2 \langle \bar{n}_u, n_v \rangle - (1 + \frac{z_t}{\Lambda}) [\langle a, n_v \rangle dz_t(u)a + \langle b, n_v \rangle dz_t(u)b] \\
&= (1 + \frac{z_t}{\Lambda}) (1 + \frac{\varepsilon s}{2\Lambda} \langle n_v, \mathbf{e} \rangle) + \frac{\varepsilon s}{2\Lambda} (1 + \frac{z_t}{\Lambda}) (\langle a, \mathbf{e} \rangle dz_t(u)a + \langle b, \mathbf{e} \rangle dz_t(u)b) \\
&= (1 + \frac{z_t}{\Lambda}) (1 + \frac{\varepsilon s}{2\Lambda} \langle n_v, \mathbf{e} \rangle) + \frac{\varepsilon s}{2\Lambda} \left( 1 - \frac{(\langle n_v, \mathbf{e} \rangle + \frac{\varepsilon s}{2\Lambda})^2}{(1 + \frac{z_t}{\Lambda})^2} \right) \times \\
&\quad \frac{1 + \frac{z_t}{\Lambda}}{1 + \frac{z_t}{\Lambda} - \frac{\varepsilon s}{2\Lambda(1 + \frac{z_t}{\Lambda})} (\langle n_v, \mathbf{e} \rangle + \frac{\varepsilon s}{2\Lambda})} \\
&= 1 + \frac{\varepsilon s}{2\Lambda} \left[ 1 - \frac{(\langle n_v, \mathbf{e} \rangle + \frac{\varepsilon s}{2\Lambda})^2}{(1 + \frac{z_t}{\Lambda})^2} + \langle n_v, \mathbf{e} \rangle \left( \frac{1}{\Lambda} + 1 \right) \right] + \mathcal{O}_{\delta r \rightarrow 0}((\delta r)^2).
\end{aligned}$$

- As  $u = v - z_t \bar{n}_u = v + \frac{\varepsilon s}{2} \langle n_v, \mathbf{e} \rangle n_{v_j} + \mathcal{O}_{\delta r \rightarrow 0}((\delta r)^2)$ , we get:

$$\begin{aligned}
\int_{S_\varepsilon(t)} du_j \tilde{G}_{J,j}(u_j) r &= \int_{\Phi_t(S_\varepsilon(t))} dv_j \tilde{G}_{J,j}(v_j + \frac{\varepsilon s}{2} \langle n_{v_j}, \mathbf{e} \rangle n_{v_j} + \mathcal{O}_{\delta r \rightarrow 0}((\delta r)^2)) \times \\
&\quad \left( 1 - \frac{\varepsilon s}{2\Lambda} \left[ 1 - \frac{(\langle n_v, \mathbf{e} \rangle + \frac{\varepsilon s}{2\Lambda})^2}{(1 + \frac{z_t}{\Lambda})^2} + \langle n_v, \mathbf{e} \rangle \left( \frac{1}{\Lambda} + 1 \right) \right] + \mathcal{O}_{\delta r \rightarrow 0}((\delta r)^2) \right) dv.
\end{aligned} \tag{142}$$

We have  $\Phi_t(S_\varepsilon(t)) \neq S(r)$ . In spherical coordinates  $(r, \theta, \phi)$  with respect to  $-\varepsilon \frac{r}{2} \mathbf{e}$  and positive vertical half-line  $\mathbb{R}_+^3 \varepsilon \mathbf{e}$  we have

$$\Phi_t(S_\varepsilon(t)) \simeq \left\{ (\Lambda, \theta, \phi), \frac{rs}{2\Lambda \sqrt{1 - \frac{rs}{2\Lambda^2} + \frac{s^2}{4\Lambda^2}}} = \cos(\theta_t) \leq \cos(\theta) \leq 1 \right\}, \tag{143}$$

and  $\cos(\theta_t) = \frac{r}{2\Lambda} - \frac{r^2}{8\Lambda^2} s + \mathcal{O}_{\delta r \rightarrow 0}((\delta r)^2)$ .

At this point, we need to differentiate  $\tilde{G}_{J,j}$ : we have

$$\tilde{G}_{J,j}(u_j) = \frac{|\langle n_{u_j}, \mathbf{e} \rangle|}{2} \int_{B(r; u_j)} du_0 \cdots \widehat{du_j} \cdots du_J \frac{G_J^0(u_\ell \pm \frac{k}{2})}{\prod_{1 \leq i \leq J} |u_i - u_{i-1}|^2}.$$

We change variables as follows:  $v_i := u_i - u_j$ , this enables us to remove  $u_j$  from the term  $|u_j - u_{j \pm 1}|^{-2}$ . Writing  $B_\varepsilon(r, u_j) := \{v : |v + u_j + \varepsilon \frac{k}{2}| < \Lambda\}$ ,  $B(r; u_j) = B_+(r; u_j) \cup B_-(r; u_j)$  and  $S_\varepsilon(r, u_j) := \partial B_\varepsilon(r, u_j)$ , we have

$$\tilde{G}_{J,j}(u_j) = \frac{|\langle n_{u_j}, \mathbf{e} \rangle|}{2} \int_{B(r; u_j)^J} dv_0 \cdots \widehat{dv_j} \cdots dv_J \frac{G_J^0(v_\ell + u_j \pm \frac{k}{2})}{\prod_{1 \leq i \leq J} |v_i - v_{i+1}|^2}, \tag{144}$$

with the convention  $v_j = 0$ . We differentiate the formula (144):  $u_j$  appears in the integrand and in the domains  $B(r; u_j)$ . We deal with the terms corresponding to differentiation of the integrand as before. Then we have for any integrable function  $F$  and small displacement  $\delta u \in \mathbb{R}^3$ :

$$\int_{B_\varepsilon(r, u_j + \delta u)} F(v) dv - \int_{B_\varepsilon(r, u_j)} F(v) dv = \int_{S_\varepsilon(r; u_j)} F(v) (\langle n(v - u_j), \delta u \rangle + \mathcal{O}_{\delta u \rightarrow 0}(|\delta u|^2)) dv, \tag{145}$$

where  $n(v - u_j)$  is the outward normal of  $S_\varepsilon(r, u_j)$  at  $v$ . Substituting in (142), as in the case 2.2. we get terms which are  $\mathcal{O}_{\delta u \rightarrow 0}(|\delta u|(1 - \log|\delta u|))$ . Writing  $u_j = u$  we have

$$\begin{aligned}
\tilde{G}_{J,j}(v - \frac{\varepsilon s}{2} \langle n_v, \mathbf{e} \rangle n_v + \mathcal{O}_{\delta r \rightarrow 0}((\delta r)^2)) &= \mathcal{O}_{\delta r \rightarrow 0}(-(\delta r)^2 \log(\delta r)) \\
+ \tilde{G}_{J,j}(v) - \frac{\varepsilon s \langle n_v, \mathbf{e} \rangle}{2} \sum_{i \neq j} &\int_{\substack{v_i \in S(r; v) \\ \mathbf{v}'_i \in B(r, v)^J}} \frac{du_i d\mathbf{v}'_i}{\prod_{1 \leq \ell \leq J} |v_\ell - v_{\ell-1}|^2} G_J^0(v_\ell \pm \frac{r}{2} \mathbf{e}) \langle n(v_i - u_j), n_v \rangle.
\end{aligned} \tag{146}$$

We write  $\mathcal{C}(r) := S_+(r) \cap S_-(r)$  (this is a curve): the integration of  $\widetilde{G}_{J,j}(v_j)$  over  $S_\varepsilon(r) \Delta \Phi_{\delta r}(S_\varepsilon(\delta r))$  gives rise to a term:

$$-\frac{2r^2}{8\Lambda^2} \int_{u_j \in \mathcal{C}(r), (u_i)_{i \neq j} \in B(r)^J} \frac{r}{4\Lambda} du_0 \cdots du_J G_J(u_\ell \pm \frac{k}{2}) + \mathcal{O}_{\delta r \rightarrow 0}((\delta r)^2).$$

Thus we get a term of order

$$-\frac{2r^2}{8\Lambda^2} \int_{u_j \in \mathcal{C}(r), (u_i)_{i \neq j} \in B(r)^J} \frac{r}{4\Lambda} du_0 \cdots du_J G_J(u_\ell \pm \frac{k}{2}) = \mathcal{O}\left(\frac{(\alpha K)^{J+1}}{\Lambda^2}\right).$$

By integrating the term  $\widetilde{G}_{J,j}(v) \times (J(\Phi_t; u_j)^{-1} - 1)$ , we get a well defined number in the limit  $\delta r \rightarrow 0$ . Furthermore this term is

$$\mathcal{O}\left(\frac{1}{\Lambda} \int_{u_j \in S(r)} \int \cdots \int du_0 \cdots du_J |G_J(u_\ell \pm \frac{k}{2})|\right) = \mathcal{O}\left(\frac{(\alpha K)^{J+1} \log(\Lambda)}{\Lambda^2} \chi_{r < 2\Lambda}\right).$$

– To conclude, we consider  $\widetilde{G}_{J,j}(\Phi_t^{-1}(v)) - \widetilde{G}_{J,j}(v)$  to deal with the problem of case 2.2.

Up to a term  $-\delta^2 \log(\delta r) = \underset{\delta r \rightarrow 0}{o}(\delta r)$ , we can take  $S(r)$  instead of  $\Phi_t(S_\varepsilon(t))$  and 1 instead of the full jacobian  $J(\Phi_t; u)$ . We have  $\varepsilon \langle n_v, \mathbf{e} \rangle = |\langle n_v, \mathbf{e} \rangle|$ . In (146) we take back the previous variables  $u_i = v + v_j$ , this gives

$$\delta r \int_{v \in S_\varepsilon(r)} \sum_{i \neq j} \int_{(u_i, \mathbf{u}') \in S(r) \times B(r)^{J-1}} du_0 \cdots du_J \frac{|\langle n_v, \mathbf{e} \rangle|}{2} \left( -\frac{|\langle n_v, \mathbf{e} \rangle \langle n_v, n_{u_i} \rangle|}{2} \right) G_J(u_\ell \pm \frac{k}{2}).$$

When we sum this term with that of (131), for each  $i \neq j$  we have

$$\begin{aligned} \left| |\langle n_{u_i}, \mathbf{e} \rangle| - |\langle n_v, \mathbf{e} \rangle \langle n_v, n_{u_i} \rangle| \right| &= \left| \varepsilon(u_i) \langle n_{u_i}, \mathbf{e} \rangle - \varepsilon(v) \langle n_v, \mathbf{e} \rangle \times \varepsilon(v) \varepsilon(u_i) \langle n_v, n_{u_i} \rangle \right|, \\ &\leq \min(\sqrt{2} |n_{u_i} - n_v|, 2). \end{aligned}$$

Thus there is no more logarithmic divergence: for  $u = u_j$  and  $v = u_{j-1}$  or  $v = u_{j+1}$ , we use the same method as that for (131) and get

$$\iint_{S(r) \times S(r)} \frac{|n_u - n_v| |\langle n_u, \mathbf{e} \rangle| du dv}{|u - v|^2} \frac{1}{\widetilde{E}(\Lambda)^2 \Lambda^2}.$$

We split the domain in 4:  $S_\varepsilon(r) \times S_{\varepsilon'}(r)$ : the case  $\varepsilon = \varepsilon'$  gives finite number. Indeed if we use spherical coordinates with respect to  $-\frac{\varepsilon r}{2} \mathbf{e}$ , we have  $|n_u - n_v| \leq \frac{|u-v|}{\Lambda}$ , and the integral is

$$\mathcal{O}\left(\int_{\mathbb{S}^2} \frac{du}{\Lambda^2 |u - \mathbf{e}|}\right) = \mathcal{O}\left(\frac{1}{\Lambda^2}\right).$$

The integration over  $S_+(r) \times S_-(r)$  is also finite. To see this we proceed as follows.

For convenience we write  $x := \frac{r}{2\Lambda}$ ,  $\theta_1^0 = \arccos(x)$ ,  $\theta_{-1}^0 = \arccos(-x)$  and  $s(\cdot)$  (resp.  $c(\cdot)$ ) for  $\sin$  (resp.  $\cos$ ). We take spherical coordinates with respect to  $-\varepsilon \frac{r}{2} \mathbf{e}$  for any  $S_\varepsilon(r)$  and obtain:

$$\begin{aligned} &\frac{2\pi}{\Lambda^2} \iiint_{(\theta_1, \theta_{-1}, \phi) \in (0, \theta_1^0) \times (-\pi, \theta_{-1}^0) \times (-\pi, \pi)} \frac{s(\theta_1) s(\theta_{-1}) d\theta_1 d\theta_{-1} d\phi}{(c(\theta_1) - c(\theta_{-1}) - 2x)^2 + s(\theta_{-1})^2 s_\phi^2 + (s(\theta_1) - s(\theta_{-1}) c_\phi)^2} \\ &\lesssim \frac{1}{\Lambda^2} \iiint_{(\theta_1, \theta_{-1}, \phi) \in (0, \theta_1^0) \times (-\pi, \theta_{-1}^0) \times (-\pi, \pi)} \frac{s(\theta_1) s(\theta_{-1}) d\theta_1 d\theta_{-1} d\phi}{(c(\theta_1) - c(\theta_{-1}) - 2x)^2 + c(\theta_{-1})^2 \phi^2} =: \frac{A}{\Lambda^2}. \end{aligned}$$

We write  $\theta_\varepsilon = \theta_\varepsilon^0 - \varepsilon\phi_\varepsilon$ : there holds

$$\begin{aligned} \varepsilon c(\theta_\varepsilon) - x &= x(c(\phi_\varepsilon) - 1) + \sqrt{1-x^2}s(\phi_\varepsilon), \\ x(c(\phi_\varepsilon) - 1) + \sqrt{1-x^2}s(\phi_\varepsilon) &\geq \phi_\varepsilon \left( \frac{2}{\pi} \sqrt{1-x^2} - x \frac{\phi_\varepsilon}{2} \right), \\ &\geq \frac{2\phi_\varepsilon}{\pi} \left( \sqrt{1-x^2} - \frac{\pi}{4} x \arccos(x) \right) \geq \frac{2\phi_\varepsilon \sqrt{1-x^2}}{\pi} \left( 1 - \frac{x\pi}{4} \right) \\ &\geq \frac{2\phi_\varepsilon \sqrt{1-x^2}}{\pi} \left( 1 - \frac{\pi}{4} \right) \geq \frac{\sqrt{1-x^2}\phi_\varepsilon}{\pi} \left( 1 - \frac{\pi}{4} \right). \end{aligned}$$

Thus we have

$$\begin{aligned} A &\lesssim \iint_{\phi_1, \phi_{-1} \in (0, \theta_1^0)} \frac{\sin(\theta_1) d\phi_1 d\phi_{-1}}{\sqrt{1-x^2} \sqrt{\phi_1^2 + \phi_{-1}^2}} \\ &\lesssim \int_{\phi_1 \in (0, \theta_1^0)} \frac{d\phi_1}{\sqrt{1-x^2}} \log \left( 1 + \frac{\arccos(x)}{\phi_1} \right) \\ &\lesssim \int_{\phi \in (0, 1)} \log(1 + \phi^{-1}) d\phi. \end{aligned}$$

**Conclusion** We obtain at last the following upper bound for the terms of cases 2. and 3.:

$$J^2 \frac{(\alpha K)^{J+1} \log(\Lambda)}{\Lambda^2}.$$

It is possible to show that the function  $\partial_r^2 f_\Lambda(x)$  tends to zero as  $|x|$  tends to  $2\Lambda$ , this is proved in the thesis of the author (to appear in 2014).  $\square$

**Alternative  $F_\Lambda$**  In the proof of Theorem 3, one is lead to consider a perturbative self-consistent equation with  $\mathcal{D}^0$  replaced by  $\mathcal{D}^0 + \frac{2}{\lambda} \frac{\mathcal{D}^0}{|\mathcal{D}^0|}$ . In particular we need Lemma 15 below for the proof of Lemma 12. We can write

$$\mathcal{D}^0 + \frac{2}{\lambda} \frac{\mathcal{D}^0}{|\mathcal{D}^0|} = \beta \tilde{w}_0(-i\nabla) + \alpha \cdot \frac{-i\nabla}{|-i\nabla|} \tilde{w}_1(-i\nabla).$$

The formulae are the same with  $g_0, g_1$  replaced by  $\tilde{w}_0, \tilde{w}_1$ , estimates of the same kind hold. The alternative functions are marked with a tilde:  $\tilde{B}_\Lambda$  and  $\tilde{g}_\Lambda$ .

We can easily estimate  $\int_{|x| \geq R} |\mathcal{F}^{-1}(\tilde{F}_\Lambda)(x)| dx$  for  $R \geq 1$ : writing  $f_\Lambda := \mathcal{F}^{-1}(\tilde{F}_\Lambda)$  we have the following Lemma:

**Lemma 15.** *For  $\lambda, \Lambda \gg 1$  we have:*

$$\int_{|x| \geq R} |f_\Lambda(x)| dx \leq \|-\Delta \tilde{F}_\Lambda\|_{L^2} \sqrt{4\pi R^{-1}} = \mathcal{O}(LR^{-1/2}). \quad (147)$$

**Acknowledgment** The author wishes to thank É. Séré for useful discussions. This work was partially supported by the Grant ANR-10-BLAN 0101 of the French Ministry of Research.

## References

- [1] V. Bach, J.-M. Barbaroux, B. Helffer, and H. Siedentop. On the stability of the relativistic electron-positron field. *Comm. Math. Phys.*, 201:445–460, 1998.
- [2] P. Chaix and D. Iracane. From quantum electrodynamics to mean-field theory: I. the Bogoliubov-Dirac-Fock formalism. *J. Phys. B: At. Mol. Opt. Phys.*, 22:3791–3814, 1989.
- [3] L. C. Evans and R. F. Gariepy. *Measure Theory and Fine Properties of Functions*. CRC Press, Boca Rota, 1992.
- [4] R. L. Frank, E. H. Lieb, R. Seiringer, and L. Thomas. Stability and absence of binding for multi-polaron systems. *Publ. Math. IHES*, 113(1):39–67, 2011.

- [5] W. H. Furry. A symmetry theorem in the positron theory. *Phys. Rev.*, 51:125–129, 1937.
- [6] P. Gravejat, M. Lewin, and É. Séré. Renormalization and asymptotic expansion of Dirac’s polarized vacuum. *Comm. Math. Phys.*, 306(1):1–33, 2011.
- [7] Ph. Gravejat, M. Lewin, and É. Séré. Ground state and charge renormalization in a nonlinear model of relativistic atoms. *Comm. Math. Phys.*, 286, 2009.
- [8] C. Hainzl, M. Lewin, and É. Séré. Existence of a stable polarized vacuum in the Bogoliubov-Dirac-Fock approximation. *Comm. Math. Phys.*, 257, 2005.
- [9] C. Hainzl, M. Lewin, and É. Séré. Self-consistent solution for the polarized vacuum in a no-photon QED model. *J. Phys. A: Math and Gen.*, 38(20):4483–4499, 2005.
- [10] C. Hainzl, M. Lewin, and É. Séré. Existence of atoms and molecules in the mean-field approximation of no-photon quantum electrodynamics. *Arch. Rational Mech. Anal.*, 192(3):453–499, 2009.
- [11] C. Hainzl, M. Lewin, and J. P. Solovej. The mean-field approximation in quantum electrodynamics. the no-photon case. *Comm. Pure Appl. Math.*, 60(4):546–596, 2007.
- [12] C. Hainzl and H. Siedentop. Non-perturbative mass and charge renormalization in relativistic no-photon quantum electrodynamics. *Comm. Math. Phys.*, 243(2):241–260, 2003.
- [13] E. Lenzmann and M. Lewin. Minimizers for the Hartree-Fock-Bogoliubov theory of neutron stars and white dwarfs. *Duke Math. Journal*, 152(2):257–315, 2010.
- [14] M. Lewin. Geometric methods for nonlinear many-body quantum systems. *J. Funct. Anal.*, 260:3535–3595, 2011.
- [15] E. H. Lieb. Existence and uniqueness of the minimizing solution of Choquard’s nonlinear equation. *Studies in Applied Mathematics*, 57:93–105, October 1977.
- [16] E. H. Lieb and M. Loss. *Analysis*. AMS, 1997.
- [17] E. H. Lieb and R. Seiringer. *The stability of matter in quantum mechanics*. Cambridge University Press, Cambridge, 2010.
- [18] E. H. Lieb and H. Siedentop. Renormalization of the regularized relativistic electron-positron field. *Comm. Math. Phys.*, 213(3):673–683, 2000.
- [19] E. H. Lieb and B. Simon. The Hartree-Fock theory for coulomb systems. *Comm. Math. Phys.*, 53:185–194, 1977.
- [20] P.-L. Lions. Solutions of Hartree-Fock equations in coulomb system. *Comm. Math. Phys.*, 109:33–97, 1987.
- [21] M. Reed and B. Simon. *Methods of Modern Mathematical Physics*, volume I-II. Academic Press Inc., 1975.
- [22] B. Simon. *Trace Ideals and their Applications*, volume 35 of *London Mathematical Society Lecture Notes Series*. Cambridge University Press, 1079.
- [23] J. Sok. Existence of ground state of an electron in the BDF approximation. To appear in RMP, <http://arxiv.org/abs/1211.3830>, 2012.
- [24] B. Thaller. *The Dirac Equation*. Springer Verlag, 1992.