



HAL
open science

An axiomatic analysis of concordance-discordance relations

Denis Bouyssou, Marc Pirlot

► **To cite this version:**

Denis Bouyssou, Marc Pirlot. An axiomatic analysis of concordance-discordance relations. 2007.
hal-00909473

HAL Id: hal-00909473

<https://hal.science/hal-00909473>

Preprint submitted on 26 Nov 2013

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

CAHIER DU LAMSADE

269

Août 2007

An axiomatic analysis of concordance-discordance relations

Denis Bouyssou, Marc Pirlot

An axiomatic analysis of concordance-discordance relations¹

Denis Bouyssou²
CNRS–LAMSADE

Marc Pirlot³
Faculté Polytechnique de Mons

22 August 2007

¹ Part of this work was accomplished while Denis Bouyssou was visiting the *Service de Mathématique de la Gestion* at the *Université Libre de Bruxelles* (Brussels, Belgium). The warm hospitality of the *Service de Mathématique de la Gestion*, the support of the Belgian *Fonds National de la Recherche Scientifique* are gratefully acknowledged.

² LAMSADE, Université Paris Dauphine, Place du Maréchal de Lattre de Tassigny, F-75775 Paris Cedex 16, France, tel: +33 1 44 05 48 98, fax: +33 1 44 05 40 91, e-mail: bouyssou@lamsade.dauphine.fr

³ Faculté Polytechnique de Mons, 9, rue de Houdain, B-7000 Mons, Belgium, tel: +32 65 374682, fax: +32 65 374689, e-mail: marc.pirlot@fpms.ac.be. **Corresponding author.**

Abstract

Outranking methods propose an original way to build a preference relation between alternatives evaluated on several attributes that has a definite ordinal flavor. Indeed, most of them appeal the concordance / non-discordance principle that leads to declaring that an alternative is “superior” to another, if the coalition of attributes supporting this proposition is “sufficiently important” (concordance condition) and if there is no attribute that “strongly rejects” it (non-discordance condition). Such a way of comparing alternatives is rather natural and does not require a detailed analysis of tradeoffs between the various attributes. However, it is well known that it may produce binary relations that do not possess any remarkable property of transitivity or completeness. This explains why the axiomatic foundations of outranking methods have not been much investigated, which is often seen as one of their important weaknesses. This paper uses conjoint measurement techniques to obtain an axiomatic characterization of preference relations that can be obtained on the basis of the concordance / non-discordance principle. It emphasizes their main distinctive feature, i.e., their very crude way to distinguish various levels of preference differences on each attribute. We focus on outranking methods, such as ELECTRE I, that produce a reflexive relation, interpreted as an “at least as good as” preference relation. The results in this paper may be seen as an attempt to give such outranking methods a sound axiomatic foundation based on conjoint measurement.

Keywords: Multiple criteria analysis, Concordance, Discordance, Outranking methods, Conjoint measurement, Nontransitive preferences

Contents

1	Introduction	1
2	The setting	2
2.1	Binary relations	2
2.2	Notation and definitions	4
2.3	Concordance relations	4
2.4	Concordance-discordance relations	6
3	Background	8
3.1	Conjoint measurement framework	8
3.2	Characterization of concordance relations	10
4	Characterization of reflexive concordance-discordance relations	12
5	Concordance-discordance relations with attribute transitivity	19
5.1	Conjoint measurement framework continued	19
5.2	R-CDR with attribute transitivity	22
6	Strict and non-strict preference models: discordance vs bonus	30
6.1	Strict and non-strict concordance relations	30
6.2	Characterizations of strict and non-strict concordance relations	33
6.3	Strict concordance-discordance relations	37
6.4	The asymmetric part of an R-CDR	40
7	Discussion	41
7.1	Limitations and directions for future work	42
7.2	Relation to the literature	43
7.2.1	The approach using noncompensation	44
7.2.2	The approach of Greco et al. (2001a)	46
	References	48

1 Introduction

Building a preference relation on a set of alternatives evaluated on several attributes is the focal point of Multiple Criteria Decision Analysis (MCDA). The classical approach to achieve this goal consists in building a value function on the set of alternatives (see Keeney and Raiffa 1976). Since this approach requires a detailed analysis of the tradeoffs between attributes and is demanding in terms of time and information, it cannot always be used in practice. For this reason, alternative methods have been proposed among which the *outranking methods* that compare alternatives in a pairwise manner and decide which is preferred on the basis of their evaluations on the several attributes. They do not require a detailed analysis of tradeoffs and mainly rest on “ordinal” considerations (for detailed presentations of these methods, we refer to Bouyssou et al. 2006; Roy 1991; Roy and Bouyssou 1993; Vincke 1992, 1999).

Most outranking methods, including the well known ELECTRE methods (Roy 1968; Roy and Bertier 1973), base the comparison of alternatives on the so-called *concordance / non-discordance* principle. It leads to accepting the proposition that an alternative is “superior” to another if:

- *concordance condition*: the coalition of attributes supporting it is “sufficiently important”,
- *non-discordance condition*: there is no attribute that “strongly rejects” it.

The fact that an alternative is “superior” to another can mean at least two different things. In the ELECTRE methods, superior means “not worse”. Such methods aim at building a reflexive preference relation that is interpreted as an “at least as good as” relation. In general, these relations may lack nice transitivity or completeness properties (on these issues, see Bouyssou 1992, 1996). Our main goal in this paper is to characterize the reflexive binary relations that can be obtained on the basis of the concordance-discordance principle like in the ELECTRE I (Roy 1968) and ELECTRE II methods (Roy and Bertier 1973). In doing so, we build on previous works aiming at characterizing the relations that can be obtained only using the first part of the concordance / non-discordance principle, i.e., the concordance condition (Bouyssou and Pirlot 2005a, 2007).

In other methods, like the TACTIC method (Vansnick 1986), superior means “strictly better than”. Such methods build an asymmetric relation that is interpreted as strict preference. As in the reflexive case, irreflexive (or asymmetric) relations that can be obtained only using the concordance

condition have been previously characterized in Bouyssou and Pirlot (2002a, 2005b). While the correspondence between reflexive and irreflexive concordance relations is straightforward (they are codual of each other), this is no longer the case when the non-discordance condition comes into play. We examine this issue below, which will give us the opportunity to extend and integrate previous results on the subject.

As first presented in Bouyssou et al. (1997), the general strategy followed in this paper is to view outranking relations as a particular case of relations having a representation in the nontransitive decomposable models introduced in Bouyssou and Pirlot (1999, 2002b, 2004a); this was indeed our initial motivation for developing them. This particular case obtains when only a few distinct levels of preference differences are distinguished. This roughly leads to analysing “ordinal aggregation” as an aggregation in which there are only three types of preferences differences: positive, null and negative ones. This paper expands on this simple idea. It is organized as follows. Our setting is introduced in Section 2. Section 3 recalls what is needed of our previous results on the axiomatic characterization of concordance relations. We propose a characterization of reflexive concordance-discordance relations in Section 4. In Section 5, we consider the special case of reflexive concordance-discordance relations with attribute transitivity. In these relations, a transitive preference structure is postulated on each attribute as is the case in most methods used in practice. In Section 6, we investigate how our results may be used, through the use of coduality, to characterize asymmetric outranking relations produced by methods such as TACTIC. A final section discusses our results and positions them with respect to the existing literature on the subject, most notably the “noncompensation approach” developed in Bouyssou and Vansnick (1986) and the approach of Greco et al. (2001a).

With this paper, we more or less put an end to a research program started more than ten years ago (see Bouyssou et al. 1997) and aiming at obtaining axiomatic foundations for outranking methods. In the course of this research, many intermediate results were obtained. This explains the large number of citations to some of our earlier papers. We beg the reader’s leniency for that.

2 The setting

2.1 Binary relations

A *binary relation* \mathcal{R} on a set A is a subset of $A \times A$. We mostly write $a \mathcal{R} b$ instead of $(a, b) \in \mathcal{R}$. A binary relation \mathcal{R} on A is said to be:

- *reflexive* if $[a \mathcal{R} a]$,
- *complete* if $[a \mathcal{R} b \text{ or } b \mathcal{R} a]$,
- *symmetric* if $[a \mathcal{R} b] \Rightarrow [b \mathcal{R} a]$,
- *asymmetric* if $[a \mathcal{R} b] \Rightarrow [Not[b \mathcal{R} a]]$,
- *transitive* if $[a \mathcal{R} b \text{ and } b \mathcal{R} c] \Rightarrow [a \mathcal{R} c]$,
- *Ferrers* if $[(a \mathcal{R} b \text{ and } c \mathcal{R} d) \Rightarrow (a \mathcal{R} d \text{ or } c \mathcal{R} b)]$,
- *semi-transitive* if $[(a \mathcal{R} b \text{ and } b \mathcal{R} c) \Rightarrow (a \mathcal{R} d \text{ or } d \mathcal{R} c)]$

for all $a, b, c, d \in A$.

The *codual* of relation \mathcal{R} on A is the relation \mathcal{R}^{cd} on A defined by:

$$a \mathcal{R}^{cd} b \Leftrightarrow Not[b \mathcal{R} a], \quad (1)$$

for all $a, b \in A$ (we often write $b \mathcal{R} a$ instead of $Not[b \mathcal{R} a]$). Taking the codual of a relation is an involutive operation, i.e., the codual of the codual of a relation is this relation. The codual of a complete relation is an asymmetric relation and conversely. Our use of the term ‘‘codual’’ follows Aleskerov and Monjardet (2002) (Roubens and Vincke 1985, use the term ‘‘dual’’ for the same relation).

The *asymmetric part* of a relation \mathcal{R} on A is the binary relation $\alpha[\mathcal{R}]$ on A such that:

$$\begin{aligned} a \alpha[\mathcal{R}] b &\Leftrightarrow a \mathcal{R} b \text{ and } Not[b \mathcal{R} a] \\ &\Leftrightarrow a \mathcal{R}^{cd} b \text{ and } Not[b \mathcal{R}^{cd} a], \end{aligned} \quad (2)$$

for all $a, b \in A$. Let us note that the asymmetric part of a relation is identical to the asymmetric part of its codual.

The *symmetric part* of a relation \mathcal{R} on A is the binary relation $\sigma[\mathcal{R}]$ on A defined by:

$$a \sigma[\mathcal{R}] b \Leftrightarrow a \mathcal{R} b \text{ and } b \mathcal{R} a \quad (3)$$

The *completion* of a relation \mathcal{R} on A is the binary relation $\overline{\mathcal{R}}$ on A such that:

$$\begin{aligned} a \overline{\mathcal{R}} b &\Leftrightarrow a \mathcal{R} b \text{ or } [Not[a \mathcal{R} b] \text{ and } Not[b \mathcal{R} a]] \\ &\Leftrightarrow a \mathcal{R} b \text{ or } a \mathcal{R}^{cd} b, \end{aligned} \quad (4)$$

for all $a, b \in A$. The completion of a relation is identical to the completion of its codual.

A *weak order* (resp. an *equivalence*) is a complete and transitive (resp. reflexive, symmetric and transitive) binary relation. If \mathcal{R} is an equivalence

on A , A/\mathcal{R} will denote the set of equivalence classes of \mathcal{R} on A . When \mathcal{R} is a weak order, it is clear that $\sigma[\mathcal{R}]$ is an equivalence. We often abuse terminology and speak of the equivalence classes of \mathcal{R} to mean the equivalence classes of $\sigma[\mathcal{R}]$.

A *semiorder* is a complete, Ferrers, semi-transitive binary relation. A *strict semiorder* is an asymmetric Ferrers, semi-transitive binary relation. It is easy to check that the asymmetric part of a semiorder is a strict semiorder. As first observed by Luce (1956), any semiorder \mathcal{R} on A induces a unique weak order \mathcal{R}^{wo} on A that is defined as follows:

$$a \mathcal{R}^{wo} b \text{ if } \forall c \in A, [b \mathcal{R} c \Rightarrow a \mathcal{R} c] \text{ and } [c \mathcal{R} a \Rightarrow c \mathcal{R} b]. \quad (5)$$

2.2 Notation and definitions

In this paper \succsim will always denote a *reflexive* binary relation on a set $X = \prod_{i=1}^n X_i$ with $n \geq 2$. Elements of X will be interpreted as alternatives evaluated on a set $N = \{1, 2, \dots, n\}$ of attributes and \succsim as an “at least as good as” relation between these alternatives. We denote by \succ (resp. \sim) the asymmetric (resp. symmetric) part of \succsim . A similar convention holds when \succsim is starred, superscripted and/or subscripted.

For any nonempty subset J of the set of attributes N , we denote by X_J (resp. X_{-J}) the set $\prod_{i \in J} X_i$ (resp. $\prod_{i \notin J} X_i$). With customary abuse of notation, (x_J, y_{-J}) will denote the element $w \in X$ such that $w_i = x_i$ if $i \in J$ and $w_i = y_i$ otherwise. We sometime omit braces around sets. For instance, when $J = \{i\}$ we write X_{-i} and (x_i, y_{-i}) .

We say that attribute $i \in N$ is *influential* (for \succsim) if there are $x_i, y_i, z_i, w_i \in X_i$ and $x_{-i}, y_{-i} \in X_{-i}$ such that $(x_i, x_{-i}) \succsim (y_i, y_{-i})$ and $(z_i, x_{-i}) \not\succeq (w_i, y_{-i})$ and *degenerate* otherwise. A degenerate attribute has no influence whatsoever on the comparison of the elements of X and may be suppressed from N . As in Bouyssou and Pirlot (2005a), in order to avoid unnecessary minor complications, we suppose henceforth that *all attributes in N are influential*.

2.3 Concordance relations

In Bouyssou and Pirlot (2005a), we have given a general definition of concordance relations and have shown that the preference relations produced by most of outranking methods fit into this framework, provided that no veto effect occurs (i.e., when only the concordance part of the concordance / non-discordance principle is used). We recall this definition below, adding the term “reflexive” to it, since we shall consider irreflexive (or asymmetric) concordance-discordance relations in Section 6.

Definition 1 (Reflexive concordance relation)

Let \succsim be a reflexive binary relation on $X = \prod_{i=1}^n X_i$. We say that \succsim is a reflexive concordance relation (or, more briefly, that \succsim is an R-CR) if there are:

- a complete binary relation S_i on each X_i ($i = 1, 2, \dots, n$),
- a binary relation \supseteq between subsets of N having N for union that is monotonic w.r.t. inclusion, i.e., for all $A, B, C, D \subseteq N$ such that $A \cup B = N$ and $C \cup D = N$,

$$[A \supseteq B, C \supseteq A, B \supseteq D] \Rightarrow C \supseteq D, \quad (6)$$

such that, for all $x, y \in X$,

$$x \succsim y \Leftrightarrow S(x, y) \supseteq S(y, x), \quad (7)$$

where $S(x, y) = \{i \in N : x_i S_i y_i\}$. We say that $\langle \supseteq, S_i \rangle$ is a representation of \succsim as an R-CR. Throughout the paper, P_i (resp. I_i) will denote the asymmetric (resp. symmetric) part of S_i . Let $A, B \subseteq N$ such that $A \cup B = N$. We abbreviate $[A \supseteq B \text{ and } B \supseteq A]$ as $A \trianglelefteq B$ and $[A \supseteq B \text{ and } \text{Not}[B \supseteq A]]$ as $A \triangleright B$.

Let us illustrate this definition by some classic examples.

Example 2 (Simple majority preferences)

The binary relation \succsim is a simple majority preference relation if there is a weak order S_i on each X_i such that:

$$x \succsim y \Leftrightarrow |\{i \in N : x_i S_i y_i\}| \geq |\{i \in N : y_i S_i x_i\}|.$$

A simple majority preference relation is easily seen to be an R-CR defining \supseteq letting, for all $A, B \subseteq N$ such that $A \cup B = N$,

$$A \supseteq B \Leftrightarrow |A| \geq |B|. \quad \diamond$$

Example 3 (Semiordeed weighted majority)

The binary relation \succsim is a semiordeed weighted majority preference relation if there are a real number $\varepsilon \geq 0$ and, for all $i \in N$,

- a semiordeed S_i on X_i ,
- a real number $w_i > 0$,

such that:

$$x \succsim y \Leftrightarrow \sum_{i \in S(x,y)} w_i \geq \sum_{j \in S(y,x)} w_j - \varepsilon.$$

Such a relation is easily seen to be a complete R-CR defining \supseteq letting, for all $A, B \subseteq N$ such that $A \cup B = N$:

$$A \supseteq B \Leftrightarrow \sum_{i \in A} w_i \geq \sum_{j \in B} w_j - \varepsilon. \quad \diamond$$

2.4 Concordance-discordance relations

ELECTRE I builds a reflexive concordance relation that is subsequently “censored” by imposing on it the non-discordance condition. This is illustrated below.

Example 4 (ELECTRE I, Roy 1968)

The binary relation \succsim is an ELECTRE I preference relation if there are a real number $s \in [1/2, 1]$ and, for all $i \in N$,

- a semiorder S_i on X_i ,
- a strict semiorder V_i on X_i such that $V_i \subseteq P_i$,
- a positive real number $w_i > 0$,

such that, for all $x, y \in X$,

$$x \succsim y \Leftrightarrow \frac{\sum_{i \in S(x,y)} w_i}{\sum_{j \in N} w_j} \geq s \text{ and } V(y, x) = \emptyset,$$

where $V(y, x) = \{i \in N : y_i V_i x_i\}$.

The “concordance part” of the ELECTRE I preference is what we obtain if there is no veto or, in other words, if the relation V_i is empty. The concordance part of an ELECTRE I preference relation is easily seen to be an R-CR defining \supseteq letting, for all $A, B \subseteq N$ such that $A \cup B = N$,

$$A \supseteq B \Leftrightarrow \frac{\sum_{i \in A} w_i}{\sum_{j \in N} w_j} \geq s.$$

In practice, the relations S_i and V_i are usually obtained as follows, using a real-valued function u_i defined on X_i and a pair of positive thresholds pt_i and vt_i , with $pt_i \leq vt_i$. We have, for all $x_i, y_i \in X_i$,

$$x_i S_i y_i \Leftrightarrow u_i(x_i) \geq u_i(y_i) - pt_i \tag{8}$$

$$x_i V_i y_i \Leftrightarrow u_i(x_i) > u_i(y_i) + vt_i. \tag{9}$$

The set $V(y, x)$ is then the set of attributes on which $u_i(y_i) > u_i(x_i) + vt_i$, i.e., those on which y is “so much better” than x that it is excluded to conclude that $x \succsim y$. An intuitive interpretation of the ELECTRE I preference relation could therefore run as follows: x is at least as good as y if there is a majority of attributes on which x is at least as good as y (the attributes belonging to $S(x, y)$ should be “sufficiently important”) and there is no attribute on which x is too much worse than y (the set $V(y, x)$ should be empty). \diamond

ELECTRE I is an example of what we call a *reflexive concordance-discordance relation*, a precise definition of which follows.

Definition 5 (Reflexive concordance-discordance relation)

Let \succsim be a reflexive binary relation on $X = \prod_{i=1}^n X_i$. We say that \succsim is a reflexive concordance-discordance relation (or, more briefly, that \succsim is an R-CDR) if there are:

- a complete binary relation S_i on each X_i ($i = 1, 2, \dots, n$) (with asymmetric part P_i and symmetric part I_i),
- an asymmetric binary relation V_i on each X_i ($i = 1, 2, \dots, n$) such that $V_i \subseteq P_i$,
- a binary relation \supseteq between subsets of N having N for union that is monotonic w.r.t. inclusion, i.e., such that (6) holds,

such that, for all $x, y \in X$,

$$x \succsim y \Leftrightarrow [S(x, y) \supseteq S(y, x) \text{ and } V(y, x) = \emptyset], \quad (10)$$

where $S(x, y) = \{i \in N : x_i S_i y_i\}$ and $V(y, x) = \{i \in N : y_i V_i x_i\}$. We say that $\langle \supseteq, S_i, V_i \rangle$ is a representation of \succsim as an R-CDR.

In the ELECTRE I method, when relations S_i and V_i are defined respectively using equations (8) and (9), we have $V_i \subseteq P_i$. It is also clear that S_i is a semiorder. Similarly, the relation V_i is a strict semiorder that is the asymmetric part of a semiorder U_i that obtains as the codual of V_i , i.e., is such that: for all $x_i, y_i \in X_i$,

$$x_i U_i y_i \text{ iff } \text{Not}[y_i V_i x_i] \text{ iff } u_i(x_i) \geq u_i(y_i) - vt_i.$$

Since $vt_i \geq pt_i$, it is clear that $S_i \subseteq U_i$ (so that their asymmetric parts are related by the opposite inclusion: $P_i \supseteq V_i$). We say that (S_i, U_i) form a *nested chain* of semiorders (ordered by inclusion).

The nested chain (S_i, U_i) has an additional crucial property. Using (5), consider the weak order S_i^{wo} associated to the semiorder S_i and the weak order U_i^{wo} associated to the semiorder U_i . From (8) and (9), it is clear that their intersection $T_i = U_i^{wo} \cap S_i^{wo}$ is again a weak order. This additional property transforms the nested chain of semiorders (S_i, U_i) into a *homogeneous* nested chain of semiorders.

Nested chains of two semiorders were first studied by Cozzens and Roberts (1982). They appear as a particular case in a more general framework, analyzed in Doignon et al. (1988).

Remark 6

It is easy to add a non-discordance condition to the definition of the relations in Examples 2 and 3, yielding respectively a simple majority preference relation with veto and a semiordered weighted majority preference with veto. •

3 Background

In this section, we briefly present the axiomatic framework and the previously obtained characterization of reflexive concordance relations within this framework.

3.1 Conjoint measurement framework

Concordance relations rely on comparing alternatives in pairwise manner on the basis of preference differences on each attribute. The relations defined below were introduced in Bouyssou and Pirlot (2002b). They will play a fundamental rôle in the sequel.

Definition 7 (Relations comparing preference differences)

Let \succsim be a binary relation on a set $X = \prod_{i=1}^n X_i$. We define the binary relations \succsim_i^* and \succsim_i^{**} on X_i^2 letting, for all $x_i, y_i, z_i, w_i \in X_i$,

$$\begin{aligned} (x_i, y_i) \succsim_i^* (z_i, w_i) &\Leftrightarrow \\ &[\text{for all } a_{-i}, b_{-i} \in X_{-i}, (z_i, a_{-i}) \succ (w_i, b_{-i}) \Rightarrow (x_i, a_{-i}) \succ (y_i, b_{-i})], \\ (x_i, y_i) \succsim_i^{**} (z_i, w_i) &\Leftrightarrow [(x_i, y_i) \succsim_i^* (z_i, w_i) \text{ and } (w_i, z_i) \succsim_i^* (y_i, x_i)]. \end{aligned}$$

The definition of \succsim_i^* suggests that $(x_i, y_i) \succsim_i^* (z_i, w_i)$ can be interpreted as saying that the preference difference between x_i and y_i is at least as large as the preference difference between z_i and w_i . The definition of \succsim_i^* does not imply that the two “opposite” differences (x_i, y_i) and (y_i, x_i) are linked.

This is at variance with the intuition concerning preference differences and motivates the introduction of the relation \succsim_i^{**} . By construction, \succsim_i^* and \succsim_i^{**} are always reflexive and transitive. The following axioms are related to further important properties of relations \succsim_i^* and \succsim_i^{**} .

Definition 8 (Conditions *RC1* and *RC2*)

Let \succsim be a binary relation on a set $X = \prod_{i=1}^n X_i$. This relation is said to satisfy:

$$RC1_i \text{ if } \left. \begin{array}{l} (x_i, a_{-i}) \succsim (y_i, b_{-i}) \\ \text{and} \\ (z_i, c_{-i}) \succsim (w_i, d_{-i}) \end{array} \right\} \Rightarrow \left\{ \begin{array}{l} (x_i, c_{-i}) \succsim (y_i, d_{-i}) \\ \text{or} \\ (z_i, a_{-i}) \succsim (w_i, b_{-i}), \end{array} \right.$$

$$RC2_i \text{ if } \left. \begin{array}{l} (x_i, a_{-i}) \succsim (y_i, b_{-i}) \\ \text{and} \\ (y_i, c_{-i}) \succsim (x_i, d_{-i}) \end{array} \right\} \Rightarrow \left\{ \begin{array}{l} (z_i, a_{-i}) \succsim (w_i, b_{-i}) \\ \text{or} \\ (w_i, c_{-i}) \succsim (z_i, d_{-i}), \end{array} \right.$$

for all $x_i, y_i, z_i, w_i \in X_i$ and all $a_{-i}, b_{-i}, c_{-i}, d_{-i} \in X_{-i}$. We say that \succsim satisfies *RC1* (resp. *RC2*) if it satisfies $RC1_i$ (resp. $RC2_i$) for all $i \in N$.

Condition $RC1_i$ is equivalent to requiring that any two preference differences are comparable in terms of \succsim_i^* . Condition $RC2_i$ imposes a “mirror effect” on the comparison of preference differences. This is summarized in the following:

Lemma 9 (Bouyssou and Pirlot 2002b, Lemma 1)

1. $RC1_i \Leftrightarrow [\succsim_i^* \text{ is complete}]$.
2. $RC2_i \Leftrightarrow$
[for all $x_i, y_i, z_i, w_i \in X_i, (x_i, y_i) \not\sucsim_i^* (z_i, w_i) \Rightarrow (y_i, x_i) \succsim_i^* (w_i, z_i)$].
3. $[RC1_i \text{ and } RC2_i] \Leftrightarrow [\succsim_i^{**} \text{ is complete}]$.
4. In the class of reflexive relations, *RC1* and *RC2* are independent conditions.

We consider binary relations \succsim on X that can be represented in the following model introduced in Bouyssou and Pirlot (2002b):

$$x \succsim y \Leftrightarrow F(p_1(x_1, y_1), p_2(x_2, y_2), \dots, p_n(x_n, y_n)) \geq 0, \quad (M)$$

where p_i are real-valued functions on X_i^2 that are *skew symmetric* (i.e., such that $p_i(x_i, y_i) = -p_i(y_i, x_i)$, for all $x_i, y_i \in X_i$) and F is a real-valued function on $\prod_{i=1}^n p_i(X_i^2)$ being *nondecreasing* in all its arguments and such that, abusing notation, $F(\mathbf{0}) \geq 0$. This model extends the additive nontransitive model studied in Fishburn (1990, 1991), in which F is a sum.

It is useful to interpret p_i as a function measuring preference differences between levels on attribute $i \in N$. The fact that the functions p_i are supposed to be skew symmetric means that the preference difference between x_i and y_i is the opposite of the preference difference between y_i and x_i , which seems a reasonable hypothesis. In order to compare alternatives x and y , model (M) proceeds as follows. On each attribute $i \in N$, the preference difference between x_i and y_i is measured using p_i . The synthesis of these preference differences is performed applying the function F to the $p_i(x_i, y_i)$'s. We then conclude that $x \succsim y$ when this synthesis is nonnegative. Given this interpretation, it seems reasonable to suppose that F is nondecreasing in each of its arguments. The fact that $F(\mathbf{0}) \geq 0$ simply means that the synthesis of null preference differences on each attribute should be nonnegative; this ensures that \succsim will be reflexive.

For finite or countably infinite sets, conditions *RC1* and *RC2* together with reflexivity are all that is needed in order to characterize model (M). We have:

Theorem 10 (Bouyssou and Pirlot 2002b, Theorem 1)

*Let \succsim be a binary relation on $X = \prod_{i=1}^n X_i$. If, for all $i \in N$, X_i^2 / \sim_i^{**} is finite or countably infinite then \succsim has a representation (M) if and only if (iff) it is reflexive and satisfies *RC1* and *RC2*.*

The extension of this result to the general case is easy but will not be useful here.

3.2 Characterization of concordance relations

The general strategy used in Bouyssou and Pirlot (2005a, 2007) to characterize concordance relations is to use model (M) as a building block adding additional conditions ensuring that all functions p_i take at most three distinct values. Hence, the “ordinal” character of the aggregation at work in concordance relations is modelled by saying that in an ordinal method, there can be at most three distinct types of preference differences: positive, null and negative ones. The additional conditions used in Bouyssou and Pirlot (2007) to capture concordance relations are as follows.

Definition 11 (Conditions *M1* and *M2*)

Let \succsim be a binary relation on a set $X = \prod_{i=1}^n X_i$. This relation is said to

satisfy:

$$\begin{aligned}
 M1_i \text{ if } \left. \begin{array}{l} (x_i, a_{-i}) \succsim (y_i, b_{-i}) \\ \text{and} \\ (z_i, c_{-i}) \succsim (w_i, d_{-i}) \end{array} \right\} &\Rightarrow \left\{ \begin{array}{l} (y_i, a_{-i}) \succ (x_i, b_{-i}) \\ \text{or} \\ (w_i, a_{-i}) \succ (z_i, b_{-i}) \\ \text{or} \\ (x_i, c_{-i}) \succ (y_i, d_{-i}), \end{array} \right. \\
 M2_i \text{ if } \left. \begin{array}{l} (x_i, a_{-i}) \succsim (y_i, b_{-i}) \\ \text{and} \\ (y_i, c_{-i}) \succsim (x_i, d_{-i}) \end{array} \right\} &\Rightarrow \left\{ \begin{array}{l} (y_i, a_{-i}) \succ (x_i, b_{-i}) \\ \text{or} \\ (z_i, a_{-i}) \succ (w_i, b_{-i}) \\ \text{or} \\ (z_i, c_{-i}) \succ (w_i, d_{-i}), \end{array} \right.
 \end{aligned}$$

for all $x_i, y_i, z_i, w_i \in X_i$ and all $a_{-i}, b_{-i}, c_{-i}, d_{-i} \in X_{-i}$. We say that $M1$ (resp. $M2$) holds if $M1_i$ (resp. $M2_i$) holds for all $i \in N$.

It is not difficult to see that $M1_i$ and $M2_i$ drastically limit the possibility of distinguishing several classes of preference differences on each attribute using \succsim_i^* . Suppose for instance that the premises of $M1_i$ holds and that its first conclusion is false. Because $(x_i, a_{-i}) \succsim (y_i, b_{-i})$ and $(y_i, a_{-i}) \not\succeq (x_i, b_{-i})$, it is clear that the preference difference (y_i, x_i) is not larger (w.r.t. the relation \succsim_i^*) than its opposite preference difference (x_i, y_i) . In an R-CR, this can only happen if the difference (x_i, y_i) is “positive” and, thus, the difference (y_i, x_i) is “negative”. But if the difference (x_i, y_i) is “positive”, there cannot exist a difference larger than (x_i, y_i) . Therefore if $(z_i, c_{-i}) \succsim (w_i, d_{-i})$, we should obtain $(x_i, c_{-i}) \succsim (y_i, d_{-i})$. This is what is required by $M1_i$ (disregarding its second possible conclusion that only ensures that the condition will be independent from the ones used to characterize model (M)). Condition $M2_i$ has a dual interpretation: if (y_i, x_i) is not larger than its opposite preference difference (x_i, y_i) then there can be no difference smaller than (y_i, x_i) .

Remark 12

In Bouyssou and Pirlot (2005a), we used two conditions that are stronger than $M1_i$ and $M2_i$. It will prove useful to introduce them. The relation \succsim is said to satisfy:

$$UC_i \text{ if } \left. \begin{array}{l} (x_i, a_{-i}) \succsim (y_i, b_{-i}) \\ \text{and} \\ (z_i, c_{-i}) \succsim (w_i, d_{-i}) \end{array} \right\} \Rightarrow \left\{ \begin{array}{l} (y_i, a_{-i}) \succ (x_i, b_{-i}) \\ \text{or} \\ (x_i, c_{-i}) \succ (y_i, d_{-i}), \end{array} \right.$$

$$LC_i \text{ if } \left. \begin{array}{l} (x_i, a_{-i}) \succsim (y_i, b_{-i}) \\ \text{and} \\ (y_i, c_{-i}) \succsim (x_i, d_{-i}) \end{array} \right\} \Rightarrow \left\{ \begin{array}{l} (y_i, a_{-i}) \succsim (x_i, b_{-i}) \\ \text{or} \\ (z_i, c_{-i}) \succsim (w_i, d_{-i}), \end{array} \right.$$

for all $x_i, y_i, z_i, w_i \in X_i$ and all $a_{-i}, b_{-i}, c_{-i}, d_{-i} \in X_{-i}$.

Bouyssou and Pirlot (2005a, Lemma 16) and Bouyssou and Pirlot (2007, Lemma 11) show that:

1. $UC_i \Rightarrow M1_i$.
2. $LC_i \Rightarrow M2_i$.
3. $UC_i \Leftrightarrow [(y_i, x_i) \succsim_i^* (x_i, y_i) \Rightarrow (x_i, y_i) \succsim_i^* (z_i, w_i), \text{ for all } x_i, y_i, z_i, w_i \in X_i]$.
4. $LC_i \Leftrightarrow [(y_i, x_i) \succsim_i^* (x_i, y_i) \Rightarrow (z_i, w_i) \succsim_i^* (y_i, x_i), \text{ for all } x_i, y_i, z_i, w_i \in X_i]$.

The problem with UC and LC is that they interact with $RC1$ and $RC2$ (see Bouyssou and Pirlot 2005a, Lemma 16). This explains our use of the slightly more involved conditions $M1$ and $M2$. •

The following appears as Theorem 13 in Bouyssou and Pirlot (2007).

Theorem 13

Let \succsim be a binary relation on $X = \prod_{i=1}^n X_i$. Then \succsim is an R-CR iff it is reflexive and satisfies RC1, RC2, M1 and M2. In the class of reflexive relations, conditions RC1, RC2, M1 and M2 are independent.

The above result demonstrates that the framework of model (M) is adequate for analyzing R-CR. We show below that this is also the case for R-CDR.

4 Characterization of reflexive concordance-discordance relations

The definition of a reflexive concordance-discordance relation has been given in Section 2.4 (Definition 5). In an R-CDR the concordance condition is tempered by a non-discordance condition forbidding to have $x \succsim y$ when there is one attribute on which y is far better than x . In terms of preference differences, this adds the possibility of having two categories of negative differences: normal ones (acting as in an R-CR) and intolerable ones (corresponding to a veto).

The following lemma shows that an R-CDR is always a particular case of model (M) (i.e., conditions *RC1*, *RC2* hold) and that, as in an R-CR, there can be only one type of positive differences (condition *M1* holds).

Lemma 14

If \succsim is an R-CDR then it satisfies RC1, RC2 and M1.

PROOF

[*RC1*] Let $\langle \triangleright, S_i, V_i \rangle$ be a representation of \succsim . Suppose that $(x_i, a_{-i}) \succsim (y_i, b_{-i})$, $(z_i, c_{-i}) \succsim (w_i, d_{-i})$. This implies that $\text{Not}[y_i V_i x_i]$ and $\text{Not}[w_i V_i z_i]$. Suppose that $y_i P_i x_i$. The definition of an R-CDR implies that $(z_i, a_{-i}) \succsim (w_i, b_{-i})$. If $x_i P_i y_i$, the definition of an R-CDR implies that $(x_i, c_{-i}) \succsim (y_i, d_{-i})$. Suppose now that $x_i I_i y_i$. If $z_i S_i w_i$, we obtain, using the definition of an R-CDR, $(z_i, a_{-i}) \succsim (w_i, b_{-i})$. If $w_i P_i z_i$, using the definition of an R-CDR leads to $(x_i, c_{-i}) \succsim (y_i, d_{-i})$.

[*RC2*] Suppose that $(x_i, a_{-i}) \succsim (y_i, b_{-i})$, $(y_i, c_{-i}) \succsim (x_i, d_{-i})$. This implies that $\text{Not}[y_i V_i x_i]$ and $\text{Not}[x_i V_i y_i]$.

Suppose that $x_i S_i y_i$. If $z_i S_i w_i$, we know that $\text{Not}[w_i V_i z_i]$ and the definition of an R-CDR leads to $(z_i, a_{-i}) \succsim (w_i, b_{-i})$. If $w_i S_i z_i$, we know that $\text{Not}[z_i V_i w_i]$. The definition of an R-CDR leads to $(w_i, c_{-i}) \succsim (z_i, d_{-i})$. The proof is similar if we suppose that $y_i S_i x_i$.

[*M1*] Suppose that $(x_i, a_{-i}) \succsim (y_i, b_{-i})$ and $(z_i, c_{-i}) \succsim (w_i, d_{-i})$. This implies that $\text{Not}[y_i V_i x_i]$ and $\text{Not}[w_i V_i z_i]$.

If $y_i S_i x_i$, we know that $\text{Not}[x_i V_i y_i]$ so that we have $(y_i, a_{-i}) \succsim (x_i, b_{-i})$, using the definition of an R-CDR. If $x_i P_i y_i$, we know that $\text{Not}[y_i V_i x_i]$ so that we have $(x_i, c_{-i}) \succsim (y_i, d_{-i})$, using the definition of an R-CDR. \square

With Lemma 14 at hand, it is clear, in view of Theorem 13 that characterizes R-CR, that condition *M2* is, in general, not satisfied by R-CDR. This is due to the possible presence of a veto effect: it may happen that the premise of *M2_i* is fulfilled while the conclusion is false because the pair (w_i, z_i) belongs to V_i . This motivates the introduction of the following condition that is satisfied by R-CDR as we shall show in the sequel. The work of Greco et al. (2001a) has been inspiring in devising it.

Definition 15

Let \succsim be a binary relation on a set $X = \prod_{i=1}^n X_i$. This relation is said to satisfy:

$$M3_i \text{ if } \left. \begin{array}{l} (x_i, a_{-i}) \succsim (y_i, b_{-i}) \\ \text{and} \\ (y_i, c_{-i}) \succsim (x_i, d_{-i}) \\ \text{and} \\ (z_i, e_{-i}) \succsim (w_i, f_{-i}) \end{array} \right\} \Rightarrow \left\{ \begin{array}{l} (y_i, a_{-i}) \succsim (x_i, b_{-i}) \\ \text{or} \\ (z_i, a_{-i}) \succsim (w_i, b_{-i}) \\ \text{or} \\ (z_i, c_{-i}) \succsim (w_i, d_{-i}), \end{array} \right.$$

for all $x_i, y_i, z_i, w_i \in X_i$ and all $a_{-i}, b_{-i}, c_{-i}, d_{-i}, e_{-i}, f_{-i} \in X_{-i}$. We say that \succsim satisfies $M3$ if it satisfies $M3_i$ for all $i \in N$.

The meaning of $M3_i$ can be intuitively expressed as follows. Suppose that the three premises of $M3_i$ hold and that its first conclusion is false (we disregard the second possible conclusion of $M3_i$, the rôle of which is to ensure that the condition is independent from the ones needed to characterize model (M)). This implies that the difference (y_i, x_i) is “negative” since it is not larger than its opposite difference (x_i, y_i) . In an R-CDR a negative preference difference is the smallest among all preference difference that do not correspond to a veto. Because $(z_i, e_{-i}) \succsim (w_i, f_{-i})$, we know that $\text{Not}[w_i V_i z_i]$. Hence, the difference (z_i, w_i) is not smaller than the difference (y_i, x_i) , so that $(y_i, c_{-i}) \succsim (x_i, d_{-i})$ implies $(z_i, c_{-i}) \succsim (w_i, d_{-i})$. The following formalizes the above reasoning showing that condition $M3$ holds in an R-CDR.

Lemma 16

If \succsim is an R-CDR then it satisfies $M3$.

PROOF

Suppose that $(x_i, a_{-i}) \succsim (y_i, b_{-i})$, $(y_i, c_{-i}) \succsim (x_i, d_{-i})$ and $(z_i, e_{-i}) \succsim (w_i, f_{-i})$. By construction, we know that $\text{Not}[x_i V_i y_i]$, $\text{Not}[y_i V_i x_i]$ and $\text{Not}[w_i V_i z_i]$. If $y_i S_i x_i$, the definition of an R-CDR implies $(y_i, a_{-i}) \succsim (x_i, b_{-i})$. If $x_i P_i y_i$, the definition of an R-CDR implies $(z_i, c_{-i}) \succsim (w_i, d_{-i})$. Hence $M3_i$ is fulfilled. \square

The following lemma analyzes the structure of the relation \succsim_i^* under $RC1$, $RC2$, $M1$ and $M3$. It shows that when there are two distinct types of negative differences, the smallest ones can be interpreted as a veto.

Lemma 17

Let \succsim be a binary relation on $X = \prod_{i=1}^n X_i$. If \succsim satisfies $RC1$, $RC2$, $M1$ and $M3$, then, for all $x_i, y_i, z_i, w_i, r_i, s_i \in X_i$,

1. $(x_i, y_i) \succ_i^* (y_i, x_i) \Rightarrow (x_i, y_i) \succsim_i^* (z_i, w_i)$.
2. $[(x_i, y_i) \succ_i^* (y_i, x_i) \succ_i^* (z_i, w_i)] \Rightarrow (r_i, s_i) \succsim_i^* (z_i, w_i)$. Furthermore, we have $(z_i, a_{-i}) \not\prec (w_i, b_{-i})$, for all $a_{-i}, b_{-i} \in X_{-i}$.

PROOF

Part 1 follows from results obtained in Bouyssou and Pirlot (2007): by Lemma 11.3 in this paper, we know that $RC2_i$ and $M1_i$ imply UC_i , i.e., that $(y_i, x_i) \not\prec_i^* (x_i, y_i) \Rightarrow (x_i, y_i) \succsim_i^* (z_i, w_i)$, for all $x_i, y_i, z_i, w_i \in X_i$ (see Remark 12). Since $RC1_i$ is equivalent to the fact that \succsim_i^* is complete, $(y_i, x_i) \not\prec_i^* (x_i, y_i)$ implies $(x_i, y_i) \succ_i^* (y_i, x_i)$, which proves Part 1.

Part 2. Suppose that, for some $x_i, y_i, z_i, w_i, r_i, s_i \in X_i$, we have $(x_i, y_i) \succ_i^* (y_i, x_i) \succ_i^* (z_i, w_i)$ and $(z_i, w_i) \succ_i^* (r_i, s_i)$. This implies $(x_i, a_{-i}) \succsim (y_i, b_{-i})$, $(y_i, a_{-i}) \not\succeq (x_i, b_{-i})$, $(y_i, c_{-i}) \succsim (x_i, d_{-i})$, $(z_i, c_{-i}) \not\succeq (w_i, d_{-i})$ and $(z_i, e_{-i}) \succsim (w_i, f_{-i})$, $(r_i, e_{-i}) \not\succeq (s_i, f_{-i})$, for some $a_{-i}, b_{-i}, c_{-i}, d_{-i}, e_{-i}, f_{-i} \in X_{-i}$. Using $M3_i$, $(x_i, a_{-i}) \succsim (y_i, b_{-i})$, $(y_i, c_{-i}) \succsim (x_i, d_{-i})$, $(z_i, e_{-i}) \succsim (w_i, f_{-i})$, $(y_i, a_{-i}) \not\succeq (x_i, b_{-i})$ and $(z_i, c_{-i}) \not\succeq (w_i, d_{-i})$ imply $(z_i, a_{-i}) \succsim (w_i, b_{-i})$. This leads to $(y_i, c_{-i}) \succsim (x_i, d_{-i})$, $(z_i, c_{-i}) \not\succeq (w_i, d_{-i})$, $(z_i, a_{-i}) \succsim (w_i, b_{-i})$ and $(y_i, a_{-i}) \not\succeq (x_i, b_{-i})$, contradicting the completeness of \succsim_i^* that follows from $RC1_i$. Note that the contradiction is obtained as soon as $(z_i, e_{-i}) \succsim (w_i, f_{-i})$, for some $e_{-i}, f_{-i} \in X_{-i}$. This proves the second part of the assertion. \square

Remark 18

It is clear that $M2_i$ implies $M3_i$. Indeed, $M3_i$ is obtained from $M2_i$ by adding the premise $(z_i, e_{-i}) \succsim (w_i, f_{-i})$. Example 38 below will show that there are reflexive relations satisfying $RC1$, $RC2$, $M1$ on all attributes and $M3_i$ on all but one attribute. In view of Theorem 13, this shows that conditions $RC1$, $RC2$, $M1$ and $M3$ are independent in the class of reflexive relations. \bullet

For relations satisfying $RC1$, $RC2$, $M1$ and $M3$, we define a relation S_i on X_i , using \succsim_i^* for comparing each difference (x_i, y_i) to its ‘‘opposite’’ difference (y_i, x_i) . Similarly, we introduce the veto relation V_i when a difference is strictly smaller (in terms of \succ_i^*) than two opposite differences. The properties of these relations are studied in the following lemma.

Lemma 19

Let \succsim be a binary relation on $X = \prod_{i=1}^n X_i$ satisfying $RC1$, $RC2$, $M1$ and $M3$.

1. The relation S_i on X_i defined letting $x_i S_i y_i$ iff $(x_i, y_i) \succsim_i^* (y_i, x_i)$ is complete. Furthermore, letting P_i denote the asymmetric part of S_i , we have that $z_i P_i w_i$ and $x_i P_i y_i$ imply $(z_i, w_i) \sim_i^* (x_i, y_i)$.
2. If $x_i I_i y_i$ and $z_i I_i w_i$, where I_i is the symmetric part of S_i , then $(x_i, y_i) \sim_i^* (z_i, w_i) \sim_i^* (y_i, x_i) \sim_i^* (w_i, z_i) \sim_i^* (a_i, a_i)$ for all $a_i \in X_i$.
3. Define the relation V_i on X_i letting $x_i V_i y_i$ if, for some $z_i, w_i \in X_i$, $(z_i, w_i) \succ_i^* (w_i, z_i) \succ_i^* (y_i, x_i)$. We have:
 - (a) $V_i \subseteq P_i$,
 - (b) $[z_i V_i w_i \text{ and } x_i V_i y_i] \Rightarrow (w_i, z_i) \sim_i^* (y_i, x_i)$,
 - (c) $[x_i P_i y_i, z_i P_i w_i, \text{Not}[x_i V_i y_i] \text{ and } \text{Not}[z_i V_i w_i]] \Rightarrow (y_i, x_i) \sim_i^* (w_i, z_i)$.

PROOF

Part 1. We have $x_i S_i y_i$ iff $(x_i, y_i) \succsim_i^* (y_i, x_i)$. Since \succsim_i^* is complete due to *RC1*, it follows that S_i is complete. Suppose now that $z_i P_i w_i$ and $x_i P_i y_i$. Using Lemma 17, we have $(z_i, w_i) \succsim_i^* (x_i, y_i)$ and $(x_i, y_i) \succsim_i^* (z_i, w_i)$ so that $(z_i, w_i) \sim_i^* (x_i, y_i)$.

Part 2. Using the definition of S_i , $x_i I_i y_i$ and $z_i I_i w_i$ is equivalent to $(x_i, y_i) \sim_i^* (y_i, x_i)$ and $(z_i, w_i) \sim_i^* (w_i, z_i)$. The conclusion follows from *RC2*.

Part 3a. We have $x_i V_i y_i$ iff $(z_i, w_i) \succ_i^* (w_i, z_i) \succ_i^* (y_i, x_i)$. Suppose that $\text{Not}[x_i P_i y_i]$ so that $(y_i, x_i) \succsim_i^* (x_i, y_i)$. Using *RC1* and *RC2*, it is easy to check that $(z_i, w_i) \succ_i^* (w_i, z_i)$ implies $(z_i, w_i) \succsim_i^* (a_i, a_i) \succsim_i^* (w_i, z_i)$, for all $a_i \in X_i$. We therefore obtain $(a_i, a_i) \succ_i^* (y_i, x_i) \succsim_i^* (x_i, y_i)$. This contradicts *RC2*.

Part 3b. Suppose that $x_i V_i y_i$ and $z_i V_i w_i$. Using Lemma 17.2, we have $(w_i, z_i) \succsim_i^* (y_i, x_i)$ and $(y_i, x_i) \succsim_i^* (w_i, z_i)$ so that $(w_i, z_i) \sim_i^* (y_i, x_i)$.

Part 3c. By definition, we have $(x_i, y_i) \succ_i^* (y_i, x_i)$ and $(z_i, w_i) \succ_i^* (w_i, z_i)$. Suppose that $(y_i, x_i) \succ_i^* (w_i, z_i)$. This would imply $(x_i, y_i) \succ_i^* (y_i, x_i) \succ_i^* (w_i, z_i)$, contradicting the fact that $\text{Not}[z_i V_i w_i]$. Similarly it is impossible that $(w_i, z_i) \succ_i^* (y_i, x_i)$. Hence, we have $(y_i, x_i) \sim_i^* (w_i, z_i)$. \square

Remark 20

With our definition of S_i and V_i , for any pair (x_i, y_i) , there are at most five possible situations:

- $x_i P_i y_i$ and $x_i V_i y_i$,
- $x_i P_i y_i$ and $\text{Not}[x_i V_i y_i]$,
- $x_i I_i y_i$,

and the two remaining cases correspond to the first two cases in which the rôles of x_i and y_i have been exchanged. In terms of \succsim_i^* , we have shown that, for all $z_i, w_i \in X_i$:

- In the first case, we have $(x_i, y_i) \succsim_i^* (z_i, w_i)$ and $(z_i, w_i) \succsim_i^* (y_i, x_i)$, for all $z_i, w_i \in X_i$. Furthermore, it is never true that $(y_i, a_{-i}) \succ (x_i, b_{-i})$.
- In the second case, we have $(x_i, y_i) \succsim_i^* (z_i, w_i)$, for all $z_i, w_i \in X_i$.
- In the third case we have $(x_i, y_i) \sim_i^* (y_i, x_i) \sim_i^* (z_i, z_i)$, for all $z_i \in X_i$.

Since we know by Lemmas 9 and 14 that \succsim_i^* is a weak order, this implies that \succsim_i^* has at most the following four equivalence classes, ordered in decreasing order by \succsim_i^* : (x_i, y_i) belongs to

- class 1 iff $x_i P_i y_i$

- class 2 iff $x_i I_i y_i$
- class 3 iff $y_i P_i x_i$ and $\text{Not}[y_i V_i x_i]$
- class 4 iff $y_i P_i x_i$ and $y_i V_i x_i$.

The relation \succsim_i^{**} , which is also a weak order, has at most five classes, the first class of \succsim_i^* being split into two subclasses depending on whether $x_i V_i y_i$ or not. •

This leads to our characterization of R-CDR.

Theorem 21

Let \succsim be a reflexive binary relation on $X = \prod_{i=1}^n X_i$. Then \succsim is an R-CDR iff it satisfies RC1, RC2, M1 and M3. These axioms are independent in the class of reflexive binary relations.

PROOF

Necessity results from Lemmas 14 and 16. The independence of the axioms results from Remark 18. We show sufficiency.

Define the relation S_i on X_i letting $x_i S_i y_i$ if $(x_i, y_i) \succsim_i^* (y_i, x_i)$. Let P_i and I_i respectively denote the asymmetric and the symmetric parts of S_i . Using Lemma 19.1, we know that S_i is complete. Note that, since all attributes have been supposed influent, P_i is not empty. Indeed, \succsim_i^* being complete, the influence of $i \in N$ implies that there are $x_i, y_i, z_i, w_i \in X_i$ such that $(x_i, y_i) \succ_i^* (z_i, w_i)$. If P_i is empty, x_i, y_i, z_i, w_i fulfill the conditions of Lemma 19.2; hence $(x_i, y_i) \sim_i^* (z_i, w_i)$, a contradiction.

Define the relation V_i on X_i letting $x_i V_i y_i$ if, for some $z_i, w_i \in X_i$, $(z_i, w_i) \succ_i^* (w_i, z_i) \succ_i^* (y_i, x_i)$. Using Lemma 19.3, we know that V_i is included in P_i .

Consider two subsets $A, B \subseteq N$ such that $A \cup B = N$ and let:

$$A \supseteq B \Leftrightarrow [x \succsim y, \text{ for some } x, y \in X \text{ such that } S(x, y) = A \text{ and } S(y, x) = B].$$

Suppose that $x \succsim y$. Using Lemma 17.2, we must have $V(y, x) = \emptyset$. By construction, we have $S(x, y) \supseteq S(y, x)$.

Suppose now that $V(y, x) = \emptyset$ and $S(x, y) \supseteq S(y, x)$. Let us show that we have $x \succsim y$. By construction, $S(x, y) \supseteq S(y, x)$ implies that there are $z, w \in X$ such that $z \succsim w$, $S(x, y) = S(z, w)$ and $S(y, x) = S(w, z)$. For all $i \in N$ such that $z_i I_i w_i$ we have $x_i I_i y_i$ so that, using Lemma 19.2, $(x_i, y_i) \sim_i^* (z_i, w_i)$. For all $i \in N$ such that $z_i P_i w_i$ we have $x_i P_i y_i$ so that, using Lemma 19.1, $(x_i, y_i) \sim_i^* (z_i, w_i)$. For all $i \in N$ such that $w_i P_i z_i$, we

have $y_i P_i x_i$. By hypothesis, we have $\text{Not}[y_i V_i x_i]$. Because $z \succsim w$, we have $\text{Not}[w_i V_i z_i]$. Using Lemma 19.3c, we know that $(x_i, y_i) \sim_i^* (z_i, w_i)$. Hence, we have $(x_i, y_i) \sim_i^* (z_i, w_i)$, for all $i \in N$ so that $z \succsim w$ implies $x \succsim y$.

It remains to show that \supseteq is monotonic. Suppose that $A \supseteq B$, so that, for some $x, y \in X$, $S(x, y) = A$, $S(y, x) = B$ and $x \succsim y$. Since $x \succsim y$, we know that $\text{Not}[y_i V_i x_i]$, for all $i \in N$. Suppose that $C \supseteq A$, $B \supseteq D$ such that $C \cup D = N$. We first show that $C \supseteq B$. Let $E = C \setminus A$; we have $B \supseteq E$. We build $z, w \in X$ with $S(z, w) = C \supseteq S(w, z) = B$. We know that for all $a_i \in X_i$, we have $a_i I_i a_i$. Using such pairs, define $z, w \in X$ as follows:

	A	E	$B \setminus E$
z	x_i	a_i	x_i
w	y_i	a_i	y_i

One verifies that $V(w, z) = \emptyset$, $S(z, w) = A \cup E = C$ and $S(w, z) = B$. For all $i \in E \subseteq B$, $(a_i, a_i) \succsim_i^* (x_i, y_i)$. Hence, using the definition of \succsim_i^* , we have $x \succsim y$ implies $z \succsim w$. This implies that $C \supseteq B$.

Starting from z and w and $C \supseteq B$, we show that $C \supseteq D$ for $D \subseteq B$. Since $C \cup D = N$, we have $B \setminus D \subseteq C$. Let $F = B \setminus D$. For all $i \in N$, P_i is not empty so that we can take, for all $i \in E$ any $a_i, b_i \in X_i$ such that $a_i P_i b_i$. Using such pairs, define $r, s \in X$ as follows:

	$C \setminus F$	F	D
r	z_i	a_i	z_i
s	w_i	b_i	w_i

One verifies that $V(s, r) = \emptyset$, $S(r, s) = C$ and $S(s, r) = D$. For all $i \in F \subseteq C$, we have $(a_i, b_i) \succsim_i^* (z_i, w_i)$. Hence, using the definition of \succsim_i^* , we have $z \succsim w$ implies $r \succsim s$. This implies that $C \supseteq D$, which completes the proof. \square

Remark 22

In Bouyssou and Pirlot (2005a, Lemma 2), we showed that, when all attributes are influent, the representation $\langle \supseteq, S_i \rangle$ of an R-CR is always unique. It is important here to note that such a result does not hold for the representation $\langle \supseteq, S_i, V_i \rangle$ of an R-CDR. Indeed suppose that on some attribute $i \in N$, it is impossible that $x \succsim y$ as soon as $y_i P_i x_i$, i.e., $(y_i, x_i) \succ_i^* (x_i, y_i)$. This can be represented saying that it is never true that $A \supseteq B$ when $i \in B$ together with $V_i = \emptyset$. Alternatively, we may take the relation V_i to hold as soon as $(y_i, x_i) \succ_i^* (x_i, y_i)$.

The first option has been taken in the proof of Theorem 21. This leads to building a representation $\langle \supseteq, S_i, V_i \rangle$ of an R-CDR that uses a minimal amount of veto. We do not investigate here the additional conditions under which the representation $\langle \supseteq, S_i, V_i \rangle$ would be unique since this does not seem

to add insight on the nature of these relations. These conditions have to do with the fact that for all attributes there are levels linked by S_i but not by V_i . Note however that assessment methods designed for R-CDR should be prepared to deal with this lack of uniqueness. •

5 Concordance-discordance relations with attribute transitivity

Our definition of R-CDR in Section 4 does not require the relations S_i or V_i to possess any remarkable property besides the completeness of S_i and the fact that $V_i \subseteq P_i$. This is at variance with what is done in most outranking methods (see the examples in Section 2.4). In this section, we show here how to characterize R-CDR with the following additional requirements:

- the relation S_i is a semiorder,
- the veto relation V_i is a strict semiorder such that $V_i \subseteq P_i$,
- (S_i, U_i) is a homogeneous nested chain of semiorders, where U_i denotes the codual of V_i .

for all $i \in N$. As discussed in Section 2.4, this will bring us quite close to the models that are used in practice.

As a first step, we show how to refine the framework provided by model (M).

5.1 Conjoint measurement framework continued

We first show, following Bouyssou and Pirlot (2004a), how to introduce a linear arrangement of the elements of each X_i within the framework of model (M). The following relations play a fundamental rôle.

Definition 23 (Relations comparing the levels on each attribute)

Let \succsim be a binary relation on a set $X = \prod_{i=1}^n X_i$. We define the binary relations \succsim_i^+ , \succsim_i^- and \succsim_i^\pm on X_i letting, for all $x_i, y_i \in X_i$,

$$x_i \succsim_i^+ y_i \Leftrightarrow \forall a_{-i} \in X_{-i}, b \in X, [(y_i, a_{-i}) \succsim b \Rightarrow (x_i, a_{-i}) \succsim b], \quad (11)$$

$$x_i \succsim_i^- y_i \Leftrightarrow \forall a \in X, b_{-i} \in X_{-i}, [a \succsim (x_i, b_{-i}) \Rightarrow a \succsim (y_i, b_{-i})], \quad (12)$$

$$x_i \succsim_i^\pm y_i \Leftrightarrow x_i \succsim_i^+ y_i \text{ and } x_i \succsim_i^- y_i. \quad (13)$$

The definition of \succsim_i^\pm suggests that there is a linear arrangement of the elements of X_i . The relation \succsim reacts in a monotonic way when an element is substituted by another that is “above” the latter in the arrangement. However, nothing in model (M) ensures that the relation \succsim_i^\pm is complete. This will indeed require extra conditions.

Remark 24

Let us note that, unsurprisingly, the relation \succsim_i^\pm has strong relationships with \succsim_i^* and \succsim_i^{**} . Using the definition of these relations, one checks directly that \succsim_i^\pm is the trace of \succsim_i^* , i.e., we have, for all $x_i, y_i \in X_i$:

$$\begin{aligned} x_i \succsim_i^+ y_i &\Leftrightarrow [\forall z_i \in X_i, (x_i, z_i) \succsim_i^* (y_i, z_i)], \\ x_i \succsim_i^- y_i &\Leftrightarrow [\forall w_i \in X_i, (w_i, y_i) \succsim_i^* (w_i, x_i)], \\ x_i \succsim_i^\pm y_i &\Leftrightarrow \end{aligned} \tag{14}$$

$$\forall z_i \in X_i, (x_i, z_i) \succsim_i^* (y_i, z_i) \text{ and } \forall w_i \in X_i, (w_i, y_i) \succsim_i^* (w_i, x_i).$$

It is easy to see that \succsim_i^\pm is also the trace of \succsim_i^{**} . •

By construction, the relations \succsim_i^+ , \succsim_i^- and \succsim_i^\pm are always reflexive and transitive. Their completeness is related to the following axioms.

Definition 25 (Conditions AC1, AC2 and AC3)

We say that \succsim satisfies:

$$AC1_i \text{ if } \left. \begin{array}{l} x \succsim y \\ \text{and} \\ z \succsim w \end{array} \right\} \Rightarrow \left\{ \begin{array}{l} (z_i, x_{-i}) \succsim y \\ \text{or} \\ (x_i, z_{-i}) \succsim w, \end{array} \right.$$

$$AC2_i \text{ if } \left. \begin{array}{l} x \succsim y \\ \text{and} \\ z \succsim w \end{array} \right\} \Rightarrow \left\{ \begin{array}{l} x \succsim (w_i, y_{-i}) \\ \text{or} \\ z \succsim (y_i, w_{-i}), \end{array} \right.$$

$$AC3_i \text{ if } \left. \begin{array}{l} z \succsim (c_i, a_{-i}) \\ \text{and} \\ (c_i, b_{-i}) \succsim y \end{array} \right\} \Rightarrow \left\{ \begin{array}{l} z \succsim (d_i, a_{-i}) \\ \text{or} \\ (d_i, b_{-i}) \succsim y, \end{array} \right.$$

for all $x, y, z, w \in X$, all $a_{-i}, b_{-i} \in X_{-i}$ and all $c_i, d_i \in X_i$. We say that \succsim satisfies AC1 (resp. AC2, AC3) if it satisfies $AC1_i$ (resp. $AC2_i$, $AC3_i$) for all $i \in N$.

These three conditions are transparent variations on the theme of the Ferrers ($AC1$ and $AC2$) and semi-transitivity ($AC3$) conditions that are made possible by the product structure of X . They are directly related to properties of relations \succsim_i^+ , \succsim_i^- and \succsim_i^\pm as stated in the next lemma.

Lemma 26

1. $AC1_i \Leftrightarrow \succsim_i^+$ is complete.
2. $AC2_i \Leftrightarrow \succsim_i^-$ is complete.
3. $AC3_i \Leftrightarrow [Not[x_i \succsim_i^+ y_i] \Rightarrow y_i \succsim_i^- x_i] \Leftrightarrow [Not[x_i \succsim_i^- y_i] \Rightarrow y_i \succsim_i^+ x_i]$.
4. $[AC1_i, AC2_i \text{ and } AC3_i] \Leftrightarrow \succsim_i^\pm$ is complete.
5. In the class of reflexive relations satisfying $RC1$ and $RC2$, $AC1$, $AC2$ and $AC3$ are independent conditions.

PROOF

See Bouyssou and Pirlot (2004b, Lemma 3) for Parts 1 to 4 and Bouyssou and Pirlot (2004a, Section 5.1.2) for Part 5. \square

In Bouyssou and Pirlot (2004a), we consider binary relations \succsim on X that can be represented as:

$$x \succsim y \Leftrightarrow F(\varphi_1(u_1(x_1), u_1(y_1)), \dots, \varphi_n(u_n(x_n), u_n(y_n))) \geq 0, \quad (M^*)$$

where u_i are real-valued functions on X_i , φ_i are real-valued functions on $u_i(X_i)^2$ that are *skew symmetric*, *nondecreasing* in their first argument (and, therefore, *nonincreasing* in their second argument) and F is a real-valued function on $\prod_{i=1}^n \varphi_i(u_i(X_i)^2)$ being *nondecreasing* in all its arguments and such that $F(\mathbf{0}) \geq 0$.

Going from model (M) to model (M*) amounts to requiring that each function p_i measuring preference differences can be factorized as $\varphi_i(u_i(x_i), u_i(y_i))$ thereby reflecting an underlying linear arrangement of the elements of X_i .

The conditions introduced so far allow to characterize model (M*) when each X_i is at most countably infinite. We have:

Theorem 27 (Bouyssou and Pirlot 2004a, Theorem 2)

Let \succsim be a binary relation on a finite or countably infinite set $X = \prod_{i=1}^n X_i$. Then \succsim has a representation (M*) if and only if it is reflexive and satisfies $RC1$, $RC2$, $AC1$, $AC2$ and $AC3$.

Remark 28

Note that, contrary to Theorem 21, Theorem 27 is only stated here for finite or countably infinite sets X . This is no mistake: we refer to Bouyssou and Pirlot (2004a) for details and for the analysis of the extension of this result to the general case. \bullet

Although model (M^*) is a particular case of model (M) , it is still flexible enough to contain as particular cases models like the additive value function model (Debreu 1960; Krantz et al. 1971) or Tversky's additive difference model (Tversky 1969). We show below that it also contains all R-CDR in which the relations S_i and U_i are semiorders and (S_i, U_i) form a homogeneous nested chain of semiorders.

5.2 R-CDR with attribute transitivity

Let us first precisely define what we will call *R-CDR with attribute transitivity* (R-CDR-AT).

Definition 29 (R-CDR with attribute transitivity)

An R-CDR with attribute transitivity (R-CDR-AT) is an R-CDR for which, for all $i \in N$:

- S_i is a semiorder with asymmetric part P_i ,
- V_i is the asymmetric part of a semiorder U_i with $U_i \supseteq S_i$ and, hence, $V_i \subseteq P_i$,
- (S_i, U_i) form a homogeneous chain of semiorders, i.e., there is a weak order R_i on X_i such that:

$$x_i R_i y_i \Rightarrow \forall z_i \in X_i, [y_i S_i z_i \Rightarrow x_i S_i z_i] \text{ and } [z_i S_i x_i \Rightarrow z_i S_i y_i], \quad (15)$$

and

$$x_i R_i y_i \Rightarrow \forall z_i \in X_i, [y_i U_i z_i \Rightarrow x_i U_i z_i] \text{ and } [z_i U_i x_i \Rightarrow z_i U_i y_i]. \quad (16)$$

Remark 30

It is easy to check that when R_i satisfies (15) and (16), we also have that:

$$x_i R_i y_i \Rightarrow \forall z_i \in X_i, [y_i P_i z_i \Rightarrow x_i P_i z_i] \text{ and } [z_i P_i x_i \Rightarrow z_i P_i y_i] \quad (17)$$

and

$$x_i R_i y_i \Rightarrow \forall z_i \in X_i, [y_i V_i z_i \Rightarrow x_i V_i z_i] \text{ and } [z_i V_i x_i \Rightarrow z_i V_i y_i]. \quad (18)$$

This will be useful later. •

The following lemma shows that all R-CDR-AT have a representation in model (M^*) .

Lemma 31

If \succsim is an R-CDR-AT then \succsim satisfies AC1, AC2 and AC3.

PROOF

Let $\langle \succeq, S_i, V_i \rangle$ be a representation of the R-CDR-AT \succsim and let R_i be the weak order obtained by intersecting the weak orders induced by the semiorders S_i and U_i .

[AC1_i]. Suppose that $(x_i, x_{-i}) \succsim (y_i, y_{-i})$ and $(z_i, z_{-i}) \succsim (w_i, w_{-i})$. We want to show that either $(z_i, x_{-i}) \succsim (y_i, y_{-i})$ or $(x_i, z_{-i}) \succsim (w_i, w_{-i})$.

From the hypothesis, we get that $\text{Not}[y_i V_i x_i]$ and $\text{Not}[w_i V_i z_i]$. Since R_i is a weak order, either $x_i R_i z_i$ or $z_i R_i x_i$. Suppose $z_i R_i x_i$ and observe that this together with $\text{Not}[y_i V_i x_i]$ implies $\text{Not}[y_i V_i z_i]$ (due to (18)). We show that $(x_i, x_{-i}) \succsim (y_i, y_{-i})$ implies $(z_i, x_{-i}) \succsim (y_i, y_{-i})$. If $y_i P_i x_i$, the conclusion follows from the monotonicity of \succeq . If $x_i S_i y_i$, we also have $z_i S_i y_i$ (using $z_i R_i x_i$ and (15)), hence the conclusion follows from the monotonicity of \succeq (if $x_i I_i y_i$) or directly (if $x_i P_i y_i$). The case in which $x_i R_i z_i$ leads to proving that $(x_i, z_{-i}) \succsim (w_i, w_{-i})$ in an analogous manner.

Hence AC1_i holds. The proof for AC2_i is similar, using the fact that either $y_i R_i w_i$ (and then $(x_i, x_{-i}) \succsim (w_i, y_{-i})$) or $w_i R_i y_i$ (and then $(z_i, z_{-i}) \succsim (y_i, w_{-i})$).

[AC3_i]. Suppose that $(z_i, z_{-i}) \succsim (x_i, a_{-i})$ and $(x_i, b_{-i}) \succsim (y_i, y_{-i})$. We want to show that either $(z_i, z_{-i}) \succsim (w_i, a_{-i})$ or $(w_i, b_{-i}) \succsim (y_i, y_{-i})$.

The hypothesis implies $\text{Not}[x_i V_i z_i]$ and $\text{Not}[y_i V_i x_i]$. We have either $x_i R_i w_i$ or $w_i R_i x_i$. Assume the former. Together with $\text{Not}[x_i V_i z_i]$ this entails $\text{Not}[w_i V_i z_i]$. As for AC1_i, one easily shows that $(z_i, z_{-i}) \succsim (x_i, a_{-i})$ and $x_i R_i w_i$ yield $(z_i, z_{-i}) \succsim (w_i, a_{-i})$. The proof, assuming $w_i R_i x_i$, is similar. \square

The following lemma is an important step towards establishing a characterization of the R-CDR-AT.

Lemma 32

Let \succsim be a binary relation on $X = \prod_{i=1}^n X_i$ and assume that it is reflexive and satisfies RC1, RC2, AC1, AC2 and AC3. Let $i \in N$ and take $a_i, b_i \in X_i$ such that $(a_i, a_i) \succsim_i^* (a_i, b_i)$. The binary relation T_i on X_i defined by:

$$x_i T_i y_i \text{ if } (x_i, y_i) \succsim_i^* (a_i, b_i) \tag{19}$$

is a semiorder and for all such a_i, b_i , the weak order induced by the semiorder T_i contains the weak order \succsim_i^\pm .

PROOF

Using RC1_i and RC2_i, it is easy to show that we have $(a_i, a_i) \sim_i^* (b_i, b_i)$, for all $a_i, b_i \in X_i$. Hence, since it is supposed that $(a_i, a_i) \succsim_i^* (a_i, b_i)$, it is clear that T_i is reflexive. Lemma 9 implies that \succsim_i^* is a weak order and Lemma 26

implies that \succsim_i^\pm is also a weak order. Moreover, \succsim_i^\pm is the trace of \succsim_i^* , i.e., \succsim_i^\pm satisfies (14).

We first show that T_i has the Ferrers property, i.e., $x_i T_i y_i$ and $z_i T_i w_i$ imply $x_i T_i w_i$ or $z_i T_i y_i$. Since \succsim_i^\pm is complete, we either have $y_i \succsim_i^\pm w_i$ or the opposite. In the former case, (14) implies that $(x_i, w_i) \succsim_i^* (x_i, y_i)$. From $x_i T_i y_i$, we obtain $(x_i, y_i) \succsim_i^* (a_i, b_i)$ and, since \succsim_i^* is transitive, we get $(x_i, w_i) \succsim_i^* (a_i, b_i)$ and $x_i T_i w_i$. Starting from $w_i \succsim_i^\pm y_i$, one proves similarly that $z_i T_i y_i$.

Establishing the semi-transitivity of T_i amounts to show that $x_i T_i y_i$ and $y_i T_i z_i$ imply $x_i T_i w_i$ or $w_i T_i z_i$ for all $x_i, y_i, z_i, w_i \in X_i$. Since \succsim_i^\pm is complete, we either have $y_i \succsim_i^\pm w_i$ or the opposite. In the latter case, from $w_i \succsim_i^\pm y_i$ and equation (14), we get $(w_i, z_i) \succsim_i^* (y_i, z_i)$. Using $y_i T_i z_i$, we have $(y_i, z_i) \succsim_i^* (a_i, b_i)$ and, since \succsim_i^* is transitive, $(w_i, z_i) \succsim_i^* (a_i, b_i)$, hence $w_i T_i z_i$. Starting from $y_i \succsim_i^\pm w_i$, one proves similarly that $x_i T_i w_i$.

We now prove that the weak order \succsim_i^\pm is included in the weak order T_i^{wo} induced by T_i , that is defined by:

$$x_i T_i^{wo} y_i \text{ if } \forall z_i \in X_i, [y_i T_i z_i \Rightarrow x_i T_i z_i] \text{ and } [z_i T_i x_i \Rightarrow z_i T_i y_i].$$

In view of (14), it is clear that $x_i \succsim_i^\pm y_i$ implies the condition that defines T_i^{wo} . It follows that $\succsim_i^\pm \subseteq T_i^{wo}$. \square

Applying the previous lemma to an R-CDR that satisfies *AC1*, *AC2* and *AC3* yields the following result.

Lemma 33

If \succsim is an R-CDR that satisfies AC1, AC2 and AC3 then it is an R-CDR-AT.

PROOF

By Theorem 21, we know that \succsim satisfies *RC1*, *RC2*. We are thus in the conditions for applying Lemma 32. With S_i as defined in Lemma 19.2 and using Remark 20, we see that S_i is just the set of pairs belonging to the first two classes of \succsim_i^* , i.e., the set of pairs (x_i, y_i) such that $(x_i, y_i) \succsim_i^* (x_i, x_i)$. Applying Lemma 32 yields that S_i is a semiorder.

In a similar way, using the definition of V_i given in Lemma 19.3, we see that V_i is formed of the pairs (x_i, y_i) such that (y_i, x_i) is in the fourth and last class of \succsim_i^* . It is thus a subset of the first class of \succsim_i^* (more precisely, it is the first class of \succsim_i^{**}). The relation V_i is the asymmetric part of its codual U_i which is easily seen to be formed of the three first classes of \succsim_i^* . Applying Lemma 32, we have that U_i is a semiorder and that V_i is the asymmetric part of a semiorder. By Lemma 19.3a, we know that V_i is included in P_i , so that S_i is included in U_i . Finally Lemma 32 guarantees that the intersection

of the weak orders induced by the semiorders S_i and U_i contains the weak order \succsim_i^\pm . It is thus complete, hence a weak order. We can thus take R_i in Definition 29 to be \succsim_i^\pm . This shows that $\langle \supseteq, S_i, V_i \rangle$ is a representation of the R-CDR \succsim with (S_i, U_i) a nested homogeneous chain of semiorders. \square

This leads to our characterization of R-CDR-AT.

Theorem 34

Let \succsim be a binary relation on $X = \prod_{i=1}^n X_i$. Then \succsim is an R-CDR-AT iff it is reflexive and satisfies RC1, RC2, M1, M3, AC1, AC2 and AC3. In the class of reflexive relations, these conditions are independent.

PROOF

The characterization of R-CDR-AT results immediately from Lemma 31, Theorem 21 and Lemma 33. In view of proving the independence of the axioms, we provide below the seven required examples (in all these examples, one of the axioms is false on the first attribute. It is easy to adapt the examples to show that the axiom can be falsified on any attribute).

Example 35 (Not[AC2₁])

Let $X = X_1 \times X_2$ with $X_1 = \{x, y, z, w\}$ and $X_2 = \{a, b\}$. We build an R-CDR on X with:

- $w P_1 z, x P_1 z, y P_1 z, w P_1 x, y I_1 w, y I_1 x$ (and all I_1 loops)
- V_1 is empty except that $y V_1 z$,
- $a P_2 b$ (and all I_2 loops) and the relation V_2 is empty,
- $\{1, 2\} \triangleright \emptyset, \{1, 2\} \triangleq \{2\}, \{1, 2\} \triangleq \{1\}$ and $\{1\} \triangleq \{2\}$.

Observe that S_1 is a semiorder (the weak order it induces ranks the elements of X_1 in the following order: w, y, x, z). The relation V_1 is a strict semiorder that is included in P_1 . But (S_1, U_1) is not a homogeneous family of semiorders on X_1 since the weak order induced by U_1 ranks y before w , while the weak order induced by S_1 does the opposite.

By construction, \succsim is an R-CDR. Hence, it satisfies RC1, RC2, M1 and M3. The relation \succsim contains all pairs in $X \times X$ except the following ones:

- $(z, b) \not\prec (w, a), (z, b) \not\prec (x, a), (z, b) \not\prec (y, a), (x, b) \not\prec (w, a)$, due to the fact that $\text{Not}[\emptyset \supseteq \{1, 2\}]$, and
- $(z, a) \not\prec (y, a), (z, a) \not\prec (y, b), (z, b) \not\prec (y, a), (z, b) \not\prec (y, b)$, due to the fact that $y V_1 z$.

One pair is common to these two series of four pairs, so that \succsim is equal to $X \times X$ minus the seven distinct pairs in the lists above.

On X_2 , it is easy to check that we have $a \succ_2^\pm b$, so that $AC1_2$, $AC2_2$ and $AC3_2$ hold.

On X_1 , it is easy to check that \succsim_1^+ is complete. We indeed have that:

$$[y \sim_1^+ w] \succ_1^+ x \succ_1^+ z.$$

The relation \succsim_1^- is not complete. We have $w \succ_1^- x$, $y \succ_1^- x$ and $x \succ_1^- z$ but neither $y \succsim_1^- w$ nor $w \succsim_1^- y$ since $(z, a) \succ (w, a)$ but $(z, a) \not\succeq (y, a)$ and $(x, b) \succ (y, a)$ but $(x, b) \not\succeq (w, a)$. This shows that $AC2_1$ is violated. Condition $AC3_1$ holds since \succsim_1^+ and \succsim_1^- are not incompatible. \diamond

Example 36 (Not[AC1₁])

This example is a slight variant of the above one obtained by reversing all relations S_i and V_i . Let $X = X_1 \times X_2$ with $X_1 = \{x, y, z, w\}$ and $X_2 = \{a, b\}$. We build an R-CDR on X with:

- $z P_1 x, z P_1 y, z P_1 w, x P_1 w, x I_1 y, y I_1 w,$
- the relation V_1 is empty except that $z V_1 y,$
- $b P_2 a,$
- the relation V_2 is empty,
- $\{1, 2\} \triangleright \emptyset, \{1, 2\} \triangleq \{2\}, \{1, 2\} \triangleq \{1\}$ and $\{1\} \triangleq \{2\}.$

By construction, \succsim is an R-CDR. Hence, it satisfies $RC1$, $RC2$, $M1$ and $M3$. The relation \succsim contains all pairs in $X \times X$ except the following ones:

- $(x, b) \not\succeq (z, a), (y, b) \not\succeq (z, a), (w, b) \not\succeq (z, a), (w, b) \not\succeq (x, a),$ due to the fact that $Not[\emptyset \supseteq \{1, 2\}]$, and
- $(y, a) \not\succeq (z, a), (y, b) \not\succeq (z, b), (y, b) \not\succeq (z, a), (y, a) \not\succeq (z, b),$ due to the fact that $z V_1 y.$

One pair is common to these two series of four pairs, so that \succsim is equal to $X \times X$ minus the seven distinct pairs in the lists above.

On X_2 , it is easy to check that we have $b \succ_2^\pm a$, so that $AC1_2$, $AC2_2$ and $AC3_2$ hold.

On X_1 , it is easy to check that \succsim_1^- is complete. We indeed have that:

$$z \succ_1^- x \succ_1^- [y \sim_1^- w].$$

The relation \succsim_1^+ is not complete. We have $z \succsim_1^+ x$, $x \succsim_1^+ y$ and $x \succsim_1^+ w$ but neither $y \succsim_1^+ w$ nor $w \succsim_1^+ y$ since $(y, b) \succsim (x, a)$ but $(w, b) \not\succeq (x, a)$ and $(w, a) \succsim (z, a)$ but $(y, a) \not\succeq (z, a)$. This shows that $AC1_1$ is violated. Condition $AC3_1$ holds since \succsim_1^+ and \succsim_1^- are not incompatible. \diamond

Example 37 (Not[AC3₁])

Consider Example 20 in Bouyssou and Pirlot (2007) (see also Example 35 in Bouyssou and Pirlot (2005a)). It is shown in Bouyssou and Pirlot (2007) that Example 20 satisfies $RC1$, $RC2$, $M1$, $M2$, $AC1$, $AC3_j$, $j \neq 1$, $Not[AC3_1]$. This example satisfies $M3$ since we know that $M2$ implies $M3$. Since in presence of $RC1$, $RC2$, $M1$, $M2$, conditions $AC1$ and $AC2$ are equivalent (Lemma 27 Bouyssou and Pirlot 2005a), \succsim also satisfies $AC2$ (for instance, \succsim_1^- is the weak order $a \succsim_1^- b \succsim_1^- [c \sim_1^- d]$). \diamond

Example 38 (Not[M3₁])

Let $X = X_1 \times X_2 \times X_3$ with $X_1 = \{x, y, z\}$, $X_2 = \{a, b\}$ and $X_3 = \{p, q\}$. Let us consider the relation \succsim such that:

$$x \succsim y \Leftrightarrow \sum_{i=1}^3 p_i(x_i, y_i) \geq 0,$$

the functions p_i being such that:

$$\begin{aligned} p_1(x, y) &= p_1(x, z) = p_1(y, z) = p_1(x, x) = p_1(y, y) = p_1(z, z) = 4, \\ p_1(y, x) &= p_1(z, y) = -1, p_1(z, x) = -4 \\ p_2(a, b) &= 2, p_2(a, a) = p_2(b, b) = 0, p_2(b, a) = -2 \\ p_3(p, q) &= 2, p_3(p, p) = p_3(q, q) = 0, p_3(q, p) = -2. \end{aligned}$$

It is easily checked that we have:

$$\begin{aligned} [(x, y) \sim_1^* (x, z) \sim_1^* (y, z) \sim_1^* (x, x) \sim_1^* (y, y) \sim_1^* (z, z)] \succsim_1^* [(y, x) \sim_1^* (z, y)] \succsim_1^* (z, x) \\ x \succsim_1^\pm y \succsim_1^\pm z, \\ (a, b) \succsim_2^* [(a, a) \sim_2^* (b, b)] \succsim_2^* (b, a), \\ a \succsim_2^\pm b, \\ (p, q) \succsim_3^* [(p, p) \sim_2^* (q, q)] \succsim_2^* (q, p), \\ p \succsim_3^\pm q. \end{aligned}$$

This shows that $RC1$, $RC2$, $AC1$, $AC2$ and $AC3$ are satisfied. Using the fact that UC_i implies $M1_i$ (see Remark 12) shows that $M1$ holds. Similarly, using the fact that LC_i implies $M2_i$ and that $M2_i$ implies $M3_i$ shows that $M3_2$ and $M3_3$ hold. Condition $M3_1$ is violated since $(x, b, q) \succsim (y, a, p)$, $(y, a, q) \succsim (x, b, q)$ and $(z, a, p) \succsim (x, b, q)$ while $(y, b, q) \not\succeq (x, a, p)$, $(z, b, q) \not\succeq (x, a, p)$ and $(z, a, q) \not\succeq (x, b, q)$. \diamond

Example 39 (*Not*[$M1_1$])

Example 23 in Bouyssou and Pirlot (2007) (see also Example 33 in Bouyssou and Pirlot (2005a)) satisfies the required properties. It is shown in Bouyssou and Pirlot (2007) that this example satisfies $RC1$, $RC2$, $M1_j$, $j \neq 1$, $Not[M1_1]$, $M2$, $AC1$ and $AC3$. Since $M2$ entails $M3$, it satisfies $M3$. One easily checks that it satisfies $AC2$. \diamond

Example 40 (*Not*[$RC2_1$])

Example 25 in Bouyssou and Pirlot (2007) is appropriate, taking into account that $M2$ implies $M3$ and that this example can be shown to satisfy $AC2$. \diamond

Example 41 (*Not*[$RC1_1$])

Let $N = \{1, 2, 3\}$ and $X = \{x_1, y_1, z_1, w_1\} \times \{x_2, y_2\} \times \{x_3, y_3\}$. Let \succsim on X be identical to X^2 except that, for all $a_1, b_1 \in X_1$, all $a_2, b_2 \in X_2$ and all $a_3, b_3 \in X_3$ the following pairs are *missing*:

$$\begin{array}{ll} (a_1, x_2, x_3) \not\succeq (b_1, y_2, y_3), & (z_1, a_2, a_3) \not\succeq (y_1, b_2, b_3), \\ (x_1, x_2, x_3) \not\succeq (y_1, x_2, y_3), & (x_1, y_2, x_3) \not\succeq (y_1, y_2, y_3), \\ (z_1, x_2, x_3) \not\succeq (w_1, y_2, x_3), & (z_1, x_2, y_3) \not\succeq (w_1, y_2, y_3), \end{array}$$

(there is a total of 35 such pairs). It is clear that \succsim is reflexive.

For $i \in \{2, 3\}$, it is easy to check that we have:

$$[(y_i, x_i), (x_i, x_i), (y_i, y_i)] \succ_i^* (x_i, y_i),$$

which shows that $RC1_2$, $RC1_3$, $RC2_2$ and $RC2_3$ hold.

For $i \in \{2, 3\}$, we also have:

$$y_i \succ_i^+ x_i \text{ and } y_i \succ_i^- x_i,$$

which shows that $AC1_2$, $AC1_3$, $AC2_2$, $AC2_3$, $AC3_2$ and $AC3_3$ hold.

By Lemma 16 in Bouyssou and Pirlot (2005a), it is clear that LC_2 , LC_3 , UC_2 and UC_3 hold, which, by Lemma 11 in Bouyssou and Pirlot (2007), implies that $M1_2$, $M1_3$, $M2_2$ and $M2_3$ hold. Because $M2_i$ implies $M3_i$, we know that $M3_2$ and $M3_3$ hold.

On attribute 1, it is easy to check that we have:

$$\begin{array}{l} (c_1, d_1) \succ_1^* (x_1, y_1), \\ (c_1, d_1) \succ_1^* (z_1, w_1), \\ (x_1, y_1) \succ_1^* (z_1, y_1), \\ (z_1, w_1) \succ_1^* (z_1, y_1), \end{array}$$

for all $(c_1, d_1) \in \Gamma = \{(x_1, x_1), (x_1, z_1), (x_1, w_1), (y_1, x_1), (y_1, y_1), (y_1, z_1), (y_1, w_1), (z_1, x_1), (z_1, z_1), (w_1, x_1), (w_1, y_1), (w_1, z_1), (w_1, w_1)\}$. The pairs (x_1, y_1) and (z_1, w_1) are not comparable in terms of \succsim_1^* since $(x_1, x_2, x_3) \succsim (y_1, y_2, x_3)$ and $(z_1, x_2, x_3) \not\succeq (w_1, y_2, x_3)$, while $(z_1, x_2, x_3) \succsim (w_1, x_2, y_3)$ and $(x_1, x_2, x_3) \not\succeq (y_1, x_2, y_3)$. Hence, $RC1_1$ is violated.

It is easy to check that $RC2_1$ and, using Lemma 16 in Bouyssou and Pirlot (2005a), that UC_1 holds. Hence, we know that $M1_1$ holds (Bouyssou and Pirlot 2007, Lemma 11). We have:

$$\begin{aligned} [y_1 \sim_1^+ w_1] \succ_1^+ x_1 \succ_1^+ z_1, \\ y_1 \succ_1^- w_1 \succ_1^- [x_1 \sim_1^- z_1], \end{aligned}$$

which shows that $AC1_1$, $AC2_1$ and $AC3_1$ hold.

It remains to check that $M3_1$ holds. The three premises of $M3_1$ are that $(a_1, a_{-1}) \succsim (b_1, b_{-1})$, $(b_1, c_{-1}) \succsim (a_1, d_{-1})$ and $(c_1, e_{-1}) \succsim (d_1, f_{-1})$. The three possible conclusions of $M3_1$ are that $(b_1, a_{-1}) \succsim (a_1, b_{-1})$ or $(c_1, a_{-1}) \succsim (d_1, b_{-1})$ or $(c_1, c_{-1}) \succsim (d_1, d_{-1})$.

Suppose first that $(b_1, a_1) \in \Gamma$. In this case, we have $(b_1, a_1) \succsim_1^* (a_1, b_1)$, so that $(a_1, a_{-1}) \succsim (b_1, b_{-1})$ implies $(b_1, a_{-1}) \succsim (a_1, b_{-1})$. Hence, the first conclusion of $M3_1$ holds.

If $(b_1, a_1) = (z_1, y_1)$, the second premise of $M3_1$ is never satisfied, so that the condition trivially holds.

Suppose now that $(b_1, a_1) = (x_1, y_1)$. If (c_1, d_1) is distinct from (z_1, w_1) and (z_1, y_1) , we have $(c_1, d_1) \succsim_1^* (x_1, y_1)$, so that $(b_1, c_{-1}) \succsim (a_1, d_{-1})$ implies $(c_1, c_{-1}) \succsim (d_1, d_{-1})$ and the third conclusion of $M3_1$ holds.

If $(c_1, d_1) = (z_1, y_1)$, the third premise of $M3_1$ is never satisfied, so that the condition trivially holds.

If $(c_1, d_1) = (z_1, w_1)$, it is easy to check that there are no $a_{-1}, b_{-1} \in X_{-1}$ such that $(y_1, a_{-1}) \succsim (x_1, b_{-1})$, $(x_1, a_{-1}) \not\succeq (y_1, b_{-1})$ and $(z_1, a_{-1}) \not\succeq (w_1, b_{-1})$, so that no violation of $M3_1$ is possible in this case.

A similar reasoning shows that the same is true if it is supposed that $(b_1, a_1) = (z_1, w_1)$. Hence, $M3_1$ holds. \square

In Bouyssou and Pirlot (2005a, 2007), we have studied reflexive concordance relations with attribute transitivity (R-CR-AT). These relations are just R-CR admitting a representation $\langle \succeq, S_i \rangle$ in which all S_i are semiorders. These are clearly a special case of R-CDR-AT (Definition 29) in which all V_i are empty relations. For further reference, we recall the characterization of R-CR-AT obtained as Theorem 26 in Bouyssou and Pirlot (2007).

Theorem 42 (Characterization of R-CR-AT)

A relation \succsim on X is an R-CR-AT iff it is reflexive and satisfies RC1, RC2, M1, M2, AC1, and AC3. In the class of reflexive relations, these axioms are independent.

The characterization of R-CR-AT only differs from that of R-CDR-AT by the substitution of axiom M3 by axiom M2 and the omission of AC2, which, in presence of the other conditions is equivalent to AC1 (see Bouyssou and Pirlot 2005a, Lemma 27).

6 Strict and non-strict preference models: discordance vs bonus

Models of preference aim at capturing either *strict* or *non-strict* preference. The latter usually corresponds to an “at least as good” relation while strict preference refers to a “better than” relation. For the special case of concordance relations, we have studied and characterized strict (asymmetric) concordance relations (see Bouyssou and Pirlot 2002a, 2005b) and non-strict concordance relations (see Bouyssou and Pirlot 2005a, 2007). The purpose of this section is to extend and unify these results.

One would expect that strict and non-strict concordance relations are related. More precisely, it would be natural that a strict concordance relation is the asymmetric part of a non-strict concordance relation. We shall see below that this is actually the case and that their axiomatic characterizations are related as well.

This picture changes however when discordance comes into play. The relationship between non-strict concordance-discordance preference models, like ELECTRE I (see Example 4) and strict concordance-discordance preference models like TACTIC (Vansnick 1986 and Example 53 below) is not entirely straightforward, as we shall see.

In our analysis of the correspondence between strict and non-strict preference models, coduality, as defined by (1), will play an important rôle.

6.1 Strict and non-strict concordance relations

We start by recalling the definition of a strict (asymmetric) concordance relation (as introduced in Bouyssou and Pirlot 2002a). We present it here as a special case of a more general “irreflexive concordance relation”. Although this more general concept is clearly of little practical interest, it will be useful for discussing relationships with reflexive concordance relations (R-CR).

Definition 43 (Irreflexive concordance relation (I-CR))

Let \mathcal{P} be an irreflexive binary relation on $X = \prod_{i=1}^n X_i$. We say that \mathcal{P} is an irreflexive concordance relation (or, more briefly, that \mathcal{P} is an I-CR) if there are:

- an asymmetric binary relation P_i° on each X_i ($i = 1, 2, \dots, n$),
- a binary relation \triangleright° between disjoint subsets of N that is monotonic w.r.t. inclusion, i.e., such that (6) holds,

such that, for all $x, y \in X$,

$$x \mathcal{P} y \Leftrightarrow P^\circ(x, y) \triangleright^\circ P^\circ(y, x), \quad (20)$$

where $P^\circ(x, y) = \{i \in N : x_i P_i^\circ y_i\}$.

A strict concordance relation is an irreflexive concordance relation that is asymmetric. We also call such a relation an asymmetric concordance relation (A-CR).

If an R-CR \succsim is a complete relation, then its asymmetric part is also the codual of \succsim since, by definition of the codual (1), we have $x \succsim^{cd} y$ iff $y \not\succeq x$. We illustrate the coduality between asymmetric I-CR and complete R-CR by the following example.

Example 44 (TACTIC, Vansnick 1986)

The binary relation \mathcal{P} is an asymmetric semioordered weighted majority preference relation if there are a real number $\varepsilon \geq 0$ and, for all $i \in N$,

- the asymmetric part P_i° of a semioorder S_i° on X_i ,
- a real number $w_i > 0$,

such that:

$$x \mathcal{P} y \Leftrightarrow \sum_{i \in P^\circ(x, y)} w_i > \sum_{j \in P^\circ(y, x)} w_j + \varepsilon,$$

where $P^\circ(x, y) = \{i \in N : x_i P_i^\circ y_i\}$.

Clearly, the relation \mathcal{P} just defined is the codual of the one in Example 3. Indeed, $\text{Not}[y \mathcal{P} x]$ is equivalent to $\sum_{j \in P^\circ(y, x)} w_j \leq \sum_{i \in P^\circ(x, y)} w_i + \varepsilon$, hence, $\sum_{i \in P^\circ(x, y)} w_i \geq \sum_{j \in P^\circ(y, x)} w_j - \varepsilon$. Adding the term $\sum_{\ell \in I^\circ(x, y)} w_\ell$ (where $I^\circ(x, y) = \{\ell \in N : x_\ell I_\ell^\circ y_\ell\}$ and I_ℓ° is the symmetric part of S_ℓ°) to both sides, one gets the definition of $x \succsim y$ as given in Example 3. \diamond

If a relation \succsim is not complete it is no longer the case that its asymmetric part is its codual. However, we cannot make the hypothesis that \succsim is complete without excluding interesting reflexive concordance relations (consider, e.g., Example 4 with $V_i = \emptyset$ and $s \gg \frac{1}{2}$). In the rest of this section, we try to clarify the relationship between strict and non-strict concordance relations in the general case of non-necessarily complete relations. We start with two lemmas.

Lemma 45

The codual of an R-CR is an I-CR and conversely.

PROOF

Let \succsim be an R-CR and $\langle \succeq, S_i \rangle$ a representation of \succsim . We obtain a representation of \succsim^{cd} as an I-CR by taking:

- $P_i^\circ = S_i^{cd}$ (P_i° is asymmetric since S_i is complete, by hypothesis),
- for all $A, B \subseteq N$ such that $A \cap B = \emptyset$,

$$A \triangleright^\circ B \text{ if } [N \setminus B] \succeq [N \setminus A], \quad (21)$$

noticing that $[N \setminus A] \cup [N \setminus B] = N$ implies $A \cap B = \emptyset$.

The required monotonicity property of \triangleright° directly follows from that of \succeq .

Conversely, starting from an I-CR, \mathcal{P} , and its representation $\langle \triangleright^\circ, P_i^\circ \rangle$, one obtains a representation $\langle \succeq, S_i \rangle$ of \mathcal{P}^{cd} by putting:

- $S_i = P_i^{\circ cd}$ (S_i is complete since P_i° is asymmetric, by hypothesis);
- for all $A, B \subseteq N$ such that $A \cup B = N$,

$$A \succeq B \text{ if } [N \setminus B] \triangleright^\circ [N \setminus A], \quad (22)$$

noticing that $[N \setminus A] \cap [N \setminus B] = \emptyset$ implies $A \cup B = N$.

The required monotonicity property of \succeq follows from that of \triangleright° . □

Lemma 46

1. *The completion of an R-CR is an R-CR;*
2. *The asymmetric part of an I-CR is an I-CR and, hence, an A-CR.*

PROOF

(1) Let \succsim be an R-CR with a representation $\langle \succeq, S_i \rangle$. The completion $\overline{\succsim}$ of \succsim is defined by $x \overline{\succsim} y$ iff $x \succsim y$ or $[x \not\succeq y \text{ and } y \not\succeq x]$. We obtain a representation

$\langle \bar{\supseteq}, S_i \rangle$ of $\bar{\succsim}$ as an R-CR by defining $\bar{\supseteq}$ as follows: for all $A, B \subseteq N$ such that $A \cup B = N$,

$$A \bar{\supseteq} B \text{ if } [A \supseteq B \text{ or } \text{Not}[B \supseteq A]]. \quad (23)$$

Using this definition, we prove that $\bar{\supseteq}$ is monotone. Assume that $A \bar{\supseteq} B$, $C \supseteq A$ and $D \subseteq B$. If $A \supseteq B$, using the monotonicity of \supseteq , we have $C \supseteq D$, hence $C \bar{\supseteq} D$. If $\text{Not}[B \supseteq A]$, then $\text{Not}[D \supseteq C]$ (by monotonicity of \supseteq). We have either $C \supseteq D$ or $\text{Not}[D \supseteq C]$, which, in both cases, yields $C \bar{\supseteq} D$. The R-CR relation having this representation is clearly the completion of $\bar{\succsim}$.

(2) Let \mathcal{P} be an I-CR with a representation $\langle \triangleright^\circ, P_i^\circ \rangle$. A representation as an I-CR of the asymmetric part $\alpha[\mathcal{P}]$ of \mathcal{P} is obtained by taking, for all $A, B \subseteq N$ such that $A \cap B = \emptyset$,

$$A \triangleright^{\circ'} B \text{ if } A \triangleright^\circ B \text{ and } \text{Not}[B \triangleright^\circ A]. \quad (24)$$

The monotonicity of $\triangleright^{\circ'}$ is easily obtained using the monotonicity of \triangleright° . Indeed, assume that $A \triangleright^{\circ'} B$, $C \supseteq A$ and $D \subseteq B$. By definition, we have $A \triangleright^\circ B$ and $\text{Not}[B \triangleright^\circ A]$. Clearly, $A \triangleright^\circ B$ implies $C \triangleright^\circ D$ and $\text{Not}[B \triangleright^\circ A]$ implies $\text{Not}[D \triangleright^\circ C]$. Hence $A \triangleright^{\circ'} B$ implies $C \triangleright^{\circ'} D$. It is easy to check that the relation having $\langle \triangleright^{\circ'}, P_i^\circ \rangle$ as a representation is indeed an asymmetric I-CR, i.e., an A-CR. \square

Using the above lemmas, we can easily prove the expected relationships between R-CR and I-CR.

Proposition 47

1. *The asymmetric part of an R-CR is an I-CR (and, hence, an A-CR).*
2. *The completion of an I-CR is an R-CR.*

PROOF

(1). Let $\bar{\succsim}$ be an R-CR with a representation $\langle \bar{\supseteq}, S_i \rangle$. Its asymmetric part \succ is also the asymmetric part of its codual $\bar{\succsim}^{cd}$. We know by Lemma 45 that $\bar{\succsim}^{cd}$ is an I-CR and by Lemma 46.2 that the asymmetric part of it, which is \succ , is an I-CR.

(2). Let \mathcal{P} be an I-CR with a representation $\langle \triangleright^\circ, P_i^\circ \rangle$. The completion of \mathcal{P} is identical to the completion of its codual \mathcal{P}^{cd} , which is an R-CR by Lemma 45. Since the completion of an R-CR is an R-CR (Lemma 46.1), the result follows. \square

6.2 Characterizations of strict and non-strict concordance relations

The correspondence through coduality between reflexive and irreflexive concordance relations (Lemma 45) is reflected in the axiomatizations of strict and

non-strict concordance relations, which we discussed respectively in Bouyssou and Pirlot (2002a, 2005b) (strict (asymmetric) concordance relations) and Bouyssou and Pirlot (2005a, 2007) (non-strict (reflexive) concordance relations). In this section, we examine the relationships between the axioms that characterize reflexive and irreflexive concordance relations.

Consider an R-CR \succsim . Its codual, \succsim^{cd} , which is not necessarily asymmetric, is an I-CR. From Theorem 13, we know that \succsim is an R-CR if and only if it is reflexive and satisfies $RC1$, $RC2$, $M1$ and $M2$. By Theorem 42, we know that \succsim satisfies, in addition, attribute transitivity (i.e., \succsim is an R-CR-AT) iff it also satisfies $AC1$ and $AC3$ (or $AC2$ and $AC3$, since in presence of the other conditions, $AC1$ and $AC2$ are equivalent). Since we want to characterize the codual of \succsim , it is interesting to observe the behavior of the axioms $RC1$, $RC2$, $M1$, $M2$, $AC1$, $AC2$ and $AC3$ through contraposition. In fact, our discussion will not depend on whether we start with a reflexive relation \succsim or an irreflexive relation \succsim^{cd} and impose on the latter or the former one of the axioms above. Our results are valid for any relation even non-reflexive or non-irreflexive. Therefore, in the following discussion, we express the axioms using the generic notation \mathcal{R} for denoting any relation on X , and \mathcal{R}^{cd} , for denoting its codual.

Consider a relation \mathcal{R} on X that is assumed to satisfy $RC1_i$. The contraposition of $RC1_i$ is a condition naturally expressed in terms of the codual of \mathcal{R} . We have:

$$\begin{aligned}
& \mathcal{R} \text{ satisfies } RC1_i \\
& \Leftrightarrow \\
& \left. \begin{array}{l} \text{Not}[(x_i, c_{-i}) \mathcal{R} (y_i, d_{-i})] \\ \text{and} \\ \text{Not}[(z_i, a_{-i}) \mathcal{R} (w_i, b_{-i})] \end{array} \right\} \Rightarrow \left\{ \begin{array}{l} \text{Not}[(x_i, a_{-i}) \mathcal{R} (y_i, b_{-i})] \\ \text{or} \\ \text{Not}[(z_i, c_{-i}) \mathcal{R} (w_i, d_{-i})] \end{array} \right. \\
& \Leftrightarrow \\
& \left. \begin{array}{l} (y_i, d_{-i}) \mathcal{R}^{cd} (x_i, c_{-i}) \\ \text{and} \\ (w_i, b_{-i}) \mathcal{R}^{cd} (z_i, a_{-i}) \end{array} \right\} \Rightarrow \left\{ \begin{array}{l} (y_i, b_{-i}) \mathcal{R}^{cd} (x_i, a_{-i}) \\ \text{or} \\ (w_i, d_{-i}) \mathcal{R}^{cd} (z_i, c_{-i}), \end{array} \right. \quad (25)
\end{aligned}$$

for all $x_i, y_i, z_i, w_i \in X_i$ and all $a_{-i}, b_{-i}, c_{-i}, d_{-i} \in X_{-i}$.

Clearly, (25) is axiom $RC1_i$ expressed in terms of relation \mathcal{R}^{cd} . Hence, we have that \mathcal{R} satisfies $RC1_1$ iff its codual does. One shows similarly that it is also the case for $RC2_i$. We refer to this property saying that axioms $RC1$ and $RC2$ are *self-dual*.

The picture is not exactly the same with $M1$ and $M2$. Contraposition of

$M1_i$ yields:

$$\begin{aligned} & \mathcal{R} \text{ satisfies } M1_i \\ & \Leftrightarrow \\ & \left. \begin{array}{l} (x_i, b_{-i}) \mathcal{R}^{cd} (y_i, a_{-i}) \\ \text{and} \\ (z_i, b_{-i}) \mathcal{R}^{cd} (w_i, a_{-i}) \\ \text{and} \\ (y_i, d_{-i}) \mathcal{R}^{cd} (x_i, c_{-i}) \end{array} \right\} \Rightarrow \left\{ \begin{array}{l} (y_i, b_{-i}) \mathcal{R}^{cd} (x_i, a_{-i}) \\ \text{or} \\ (w_i, d_{-i}) \mathcal{R}^{cd} (z_i, c_{-i}), \end{array} \right. \quad (26) \end{aligned}$$

for all $x_i, y_i, z_i, w_i \in X_i$ and all $a_{-i}, b_{-i}, c_{-i}, d_{-i} \in X_{-i}$.

Condition (26) is exactly axiom $Maj2_i$ in Bouyssou and Pirlot (2005b), imposed to relation \mathcal{R}^{cd} . Paraphrasing this result, imposing $M1_i$ to \mathcal{R} is tantamount to imposing $Maj2_i$ to its codual \mathcal{R}^{cd} and conversely. In a similar way, starting from condition $M2_i$ imposed on \mathcal{R} , we obtain the following equivalent condition on \mathcal{R}^{cd} : for all $x_i, y_i, z_i, w_i \in X_i$ and all $a_{-i}, b_{-i}, c_{-i}, d_{-i} \in X_{-i}$,

$$\left. \begin{array}{l} (x_i, b_{-i}) \mathcal{R}^{cd} (y_i, a_{-i}) \\ \text{and} \\ (w_i, b_{-i}) \mathcal{R}^{cd} (z_i, a_{-i}) \\ \text{and} \\ (w_i, d_{-i}) \mathcal{R}^{cd} (z_i, c_{-i}) \end{array} \right\} \Rightarrow \left\{ \begin{array}{l} (y_i, b_{-i}) \mathcal{R}^{cd} (x_i, a_{-i}) \\ \text{or} \\ (x_i, d_{-i}) \mathcal{R}^{cd} (y_i, c_{-i}). \end{array} \right. \quad (27)$$

Condition (27) is exactly axiom $Maj1_i$ in Bouyssou and Pirlot (2005b), imposed to relation \mathcal{R}^{cd} . Again, imposing $M2_i$ to \mathcal{R} is equivalent to imposing $Maj1_i$ to its codual \mathcal{R}^{cd} and conversely.

Dealing in the same way with axioms $AC1$, $AC2$ and $AC3$, it is easy to show that \mathcal{R} has property $AC1_i$ (resp. $AC2_i$) iff its codual satisfies $AC2_i$ (resp. $AC1_i$). Condition $AC3_i$ is self-dual.

We collect our findings in the next lemma. Its proof results from the above observations.

Lemma 48

Let \mathcal{R} be any relation on X and \mathcal{R}^{cd} its codual. We have the following equivalences, for all $i \in N$:

1. \mathcal{R} satisfies $RC1_i$ iff \mathcal{R}^{cd} satisfies $RC1_i$;
2. \mathcal{R} satisfies $RC2_i$ iff \mathcal{R}^{cd} satisfies $RC2_i$;
3. \mathcal{R} satisfies $M1_i$ iff \mathcal{R}^{cd} satisfies $Maj2_i$ (i.e., condition (26));
4. \mathcal{R} satisfies $M2_i$ iff \mathcal{R}^{cd} satisfies $Maj1_i$ (i.e., condition (27));

5. \mathcal{R} satisfies $AC1_i$ iff \mathcal{R}^{cd} satisfies $AC2_i$;

6. \mathcal{R} satisfies $AC3_i$ iff \mathcal{R}^{cd} satisfies $AC3_i$.

Using Lemma 48 it is straightforward to derive the following characterization of an irreflexive concordance relation I-CR (resp. of an I-CR with attribute transitivity I-CR-AT) from Theorem 13 (resp. from Theorem 42).

Proposition 49 (Characterization of I-CR and I-CR-AT)

A relation \mathcal{P} on X is an I-CR iff it is irreflexive and satisfies $RC1$, $RC2$, $Maj1$, and $Maj2$. A relation \mathcal{P} is an I-CR-AT iff it is irreflexive and satisfies $RC1$, $RC2$, $Maj1$, $Maj2$, $AC2$ and $AC3$. In the class of irreflexive relations, $RC1$, $RC2$, $Maj1$, $Maj2$, $AC2$ and $AC3$ are independent.

PROOF

A relation is an I-CR iff its codual is an R-CR (Lemma 45). Using the characterization of an R-CR (Theorem 13) and Lemma 48, we get the result. Obviously, if a relation is an I-CR-AT, its codual is an R-CR-AT and we obtain the result using Theorem 42 and Lemma 48. The independence of the axioms results from coduality and the fact that $RC1$, $RC2$, $M1$, $M2$, $AC1$ and $AC3$ are independent axioms (Theorem 42). \square

Remark 50

The first part of Proposition 49 has been obtained, for asymmetric relations, in Bouyssou and Pirlot (2006, Theorem 2). The characterization of I-CR-AT has not appeared before. A variant of this characterization is obtained by substituting axiom $AC2$ by axiom $AC1$. It indeed results from Bouyssou and Pirlot (2005a, Lemma 27) that $AC1$ and $AC2$ are equivalent in the class of R-CR. \bullet

Remark 51

Proposition 49 also offers a characterization of strict (asymmetric) concordance relations and of strict concordance relations with attribute transitivity. The independence of axioms $RC1$, $RC2$, $Maj1$ and $Maj2$ in the class of asymmetric relations has been established in Bouyssou and Pirlot (2006). Axioms $RC1$, $RC2$, $Maj1$, $Maj2$, $AC1$ and $AC3$ are independent in the class of asymmetric relations since axioms $RC1$, $RC2$, $M1$, $M2$, $AC1$ and $AC3$ are independent in the class of complete relations: the examples used to show the independence of the latter in Bouyssou and Pirlot (2005a, 2007) were all complete relations. \bullet

6.3 Strict concordance-discordance relations

In Bouyssou and Pirlot (2006), we have studied a model of strict (asymmetric) concordance-discordance relation that provides a framework for outranking methods, such as TACTIC (Vansnick 1986), building an asymmetric relation interpreted as strict preference. We recall here the definition of an asymmetric concordance-discordance relation (A-CDR). As we did for I-CR in Section 6.2, we define the more general irreflexive concordance discordance relations I-CDR, while A-CDR appears as a particular case of it.

Definition 52 (Irreflexive concordance-discordance relation)

Let \mathcal{P} be an irreflexive binary relation on $X = \prod_{i=1}^n X_i$. We say that \mathcal{P} is an irreflexive concordance-discordance relation (or, more briefly, that \mathcal{P} is an I-CDR) if there are:

- an asymmetric binary relation P_i° on each X_i ($i = 1, 2, \dots, n$),
- an asymmetric binary relation V_i° on each X_i ($i = 1, 2, \dots, n$), with $V_i^\circ \subseteq P_i^\circ$,
- a binary relation \triangleright° between disjoint subsets of N that is monotonic w.r.t. inclusion, i.e., such that (6) holds,

such that, for all $x, y \in X$,

$$x \mathcal{P} y \Leftrightarrow P^\circ(x, y) \triangleright P^\circ(y, x) \text{ and } V^\circ(y, x) = \emptyset, \quad (28)$$

where $P^\circ(x, y) = \{i \in N : x_i P_i^\circ y_i\}$ and $V^\circ(y, x) = \{i \in N : y_i V_i^\circ x_i\}$. A strict concordance-discordance relation is an asymmetric I-CDR. We also call strict concordance-discordance relations asymmetric concordance-discordance relations (A-CDR).

An example of an A-CDR can be obtained by adding a non-veto condition to the relation defined in Example 44. The result is a version of the outranking relation obtained using the TACTIC method proposed by Vansnick (1986).

Example 53 (TACTIC, Vansnick 1986)

The binary relation \mathcal{P} is a TACTIC outranking relation (with additive threshold) if there are $\varepsilon, P_i^\circ, w_i$ like in Example 44 and, for each $i \in N$, a strict semiorder V_i° such that $V_i^\circ \subseteq P_i^\circ$ and (V_i°, P_i°) forming a nested homogeneous chain of strict semiorders, such that:

$$x \mathcal{P} y \Leftrightarrow \sum_{i \in P^\circ(x, y)} w_i > \sum_{j \in P^\circ(y, x)} w_j + \varepsilon \text{ and } V^\circ(y, x) = \emptyset$$

where $V^\circ(y, x) = \{i \in N : y_i V_i^\circ x_i\}$. ◇

Clearly, a TACTIC outranking relation is an A-CDR. We recall the characterization of A-CDR obtained in Bouyssou and Pirlot (2006). It is easy to check that this result also characterizes I-CDR, when the relation is not supposed to be asymmetric.

Theorem 54 (Characterization of I-CDR and A-CDR)

A relation \mathcal{P} on X is an I-CDR iff it is irreflexive and satisfies RC1, RC2, Maj1, and Maj3, where Maj3 is satisfied as soon as the following condition, Maj3_{*i*}, is satisfied for all $i \in N$:

$$\left. \begin{array}{l} (x_i, a_{-i}) \mathcal{P} (y_i, b_{-i}) \\ \text{and} \\ (w_i, a_{-i}) \mathcal{P} (z_i, b_{-i}) \\ \text{and} \\ (y_i, c_{-i}) \mathcal{P} (x_i, d_{-i}) \\ \text{and} \\ (z_i, e_{-i}) \mathcal{P} (w_i, f_{-i}) \end{array} \right\} \Rightarrow \left\{ \begin{array}{l} (y_i, a_{-i}) \mathcal{P} (x_i, b_{-i}) \\ \text{or} \\ (z_i, c_{-i}) \mathcal{P} (w_i, d_{-i}), \end{array} \right. \quad (29)$$

for all $x_i, y_i, z_i, w_i \in X_i$ and all $a_{-i}, b_{-i}, c_{-i}, d_{-i}, e_{-i}, f_{-i} \in X_{-i}$. A relation \mathcal{P} is an A-CDR iff it is an asymmetric I-CDR. In the class of asymmetric relations (and, therefore in the class of irreflexive relations), conditions RC1, RC2, Maj1 and Maj3 are independent.

The only difference in the characterization of I-CDR w.r.t. I-CR is the substitution of Maj2 by Maj3, which, obviously, is a weakening of Maj2. Condition Maj3 is obtained from Maj2 exactly in the same way as M3 is obtained from M2 (see Remark 18 in Section 4 above).

In Section 6.1, we have established that the codual of an R-CR is an I-CR (and conversely) and that the asymmetric part of an R-CR is an I-CR. The situation is not the same for R-CDR and I-CDR. It is not hard to see that the codual of an I-CDR is an R-CR *with bonus* as defined below.

Definition 55 (Reflexive concordance relation with bonus R-CRB)

Let \succsim be a reflexive binary relation on $X = \prod_{i=1}^n X_i$. We say that \succsim is a reflexive concordance relation with bonus (or, more briefly, that \succsim is an R-CRB) if there are:

- a complete binary relation S_i on each X_i ($i = 1, 2, \dots, n$),
- an asymmetric binary relation V_i on each X_i ($i = 1, 2, \dots, n$), with $V_i \subseteq P_i$,
- a binary relation \supseteq between subsets of N having N for union that is monotonic w.r.t. inclusion, i.e., such that (6) holds,

such that, for all $x, y \in X$,

$$x \succsim y \Leftrightarrow S(x, y) \supseteq S(y, x) \text{ or } V(x, y) \neq \emptyset \quad (30)$$

where $S(x, y) = \{i \in N : x_i S_i y_i\}$ and $V(x, y) = \{i \in N : x_i V_i y_i\}$.

The correspondence between I-CDR and R-CRB is straightforward. If $\langle \triangleright^\circ, P_i^\circ, V_i^\circ \rangle$ is a representation of \mathcal{P} as an I-CDR, then a representation, $\langle \supseteq, S_i, V_i \rangle$, of its codual \mathcal{P}^{cd} as an R-CRB is obtained by defining S_i as the codual of P_i , \supseteq from \triangleright through (22), and taking $V_i = V_i^\circ$.

To illustrate the coduality of I-CDR and R-CRB, consider the relation \mathcal{P} defined in Example 53. Its codual \mathcal{P}^{cd} is as follows:

$$x \mathcal{P}^{cd} y \Leftrightarrow \sum_{i \in S(x, y)} w_i \geq \sum_{j \in S(y, x)} w_j - \varepsilon \text{ or } V^\circ(x, y) \neq \emptyset.$$

The first part of the above condition corresponds the definition of a semi-ordered weighted majority relation (Example 3) with S_i the codual of P_i° and $S(x, y) = \{i \in N : x_i S_i y_i\}$. The second branch of the alternative has an interpretation that is in sharp contrast with the concept of a veto: it says that x can be declared at least as good as y as soon as x is ‘‘much better’’ than y on some attribute (i.e., there is some attribute i on which $x_i V_i^\circ y_i$). Clearly, relation \mathcal{P}^{cd} is an R-CRB.

Remark 56

Definition 55 introduces a new preference structure that deserves attention in its own right. It is quite easy to derive a characterization of R-CRB using coduality and the characterization of I-CDR. We know from Lemma 48 that imposing $RC1$ (resp. $RC2$, $Maj1$) to a relation amounts to imposing $RC1$ (resp. $RC2$, $M2$) to its codual. It is readily seen that imposing $Maj3$, i.e., (29) (for all $i \in N$), to a relation \mathcal{R} amounts to imposing the following axiom, referred to as $M4$, to its codual \mathcal{R}^{cd} . We say that \mathcal{R}^{cd} satisfies $M4$ if, for all $i \in N$, for all $x_i, y_i, z_i, w_i \in X_i$ and all $a_{-i}, b_{-i}, c_{-i}, d_{-i}, e_{-i}, f_{-i} \in X_{-i}$, we have:

$$\left. \begin{array}{l} (x_i, b_{-i}) \mathcal{R}^{cd} (y_i, a_{-i}) \\ \text{and} \\ (w_i, d_{-i}) \mathcal{R}^{cd} (z_i, c_{-i}) \end{array} \right\} \Rightarrow \left\{ \begin{array}{l} (y_i, b_{-i}) \mathcal{R}^{cd} (x_i, a_{-i}) \\ \text{or} \\ (z_i, b_{-i}) \mathcal{R}^{cd} (w_i, a_{-i}) \\ \text{or} \\ (x_i, d_{-i}) \mathcal{R}^{cd} (y_i, c_{-i}) \\ \text{or} \\ (w_i, f_{-i}) \mathcal{R}^{cd} (z_i, e_{-i}). \end{array} \right. \quad (31)$$

Note that this axiom is clearly a weakened variant of $M1$ that simply adds to it one possible conclusion. Using this new axiom, we directly derive a characterization of an R-CRB from Theorem 54.

Proposition 57 (Characterization of R-CRB)

A relation \succsim on X is a reflexive concordance relation with bonus (R-CRB) iff it is reflexive and satisfies RC1, RC2, M2, and M4.

We leave to the reader the study of the particular case of R-CRB with attribute transitivity. •

6.4 The asymmetric part of an R-CDR

The asymmetric part of an R-CDR is not always an I-CDR, as shown by the following example.

Example 58

Let $N = \{1, 2, 3, 4\}$ and $X = \{x_1, y_1, z_1\} \times \{x_2, y_2\} \times \{x_3, y_3\} \times \{x_4, y_4\}$. Let \succsim on X be such that

$$x \succsim y \Leftrightarrow |S(x, y)| \geq |S(y, x)| \text{ and } V(y, x) = \emptyset,$$

where $S(x, y) = \{i \in N : x_i S_i y_i\}$ and $V(y, x) = \{i \in N : y_i V_i x_i\}$ with:

$$\begin{aligned} x_1 P_1 y_1 P_1 z_1, \\ x_2 P_2 y_2, x_3 P_3 y_3, x_4 P_4 y_4, \\ V_i = \emptyset, \text{ for all } i \in \{2, 3, 4\}, x_1 V_1 z_1. \end{aligned}$$

(where P_i denotes the asymmetric part of the complete relation S_i).

By construction, \succsim is an R-CDR. Denoting by \succ the asymmetric part of \succsim , it is easy to check that we have:

$$\begin{aligned} (x_1, x_2, x_3, y_4) \succ (z_1, x_2, x_3, x_4) \text{ and } \text{Not}[(x_1, x_2, x_3, y_4) \succ (y_1, x_2, x_3, x_4)], \\ (x_1, x_2, x_3, x_4) \succ (y_1, y_2, x_3, x_4) \text{ and } \text{Not}[(y_1, x_2, x_3, x_4) \succ (x_1, y_2, x_3, x_4)]. \end{aligned}$$

The first line implies that, w.r.t. \succ , the difference (x_1, z_1) is strictly larger than the difference (x_1, y_1) . The second line shows that the difference (x_1, y_1) is strictly larger than its opposite (y_1, x_1) . It is easy to see that this is impossible in an I-CDR (see Bouyssou and Pirlot 2006, Lemma 9, for details). \diamond

Let \succsim be an R-CDR with its representation $\langle \supseteq, S_i, V_i \rangle$. Using the definition of an R-CDR, we can say that its asymmetric part, denoted by \succ , is such that:

$$\begin{aligned} x \succ y \Leftrightarrow [x \succsim y \text{ and } y \not\prec x] \\ \Leftrightarrow \begin{cases} [S(x, y) \supseteq S(y, x) \text{ and } V(y, x) = \emptyset] \\ \text{and} \\ [\text{Not}[S(y, x) \supseteq S(x, y)] \text{ or } V(x, y) \neq \emptyset]. \end{cases} \end{aligned}$$

Let us denote by \succsim^c , the “concordance part” of the R-CDR \succsim , i.e., \succsim^c is the relation defined by:

$$x \succsim^c y \text{ iff } S(x, y) \supseteq S(y, x).$$

Clearly, \succsim^c is an R-CR with representation $\langle \supseteq, S_i \rangle$; we denote by \succ^c its asymmetric part (it is an A-CR) and by \sim^c its symmetric part. An alternative definition of \succ is the following:

$$x \succ y \text{ iff } \begin{cases} x \succ^c y \text{ and } V(y, x) = \emptyset \\ \text{or} \\ x \sim^c y, V(y, x) = \emptyset \text{ and } V(x, y) \neq \emptyset. \end{cases} \quad (32)$$

This means that the asymmetric part of \succ originates either from the asymmetric part of the concordance relation, provided there is no veto, or from the symmetric part of the concordance relation provided that there is no attribute on which $y_i V_i x_i$ and that there is at least one attribute for which x is “much better” than y (i.e., $x_i V_i y_i$).

As a conclusion, in the asymmetric part of an R-CDR, V_i may play the rôle of a veto (discarding a pair belonging to the asymmetric part of the concordance relation) or, on the contrary, play the rôle of a *bonus* (transforming an indifference, w.r.t. the concordance relation, into a strict preference).

Hence, the asymmetric part of an R-CDR is a more complex object than an R-CDR. It is not directly based on the application of concordance / non-discordance principle. This was already observed in Bouyssou and Marchant (2007a,b) for the analysis of the optimistic version of the ELECTRE TRI procedure that make use of the asymmetric part of an R-CDR. Combining the conditions introduced above giving rise to veto effects (29) and bonus effects (31), a characterization of such relations does not seem out of reach. We do not develop this point here.

7 Discussion

In this paper, we have given several axiomatic characterizations of outranking relations based on the concordance / non-discordance principle. This was done in a traditional conjoint measurement setting, i.e., using a binary relation defined on a Cartesian product as the only primitive. This leads to conditions entirely phrased in terms of \succsim that could be subject to empirical tests. Furthermore, the proofs of our results are constructive, which may be useful to devise assessment protocols. Different primitives are used in Pirlot (1997) who uses concepts from social choice and, hence, views outranking methods as techniques aggregating information available on each attribute

(see also Marchant 2007). For a detailed comparison of the conjoint measurement approach followed here and the approach based on social choice concepts, we refer to Bouyssou et al. (2006, Ch. 4–6).

We would like to conclude with the mention of some limitations of the present study and its relation to the literature. We refer to Bouyssou and Pirlot (2005a) for other elements of discussion, including the important issue of the sweeping consequences of imposing nice transitivity properties to outranking relations obtained using the concordance / non-discordance principle.

7.1 Limitations and directions for future work

The practitioner of outranking methods will surely have noticed a number of limitations of the present work. We mention here what we consider to be the most important ones.

Our emphasis has been on outranking methods that leads to the construction of “crisp” preference relations. This is a severe limitation since many well known outranking methods, like ELECTRE III (Roy 1978) or the various versions of PROMETHEE (Brans and Mareschal 2002; Brans and Vincke 1985) lead to the construction of valued relations, i.e., relations in which a number is attached to each ordered pair of alternatives reflecting the credibility or intensity of the underlying preference statement. Strictly speaking, this paper does not bring anything to the study of such methods. However, let us mention that it is possible to extend models (M) and (M*) to cover the case of valued binary relations: instead of comparing the value of F to a fixed threshold (0), this value can be seen as defining the valued relation. This calls for further study that will be the subject of a subsequent paper.

With the ELECTRE methods in mind, another limitation of our work is that it does not take “weak preference” (interpreted as an hesitation between indifference and strict preference) into account. This is also a severe limitation since “weak preference” plays an important part in some outranking methods. Nevertheless, as pointed out in Tsoukiàs et al. (2002), modelling “hesitation” is not an easy task and may necessitate the use of non-classical logics. Hence, it would be accurate to say that our results only deal with “idealized” outranking methods in which “hesitation” plays no rôle. Nevertheless our feeling is that these idealized methods remain close in spirit to the real outranking methods and that our conditions capture some of the central features of the latter.

Another limitation of this work is that our model for concordance remains too general. Indeed, in most methods weights are attached to each attribute

and their sum are used to perform the concordance test. Our models do not use weights at all and are simply based on an importance relation between coalition of attributes. We do not consider this limitation as very serious. On the practical side, dispensing with additive weights is indeed feasible, e.g., making use of symbolic inference techniques inspired from the Artificial Intelligence field: this was convincingly demonstrated with the “rough set” approach to MCDA, as presented, e.g., in Greco et al. (2001b, 2005). On the theoretical side, formulating conditions ensuring that the relation \succeq has an additive representation is not a difficult task. Completing it is unlikely to lead to conditions giving much insight on the underlying methods: since N is finite, these conditions will require a denumerable scheme of conditions that cannot be truncated.

A final important limitation of our work is that it is limited to the so-called “construction phase” of outranking methods. Since the relations obtained as the result of this construction phase do not, in general, possess remarkable properties of transitivity or completeness, using them to devise a recommendation is not an easy task and calls for the application of specific techniques (this is the so-called “exploitation phase”; on such techniques, see Bouyssou et al. 2006; Roy 1991; Roy and Bouyssou 1993; Vanderpooten 1990). We have little to say on how an axiomatic analysis that would include both the construction and the exploitation phases of outranking methods could be conducted. This is all the more true that there is no general agreement on what are the “best” techniques to derive a recommendation on the basis of an outranking relation in a given problem formulation. The sorting problem formulation seems the most easy one to deal with: since it only uses the comparison of alternatives with carefully selected profiles, so that intransitivity and incompleteness are not central issues here. Results of this kind have been obtained in Bouyssou and Marchant (2007a,b). The analysis of both the construction and exploitation phases of outranking methods clearly calls for more research (first results were obtained in Bouyssou et al. 2006, Ch. 5, in a social choice framework).

7.2 Relation to the literature

This paper is not the first attempt to analyze the concordance / non-discordance principle in a conjoint measurement perspective. In what follows, we try to position our contribution w.r.t. this earlier literature.

7.2.1 The approach using noncompensation

The work of Bouyssou and Vansnick (1986) on TACTIC is probably the first attempt to tackle the problem studied here. This paper uses a classical conjoint measurement setting to analyze A-CR and A-CDR. The central condition used in this paper to characterize A-CR is a condition called “non-compensation” that was introduced in Fishburn (1975, 1976, 1978). It says that, for all $x, y, z, w \in X$,

$$\left. \begin{array}{l} \succ(x, y) = \succ(z, w) \\ \succ(y, x) = \succ(w, z) \end{array} \right\} \Rightarrow [x \succ y \Leftrightarrow z \succ w], \quad (33)$$

where $\succ(x, y) = \{i \in N : x_i \succ_i y_i\}$ and \succ_i is the marginal binary relation on X_i induced by \succ , i.e., the relation such that, for all $x_i, y_i \in X_i$,

$$x_i \succ_i y_i \Leftrightarrow [(x_i, a_{-i}) \succ (y_i, a_{-i})], \text{ for all } a_{-i} \in X_{-i}].$$

When coupled with a suitable monotonicity condition, or when strengthened as

$$\left. \begin{array}{l} \succ(x, y) \supseteq \succ(z, w) \\ \succ(y, x) \subseteq \succ(w, z) \end{array} \right\} \Rightarrow [x \succ y \Rightarrow z \succ w], \quad (34)$$

the noncompensation condition seems to offer an interesting basis to analyze the concordance principle. However, as discussed at length in Bouyssou and Pirlot (2005a, Sect. 5.2) this approach to concordance does not allow to deal with the whole variety of concordance relations. For instance, it prevents one from using a threshold ε that would be greater than the weight of one attribute in the concordance part of TACTIC (see Example 53). Furthermore, this approach is not well suited to characterize a relation \succ in which the relations \succ_i would have nice transitivity properties. Finally, the non-compensation condition is very specific to “ordinal” methods. Using it does not allow to characterize concordance relation within a broader framework that also encompasses other types of relations.

Bouyssou and Vansnick (1986) have approached the introduction of discordance via the following weakening of the noncompensation condition saying that, for all $x, y, z, w \in X$,

$$\left. \begin{array}{l} \succ(x, y) = \succ(z, w) \\ \succ(y, x) = \succ(w, z) \end{array} \right\} \Rightarrow [x \succ y \Rightarrow w \not\succeq z]. \quad (35)$$

When $\succ(x, y) = \succ(z, w)$ and $\succ(y, x) = \succ(w, z)$ and $x \succ y$, this new condition allows to have either $z \succ w$ or $z \sim w$, where \sim is seen here as the symmetric complement of \succ . It is not difficult to see that TACTIC satisfies

this new condition. Indeed, the original noncompensation condition is (up to the point made earlier) satisfied for the concordance part of the method. The effect of discordance is to transform \succ relations into \sim ones. Hence, TACTIC satisfies condition (35).

There are several problems with this approach (that motivated one of the author of the present paper to abandon it and to develop the material presented here). We already mentioned that it is not flexible enough to cover all concordance relations of interest. We also noted that it is not well suited to introduce a linear arrangement of elements on each X_i . This last problem becomes even more important when discordance comes into play and there is a need to introduce links between \succ_i and V_i . The route followed in Bouyssou and Vansnick (1986) to tackle this problem is correct but somewhat ad hoc.

A final and major limitation of this approach is that, contrary to the approach taken here, it does not generalize outside the realm of asymmetric relations. As first noted in Bouyssou (1986, 1992) it is simple to reformulate the noncompensation condition so that it becomes adapted to the treatment of “at least as good as” relations. This leads to a condition of the type:

$$\left. \begin{array}{l} \succsim(x, y) = \succsim(z, w) \\ \succsim(y, x) = \succsim(w, z) \end{array} \right\} \Rightarrow [x \succsim y \Leftrightarrow z \succsim w], \quad (36)$$

where $\succsim(x, y) = \{i \in N : x_i \succsim_i y_i\}$ and \succsim_i is the marginal binary relation on X_i induced by \succ , i.e., the relation such that, for all $x_i, y_i \in X_i$,

$$x_i \succsim_i y_i \Leftrightarrow [(x_i, a_{-i}) \succ (y_i, a_{-i}), \text{ for all } a_{-i} \in X_{-i}].$$

This approach was later developed in Fargier and Perny (2001) and Dubois et al. (2001, 2003). As was the case with the original noncompensation condition, (36) does not allow to deal with the whole variety of concordance relations (see Bouyssou and Pirlot 2005a, Examples 30 and 31), is very specific to this type of relations, and is not very well suited to introduce a linear arrangement of the elements on each X_i . Even worse, the route followed by Bouyssou and Vansnick (1986) in order to cope with veto effects does not work here. Mimicking the above approach, we could consider a condition saying that

$$\left. \begin{array}{l} \succsim(x, y) = \succsim(z, w) \\ \succsim(y, x) = \succsim(w, z) \end{array} \right\} \Rightarrow [x \succ y \Rightarrow w \not\succeq z], \quad (37)$$

but even such a weak condition fails with most outranking relations. Indeed, as discussed in Section 6, an indifference situation between two alternatives obtained on the sole basis of the concordance test may be broken in an unpredictable way by veto effects. This leads to possible violations of condition

(37). It is easy to build examples, e.g., using ELECTRE I, in which one has $\succsim(x, y) = \succsim(z, w)$, $\succsim(y, x) = \succsim(w, z)$, $x \sim y$ and $z \sim w$ when only the concordance part of the method is used. Introducing veto effects, one can easily obtain $x \succ y$ and $w \succ z$, violating (37).

The approach using variants of noncompensation is often seen to have some advantages w.r.t. the approach used here. First it has been shown (see Bouyssou 1992, or Dubois et al. 2003) to be particularly well suited to transfer “Arrow-like” results (i.e., results showing that requiring \succsim to have “nice properties” induces a very undesirable repartition of “importance” among the various attributes) to the context of MCDA. Nevertheless, we have shown in Bouyssou and Pirlot (2002a) that a similar analysis can also be performed in our more general framework, which leads to even more powerful results. Noncompensation-like conditions are also often seen as being “simpler” and “more natural” than the type of conditions used here. We disagree. Besides the fact that “simplicity” is a very subjective criterion, we would like to point out that we have strived to present conditions that are entirely phrased using our primitives, i.e., the relation \succsim on X . If one tries to reformulate the noncompensation condition with such a constraint, the result does not appear to be much simpler than our conditions.

7.2.2 The approach of Greco et al. (2001a)

Greco et al. (2001a) give results that are closely related to the ones presented here. Motivated by the ELECTRE I method, their aim was to characterize the particular class of R-CDR in which the relation \supseteq is such that, for all $A, B \subseteq N$ such that $A \cup B = N$,

$$A \supseteq B \Rightarrow A \supseteq N. \quad (38)$$

It is easy to see that this additional condition is satisfied in ELECTRE I (see Example 4).

As we did above, the approach of Greco et al. (2001a) is based on conditions limiting the number of distinct equivalence classes of \succsim_i^* (see Bouyssou et al. 1997).

When discordance is not taken into account, the central condition used by Greco et al. (2001a) is the following. We say that \succsim is *super-coarse* on attribute $i \in N$ if, for all $x_i, y_i, z_i, w_i, r_i, s_i \in X_i$ and all $a_{-i}, b_{-i}, c_{-i}, d_{-i} \in X_{-i}$,

$$\left. \begin{array}{l} (x_i, a_{-i}) \succsim (y_i, b_{-i}) \\ \text{and} \\ (z_i, c_{-i}) \succsim (w_i, d_{-i}) \end{array} \right\} \Rightarrow \left\{ \begin{array}{l} (x_i, c_{-i}) \succsim (y_i, d_{-i}) \\ \text{or} \\ (r_i, a_{-i}) \succsim (s_i, b_{-i}). \end{array} \right. \quad (39)$$

This condition is clear strengthening of $RC1$. It is not difficult to see that a \succsim is super-coarse on attribute $i \in N$ if and only if \succsim_i^* is complete and has at most two distinct equivalence classes.

It is important to notice that, on its own, super-coarseness does not imply independence (in our framework independence is ensured via the use of $RC2$). Therefore nothing prevents (x_i, x_i) and (y_i, y_i) from belonging to two distinct equivalence classes of \succsim_i^* . In order to characterize R-CR satisfying (38), Greco et al. (2001a) use super-coarseness as well as an additional condition saying that, for all $i \in N$, all $x_i, y_i, w_i \in X_i$ and all $a_{-i}, b_{-i} \in X_{-i}$,

$$(x_i, a_{-i}) \succsim (y_i, b_{-i}) \Rightarrow (w_i, a_{-i}) \succsim (w_i, b_{-i}). \quad (40)$$

This is a rather strong condition implying at the same time independence and the fact that the null preference differences (w_i, w_i) always belong to the first equivalence class of \succsim_i^* . Its intuitive content appears to be limited.

As discussed in Bouyssou and Pirlot (2005a, Sect. 5.2), the result of Greco et al. (2001a) on concordance relations deals with a particular class of R-CR, in contrast with our own results. Furthermore, as this analysis does not explicitly uses conditions $RC1$ and $RC2$, it not conducted within the broader framework of model (M), which may be viewed as a drawback in view of comparing concordance relations with other types of relations. Finally, condition (40) is not very easy to interpret.

Greco et al. (2001a) have noted that adding conditions $AC1$, $AC2$ and $AC3$ implies that the resulting relations S_i must be semiorders. Nevertheless, they have not studied the independence of these conditions with respect to conditions (39) and (40). As shown in Bouyssou and Pirlot (2005a, Lemma 27), conditions $AC1_i$ and $AC2_i$ turn out to be equivalent for R-CR. Independence issues were indeed the most delicate ones to tackle in Bouyssou and Pirlot (2005a) and Bouyssou and Pirlot (2007).

Although we already mentioned in Bouyssou et al. (1997) that model (M) offers an adequate framework to tackle R-CDR, it took us some time to devise adequate conditions that would characterize R-CDR within this framework. Greco et al. (2001a) were the first to come up with such conditions; we already mentioned that our condition $M3$ is inspired from their work. The main condition used in Greco et al. (2001a) to characterize R-CDR satisfying (38) is the following. We say that \succsim is *super-coarse with veto* on attribute $i \in N$ if, for all $x_i, y_i, z_i, w_i, r_i, s_i \in X_i$ and all $a_{-i}, b_{-i}, c_{-i}, d_{-i} \in X_{-i}$,

$$\left. \begin{array}{l} (x_i, a_{-i}) \succsim (y_i, b_{-i}) \\ \text{and} \\ (z_i, c_{-i}) \succsim (w_i, d_{-i}) \\ \text{and} \\ (r_i, e_{-i}) \succsim (s_i, f_{-i}) \end{array} \right\} \Rightarrow \left\{ \begin{array}{l} (x_i, c_{-i}) \succsim (y_i, d_{-i}) \\ \text{or} \\ (r_i, a_{-i}) \succsim (s_i, b_{-i}). \end{array} \right. \quad (41)$$

This condition is a clear weakening of (39) that have inspired our own weakening of $M2$ to obtain $M3$. The main result in Greco et al. (2001a) is that R-CDR satisfying (38) are characterized by the conjunction of conditions (40) and (41). We do not repeat here the comments made above: this result uses condition (40) and is not conducted within a broader framework, like the one provided by model (M). Nevertheless, it was the first approach that convincingly took discordance into account in a conjoint measurement setting.

References

- Aleskerov, F., Monjardet, B., 2002. *Utility Maximization, Choice and Preference*. Springer-Verlag, Heidelberg.
- Bouyssou, D., 1986. Some remarks on the notion of compensation in MCDM. *European Journal of Operational Research* 26, 150–160.
- Bouyssou, D., 1992. On some properties of outranking relations based on a concordance-discordance principle. In: Duckstein, L., Goicoechea, A., Zionts, S. (Eds.), *Multiple criteria decision making*. Springer-Verlag, Berlin, pp. 93–106.
- Bouyssou, D., 1996. Outranking relations: Do they have special properties? *Journal of Multi-Criteria Decision Analysis* 5, 99–111.
- Bouyssou, D., Marchant, Th., 2007a. An axiomatic approach to noncompensatory sorting methods in MCDM, I: The case of two categories. *European Journal of Operational Research* 178 (1), 217–245.
- Bouyssou, D., Marchant, Th., 2007b. An axiomatic approach to noncompensatory sorting methods in MCDM, II: More than two categories. *European Journal of Operational Research* 178 (1), 246–276.
- Bouyssou, D., Marchant, Th., Pirlot, M., Tsoukiàs, A., Vincke, Ph., 2006. *Evaluation and decision models: Stepping stones for the analyst*. Springer, New York.
- Bouyssou, D., Pirlot, M., 1999. Conjoint measurement without additivity and transitivity. In: Meskens, N., Roubens, M. (Eds.), *Advances in Decision Analysis*. Kluwer, Dordrecht, pp. 13–29.
- Bouyssou, D., Pirlot, M., 2002a. A characterization of strict concordance relations. In: Bouyssou, D., Jacquet-Lagrèze, É., Perny, P., Słowiński, R., Vanderpooten, D., Vincke, Ph. (Eds.), *Aiding Decisions with Multiple Criteria: Essays in Honour of Bernard Roy*. Kluwer, Dordrecht, pp. 121–145.
- Bouyssou, D., Pirlot, M., 2002b. Nontransitive decomposable conjoint measurement. *Journal of Mathematical Psychology* 46, 677–703.
- Bouyssou, D., Pirlot, M., 2004a. ‘Additive difference’ models without additivity and subtractivity. *Journal of Mathematical Psychology* 48, 263–291.

- Bouyssou, D., Pirlot, M., 2004b. Preferences for multiattributed alternatives: Traces, dominance, and numerical representations. *Journal of Mathematical Psychology* 48, 167–185.
- Bouyssou, D., Pirlot, M., 2005a. A characterization of concordance relations. *European Journal of Operational Research* 167/2, 427–443.
- Bouyssou, D., Pirlot, M., 2005b. Notes on strict concordance relations. In: Barthélemy, J.-P., Lenca, Ph. (Eds.), *Advances in Multi-Criteria Decision Aid*. ENST Bretagne, Brest, pp. 21–31, Also available at <http://lamsade.dauphine.fr/~bouyssou>.
- Bouyssou, D., Pirlot, M., 2006. An axiomatic approach to TACTIC, Cahier du LAMSADE # 238, Université Paris Dauphine. Available at <http://lamsade.dauphine.fr/~bouyssou>.
- Bouyssou, D., Pirlot, M., 2007. Further results on concordance relations. *European Journal of Operational Research* 181, 505–514.
- Bouyssou, D., Pirlot, M., Vincke, Ph., 1997. A general model of preference aggregation. In: Karwan, M. H., Spronk, J., Wallenius, J. (Eds.), *Essays in Decision Making*. Springer Verlag, Berlin, pp. 120–134.
- Bouyssou, D., Vansnick, J.-C., 1986. Noncompensatory and generalized noncompensatory preference structures. *Theory and Decision* 21, 251–266.
- Brans, J.-P., Mareschal, B., 2002. PROMETHEE-GAIA. Une méthodologie d’aide à la décision en présence de critères multiples. Éditions de l’Université de Bruxelles, Brussels.
- Brans, J.-P., Vincke, Ph., 1985. A preference ranking organisation method. (The PROMETHEE method for multiple criteria decision-making). *Management Science* 31 (6), 647–656.
- Cozzens, M. B., Roberts, F., 1982. Double semiorders and double indifference graphs. *SIAM Journal on Algebraic and Discrete Methods* 3, 566–583.
- Debreu, G., 1960. Topological methods in cardinal utility theory. In: Arrow, K. J., Karlin, S., Suppes, P. (Eds.), *Mathematical methods in the Social Sciences*. Stanford University Press, Stanford, pp. 16–26.
- Doignon, J.-P., Monjardet, B., Roubens, M., Vincke, Ph., 1988. Biorder families, valued relations and preference modelling. *Journal of Mathematical Psychology* 30, 435–480.
- Dubois, D., Fargier, H., Perny, P., Prade, H., 2001. Towards a qualitative multicriteria decision theory. In: *Proceedings of the EUROFUSE Workshop on Preference Modelling and Applications*. Granada, Spain, April 25–27, 2001, pp. 121–129.
- Dubois, D., Fargier, H., Perny, P., Prade, H., 2003. A characterization of generalized concordance rules in multicriteria decision-making. *International Journal of Intelligent Systems* 18 (7), 751–774.
- Fargier, H., Perny, P., 2001. Modélisation des préférences par une règle de concordance généralisée. In: Colorni, A., Paruccini, M., Roy, B. (Eds.), *A-MCD-A*,

- Aide Multicritère à la Décision / Multiple Criteria Decision Aid. European Commission, Joint Research Centre, pp. 99–115.
- Fishburn, P. C., 1975. Axioms for lexicographic preferences. *Review of Economic Studies* 42, 415–419.
- Fishburn, P. C., 1976. Noncompensatory preferences. *Synthese* 33, 393–403.
- Fishburn, P. C., 1978. A survey of multiattribute/multicriteria evaluation theories. In: Zionts, S. (Ed.), *Multicriteria problem solving*. Springer Verlag, Berlin, pp. 181–224.
- Fishburn, P. C., 1990. Continuous nontransitive additive conjoint measurement. *Mathematical Social Sciences* 20, 165–193.
- Fishburn, P. C., 1991. Nontransitive additive conjoint measurement. *Journal of Mathematical Psychology* 35, 1–40.
- Greco, S., Matarazzo, B., Słowiński, R., 2001a. Axiomatic basis of noncompensatory preferences, communication to *FUR X*, 30 May–2 June, Torino, Italy.
- Greco, S., Matarazzo, B., Słowiński, R., 2001b. Rough sets theory for multicriteria decision analysis. *European Journal of Operational Research* 129, 1–7.
- Greco, S., Matarazzo, B., Słowiński, R., 2005. Decision rule approach. In: Figueira, J., Greco, S., Ehrgott, M. (Eds.), *Multiple Criteria Decision Analysis: State of the Art Surveys*. Springer-Verlag, Boston, pp. 507–562.
- Keeney, R. L., Raiffa, H., 1976. *Decisions with multiple objectives: Preferences and value tradeoffs*. Wiley, New-York.
- Krantz, D. H., Luce, R. D., Suppes, P., Tversky, A., 1971. *Foundations of measurement, vol. 1: Additive and polynomial representations*. Academic Press, New-York.
- Luce, R. D., 1956. Semiorders and a theory of utility discrimination. *Econometrica* 24, 178–191.
- Marchant, Th., 2007. An axiomatic characterization of different majority concepts. *European Journal of Operational Research* 179 (1), 160–173.
- Pirlot, M., 1997. A common framework for describing some outranking procedures. *Journal of Multi-Criteria Decision Analysis* 6, 86–92.
- Roubens, M., Vincke, Ph., 1985. *Preference modelling*. Springer Verlag, Berlin.
- Roy, B., 1968. Classement et choix en présence de points de vue multiples (la méthode ELECTRE). *RIRO* 2, 57–75.
- Roy, B., 1978. ELECTRE III : un algorithme de classement fondé sur une représentation floue des préférences en présence de critères multiples. *Cahiers du CERO* 20, 3–24.
- Roy, B., 1991. The outranking approach and the foundations of ELECTRE methods. *Theory and Decision* 31, 49–73.
- Roy, B., Bertier, P., 1973. La méthode ELECTRE II : une application au media-planning. In: Ross, M. (Ed.), *OR'72*. North Holland, Amsterdam, pp. 291–302.
- Roy, B., Bouyssou, D., 1993. *Aide multicritère à la décision : Méthodes et cas*.

Economica, Paris.

- Tsoukiàs, A., Perny, P., Vincke, P., 2002. From concordance/discordance to the modelling of positive and negative reasons in decision aiding. In: Bouyssou, D., Jacquet-Lagrèze, É., Perny, P., Słowiński, R., Vanderpooten, D., Vincke, Ph. (Eds.), *Aiding Decisions with Multiple Criteria: Essays in Honour of, Bernard Roy*. Kluwer, Dordrecht, pp. 147–174.
- Tversky, A., 1969. Intransitivity of preferences. *Psychological Review* 76, 31–48.
- Vanderpooten, D., 1990. The construction of prescriptions in outranking methods. In: Bana e Costa, C. A. (Ed.), *Readings in multiple criteria decision aid*. Springer Verlag, Berlin, pp. 184–215.
- Vansnick, J.-C., 1986. On the problems of weights in MCDM (the noncompensatory approach). *European Journal of Operational Research* 24, 288–294.
- Vincke, Ph., 1992. *Multi-criteria decision aid*. Wiley, New York, original version in French “*L’aide multicritère à la décision*”, Éditions de l’Université de Bruxelles — Éditions Ellipses, Brussels, 1989.
- Vincke, Ph., 1999. Outranking approach. In: Gal, T., Stewart, T., Hanne, T. (Eds.), *Multicriteria decision making, Advances in MCDM models, algorithms, theory and applications*. Kluwer, pp. 11.1–11.29.