



**HAL**  
open science

## **Fringing Reefs of Reunion Island and Eutrophication Effects - Part 1: Long-Term Monitoring of Two Shallow Coral Reef Communities**

Catherine Tourrand, Odile Naim, Lionel Bigot, Chrisophe Cadet, Bruce Cauvin, Stuart Semple, Lucien F. Montaggioni, Pascale Chabanet, J. Henrich Bruggemann

► **To cite this version:**

Catherine Tourrand, Odile Naim, Lionel Bigot, Chrisophe Cadet, Bruce Cauvin, et al.. Fringing Reefs of Reunion Island and Eutrophication Effects - Part 1: Long-Term Monitoring of Two Shallow Coral Reef Communities. *Atoll Research Bulletin*, 2013, 596, pp.1–35. 10.5479/si.00775630.596 . hal-00909558

**HAL Id: hal-00909558**

**<https://hal.science/hal-00909558>**


Submitted on 27 Sep 2021

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial 4.0 International License



**FRINGING REEFS OF REUNION ISLAND  
AND EUTROPHICATION EFFECTS**

No. 596

**PART 1: Long-Term Monitoring of Two Shallow Coral Reef Communities**

*Catherine Tourrand, Odile Naim, Lionel Bigot, Christophe Cadet, Bruce Cauvin,  
Stuart Semple, Lucien F. Montaggioni, Pascale Chabanet and Henrich Bruggemann*

No. 597

**PART 2: Long-Term Monitoring of Primary Producers**

*Odile Naim, Catherine Tourrand, Enric Ballesteros, Stuart Semple, Lionel Bigot,  
Bruce Cauvin, Pascale Cuet, and Lucien F. Montaggioni*

No. 598

**PART 3: Long-Term Monitoring of Living Corals**

*Odile Naim, Catherine Tourrand, Gerard F. Faure, Lionel Bigot, Bruce Cauvin,  
Stuart Semple and Lucien F. Montaggioni*

---

# ATOLL RESEARCH BULLETIN

---



Smithsonian Institution  
Scholarly Press

# **FRINGING REEFS OF REUNION ISLAND AND EUTROPHICATION EFFECTS**

## **PART 1: Long-Term Monitoring of Two Shallow Coral Reef Communities**

*Catherine Tourrand, Odile Naim, Lionel Bigot, Christophe Cadet, Bruce Cauvin,  
Stuart Semple, Lucien F. Montaggioni, Pascale Chabanet and Henrich Bruggemann*

## **PART 2: Long-Term Monitoring of Primary Producers**

*Odile Naim, Catherine Tourrand, Enric Ballesteros, Stuart Semple, Lionel Bigot, Bruce Cauvin,  
Pascale Cuet, and Lucien F. Montaggioni*

## **PART 3: Long-Term Monitoring of Living Corals**

*Odile Naim, Catherine Tourrand, Gerard F. Faure, Lionel Bigot, Bruce Cauvin, Stuart Semple  
and Lucien F. Montaggion*

**Atoll Research Bulletin No. 596 ♦ 15 November 2013**



Smithsonian Institution  
Scholarly Press

All statements made in papers published in the *Atoll Research Bulletin* are the sole responsibility of the authors and do not necessarily represent the views of the Smithsonian Institution or of the editors of the Bulletin. Articles submitted for publication in the *Atoll Research Bulletin* should be original papers and must be made available by authors for open access publication. Manuscripts should be consistent with the “Author Formatting Guidelines for Publication in the Atoll Research Bulletin.” All submissions to the Bulletin are peer reviewed and, after revision, are evaluated prior to acceptance and publication through the publisher’s open access portal, Open SI (<http://opensi.si.edu>).

Published by SMITHSONIAN INSTITUTION SCHOLARLY PRESS  
P.O. Box 37012, MRC 957  
Washington, D.C. 20013-7012  
[www.scholarlypress.si.edu](http://www.scholarlypress.si.edu)

The rights to all text and images in this publication are owned either by the contributing authors or third parties. Fair use of materials is permitted for personal, educational, or noncommercial purposes. Users must cite author and source of content, must not alter or modify the content, and must comply with all other terms or restrictions that may be applicable. Users are responsible for securing permission from a rights holder for any other use.

ISSN: 0077-5630 (online)

# FRINGING REEFS OF REUNION ISLAND AND EUTROPHICATION EFFECTS. PART 1: LONG-TERM MONITORING OF TWO SHALLOW CORAL REEF COMMUNITIES

BY

CATHERINE TOURRAND,<sup>1 2\*</sup> ODILE NAIM,<sup>3 2</sup> LIONEL BIGOT,<sup>2</sup> CHRISTOPHE CADET,<sup>4</sup> BRUCE CAUVIN<sup>4</sup>, STUART SEMPLE,<sup>5</sup> LUCIEN F. MONTAGGIONI,<sup>6</sup> PASCALE CHABANET<sup>7</sup> and HENRICH BRUGGEMANN<sup>2</sup>

## ABSTRACT

This study assesses changes in subtidal benthic communities on the largest reef flat in Reunion, *Saint-Gilles La Saline*, using several types of surveys. Temporal and spatial trends are documented over a 22 year period (1987-2009), thus spanning the 1998 and 2000s bleaching events. The most plausible explanations for the observed trends are proposed. We chose two sites that are characterized by two types of community and metabolism: (1) an oligotrophic site dominated by *Acropora* corals (Site-Toboggan), where sea-urchins are numerous and macroalgae rare and (2) a dystrophic site dominated by non-*Acropora* corals, mostly massive and submassive, where macroalgae abound and sea-urchins are almost absent (Site-Planch'Alizés). Results are presented in three parts : Part 1 : general trends of the communities, part 2 : primary producers, part 3 : living corals.

Part 1 presents three surveys. Survey 1 reports status and trends across the reef flats in 1993, 1996, and 2002, with all attached benthic components reported at the level of the species where possible. Survey 2 reports composition and changes in associated sedentary organisms such as sea urchins, holothurids, and the Pomacentridae fish *Stegastes*. Survey 3 focuses on the period 1998 to 2009 on permanent transects established in 1987 and monitored periodically, partly as a contribution to the Global Coral Reef Monitoring Network (GCRMN). In general, community structures at both locations remained stable, with primary producer coverage around 5% at Toboggan (T) and around 60% at Planch'Alizés (P). Living coral coverage averaged around 17% at both sites, but it was dominated by *Acropora* on Site-T (68%) and by massive corals on Site-P (88%). Regular echinoids were very abundant at T but almost absent from P, while the opposite was true for holothurids. The territorial damselfish *Stegastes* was of comparable abundance at both sites (<1 individuals per m<sup>2</sup>). There were large changes in coral cover in some zones, especially due to growth in *Acropora* from 1993 to 2002. On the two GCRMN permanent transects, 1999 and 2004 were turning points, so results are reported in three blocks: 1987-98, 1999-2003 and 2004-09. From 1987 to 1998 primary

---

<sup>1</sup> Département de Physique, Université de la Réunion, Saint-Denis, France (DOM)

<sup>2</sup> Laboratoire ECOMAR, Université de la Réunion, Saint-Denis, France (DOM)

<sup>3</sup> Laboratoire GMGL, Université de Bretagne Sud, Vannes, France

<sup>4</sup> GIP Réserve Nationale Marine de la Réunion, France (DOM)

<sup>5</sup> Department of Life Sciences, Roehampton University, England

<sup>6</sup> UMR 6536, Université d'Aix-Marseille, Marseille, France

<sup>7</sup> IRD, Sainte-Clotilde, Réunion, France (DOM)

\* catherine.tourrand@univ-reunion.fr

producers were inconspicuous on T transect but increased after that, suggesting abiotic conditions had become more favorable to algae. On the other transect (P-one), algae began to decline from 1999 and massive corals increased. Possible causes for these changes are discussed.

*Key words:* Coral reefs, benthic community, subtidal, stability, benthic primary producers, living corals, *Acropora*, sea urchins, holothurids, *Stegastes*.

## INTRODUCTION

Forty years ago, reef workers and oceanographers claimed that the Indian Ocean was the least known of the tropical oceans (Behrman, 1981; Stoddart, 1973). A workshop held in Mombasa in 1997 put an end to this situation through releasing a synthesis of existing knowledge on major Indian Ocean coastal ecosystems (McClanahan *et al.*, eds, 2000). The widespread degradation of the coastal habitats and their associated ecosystem services was recently reported (Sheppard, 2000; Lindèn *et al.*, eds, 2002; Souter & Lindèn, eds, 2005; Obura *et al.*, eds, 2008; Wilkinson, ed., 2008; Burke *et al.*, 2011), especially as a result of previous bleaching events that occurred in 1998, 2000, 2005 and 2010.

In oceanic islands, highly dependent on coastal resources, increasing urbanization and economic development along the coastlines exacerbates ecological problems. The Mascarene Islands (e.g. Reunion, Mauritius and Rodrigues), belong to this category, suffering from fast human-derived disturbances, largely concentrated along their narrow coasts. It is difficult to estimate reef degradation since man's first arrival occurred in the 17<sup>th</sup> century. Gardiner (1936) collected corals from Mauritius and the St Brandon Bank during the Percy Sladen Expedition of 1905. Although some studies on specific taxonomic groups were undertaken in Mauritius in the intervening period, no further coral reef studies were made until the early 1970s. The studies by Pichon (1971), Faure & Montaggioni (1970, 1971a, 1971b, 1976), Faure (1974, 1975, 1977), Montaggioni (1978), Montaggioni & Faure (1980), Bouchon (1981, 1996), and Faure (1982) provided baseline data for the Mascarene coral reefs, appropriately just prior to the recent period of rapid demographic and economic development of these islands. The transformation of original coastal habitats for agricultural and urban use have accompanied these developments (Cheke, 1987) and contributed to reef degradation.

The mapping of the Mascarene reefs using satellite imagery indicates that the shallow reef ecosystems cover some 705 km<sup>2</sup> (Turner & Klaus, 2005). In Reunion, the fringing reefs are good test sites due to their small sizes (12 km<sup>2</sup> of shallow habitats) (Spalding *et al.*, 2001). Their study started from the 1970s, in particular the largest *Saint-Gilles La Saline* reef (See references herein). As a result of the increasing human population (1982: 515,814 inhabitants; 2010: 833,000 inhab., INSEE), urbanization and beach activities have greatly increased reef utilization, degradation and other impacts (Faure, 1994; Naim *et al.*, 2000). In addition, natural events, such as bleaching (McClanahan *et al.*, 2007) and extreme low tides, chronically affect the subtidal coral communities.

In Reunion, Guillaume *et al.* (1983) first recorded reef degradation. In some sectors of *Saint-Gilles La Saline*, *Acropora* assemblages died and macroalgae became dominant, and the reef degradation was attributed to water nutrient-enrichment *via* submarine groundwater discharge (Cuet *et al.*, 1988). Nowadays, in spite of local

degradation, the major part of the Reunion shallow reef flats are still dominated by *Acropora* populations (Bruggemann *et al.*, 2007; Faure, 2009), and Reunion still harbours original benthic communities mainly because it is often just outside the core area of major temperature anomalies (McClanahan *et al.*, 2007).

Phase-shifts from one persistent assemblage of species to another have become increasingly commonplace in many coral ecosystems. Coral reef science, monitoring and global assessments have focused mainly on producing detailed descriptions of reef decline, and continue to pay insufficient attention to the underlying processes causing degradation (Hughes *et al.*, 2010). Long term records on benthic assemblages are thus important to assess the resilience capacity of different communities and predict future changes (Done, 1997). Such data are rare. Using a combination of different datasets, this work focuses on the spatio-temporal evolution, from 1987 to 2009, of the major components of two major subtidal reef community structures, one classified as being in “good health” (ACR), one considered to be in “bad health” (MAS). These communities are respectively observed at Site-Toboggan (Site-T) and at Site-Planch’Alizés (Site-P).

The results will be presented in three distinct parts: (1) Part 1 (the present paper) discusses the spatio-temporal variation, from 1993 to 2002, and then the temporal variation from 1987 to 2009 experienced respectively by the two (ACR, MAS) community structures, in terms of zonation, patterns of primary producers, living and dead corals, echinoids, holothurids and sedentary species of fish; (2) Part 2 (Naim *et al.*, 2013a) deals with the temporal variation that affected the benthic primary producer community in ACR and MAS, in terms of dominant forms (cyanophytes, turfs, encrusting coralline, and macroalgae) and dominant species. The species composition, diversity and biomass of primary producers will be also analyzed at the two sites; (3) Part 3 (Naim *et al.*, 2013b) focuses on the temporal variation in coral abundance, diversity and coral species as bioindicators.

## MATERIAL AND METHODS

### General environmental setting and study sites

**Island and climatic setting.** Reunion (21°07'-19°40' S, 55°13'-61°13' E) is a steep volcanic island, regarded as not exceeding 2.1 mya in age (Bachelery, 2000), and typified by a narrow shelf. Coral reef development is limited to the drier west and south coasts, forming a series of fringing reefs intersected by gullies that transfer terrestrial run-off during tropical storms (Montaggioni & Faure, 1980). Reefs are exposed to high wave energy conditions. During the cool season, from June to August, southeast trade winds (speed: 7-11 km.h<sup>-1</sup>) generate southeast 1-2 m high swells that affect most of the coastline. During the hot season, from November to February, winds and swells are weak (wind speed: less than 6 km.h<sup>-1</sup>; swell under 0.5 m high) but cyclones occur occasionally, generating whirlwinds of 120-250 km.h<sup>-1</sup> and swells of 4-11 m amplitude, although the west coast often escapes the harshest impacts. The reef is also affected by 3-5 m-high swells, mainly from the southwest, generated by polar depressions starting in the Marion Island region (in the Roaring Forties). Tides are semi-diurnal. The tidal range, greatly modified by swells, varies from 0.10 m (neaps) to 0.95 m (springs). Reunion held two world records for 3-days precipitation events: 3.24 meters that fell during January 1980 (cyclone Hyacinthe), and 3.93 m recorded in February 2007 (cyclone Gamède). Nevertheless, on the leeward side, mean annual rainfall is only 570 mm, and rain occurs mainly between

December and March. Mean costal sea surface temperature varies from  $23.7 \pm 0.3$  °C in cold season to  $27.7 \pm 0.4$  °C in hot season (Conand *et al.*, 2007), but can reach more than 31.0°C on the reef flats (Cuet, 1989a).

The *Saint-Gilles La Saline reef*, located on the west coast, is the most developed. It is typified by two subtidal reef flats of < 1.5 m depth, 250-550 m in width, and extending over 9 km. This nearly continuous reef rim is widest in the south and slightly deeper (**Fig.1**). Narrow passes occur where freshwater runoff periodically overflows the shoreline (Ermitage and Trois Bassins). Runoff occurs intermittently in the hot and wet season, but compared to other large Reunion reefs, such as those of Saint-Leu and Saint-Pierre, the *Saint-Gilles La Saline reef* can be considered to be moderately influenced by land-derived freshwater and sediment discharge, except when the island is crossed by the trajectories of strong cyclones, like Gamède (2007) (Kolasinski *et al.*, 2011). Sediments are predominantly of biogenic origin (Montaggioni, 1978, Gabrié & Montaggioni, 1982). From the coast seawards, the studied area refers to the inner reef flat. The outermost reef flat zone was not studied.

#### **Water energy and physico-chemistry.**


Average current speed varies between 0.3 and 1.0 km.h<sup>-1</sup> and tidal currents rarely exceed 2 km.h<sup>-1</sup> (Chazottes *et al.*, 2002). At the two studied sites, Toboggan (Site-T) and Planch'Alizés (Site-P), the flow of oceanic waters is dependent on swell, tide, wind, and topography (Cordier, 2007). In the north, Site-T receives a strong flux of oceanic waters, while further south, at Site-P, a depression in the reef front serves as an outlet of reef flat waters. Thus, at Site-P, benthic communities are bathed in waters that have transited over the reef flat (Conand *et al.*, 2002a). In addition, submarine groundwater discharge occurs near shore during low tides, and is drained by the Site-P outlet (Cuet *et al.*, 1988). Thus, P waters display very variable salinities and silicate concentrations (Cuet, 1989a, 1989b). Moreover, P waters have slower flow rates, higher mean and more variable sea surface temperature (Denis *et al.*, 2009). Thus, physico-chemical parameters of waters at Site-P greatly differ from those measured at Site-T (see discussions in parts II and III, Naim *et al.*, 2013a, 2013b).

#### **Background**

Selection of Site-Toboggan and Site-Planch'Alizés is based on four previous studies carried out on the *Saint-Gilles La Saline* reef flat: (1) Cuet *et al.* (1988) mapped hydrographic conditions, silicate concentrations and reef degradation from data obtained along the 17 radial lines on the entire reef flat; (2) Naim (2006) described the *Saint-Gilles La Saline* reef community structure through the analysis of the benthic components along the same 17 radial lines. The benthic community structure appeared to be typified by two distinct assemblages: (a) a dominant assemblage, labelled ACR, made up of branching *Acropora* corals, with a very low primary producer coverage, abundant sea urchins and a low number of holothurids; (b) an assemblage, labelled MAS, dominated by massive or submassive coral species, together with abundant cyanophytes and soft macro-algae, few sea urchins and abundant holothurids. In 1987, ACR assemblages were observed in reef-flat sites subjected to ingoing oceanic water flow (*inflow sites*), such as Site-Toboggan (Site-T) and Site-Petit Trou d'eau, while the MAS assemblages characterized subtidal areas where the reef flat waters come back to the ocean, after having passed over the reef



flat (*outflow sites*). These MAS assemblages were only observed at Site-Planch'alizés (Site-P) and Site-Club Med; (3 & 4) Mioche (1998), Mioche & Cuet (2002) and Mioche *et al.* (2002), in two studies conducted in 1996 and 2002 on community metabolism, demonstrated that communities at Site-Toboggan and Site-Petit Trou d'eau (inflow sites, ACR) have an autotrophic functioning, thus indicating a "healthy" state for the reef ecosystem.



**Figure 1:** The *Saint-Gilles La Saline* reef. The two studied sites. T180 and P270 are the Global Coral Reef Monitoring Network (GCRMN) stations.

By contrast, communities at Site-Planch'Alizés and Site-Club Med (outflow sites, MAS) are significantly altered to heterotrophy, indicating depletion of oxygen and eventual anaerobic conditions at the end of the night. Further investigations on reef hydrology (Cordier, 2007) and on sediment metabolism at Site-P (Clavier *et al.*, 2008) were in accordance with these previous results and observations.

In 1987, according to the high amount of macroalgae (Naim, 1993) and the lack of sewage treatment along the west coast, MAS at Site-P was regarded as resulting from the nutrient-enriched submarine groundwater discharge onto the reef flat (Cuet, 1989a; 1989b; Naim, 1989).

#### Benthic community assessment

Three kinds of surveys have been performed at the two Sites-T and P (**Table 1**).

### **Survey 1: abundance of major benthic associations (1993, 1996, 2002).**

Estimates of percent coverage and species composition of major benthic categories (including live and dead corals, primary producers, sand and rubble at each site) were made at a 1 centimeter resolution, using the Line Intercept Transect (LIT) method (Lucas & Seber, 1977; Loya, 1978). LITs, 50 m in length, parallel to the front of the reef, were run at fixed intervals of 30 m at each site from 30 m to 330 m off the beach in 1993, 1996, and 2002. At 360 m, analyses became too hazardous owing to strong waves and a reduction of the depth to 0.30 m or less. A ten meter-LIT is considered as a sample for mean calculation. The study was carried out during the hot season when algae are most fully developed (Semple, 1997). The terms “primary producers” refer to sea-grass, cyanophytes and benthic algae and the terms “bare dead corals” refer to dead substrates that are not covered by any living organism visible to the naked eye.

### **Survey 2: densities of echinids, holothurids and the territorial fish**

***Stegastes* (Pomacentridae) (1993, 1996, 2002).** On Saint-Gilles la Saline, Cuet *et al.* (1988), Naim *et al.* (1997), and Naim (2006) showed that spatial distribution of sedentary fauna (echinids, holothurids and the territorial damselfish *Stegastes*) was correlated with the physico-chemistry of reef waters and reef degradation. Thus, their occurrence in a benthic assemblage may give also an indication of the health state of the reef.

Echinids. On the Saint-Gilles la Saline reef flat, sea urchins occur mostly on the coral zone. *Echinometra mathaei* is the dominant species, followed distantly by *Tripneustes gratilla*, *Diadema setosum*, *Toxopneustes pileolus* and *Stomopneustes variolaris*. *Heterocentrotus spp* and *Cidarids* were abundant in the 1980's on the outer reef flat but are now very uncommon.

Holothurids. On the Saint-Gilles la Saline reef flat, sea cucumbers occur mostly in the back reef area. *Holothuria leucospilota* is the dominant species, followed by *H. atra*, *Synapta maculata*, *Stichopus chloronotus*, *Actinopyga echinites*, *A. mauritiana*, *Stichopus chloronotus*, *H. impatiens* (hidden under blocks). *H. nobilis* and *Bohadschia vitensis* also occur but are uncommon.

The territorial damselfish, *Stegastes*. On the Saint-Gilles la Saline reef flat, *Stegastes* is a common damselfish in the inner coral zone. *S. nigricans* is the dominant species, followed by *S. lividus* and *S. limbatus*. Two other species, *S. fasciolatus* and *S. pelicierii*, occur rarely.

Densities, at the specific level, were censused in belt-transects, 1 m strip on either side of each 10 m LIT in the back reef zone, or in 5 \* 1 m<sup>2</sup>-quadrats for 10 m-LIT in the coral zone. All animals, visible from a vertical position as well as those hiding underneath ledges and in crevices, were counted. Animals cut by the upper and left boundaries of the belt-transects were not counted. Data from surveys 1 and 2 form a matrix of 127 variables (including fixed organisms, in cm per 10m-LIT and echinoderms and fishes, in number of individuals per 10 square meter), each species identified if possible, measured on 55 \*(10m-LITs) per year at each site, *i.e.* a matrix made of 127 lines \* 330 columns in total.

**Survey 3: abundance of primary producers, living corals, abiotic coverage and *Stegastes* along two permanent transects T180m and P270m (1987, 1993, 1996, and each year from 1998 to 2009).** In 1987, two 50 m-permanent transects,

incremented every 5 meters, were established on Sites-T and P, at a distance of 180 m and of 270 m seawards from the beach. These T180 and P270 transects were monitored in survey 1 (1993, 1996, 2002). In March 1996, 24 h-variation of sea urchin density was also measured along these two permanent 50 m-LITs.

Table 1: Summary of the 3 studies. "LIT"= Line Intercept Transect. "T180" means: a permanent transect, marked off every 5 meters at Site-T, distance from the shore line : 180 m).

	<b>Aim of the study</b>	<b>Stations</b>	<b>Method</b>	<b>Period</b>
<b>Survey 1</b>	Estimate of coverage and species composition of major fixed benthic categories	At each site, T and P, LITs are parallel to the front, run at fixed intervals of 30m, from 30m to 330m off the beach	50 m-LIT separated in 10 m, 1 cm resolution	1993, 1996, 2002 hot season
<b>Survey 2</b>	Estimate of densities of sea-urchins, holothurids, and sedentary fishes	At each site, T and P, 5 belt-transects are located 1m apart each 50 m-LIT	Belt-transects (10m * 2 m)	1993, 1996, 2002 hot season
<b>Survey 3</b>	Estimate of coverage and species composition of major fixed benthic categories	T 180 and P 270	Permanent transects (2 * 20 m)	1987, 1993, 1996, 2002, and GCRMN each year from 1998 to 2009, hot season
	24 h-variation in sea urchin density	T180 and P270	30 quadrats of 1m <sup>2</sup> thrown along a 50m-LIT	1996, hot season
	<i>Stegastes</i> density	T180 and P270	Belt-transects (50m * 2.0m)*1 (50m * 2.5m)*3	1987, 1993, 1996 GCRMN from 1998 to 2002, hot season

Since 1998, the Global Coral Reef Monitoring Network (GCRMN) (Wilkinson, ed., 2008) started in Reunion (Chabanet *et al.*, 2002), and 3 \* 20m-LITs were monitored each year on each site. Two of these three 20m-LITs are located on the permanent transects T180 and P270, monitored from 1987. Then, temporal variation in abundance of primary producers, living coral and abiotic coverage was analyzed from 1987 to 2009 along the two same 20m-permanent transects. *Stegastes nigricans* density was also analyzed (3 belt transects, length: 50m, width: 2.5 m).

#### Data analysis

Due to the multi-species nature of the data and the design of the survey, the most appropriate analyses are multivariate (Clarke & Warwick 2001). To assess patterns in benthic data from all stations during all years of survey, we used non-metric multidimensional scaling (MDS). To assess temporal patterns in benthic variation, we used Kruskal-Wallis one-way analysis of variance by ranks: this non-parametric method tests equality of population medians among groups. It is identical to a one-way analysis of variance with the data replaced by their ranks and is an extension of the Mann-Whitney U test to 3 or more groups. Since it is a non-parametric method, the Kruskal-Wallis test does not assume a normal population, but it does assume an identically-shaped and scaled distribution for each group, except for any difference in medians.

Mean coral coverage and densities are given with standard deviation, SD (mean  $\pm$  SD). These are calculated as per the following example (shown only for mean calculation here): on Site-T, for a given parameter (e.g. the primary producer coverage), the mean in a subzone (e.g. the back reef, B), for a given-year (e.g. 1993) is denoted TB93, and is the average of the 20 "10m-samples" taken on the 50m-LITs located at 30m, 60m, 90m and 120m. Similarly, the TL93 mean is the average over the 25 "10m-samples" taken on the 150 to 270m LITs, and the TN93 is the average

over the 10 “10m samples” taken on the 300 and 330m LITs. The TB mean is the average over the 60 “10m-samples” taken in the back reef – 20 in each of 1993, 1996 and 2002; the T mean is the average of the 165 “10m-samples” taken – 55 in each of 1993, 1996 and 2002.


### Nomenclature

Concerning the nomenclature used in the text, the « ACR » and « MAS » terms refer to *Acropora*-dominated and massive corals-dominated community structures. The morphological zones are noted as follows, from the beach seawards: the back reef zone  $T_B$  and  $P_B$ , respectively of Sites T and P; the inner coral zone made of « large coral strips »:  $T_L$  and  $P_L$ ; the outer coral zone made of « narrow coral strips »:  $T_N$  and  $P_N$ ; the inner reef flat including the two coral strip subzones:  $T_{L+N}$  and  $P_{L+N}$ . The living coral coverage can be noted as LCcv and the primary producer coverage as PPcv.

## RESULTS

Survey 1: total reef flat, 50m LITs, 1993, 1996, 2002

**Are ACR and MAS communities stable over time?** Based on the mean abundance of species in the communities censused along each 50m-LIT across the reef flat, the MDS plot clearly shows that the data from Site-T and Site-P form two separate clusters (Fig. 2). Apparently, the community structure of ACR at Site-T and MAS at Site-P remained different during the nine years covered by this survey. A stress of 0.15 corresponds to a potentially useful ordination.




**Figure 2:** Two dimensional-MDS configuration (T: triangles, P: circles; Back reef in green, L-zone in orange and N-zone in blue), obtained with a hierarchical clustering of the 66\*(50m-LIT) (all benthic data: corals, algae, etc., echinoderms and territorial fishes included) using group-average linking of Bray-Curtis. "LIT 3P330" means: year 1993, Site-P, 330 m off the beach; similarly, "LIT 6T60" means: year 1996, Site-T, 60 m off the beach, 2P50: year 2002 etc.

Bubble plots, overlaying percent cover of benthic variables without the detail of species, further highlight these trends (**Fig. 3**). Bubble plots of living *Acropora* and sea-urchins, characterizing ACR, overlay Site-T LITs (reef flat and back reef LITs), while bubble plots of primary producers, non-*Acropora* corals, holothurids, characterizing MAS, overlay Site-P LITs (only back reef LITs for holothurids). These results show that from 1993 to 2002, ACR and MAS have kept the characteristics described previously in Naim (2006). Thus, in the following results, we will consider separately the two sites and their spatio-temporal evolution.

**Geomorphology of the reef flat.** A cross-check with a hierarchical clustering was performed based on the same data (not shown). Cluster analysis separates each site into 3 distinct groups (depth value refers to fairweather conditions): **(1)** LITs 30, 60, 90, 120 m group together in Site-T and Site-P clouds and highlight the *Back reef (B-zone)*, where depth reaches 1.50 m maximum and coral coverage is low; **(2)** LITs 150, 180, 210, 240, 270 m, grouping in separate T and P clouds, highlight the deeper zone (0.8 m) of the coral zone, where depth allows corals to organize in *Large strips* of a meter-wide or more (**L-zone**), often anastomosing shoreward where the swell decreases. **(3)** LITs 300 and 330m grouping together, in T and P clouds, are located in the shallow (0.2-0.4 m deep) coral zone where corals can only form *Narrow* (inframetric) *strips (N-zone)*. L+N zones form the “inner reef flat” or the “coral zone”.

**Primary producers (Cyanophytes and benthic algae).** The benthic algae form a very complex community (see part 2, Naim *et al.*, 2013a for further details). Algal coverage at Site-P is about 10 times higher than at Site-T (T:  $5.3 \pm 11.4\%$ ; P:  $61.3 \pm 26.6\%$ ) (**Fig. 3a**). Marine phanerogams are rare and not recorded in the survey.



**Figure 3a to 3f:** Bubble plots, overlaying LITs from figure 2 (benthic variables without the detail of species).

At Site-T, the primary producers coverage -PPcv-, which was very low in 1993, increased significantly to 2002 (**Fig. 4a, Table 2**). However, the values remained low (mean < 10%). The increase was mostly obvious at TN outer subzone where the primary producers, barely visible in 1993 ( $0.9 \pm 2.0\%$ ), increased to  $9.1 \pm 10.2\%$  in 2002 (**Fig. 4b**). At Site-P, the highest coverage was recorded in 1993 ( $81.6 \pm 19.3\%$ ). Between 1993 and 1996, and between 1996 and 2002, the primary producers declined significantly (**Table 2**), but still covered  $49.0 \pm 18.2\%$  of the substrate in 2002. PB and PL-subzones showed the same trend as the entire site. At PN subzone, although the PPcv also significantly declined between 1993 and 1996, it stabilized at  $63.7 \pm 9.0\%$  in 2002.

**Living corals, *Acropora* and non-*Acropora*.** Between 1993 and 2002, although the living coral coverage -LCcv- was similar at both sites (T:  $17.4 \pm 20.7\%$ ;

P:  $16.2 \pm 16.9\%$ ), the genus *Acropora* (*Acr*) dominated the T-coral community ( $11.9 \pm 17.4\%$ ) and the "non-*Acropora*" corals (*NAcr*) dominated to an even greater extent at Site-P ( $14.2 \pm 15.8\%$ ).

Species richness and diversity of corals is treated in part 3 of the paper (Naim *et al.*, 2013b). At Site-T, 11 *Acropora* species formed 68.4% of the coral coverage, while at Site-P, only two species of *Acropora* accounted for 12.6% of cover.

*Acropora muricata* was the dominant species at both sites (T: 76.4% of *Acropora* corals; P: 99.2%).


Among non-*Acropora* (*NAcr*) species, both *Montipora circumvallata* and *Porites (Synaraea) rus* dominated (T: 60.8% of *NAcr*; P: 89.0% of *NAcr*).

Considering spatio-temporal variation, the living coral coverage was relatively stable at Site-P while increasing at Site-T (**Table 2**). The Site-T increase was due to an intense growth of *Acropora* from  $1.9 \pm 3.3\%$  in 1993 to  $25.8 \pm 22.4\%$  in 2002 (**Fig. 4c**) while non-*Acropora* coverage remained stable through time at this site (**Fig. 4e**). In the T-coral zone (TL + TN), the LCcv increased from 1993 to 2002 (**Fig. 4d**) while remaining relatively stable in the P-coral zone (PL + PN), except for a slight increase from 1996 to 2002 in PL (**Fig. 4f**). *Acropora* growth was particularly spectacular in the outer TN zone, where the coverage, reduced to  $1.2 \pm 1.6\%$  in 1993, recovered to  $46.2 \pm 14.4\%$  in 2002. In the back reef areas, due to the rarity of coral patches and the associated high variance, the LCcv did not vary significantly.

At Site-P, *Acropora* and non-*Acropora* coverages did not vary significantly over time. At both sites, non-branching corals showed a remarkable stability over time.


**Bare dead corals.** The bare dead corals were locally abundant at Site-T but were exceedingly rare at Site-P (coverage: T:  $13.5 \pm 16.9\%$ ; P:  $1.2 \pm 3.1\%$ ).

Considering spatio-temporal variation, their coverage remained stable through time, except in TN, where it dropped from  $36.1 \pm 8.3\%$  in 1993 to  $12.4 \pm 8.8\%$  in 2002 (**Fig. 4g, 4h, Table 2**). In the same subzone, TN, considering the {living corals/(bare substrates + living corals)} ratio, it increased continually, from 20% in 1993, to 44% in 1996, and to 83% in 2002, meaning that young coral colonies settled preferentially on bare substrates.



**Figure 4a to 4f:** Spatio-temporal variation from 1993 to 2002 of the coverage of primary producers, living corals, *Acropora* and non-*Acropora*. Left : over the total site, right: in the three geomorphological zones, B= back reef, L= inner zone of Large coral strips, N= outer zone of Narrow coral strips. Error bars are standard deviation.





**Figure 4g, 4h:** Spatio-temporal variation from 1993 to 2002 in the coverage of bare substrate, the density of sea urchins and holothurids. Left : over the total site, right: in the three morphological zones, B= back reef, L= inner zone of Large coral strips, N= outer zone of Narrow coral strips.


#### Survey 2: sedentary fauna

**Sea urchins.** At Site-T, from 1993 to 2002, sea urchin densities were 100 times higher than at Site-P (T:  $70.4 \pm 69.6$  number of individuals (ind.) per  $10 \text{ m}^2$ ; P:  $0.6 \pm 1.2$ ). *Echinometra mathaei* is the dominant species, averaging 99.7% of the echinoid community at Site-T, and 77.7% at Site-P.

Looking at spatio-temporal variation, the densities varied at Site-T and remained stable at Site-P (mean always  $<1$  ind. per  $10 \text{ m}^2$ ) (**Table 2**). At Site-T, the lowest density was observed in 1993 ( $42.9 \pm 46.8$  ind. per  $10 \text{ m}^2$ ). Then the density increased in 1996, to reach in 2002  $89.9 \pm 86.7$  ind. per  $10 \text{ m}^2$  (**Fig. 4i, 4j**). Sea urchin density increased on TL and TN, but in the back reef zone, TB, density and distribution were highly variable through time as follows: very low density and regular distribution in 1993 ( $3.5 \pm 4.1$  ind. per  $10 \text{ m}^2$ ), high density and random distribution in 1996 ( $27.9 \pm 10.0$  ind. per  $10 \text{ m}^2$ ), and again low density but aggregated distribution in 2002 ( $7.6 \pm 19.4$  ind. per  $10 \text{ m}^2$ ).

**Holothurids.** Unlike sea urchins, sea cucumbers were abundant at Site-P, but rare at Site-T (T:  $1.8 \pm 2.6$  ind. per  $10 \text{ m}^2$ ; P:  $12.5 \pm 19.2$ ). Considering the subzones, no significant differences appeared at Site-T, while at Site-P, holothurids were mostly located in the back reef (PB:  $27.8 \pm 21.2$  ind. per  $10 \text{ m}^2$ ).

Considering spatio-temporal variation (**Fig. 4.k, 4.l, Table 2**), holothurid density did not vary significantly from site to site. However, in TL and TN, their numbers increased significantly in 1993-96 and 1993-2002 periods. But, due to low densities in L and N-zones ( $\leq 5$  ind. per  $10 \text{ m}^2$ ), these results prove not to be highly significant. In contrast, in PL and PN, the density decreased between 1993 and 2002. In PB, density declined just as much (on average) but the variance was too great for this decrease to be significant.



**Figure 4j to 4n:** Spatio-temporal variation in the coverage of bare substrate, the density of sea urchins and holothurids from 1993 to 2002. Left : over the total site, right: in the three geomorphological zones, B= back reef, L= inner zone of Large coral strips, N= outer zone of Narrow coral strips.

Table 3: Surveys 1 & 2 (whole sites, 1993, 1996, 2002): Kruskal-Wallis test between means ( $\mu$ ) of the variables in the 3 different years, at the two sites (T= “Toboggan”, P= “Planch’Alizés”). B= back reef, L= zone of large coral strips, N= zone of narrow coral strips). (>: significant increase, <: significant decrease, Ns: non significant).

		$\mu$ 1993- $\mu$ 1996	$\mu$ 1993- $\mu$ 2002	$\mu$ 1996- $\mu$ 2002		$\mu$ 1993- $\mu$ 1996	$\mu$ 1993- $\mu$ 2002	$\mu$ 1996- $\mu$ 2002
Primary Producers	T	Ns	Ns	>>>>	P	<<<<	<<<<	Ns
				p=0.0000		p=0.0000	p=0.0000	
	TB	Ns	Ns	Ns	PB	<<	<<<	Ns
						p= 0.0074	p= 0.0007	
	TL	Ns	Ns	Ns	PL	<	<<<	Ns
						p= 0.0200	p= 0.0009	
	TN	Ns	>>	>	PN	<	Ns	Ns
			p= 0.0060	p= 0.0201		p= 0.0247		
Living corals	T	Ns	>>>	Ns	P	Ns	Ns	Ns
			p= 0.0001					
	TB	Ns	Ns	Ns	PB	Ns	Ns	Ns
	TL	Ns	>>>>	Ns	PL	Ns	Ns	>
			p=0.0000					p= 0.023
	TN	Ns	>>>>	>>>>	PN	Ns	Ns	Ns
			p=0.0000	p=0.0000				
<i>Acropora</i>	T	>	>>>>	Ns	P	Ns	Ns	Ns
		p=0.0233	p=0.0000					
	TB	Ns	Ns	Ns	PB	Ns	Ns	Ns
	TL	Ns	>>>>	>	PL	Ns	Ns	Ns
			p=0.0000	p=0.0205				
	TN	Ns	>>>>	>>>	PN	Ns	Ns	Ns
			p=0.0000	p= 0.0001				
Non- <i>Acropora</i>	T	Ns	Ns	Ns	P	Ns	Ns	Ns
	TB	Ns	Ns	Ns	PB	Ns	Ns	Ns
	TL	Ns	Ns	Ns	PL	Ns	Ns	Ns
	TN	Ns	Ns	Ns	PN	Ns	Ns	Ns
Bare dead corals	T	Ns	Ns	Ns	P	Ns	Ns	Ns
	TB	Ns	Ns	Ns	PB	Ns	Ns	Ns
	TL	Ns	Ns	Ns	PL	Ns	Ns	Ns
	TN	Ns	<	Ns	PN	Ns	Ns	Ns
			p= 0.0103					
Sea urchins	T	>>>>	>>	Ns	P	Ns	Ns	Ns
		p=0.0000	p= 0.0092					
	TB	>>>>	Ns	<<<<	PB	Ns	Ns	Ns
		p=0.0000		p=0.0000				
	TL	Ns	>>	Ns	PL	Ns	Ns	Ns
			p= 0.0073					
	TN	Ns	>	Ns	PN	Ns	Ns	Ns
			p= 0.0440					
Holothurids	T	Ns	Ns	Ns	P	Ns	Ns	Ns
	TB	Ns	Ns	Ns	PB	Ns	Ns	Ns
	TL	>	>	Ns	PL	Ns	<	Ns
		p= 0.0408	p= 0.0280				p= 0.042	
	TN	>	>	Ns	PN	Ns	<	Ns
		p= 0.0384	p= 0.0343				p=0.0221	
<i>Stegastes</i> (all species)	T	>	>>>	Ns	P	Ns	Ns	Ns
		p=0.0162	p= 0.0002					
	TB	absent	absent	absent	PB	Ns	Ns	Ns
	TL	Ns	>>>>	Ns	PL	Ns	>>>	>>>
			p=0.0000				p= 0.0004	p= 0.0009
	TN	Ns	>>	Ns	PN	Ns	>>	>
			p= 0.0026				p= 0.0075	p=0.0213

***Stegastes*.** *Stegastes* is a territorial damselfish, that colonizes dead corals. Thus, this Pomacentridae is rare or absent from back reef areas. The density of *Stegastes* was not significantly different between Sites T and P (T:  $6.2 \pm 14.5$  ind. per

10 m<sup>2</sup>; P: 3.8 ± 8.1), but its spatio-temporal distribution was more variable at Site-T (**Table 2**). In 1993, *Stegastes* density was very low at both sites (<2 ind. per 10 m<sup>2</sup>) (**Fig. 4m**). At Site-T, a significant increase in density was recorded between 1993 and 2002 to 14.8 ± 22.1 ind. per 10 m<sup>2</sup>. This increase mainly concerned the TL area where the density increased from 1.0 ± 2.8 to 32.8 ± 30.0 ind. per 10 m<sup>2</sup> (**Fig. 4n**). In contrast, at Site-P, between 1993 and 2002, the density also significantly increased in PL and PN, but to a lesser extent.

Survey 3: the permanent transects T180m, P270m from 1987 to 2009


Do ACR and MAS structures remain stable until 2009? The primary producer coverage -PPcv- was lower on average and more variable at T180 than at P270 (T180: 23.9 ± 15.1%; P270: 40.3 ± 13.5%) whereas the living coral coverage -LCcv- was relatively stable at the two sites (T180: 36.8 ± 13.3%; P270: 42.0 ± 9.8%). Among corals, the genus *Acropora* remained dominant at T180 while scarce at P270 (relative %: T180: 73.5 ± 8.4% of LCcv, P270: 3.6% ± 5.5 of LCcv).

Considering temporal variations: (1) at **T180**, from 1987 to 1998, the primary producer coverage remained below 10%. From 1999, it clearly increased, stabilizing in the range of 20-50% (**Fig. 5a**). For living corals, the total living coral coverage (as well as the *Acropora* and non-*Acropora* coverage), did not vary significantly (**Fig. 5a, 5c, 5d**). The *Acropora* coverage went through cycles of demise (1987-1993, 1999-00, 2003-06) followed by a rapid recovery (1993-98, 2000-03, 2006-09) while the non-*Acropora* coral coverage remained remarkably stable; (2) at **P270**, from 1987 to 1998, the primary producer coverage was high (range of 40-60%). In 1999, it clearly decreased, stabilizing then in the range of 20-40% until 2008 (**Fig. 5b**). In 2009, the algal coverage increased again to 60%. The living coral coverage was not significantly different in the 1987-98 and 1998-2003 periods (28-40%). Later, it increased to 40-60%, but markedly decreased in 2009 to 26.5%, with the lowest LCcv ever measured (**Fig. 5b**). Massive corals remained dominant over time and *Acropora* represented less than 10% of the benthic coverage (**Fig. 5c, 5d**).

In conclusion, (1) at **T180** (**Fig. 5a**), from 1987 to 1999, the living corals dominated, while primary producers remained inconspicuous, the 1987 coverage (70%) being no longer achieved. Then, from 1999 to 2009, living corals and primary producers dominated alternately: the 2001-2003 period was favorable to coral development, the 2004-2006 interval was favorable to primary producers, while from 2008, the conditions were again favorable to corals. This suggests that a critical shift occurred in 1999 in the functioning of the community structure, and a lesser shift happened in 2004;

(2) at **P270** (**Fig. 5b**), in contrast to T180, from 1987 to 1998, the algal coverage was higher than living coral coverage, the year 1987 being the worst period. In 1999, living corals became dominant over primary producers, and from 2004, the situation improved further. In 2009, there was a sharp decline in living corals and related increase in primary producers, thus once more inverting the dominance ratio. In terms of living coral coverage, 2009 appears to be the worse year of the 2000s.

Considering living coral and primary producer relative dominance, two 'turning points' occurred at both transects, one in 1999 in one direction and the other in 2004 in the opposite direction. Therefore, we divided the monitoring period into 3 blocks (1987-98, 1999-2004 and 2004-2009) and compared the mean coverages of the three periods.




**Figure 5:** Yearly variation of primary producers, living corals, *Acropora* and non-*Acropora* at the two permanent transects T180 (Site-T) and P270 (Site-P).

**Comparison of 1987-98, 1999-2003, and 2004-2009 periods.** The primary producer coverage increased significantly at T180 between 1987-98 and 2004-09, and decreased significantly at P270 between 1987-98 and 1998-2003 (Fig. 6a, Table 3). The living coral coverage, as well as *Acropora* coverage, did not vary significantly at T180 from 1987-98 to 2004-09 (Fig. 6b, 6c, 6d, Table 3). By contrast, at P270, the massive coral coverage increased significantly between 1987-98 and 2004-09 (Fig. 6d, Table 3). At T180, *Stegastes* density significantly increased while it remained stable at P270 (Fig. 6e, Table 3).

Regarding PPcv and LCcv temporal trends, T180 and P270 seem to vary in opposite directions: (1) from 1987 to 1998, at T180, living corals dominate primary producers while at P270 living corals are heavily overgrown by algae (see part 2, Naim *et al.*, 2013a), then (2) since 1999, at T180, a period less favorable for coral growth began (alternate dominance of PPcv and LCcv), while at P270, a period more favourable to coral growth started. Then, in 2009, P270 provided the worst PPcv/LCcv ratio in the whole monitoring period, indicating a sharp decline in the coral health on the transect, while at T180, 2009 still was a prosperous period for

coral life. The *Acropora* coverage did not vary significantly (high variance) while the non-*Acropora* increased significantly at P270 between 1987-98 and 2004-2009.



**Figure 6:** Temporal variation of primary producer, living *Acropora* and non-*Acropora* coverages between 1987-98, 1999-2003 and 2004-2009 "year-blocks".

Table 3: Survey 3: Kruskal-Wallis test between means ( $\mu$ ) of the variables in the 3 "year-blocks", at the two permanent transects (T180= "Toboggan" 180m off shore, P= "Planch'Alizés", 270 m off shore). *N.b.* For *Stegastes*, 2001 densities are not included. (>: significant increase, <: significant decrease, Ns: non significant).

		$\mu$ (1987-1998)	$\mu$ (1999-2003)	$\mu$ (1987-1998)	$\mu$ (1987-1998)	$\mu$ (1999-2003)	$\mu$ (1987-1998)
		$\mu$ (1999-2003)	$\mu$ (2004-09)	$\mu$ (2004-09)	$\mu$ (1999-2004)	$\mu$ (2004-09)	$\mu$ (2004-09)
Primary Producers	T180	Ns	Ns	>>	P270	<	Ns
				p= 0.0019		p= 0.0373	Ns
Living corals	T180	Ns	Ns	Ns	P270	Ns	>
							p=0.0391
<i>Acropora</i>	T180	Ns	Ns	Ns	P270	Ns	Ns
Non- <i>Acropora</i>	T180	Ns	Ns	Ns	P270	Ns	>
							p=0.0361
<i>Stegastes</i>	T180	>	Ns	Ns	P270	Ns	Ns
		p=0.0298					

Concerning sea urchins, at T180 and P270, their density was measured only in 1987, 1993, 1996 and 2002 (survey 1) and 2011. In 1987, at T180, the density was the highest ever measured:  $180.2 \pm 52.1$  ind. per  $10\text{ m}^2$ , while at P270, no sea urchin was recorded. In 2011, the sea urchin density is still  $27.1 \pm 3.6$  ind. per  $10\text{ m}^2$  on T180 and  $0.5 \pm 0.6$  ind. per  $10\text{ m}^2$  on P270.

The measurement of the nycthemeral variation in the density of sea urchins showed that at T180, the density at midnight is three times higher than at noon, while at P270, no sea urchin was observed (**Table 4**).

Table 4: Nycthemeral variation in the density of sea urchins on T180 and P270, in March 1996 (in number of individuals per 10 m<sup>2</sup> ± standard deviation).

Sea urchin density	6 am	12 am	6 pm	12 pm
T180	30.1 ± 10.9	20.5 ± 22.2	30.3 ± 32.5	62.9 ± 32.9
P270	0.0 ± 0.0	0.0 ± 0.0	0.0 ± 0.0	0.0 ± 0.0

## DISCUSSION

Spatio-temporal variation of benthic communities: ACR and MAS are stable over 25 years

To our knowledge (the quantitative studies started in 1985), the Site-T and Site-P communities, labelled ACR and MAS, have preserved their principal characteristics until now, although the MAS components remain much more stable than the ACR ones.

**Primary producers.** From 1993 to 2002, at Site-T, the primary producer coverage remained stable and inconspicuous, except in the TN-subzone where it increased significantly. However, the average coverage never exceeded 10%. At Site-P, in 1993, algae were dominating the substratum (60-100% cov.) (see details in part 2). But from 1993 to 2002, their coverage steadily decreased and stabilized around 50% in 2002, except in the outer PN-subzone where the coverage remained stable. Outer N-subzones had the highest algal coverage at both sites, revealing then a greater fragility in relation to environmental factors, due certainly in part to their shallowness (0.4 m maximum depth).

From 1987 to 2009, a significant change in the amount of primary producers occurred at the two sites: at Site-T, the algal coverage increased while at Site-P it decreased. These results will be further developed and discussed in Part 2 (Naim *et al.*, 2013a).

**Living corals.** Between 1993 and 2002, the mean living coral coverage was relatively low, and not significantly different between sites (T: 17 ± 21%; P: 16 ± 17%). This might reflect a carrying capacity in living corals of the *Saint-Gilles La Saline reef* flat in extreme conditions (pure autotrophic, pure heterotrophic).

In the 1993-2002 period, the coral coverage was stable at Site-P and very unstable at Site-T. Instability observed at Site-T is linked to the dominance and fluctuations in the *Acropora* coverage. Conversely, the massive coral cover showed a remarkable 10 year-stability at both sites. The same trends were observed in the 1998-2009 GCRMN data, although massive coral coverage increased significantly at Site-P in the 2004-2008 period.

Before the creation of the marine reserve (2007), the T-coral zone (TL + TN) was a hotspot for visitors, either by snorkelers during the day, and/or by poachers at night. The great coral vitality and the fish richness (Letourneur, 1992; Chabanet, 1994; Chabanet *et al.*, 1997; Tessier *et al.*, 2008) probably explain the high touristic pressure it has suffered. In 2007, the area became a sanctuary, and such a new status may modify the coral as well as the fish carrying capacity in the future.

*Acropora, stenoecious corals, ruderals (r-strategists), sensitive but rapidly growing, structuring the ACR community.* Outer shallow coral zones (N-zones) develop on the large Reunion reef flats (La Saline, Saint-Leu, Saint-Pierre). At these particular zones, *Acropora* can form what Montaggioni and Faure (1980) and Faure (1982) called "fields of branching *Acropora*", that approach 100% coverage and in which rare sandy channels do not exceed some decimeters in width. These dense bushes appear to be the most vulnerable to environmental factors that chronically affect the reef flats (extreme low tides, bleaching events, etc.), and mortality can reach almost 100% of the coverage in one episode (Naim, pers. obs.). In this study, at TN, *Acropora* bushes were least in 1993 (1%), and recovered to more than 40% over 9 years, in spite of two small *Acropora* "declines" in 1998 and 1999.

In the inner L-coral zones, which are deeper than N-zones, *Acropora* form large colonies or coral strips that often merge shoreward. These coral formations are generally less affected by bleaching events and extreme low tides, as deeper parts of the colonies/strips remain alive. Nevertheless, at TL, between 1993 and 2002, *Acropora* coverage displayed, as in TN, a rapid recovery (5% to 30%). The GCRMN data showed that recovery can be quite rapid, increasing by up to 5-10% cover per year from 2000 to 2003, but the *Acropora* community did not reach 64% of coverage as reached in 1987 on the same permanent transect. Either this is due to lack of recovery time before a next disturbance affects the *Acropora* population, or this indicates an ecological change on the reef flat at Site-T.

The sensitivity of *Acropora* to environmental factors, as well as its ability to quickly recolonize the space, is demonstrated here. Similarly, on the Saint-Leu ville reef flat (ACR community, west Reunion coast), the *Acropora* assemblages recovered through 10 years following the impact of cyclone Firinga (Naim *et al.*, 2002; Scopélitis *et al.*, 2009). The rapid recovery of *Acropora* populations indicates that the Site-T and Site-Saint-Leu ville environments remained favorable enough for *Acropora* resettlement and survival. Similarly, in the Caribbean, after a serious degradation, reefs of the Florida Keys recovered rapidly (Shinn, 1976), but by contrast the Carysfort reef did not (Dustan & Halas, 1987).

Many studies over the past 30 years have been interested in the coral composition and distribution on the largest reef units at Reunion, including Saint-Gilles la Saline, Saint-Leu and Saint-Pierre (Faure & Montaggioni, 1970; Montaggioni, 1978; Faure, 1982; Naim, 1989; Naim *et al.*, 2002; Bruggemann *et al.*, 2007; Tessier *et al.*, 2008) and particularly on the *Saint-Gilles La Saline reef* (Bouchon, 1981, 1996; Guillaume, 1988; Naim, 2002, 2006). All show that *Acropora* is usually dominant on Reunion reef flats. Classified as ruderal, or r-strategist by Edinger and Risk (2000), *Acropora* still dominates the healthy parts of reef flats in Reunion, with both high coverage and high diversity. **Figure 5b** shows that except in 1985 and 2009, the T180-living coral coverage is still lower than the P270 one. The *Acropora* growth periods (1993-98, 2000-03, 2006-08) apparently do not compensate for periods of decline (1985-93, 1998-2000, 2003-04), some of which are very fast. Events that may cause *Acropora* degradation will be discussed further.

At Site-P, between 1993 and 2002, *Acropora* coverage did not exceed 2% ( $\pm 5.7$ ). At P270, the *Acropora* coverage was always less than 4% between 1987 and 2009, except in 2008 when it reached 9% of the total coverage.

*Massive and submassive corals, euryoecious, stress-tolerators (s-strategists), structuring the MAS community.* From 1993 to 2002, in the PN shallow outer zone,



the flattened upper part of corals was generally invaded with macroalgae. The outer “ring pad” and the small (0.3 m max) vertical walls of coral colonies were the only parts alive. At PL, in 1993, the flat subtidal top of corals was also invaded by macroalgae. Exposure appears to be an important factor of degradation at the PL and PN subzones, but does not explain the abundance of primary producers.

Massive and submassive corals are considered as stress-tolerators (s-strategists) by Edinger and Risk (2000). In this study, they show a high spatial and temporal stability over time. From 2004 to 2008, their coverage increased at Site-P, certainly due in part, to the reduction of competition for space with primary producers (see part 2, Naim *et al.*, 2013a).

**Bare dead substrates.** Measurements of bare dead coral coverage are interesting because a bare substrate indicates, because of the total absence of primary producers, a high grazing pressure. Bare substrates were rare on the reef, except at TN where the sea-urchin *Echinometra mathaei* lived in nearby sites, either hidden in the cavities or at the base of these substrates. From 1993 to 2002, the significant decrease in bare dead corals at TN can be related to the increase of *Acropora* coverage since *Acropora* juveniles colonized -and extended beyond- these dead substrates over the 9 year-period, but also to the sea-urchin bioerosion (see below). As no bare substrate was found on the two GCRMN transects, temporal evolution was not documented after 2002.

**Sea urchins.** From 1993 to 2002, the density of sea urchins was 100 times higher at Site-T than at Site-P, as confirmed by Conand *et al.* (1998) in the coral-zones, TL and TN. Minimal in 1993 the urchin density increased significantly until 2002. *Echinometra mathaei* formed more than 99.7% of the community at Site-T.

In the T back reef, the density and distribution patterns of sea urchins were highly variable over years, partially due to animal movements but mainly to swell action. Density of sea-urchins was not monitored after 2002, but a 2011 monitoring confirms that the general trends did not change (GCRMN, not published). Chazottes *et al.* (2002) found significant variations in bioerosion among ACR and MAS sites.

Circadian census performed at the TL coral subzone revealed that the sea urchin density was three times higher at night than during the day. This very high nocturnal density shows how the *Echinometra mathaei* pressure on the reef may be underestimated by measures of the day-time densities.

*Echinoids and algae.* Inconspicuousness or dominance of algae at the T and P sites were obviously linked to nutrient concentrations and/or the quantity of nutrient regeneration through organic matter depletion (Cuet *et al.*, 1988; Mioche, 1998). However, regular echinoid grazing is also a major factor determining the dominance of primary producers (for instance, see Hughes *et al.*, 1987; Littler *et al.*, 2009). So, at both sites, the abundance (or rarity) of sea urchins provides a strong (or very weak) feedback on algal biomass (See part 2, Naim *et al.*, 2013a).

*Echinoids and living corals.* In September 1992, *Echinometra mathaei* and *Acropora* were similarly affected by a high rate of mortality and from 1993 to 2002, *Acropora* coverage and *Echinometra mathaei* densities increased significantly in the same way. It is interesting to note that the 1993-2002 increase in echinoid densities has not hampered the settlement of corals, as reported by Hardman *et al.* (2007). It

has to be noted that in 1987, when *Acropora* populations were at their best (64% of the coverage), the sea urchin density was the highest measured on Site-T (180 ind. per 10 m<sup>2</sup>).

Cover & Potts (2008) documented the direct impact of *E. mathaei* on living coral fragments (mostly *Montipora* and *Porites*): most fragments of all coral species placed in plots where *E. mathaei* occurred were grazed down, often resulting in a complete removal of coral tissue and corallites.

By contrast, Sammarco's experiments in Jamaica (1982) showed that percent-cover of adult corals was enhanced in relation to the presence of *Echinometra viridis*, and greatly reduced in the absence of echinoid grazing, as a result of massive overgrowth by algae. In the same way, this author showed that the presence of *Echinometra viridis*, even at the high densities naturally present on the reef, did not deter coral settlement. He showed also that as echinoid species composition and density were altered, the species composition of benthic algae and associated invertebrates shifted.

Therefore, are echinoids coral safe or not? This is a very debatable subject. Sea urchins are very often considered as reef pests, occurring in nutrient-enriched areas (Horrill *et al.*, 2000). From our point of view, it may depend on the species considered. At Site-P, *E. mathaei* is rare in the back reef zone and totally absent from the inner and outer reef flat. More generally, in Reunion as well as in the nearby island of Mauritius, the species seems to avoid freshwater or nutrient-enriched areas (except in strong current areas): the more oceanic are the reef flat conditions, the more abundant is *E. mathaei* (Naim, 2006). In Reunion, this species was considered to be grazing on turfs and microflora and eroding the dead corals (Conand *et al.*, 1998). But *E. mathaei* is an opportunistic species: at night, it has been observed to gnaw on living polyps and their corallites located at the base of *Acropora* branching colonies, while during the day, it has also been observed out of its hole, eating rice and corn from the dirty dishes of the beach users (Prigent & Naim, pers. obs.).

In september 1992, at TN, when *Acropora* colonies were almost all terminated, the last *E. mathaei* were hiding in the remaining coral framework (20-40 cm high). During the 1993-2002 recovery, young *Acropora* colonies settled despite the increase in *E. mathaei* density and did not seem to suffer from sea urchin grazing, suggesting that *E. mathaei* found enough « non coral-food » and/or had not exactly the same micro-niche as *Acropora* juveniles. Such contradictory results between different studies suggest that echinoid impact on living corals may depend on very diverse factors, including sea urchin species, density, food availability, coral forms (massive, branching, corymbose, etc.), accessibility of the corals, spatial competition with other organisms such as *Stegastes* (Naim *et al.*, 1997) and the exact overlapping of the coral and echinoid micro-niche.

*Echinoids and bare substrates.* At TN, *Echinometra mathaei* density increases from 1993 to 2002 (1993: 8.2 ind.m<sup>-2</sup>; 1996: 13.6 ind.m<sup>-2</sup>; 2002: 15.7 ind.m<sup>-2</sup>; **Fig. 4j**). At Reunion (Petit Trou d'eau, south Saline reef, **fig.1**), Conand *et al.* (1997) showed that one *Echinometra mathaei* adult grazes 113 g CaCO<sub>3</sub>.y<sup>-1</sup> (in mean) on the N-zone of the reef flat.

Then, if we assume (1) that *E. mathaei* has a similar behavior at TN than at Petit Trou d'eau, and (2) that at TN, the population density increases *linearly* from 1993 to 1996 (1.8 ind. more per year), and from 1996 to 2002 (0.35 ind. more per year) the total bioerosion of the *E. mathaei* population can be estimated:

$$\int_0^3 0.113(1.8x + 8.2).dx + \int_0^6 0.113(0.35x + 13.6).dx = 13.6 \text{ kg.m}^{-2}$$

This approximative value of  $13 \text{ kg.m}^{-2}$  (that corresponds to the amount of  $\text{CaCO}_3$  grazed by the *E. mathaei* population in 9 years), allows us to better understand how the *E. mathaei* population contributes to the decrease of dead coral coverage in this outer zone of the reef flat.

**Holothurids.** Sea cucumbers, as deposit-feeders, collect food by swallowing large volumes of sediment requiring both microbial and non-living organic matter (Lopez & Levinton, 1987). Numerous holothurid species remain hidden during the day, and their density is also underestimated but less markedly than that of sea urchins. At Site-P, densities of holothurids are far higher than at Site-T, but, from 1993 to 2002, they decreased, in conjunction with the demise in primary producer coverage. Their abundance at Site-P appeared then to be linked to the abundance of primary producers as the bulk of detritus derives mainly from algae (Lewis, 1977). Similarly, Frouin (2000) found a significant difference in ACR and MAS soft-bottom communities.

Conand & De Ridder (1990) and Mangion *et al.* (2004) explained the high density of sea cucumbers in the P-back reef by a high amount of organic matter but also by the reproductive strategy of one of the dominant species, *Holothuria atra*. Conand (1996) reported that 20 percent of the *H. atra* population reproduce asexually in the P-back reef, by transverse fission followed by regeneration of both parts, whilst in the P-coral zone, *H. atra* is much larger and reproduces sexually. In the Western Central Pacific, *H. atra* apparently shows the same two reproductive strategies (Conand, 1998).

**Stegastes.** These opportunistic damselfish settle on dead corals (Letourneur *et al.*, 1993; Naim *et al.*, 1997; Letourneur, 2000), and are rare or absent from back reef areas. Among the *Stegastes* species recorded here, *Stegastes nigricans* was the dominant species, as demonstrated by Letourneur (1992) on Reunion reef flats. Its dominance among the *Stegastes* community can vary in time (Naim & Tourrand, unpublished).

From 1993 to 2002, the *Stegastes* density was not significantly different between Sites T and P. Very low in 1993 at both sites, the density showed a significant increase in 2002, but to a lesser extent at Site-P. Later, from 1998 to 2009, the *Stegastes* density significantly increased at Site-T and decreased at Site-P. Done *et al.* (1991) emphasized the importance of *Stegastes nigricans* and their turf territories in the dynamics of the benthic community structure. In Reunion, further investigations are needed to better understand the ecological role of this key species of the reef flats.

Primary producers and living corals: do the two sites, T and P, behave oppositely?

Despite the lack of a quantitative approach, the earlier observations are of some interest. As far as we know, in the 1970s, coral populations were luxuriant on the Reunion reef flats (Montaggioni, 1978; Faure, 1982; Bouchon, 1996). In 1983, Guillaume *et al.* reported that 80% of corals were dying: *Acropora* corals from the reef flats and *Porites* on the outer slope down to 20 m deep were particularly affected. This coral demise was at that time partly attributed to a decrease in water quality.

Year 1983 is since known to have been typified by strong positive Sea Surface Temperature (SST) anomaly in the southwest Indian Ocean (**Table 6**). Observations on Sites T and P have begun from 1985.

**Site-T.** From 1985 to 1992, the inner TL and the outer TN coral zones were characterized by a high coverage in branching corals (Naim, 1989). Data available indicate a very high coverage with a very low standard deviation: TL:  $70.0 \pm 5.2\%$  in April 1987 (Naim, 2006), TN:  $65.1 \pm 1.6\%$  in January 1992 (Chabanet, 1994).

In September 1992, the *Acropora* assemblages suffered a sharp deterioration due to an important bleaching event, explaining the very low living *Acropora* coverage recorded in 1993. The bleaching occurred mostly on the oceanic side of the reef flat and the associated mortality of *Echinometra mathaei* suggested that corals were not affected by a bleaching event induced by high sea-surface temperature nor by extreme low tides. Hypotheses of a potential contaminant supplied by the ocean waters rather than by terrigenous inputs, or of inimical environmental conditions, were advanced. The sewage treatment plant, established close to the Ermitage ravine at the end of the 1980s, was suspected to release toxins to the ocean and then to the reef. Later, *Acropora* colonies resettled, but without achieving their highest recorded coverage.

From 1985 to 1998, the primary producers remained inconspicuous and living corals dominated. Since 1999, the coverage of primary producers increased and in the 2004-2007 period, it overtook the living coral coverage suggesting that the abiotic conditions became more favorable to algae and/or unfavourable to living *Acropora* (part 2 of the paper). In the same period, the increase in *Stegastes* densities corroborates this assertion.

**Site-P.** At Site-P, in 1985-1988, corals could be overgrown by macroalgae within one month (Naim, 1993). In 1995, algal amounts dropped; although still dominant in coverage, their thickness significantly decreased and coral overgrowth by macroalgae was no longer observed (part 2, Naim *et al.*, 2013a). From 1999, the primary producer coverage began also to decline and massive corals appear to do better. Similarly, in the Caribbean (Gardner *et al.*, 2003) and Pacific Southeast Asia (Bruno & Selig, 2007), the greatest decline in coral coverage occurred in the 1980's, due to multi-stress factors such as diseases, hurricane activity and pollution.

**Sites T and P.** In summary, in the 1980s, corals at Site-T were flourishing while at Site-P, they were overgrown by macroalgae. Then, from 1999, the Site-T health was getting worse, while Site-P health was improving. In 2009, a sharp deterioration at Site-P occurred while a slight health improvement occurred at Site-T.


The two sites seem to behave oppositely. According to Join (1991), at Site-P, the sand aquifer seeps onto the reef flat at low tide and at Site-T, seeps through the reef framework to trickle down on the outer slope. This freshwater affects coral communities (Cuet *et al.*, 1988, Chabanet, 1994; Chabanet *et al.*, 1995).

At the end of the 1990s, nutrient-pollution through groundwater discharge greatly decreased at Site-P, suggesting that the connection of coastal homes to the sewage plant increased (Cuet, 2007). However, the increase in incoming sewage waters is correlated to an increase in discharge of the sewage plant into the Ermitage ravine (36%), and by infiltration in the ravine sub-flow (64%) ([www.tco.re](http://www.tco.re)). The ravine water flows then southwards and northwards to the ocean and then flows back

onto the reef (Biais & Kopp, 1981) and may bring contaminated water to the inflow Site-T.

Moreover, from 1994, a major project was intended to transfer water available in two rivers in northern and eastern shores of the island, which were well watered, to the west coast, which is naturally arid. It included two major projects: Gallery Mafate and transfer of Salazie. Mafate work started in 1989 and ended in 1999 with the commissioning of « antennas 0 and 4 » (Borca & Margaloff, 2004). In a normal year, the contribution will transfer to the West in low water 60,000 m<sup>3</sup> from Mafate and 200,000 m<sup>3</sup> from Salazie. The project will ultimately irrigate 7,300 hectares of sugar cane on the West watershed (including new 5.150 ha) (**Fig.7**); this represents a peak demand in a normal year of 260,000 m<sup>3</sup>.day<sup>-1</sup> (SAGE ouest, état des lieux 2006).

While the sewage treatment resulted in improvement in health at Site-P, the increasing freshwater flux, contaminated or not, to the Ermitage watershed may possibly account for the alteration of *Acropora assemblages* at Site-T.



**Figure 7:** The Saint-Gilles la Saline watershed

Although Tessier *et al.* (2008) showed that increase in sea surface temperature weakly influences the structure of coral populations at Reunion, the 1998 temperature anomaly can be regarded as one of the most prominent factors that are controlling regional change in coral coverage in the Indian Ocean (CORDIO, 1999). Thus, it can be hypothesized that Reunion stands of *Acropora* may have been undermined by this event.

#### Causes of coral degradation (other than nutrient-enrichment)

Climatic causes of degradation, including cyclone impacts, high SST, extreme low tides, together with crown-of-thorns outbreaks, may have also important impacts

on r-strategist *Acropora*. Nutrient-enrichment is treated in detail in part 2 of the paper (Naim *et al.*, 2013a).

**Crown-of-thorns outbreaks.** In Reunion Island, research on *Acanthaster planci* outbreaks has been sporadic. On the outer slope, aggregations of tens of individuals have been observed (Faure, Conand, non published), but on the reef flats, only a few isolated individuals have been observed. Degradation of *Acropora* cannot then be attributed to starfish predation.

**Runoff, cyclonic impacts.** The west coast of the island is generally protected from the impact of hurricanes. However, some cyclones, such as Firinga (1989), Dina (2002) and Gamède (2007) have seriously affected the west and south coasts. In 1989, Saint-Leu and Saint-Pierre coral colonies were destroyed by 99% (Letourneur *et al.*, 1993; Naim *et al.*, 2002) but the *Saint-Gilles La Saline reef* corals remained relatively unaffected.

At the beginning of the 1990s, a runoff outlet was set up at the T-shore, in order to clear out the flooding of the urbanized coastal plain. The runoff has certainly contributed to coral degradation because turbid water plumes occur in the T-coral area after heavy rains. Nevertheless, the coral area is probably relatively protected from runoff by the incoming oceanic waters and by a strong coastal current that flows southwards, to the pass.

For the Site-P, the impact of hurricanes is not really apparent. At both sites, coral breakage due to cyclonic events is negligible, compared to the anthropogenic effect.

**Sedimentation processes.** Given the strong impact of human activities on the coastal beaches, depositional processes have changed qualitatively and quantitatively on both sites. Beach erosion is intense at Toboggan and contributes to fill the back reef area.

**Global change: regional implications.** Considering regional-wide changes in composition of coral assemblages, coral reefs throughout South Asia and the central and the western Indian Ocean seem to be undergoing significant changes in coral community composition. This is a result of differential mortality among adult colonies of different genera. *Acropora*, which was easily the most abundant and diverse genus in the Indian Ocean, is now conspicuously absent or has become rare due to severe bleaching events (Obura *et al.*, eds, 2008; McClanahan *et al.*, 2007).

At *Saint-Gilles La Saline*, a number of bleaching events were recorded (**Table 6**), and *Acropora* was recognized as the most sensitive genus by Conand *et al.* (2002b). Nevertheless, after all these events, although the coverage declined since the 1980s at Site-T, and in 2009 at Site-P, both living *Acropora* at Site-T, and living massive corals at Site-P, show a coverage of 40 to 50% in 2011, which can be considered a high coverage.

ACR coral zones such as “Toboggan” and “Petit Trou d’eau” have been included into a sanctuary area in 2007. This will favour the resilience capacity of *Acropora* communities. In this context, in addition to monitoring the coral populations in the reserve after the initial state (Bruggemann *et al.*, 2007), further monitoring of the T-coral zone is required, especially focusing on benthic communities, coral biodiversity and abiotic conditions, in order to see if, in the

absence of direct human impacts, *Acropora* coverage is able to recover its previous 1980s health state or if it is still delayed by pollution effects and/or bleaching events.

Table 6: Significant bleaching events, extreme low tides recorded in Reunion and/or in the southwest Indian Ocean (due either to climatic conditions or low tides). T180: Site-T, 180m from the beach. General data come from Burke *et al.* (2011).

Year	Events
1983	Guillaume <i>et al.</i> (1983) reported a high coral bleaching and mortality event on Reunion reef flats and along outer slopes. This episode also affected the Mozambique channel (Mayotte Island, Faure <i>et al.</i> , 1983, 1984), Indonesia (PulauPari, Seribu island, Dauget, 1983), and the Pacific coast of Panama (Gulf of Chiriqui) and Galapagos (Glynn, 1983).
1987	In April, 30% of <i>Acropora muricata</i> bleached and died over the whole <i>Saint-Gilles La Saline reef flat</i> (Naim, 2006). In addition, this bleaching event affected Saint-Leu and Saint-Pierre reef flats: other <i>Acropora</i> species including <i>A. digitifera</i> , <i>A. abrotanoides</i> , <i>A. austera-hemprichi</i> , <i>A. cytherea</i> , ... but also <i>Platygyra</i> , <i>Millepora</i> were totally bleached to death.
1992	A large bleaching event was observed at Site-T in September, and around the mouth of the Ermitage ravine. This event may be related to a possible pollution from the Ermitage sewage treatment plant.
<b>1998: the extreme El Niño-La Niña switch</b>	This climatic event resulted in the most extensive coral bleaching and mortality ever recorded with approximately 16% of the world's coral reefs being destroyed (approximately 3/4 of the bleached corals recovered). The warm phase was particularly devastating to the western Indian Ocean (Quod, 1999; Goreau <i>et al.</i> , 2000, Turner <i>et al.</i> , 2002). However, while some locations, such as the Maldives, Comoros, Mayotte and the inner Seychelles suffered 75-99% mortality, in Reunion, in 1998, the coral coverage was the highest for the 1990-2000 period (GCRMN data).
1999	In 1999, Tessier <i>et al.</i> (2008) reported exceptional low tides.
2001	In March 2001, a large bleaching event occurred in the <i>Saint-Gilles La Saline reef</i> . Bleaching affected more than 45% of living corals at Site-T and 70% at Site-Livingstone (mainly <i>Acropora</i> communities) (Turquet <i>et al.</i> , 2002).
2002	Early in the year 2002, large scale bleaching affected the world's largest continuous coral reef. More than 60% of the Great Barrier Reef bleached and up to 5% was severely damaged as a result. Bleaching events have also been recorded in Rodrigues in February (Hardman <i>et al.</i> , 2007). At Site-T, <i>Acropora</i> corals were flourishing, indicating that they recovered from the 2001-bleaching event.
2003	In February 2003, a new local bleaching phenomenon is recorded in Reunion, with variable intensity. The more affected sites were situated on the North of Saint Gilles La Saline, Pointe des Chateaux and Colimaçons (St Leu reef). Since 2001, Site-Livingstone showed a very low recovery (living coral cover < to 8%) (Turquet <i>et al.</i> , 2002). In the T-coral zone, <i>Acropora</i> bleached and a month later, all colonies of <i>A.austera-hemprichi</i> were dead (Naim & Tourrand pers.obs.), explaining in part the decrease in living coral coverage in 2004 (GCRMN). In Mauritius, McClanahan <i>et al.</i> (2005) documented 24% coral death.
2004	In 2004, a bleaching event affecting Site-T caused a decrease in coral coverage from 41 to 27% at Site-T while Site-P was hardly affected (GCRMN, Tessier <i>et al.</i> , 2008).
2005	Coral bleaching reported affected western Madagascar, Mauritius, and Reunion in March-April but did not cause significant mortality.
2009	In March, corals bleached intensely at Saint-Gilles la Saline. A large number of species were affected, including <i>Pavona spp.</i> , <i>Pocillopora damicornis</i> , but also the fire coral <i>Millepora</i> , Alcyonarians. The species <i>Porites nigrescens</i> appeared to be the most resistant. Since 2004, it was the most widespread bleaching phenomenon at Reunion. Global sea surface temperature measurements (NOAA) in this period suggested a coral bleaching related to thermal stress. GCRMN: at Site-T, half of <i>Acropora muricata</i> coverage bleached but was not dead at the time of the analysis. At Site-P, Cauvin observed a reduction of 50% living coral coverage whereas this coverage remained relatively stable for 11 years. <i>Montipora circumvallata</i> and <i>Porites (Synaraea) rus</i> and the few <i>Acropora muricata</i> were affected but only <i>P. (S.) rus</i> recovered. Living <i>M. circumvallata</i> coverage dropped from 34% in 2008 to 10% in 2009.
2010	GCRMN: at Site-T, <i>Acropora muricata</i> recovered from the 2009-bleaching event, and the living coral coverage still reached 43%. At Site-P, in 2010, some <i>Montipora circumvallata</i> recovered (2009 : 10% coverage, 2010 : 15%). <i>Porites (S.) rus</i> , which has withstood the 2009-bleaching, became the dominant species (2009 : 14%, 2010 : 20%).

Climate change is now considered to be one of the greatest long-term threats to coral reefs, with some future unavoidable change despite mitigation efforts (Hughes *et al.*, 2003; Bellwood *et al.*, 2004; Wild *et al.*, 2011). Managers must therefore focus on supporting the natural resilience of reefs, requiring identification of resilient reefs and reef regions. Considering the 1998-bleaching event in the southwest Indian Ocean, coral coverage suffered negligible damage in South Africa (Obura, 2005). Similarly, in Mauritius the impact was minimal with less than 10% coral cover affected (Moothien-Pillay *et al.*, 2002; Turner & Klaus 2005) although,

later, in 2004, 24% was affected (McClanahan *et al.*, 2005). After these two events, in Mauritius, coral cover was still dominated by thermally sensitive genera such as *Acropora* and overall cover appeared to have risen since a broad survey in 1992 (McClanahan *et al.*, 2005). Graham *et al.* (2007) recorded the composition and distribution of benthic and fish communities on a reef flat in Mauritius during five surveys between 1994 and 2005. They reported a trend of increasing coral coverage and the dominance of *Acropora*, highlighting Mauritius as a regional refugium for temperature-sensitive corals and specialized species of fish.

In the southwest Indian Ocean, McClanahan *et al.* (2007) showed that, from Kenya to Mauritius, coral communities were less disturbed, with *Acropora* and *Montipora* as dominating forms. Similarly, a survey of coral along Madagascar's northeast coast by researchers from Conservation International (CI), suggests that these island's reefs may have so far escaped the damaging effects of warmer ocean temperatures attributed to global climate change.

These findings suggest that the southern Indian Ocean often escape rising temperatures. In the Mascarenes, 189 reef-building coral species were recorded and *Acropora* was considered as the most diversified genus with 39 species (Faure, 1982; Faure *et al.*, 2008). Reefs from the southern Indian Ocean region can contribute then to the maintenance of *Acropora* biodiversity in the southwest Indian Ocean. These living *Acropora* may ultimately be used to reseed damaged reefs and may also provide valuable information about how to protect corals from future damage.

## CONCLUSION

In summary, the ACR-type community, typified both by the abundance of several *Acropora* species and by dense populations of sea urchins, has occurred at Site-Tobbogan for 25 years at least. During the same time interval, Site-Planch'Alizés has been occupied by the MAS-type community as dominating features. This is typified by massive corals, abundant benthic primary producers and lack of sea urchins. There is a greater temporal variability of composition of the ACR assemblages, due to a higher sensitivity of acroporids to ambient disturbances. However, recovery of ACR is quite rapid, resulting in *Acropora* populations recovering at a rate of up to 5% per year. By contrast, the cover rate of massive corals is more stable through time, except for some exceptional bleaching events (2009). Perennial macroalgal stands which invaded the MAS community during the 1980's, declined in abundance from the beginning of the 1990's. Related to this, massive coral colonies increased in coverage. Sea urchins permanently remained absent from the MAS assemblages and, as such, undergrazing lasted until now. In 2011, ACR and MAS communities still remain apparently luxuriant on the reef flats of Reunion.

## ACKNOWLEDGMENTS

We thank M. Daniel CHOUSSY† for his outstanding assistance during fieldwork. We also thank Professor Terence J. DONE for rereading this paper. This project was funded by the « programme quadriennal laboratoire ECOMAR, université de la Réunion, Région Réunion » and the « Ministère de l'Enseignement Supérieur et de la Recherche (MESR) ».



## REFERENCES

- Bachèlery, P.  
2000. Origin of the islands (chapter: The Mascarene Islands). *In: Coral reefs of the Indian Ocean: their Ecology and Conservation* (McClanahan, T., Sheppard, C., Obura, D.O. eds), Oxford Univ. Press, Pp. 360-361.
- Bellwood, D.R., T.P. Hughes, C. Folke and M. Nystroëm  
2004. Confronting the coral reef crisis. *Nature* 429: 827–833.
- Behrman, D.  
1981. *Assault on the largest unknown. The international Indian Ocean expedition 1959-65.* UNESCO Press, 96 pp.
- Biais, G. and J. Kopp  
1981. *Etude courantologique au droit des passes du lagon de La Saline.* Rapport ISTPM / DDA Réunion / Mairie de St-Paul, 46 pp.
- Bouchon, C.  
1981. Quantitative study of the scleractinian coral communities of a fringing reef of Reunion Island (Indian Ocean). *Marine Ecology Progress Series* 4: 273-288.  
1996. *Recherche sur les peuplements de Scleractinaires Indo-Pacifiques (Mer Rouge, Océan Indien, Océan Pacifique).* Doctorat ès Sciences, Univ. Aix-Marseille II, 338 pp.
- Borca, T. and N. Margoloff  
2004. Basculement des eaux de la Réunion: Un chantier titanesque. La Reunion Island: shifting waters from East to West, a giant jobsite. *Tunnels et ouvrages souterrains*, ISSN 0399-0834, CODEN TNOSAI 182 : 71-74.
- Bruggemann, H., M.M.M. Guillaume, V. Denis, L. Bigot, O. Naim, E. Tessier, P. Durville and T. Mulochau  
2007. The recent marine reserve at Reunion Island: Initial state of the benthic communities of the coral reefs of Saint-Leu and la Saline. *5<sup>th</sup> Western Indian Ocean Marine Science Association (WIOMSA) Conference, Durban, South Africa*, abstract
- Bruno, J.F. and E.R. Selig  
2007. Regional decline of coral cover in the Indo-Pacific: timing, extent and subregional comparisons. *PLoS ONE* 2 (8): e711.
- Burke, L., K. Reytar, M. Spalding and A. Perry  
2011. *Reefs at risk revisited.* World resources Institute, Washington D.C., 114 pp.
- Chabanet, P.  
1994. *Étude des relations entre les peuplements coralliens et les peuplements ichtyologiques sur le complexe récifal de Saint-Gilles La Saline.* Doctorat de spécialité Environnement marin, Université Aix-Marseille III., France, 200 pp.
- Chabanet P., J.-L. Join, P. Cuet and O. Naim  
1995. Spatial variability in submarine groundwater discharge occurrence and benthic and fish communities patterns on St-Gilles/la Saline reef : a tentative interpretation through an hydrogeological model (Reunion Island). *International Society for Reef Studies, Annual Meeting, Marseille, France*, abstract.
- Chabanet, P., H. Ralambondrainy, M. Amanieu, G.F. Faure and R. Galzin  
1997. Relationships between coral reef substrata and fish. *Coral Reefs* 16: 93-102.
- Chabanet, P., L. Bigot, O. Naim, R. Garnier, E. Tessier and M. Moyne-Picard  
2002. Coral reef monitoring at Reunion Island (Western Indian ocean) using the GCRMN method. *Proceedings of the 9<sup>th</sup> International Coral reef Symposium, Bali* 2 : 873-878.
- Chazottes, V., T. Le Campion-Alsumard, M. Peyrot-Clausade and P. Cuet  
2002. The effects of eutrophication-related alterations to coral reef communities on agents and rates of bioerosion. *Coral reefs*, 21: 375-390.
- Cheke, A.S.  
1987. An ecological history of the Mascarene Islands with particular refernce to land vertebrates. *In : Studies of Mascarene Island birds* (Diamond, E.W., ed.), Cambridge University press, Pp. 5-89.

- Clarke, K.R. and R.M. Warwick  
2001. *Change in marine communities: an approach to statistical analysis and interpretation*. 2<sup>nd</sup> edn, PRIMER-E Ltd, Plymouth.
- Clavier, J., L. Chauvaud, P. Cuet, C. Esbelin, P. Frouin, D. Taddei and G. Thouzeau  
2008. Diel variation of benthic respiration in a coral reef sediment (Reunion Island, Indian Ocean). *Estuarine, Coastal & Shelf Science* 76 (2): 369-377.
- Conand, C.  
1996. Asexual reproduction by fission in *Holothuria atra*: variability of some parameters in populations from the tropical Indo-Pacific. *Oceanologica acta* 19 (3-4): 209-216.  
1998. Holothurians. In: *FAO Species identification guide. The marine living resources of the western Central Pacific* (Carpenter K., Niem, V., eds) vol.2, Pp. 1157-1190.
- Conand, C. and C. De Ridder  
1990. Reproduction asexuée par scission chez *Holothuria atra* (Holothuroidea) dans des populations de platiers récifaux. In: *Echinoderm research* (De Ridder, C., Dubois, P., Jangoux, M., Lehayé, M.C., eds). Balkema, Rotterdam, Pp. 71-76.
- Conand, C., P. Chabanet, P. Cuet and Y. Letourneur  
1997. The carbonate budget of a fringing reef in La Reunion Island (Indian Ocean) : external bioerosion and benthic flux of CaCO<sub>3</sub>. *Proceedings of the 8<sup>th</sup> International Coral reef Symposium*, Panama, 1: 953-958.
- Conand, C., M. Heeb, M. Peyrot-Clausade and M.-F. Fontaine  
1998. Evaluations of bioerosion by two types of the sea urchin *Echinometra mathaei*, on several sites of a fringing reef in La Reunion Island (Indian Ocean) and comparison with other sites. In: *Echinoderms: San Francisco* (Mooi, R. & Telford, M., eds). Balkema, A.A. Rotterdam, Pp. 699-615.
- Conand, C., P. Cuet, O. Naim and D. Mioche  
2002a. Des coraux sous surveillance. *Pour la Science* 298 : 75-81.
- Conand, C., J.-P. Quod, F. Conand and J. Turquet  
2002b. Bleaching in a western Indian Ocean island, La Reunion: a multi-scale approach. *Proceedings of the 9<sup>th</sup> International Coral reef Symposium*, Bali, Indonesia 2: 1155-1159.
- Conand, F., F. Marsac, E. Tessier and C. Conand  
2007. A ten-year period of daily sea surface temperature at a coastal station in Reunion Island (July 1993-April 2004). *Western Indian Ocean Journal of Marine Science*, 6: 1-16.
- Cordier, E.,  
2007. *Processus physiques et sédimentaires en milieu récifal: application aux récifs frangeants de l'Hermitage/La Saline, Ile de La Réunion*. Doctorat de spécialité, Université de la Réunion, France, 150 pp.
- CORDIO  
1999. Consequences of the 1998 coral bleaching event for the islands of the western Indian Ocean. In: *Coral Reef Degradation in the Indian Ocean. Status report and project presentations* (Lindén, O. Sporrang, N., eds). SIDA /SAREC Mar. Sci. Progr., Sweden.
- Cover, W. and D. Potts  
2008. Direct specific impacts of sea-urchins on live corals. *International Society for Reef Studies*, Annual Meeting, summary
- Cuet, P.  
1989a. *Influence des résurgences d'eau douce sur les caractéristiques physico-chimiques et métaboliques de l'écosystème récifal à La Réunion*. Doctorat de spécialité en Environnement marin, Univ. Aix-Marseille III, France, 204 pp.  
1989b. *Les platiers récifaux de l'île de la Réunion. Physico-chimie des eaux*. Rapport Univ. Réunion (Lab.Ecol.Mar) - Agence Urbanisme Réunion, 151 pp.  
2007. *Influence de l'irrigation du littoral Ouest sur la qualité du milieu naturel. Suivi du réseau qualitatif Antenne 4. Années 2004-2005 et synthèse 1999-2005 (rapport final)*. Rapport Université de la Réunion - Département de la Réunion, 77 pp.
- Cuet, P., O. Naim, G.F. Faure and J.-Y. Conan

1988. Nutrient-rich groundwater impact on benthic communities of la Saline fringing reef (Reunion Island, Indian Ocean): preliminary results. *Proceedings of the 6<sup>th</sup> International Coral Reef Symposium*, Townsville, Australia 2: 207-212.
- Dauget, J.-M.  
1983. *Application of architectural methods used in tropical botany to the corals and the coral reefs*. Report Surat Pemberitahuan Penelitian, n° 070/4074, Seribu Islands, Indonesia, 6 pp.
- Denis, V., S. De Palmas, J. Debreuil, H. Bruggemann and M.M.M. Guillaume  
2009. Resisting climate change: phenotypic plasticity and environmental imprints in *Acropora muricata* from Reunion Island. *Proceedings of the 6<sup>th</sup> Western Indian Ocean Marine Science Association (WIOMSA) Scientific Symposium*, St-Denis, La Réunion, summary, p. 89.
- Done, T.J.  
1997. Decadal changes in reef-building communities: implications for reef growth and monitoring programs. *Proceedings of the 8<sup>th</sup> International Coral Reef Symposium* 1: 411-416.
- Done, T.J., P.K. Dayton, A.E. Dayton and R. Steger  
1991. Regional and local variability in recovery of shallow coral communities: Moorea, French Polynesia and Central Great Barrier Reef. *Coral reefs* 9: 183-192.
- Dustan, P. and J.C. Halas  
1987. Changes in the reef-coral community of Carysfort reef, Key Largo, Florida: 1974 to 1982. *Coral reefs* 6: 91-106.
- Edinger, E.N. and M.J. Risk  
2000. Reef classification by coral morphology predicts coral reef conservation value. *Biological Conservation* 92: 1-13.
- Faure, G. F.  
1974. Morphology and bionomy of the coral reef discontinuities in Rodrigues Island. *Proceedings of the 2<sup>nd</sup> International Coral Reef Symposium*, Brisbane, The Great Barrier Reef Committee 2: 161-172.  
1975. Etude comparative des récifs coralliens de l'Archipel des Mascareignes (Océan Indien). *Bulletin of the Mauritius Institute* 8: 1-26.  
1977. Distribution of coral communities on reef slopes in the Mascareignes Archipelago, Indian Ocean. *Marine Research Indonesia* 17 : 73-97.  
1982. *Recherche sur les peuplements de Scléactiniaires des récifs coralliens de l'Archipel des Mascareignes (Océan Indien Occidental)*. Doct. ès Sciences, Univ. Aix-Marseille II, 2 vol.  
1994. Principales dégradations de l'écosystème récifal. In: *Environnement en Milieu tropical* (Coudray, J., Bouguerra, M.L., eds), ESTEM, Paris: pp. 86-97.
- Faure, G. F. and L.F. Montaggioni  
1970. Le récif corallien de Saint-Pierre de la Réunion (Océan Indien): géomorphologie et répartition des peuplements. *Recueil des Travaux de la Station Marine Endoume, fascicule hors série, suppl. 10*: 271-284.  
1971a. Les récifs coralliens de l'île Rodrigues (Archipel des Mascareignes, Océan Indien): géomorphologie et répartition des peuplements. *Journal of the Marine Biology Association of India* 16 (3): 653-682.  
1971b. Les récifs coralliens Sous-Le-Vent de l'île Maurice (Archipel des Mascareignes, Océan Indien): morphologie et bionomie de la pente externe. *Comptes Rendus de l'Académie des Sciences, Paris, Série II* 273: 1914-1916.  
1976. Les récifs coralliens au-vent de l'île Maurice (Archipel des Mascareignes, Océan Indien): géomorphologie et bionomie de la pente externe. *Marine Geology* 21 : M9-M16.  
2009. Biodiversité des Scléactiniaires, la Saline 2009. *Rapport non publié*, GIP Réserve nationale marine Réunion, 10 pp.
- Faure, G.F., M.M.M. Guillaume, C.E. Payri, B. Thomassin and P. Vasseur  
1983. Massive bleaching and death of scleractinian corals in Mayotte coral reef ecosystem (Mozambique Channel, West Indian Ocean). *International Society for Reef Studies, Annual Meeting*, Nice: p. 26.
- Faure, G.F., M.M.M. Guillaume, C.E. Payri, B. Thomassin, M. Van Praet and P. Vasseur

1984. Sur un phénomène remarquable de blanchissement et de mortalité massive des Madréporaires dans le complexe récifal de l'île de Mayotte (S.W. Océan Indien). *Comptes Rendus de l'Académie des Sciences*, Paris sér. III, 299 (15): 637-642.
- Faure, G.F., M. Pichon, F. Trentin, Y. Geynet, N. Conruyt, P. Gigord, P. and D. Caron  
2008. Base de Connaissance sur les Coraux des Mascareignes. Université de la Réunion, 50pp.  
<http://coraux.univ-reunion.fr>
- Frouin, P.  
2000. Anthropogenic disturbances of tropical soft-bottom benthic communities. *Marine Ecology Progress Series* 194: 39-53.
- Gabrié, C. and L.F. Montaggioni  
1982. Sediments from fringing reefs of Reunion Island, Indian Ocean. *Sedimentary Geology* 31: 281-301.
- Gardiner, J.S.  
1936. The reefs of the Western Indian Ocean. I. Chagos archipelago. II. The Mascarene region. *Transactions of Linnean Society*, London 19: Pp. 393-436.
- Gardner, T.A., I.M. Côté, J.A. Gill, A. Grant and A. Watkinson  
2003. Long-term region-wide declines in Caribbean corals. *Science* 301 : 958-960.
- Glynn, P.W.  
1983. Widespread coral bleaching and death in Panama (1983). *Coral reefs newsletter, Pacific Science Association* 12: 8-9.
- Goreau, T., T.R. McClanahan, R. Hayes and A. Strong  
2000. Conservation of coral reefs after the 1998 global bleaching event. *Conservation Biology* 14: 5-15.
- Graham, N.A.J., T.R. McClanahan, Y. Letourneur and R. Galzin  
2007. Anthropogenic stressors inter-specific competition and ENSO effects on a Mauritian coral reef. *Environmental Biology of Fishes* 78: 57-69.
- Guillaume, M.M.M.  
1988. *La croissance du squelette de Porites lutea, Scleractiniaire hermatypique, sur le récif frangeant de La Saline, île de la Reunion, Océan Indien*. Doct. Spéc. Océanographie marine, Univ. Aix-Marseille II., France, 150 pp.
- Guillaume, M.M.M., C.E. Payri and G.F. Faure  
1983. Blatant dégradation of coral reefs at la Reunion Island (West Indian Ocean). *International Society for Reef Studies*, Annual Meeting, Nice, France, 28.
- Hardman, E.R., N.S. Stampfli, L. Hunt, S. Perrine, A. Perry and J.S. Jovani Raffin  
2007. The impact of coral bleaching in Rodrigues, Western Indian Ocean. *Atoll Research Bulletin* 555: 1-12.
- Horrill, J.C., A.T. Kamukuru, Y.D. Mgaya and M. Risk  
2000. Northern Tanzania and Zanzibar. In: *Coral reefs of the Indian Ocean : their Ecology and Conservation* (McClanahan, T., Sheppard, C., Obura, D.O. eds), Oxford Univ. Press, Pp. 167-198.
- Hughes, T.P., D.C. Reed and M.-J. Boyle  
1987. Herbivory on coral reefs: community structure following mass mortalities of sea urchins. *Journal of Experimental Marine Biology and Ecology* 113: 39-59.
- Hughes, T.P., A.H. Baird, D.R. Bellwood, M. Card, S. Connolly, C. Folke, R. Grosberg, O. Hoegh-Guldberg, J.B.C. Jackson, J. Kleypas, J.M. Lough, P. Marshall, M. Nystrom, S.R. Palumbi, J.M. Pandolfi, B. Rosen and J. Roughgarden  
2003. Climate change, human impacts, and the resilience of coral reefs. *Science* 301: 929-933.
- Hughes, T.P., N.A.J. Graham, J.B.C. Jackson, P.J. Mumby and R.S. Steneck  
2010. Rising the challenge of sustaining coral reef resilience. *Trends in ecology and evolution* 25: 633-642.
- INSEE  
Institut National de la Statistique et des Etudes Economiques, France. <http://www.insee.fr/>
- Join, J.-L.

1991. *Caractérisation hydrologique du milieu volcanique insulaire, le Piton des Neiges, île de la Réunion*. Doctorat de Spécialité en Hydrogéologie, Université Montpellier II, France, 187 pp.
- Kolasinski, J., K. Rogers, P. Cuet, B. Barry and P. Frouin  
2011. Sources of particulate organic matter at ecosystem scale: a stable isotope and trace element study in a coral reef (Reunion island, Indian Ocean). *Marine Ecology Progress Series* 443: 77-93.
- Letourneur, Y.  
1992. *Dynamique des peuplements ichtyologiques des platiers récifaux de l'île de la Réunion*. Doctorat de Spécialité en Océanologie, Université Perpignan-Réunion, France, 244 pp.  
2000. Spatial and temporal variability in territoriality of a tropical benthic damselfish on a coral reef (Reunion Island). *Environmental Biology of Fishes* 57: 377-391.
- Letourneur Y., M.L. Harmelin-Vivien and R. Galzin  
1993. Impact of hurricane Firinga on fish community structure on fringing reefs of Reunion Island, S.W. Indian Ocean. *Environmental Biology of Fishes* 37: 109-120.
- Lewis, J.B.  
1977. Processes of organic production on coral reefs. *Biological Reviews* 52: 305-347.
- Lindén, O., D. Souter, D. Wilhemsson and D.O. Obura  
2002. Coral Reef Degradation in the Indian Ocean. Status Report, CORDIO, Kalmar, Sweden, 284 pp.
- Littler, M.M., D.S. Littler and B.L. Brooks  
2009. Herbivory, nutrients, stochastic events, and relative dominances of benthic indicator groups on coral reefs: a review and recommendations. *Proceedings of the Marine Science Network Symposium*, Washington D.C. 38: 401-414
- Lopez, G.R. and J.S. Levinton  
1987. Ecology of the deposit-feeding animals in marine sediments. *Quarterly Review of Biology* 62: 235-260.
- Loya, Y.  
1978. Plotless and transect methods. In: *Coral research methods* (Stoddart D.R., Johannes R.E., eds). UNESCO, Paris, Pp. 197-217.
- Lucas, H.A. and G.A.F. Seber  
1977. Estimating coverage and particle density using the line intercept method. *Biometrika* 64 (3): 618-622.
- Mangion, P., D. Taddei, P. Frouin and C. Conand  
2004. Feeding rate and impact of sediment reworking by two deposit feeders *Holothuria leucospilota* and *Holothuria atra* on a fringing reef (Reunion Island, Indian Ocean). In: *Echinoderms* (München, Heizeller & Nebelsick, eds). Taylor & Francis Group, Londres, Pp. 311-317.
- McClanahan, T.R., C.R.C. Sheppard and D.O. Obura, eds,  
2000. *Coral reefs of the Indian Ocean. Their ecology and conservation*, Oxford University Press: 525 pp.
- McClanahan, T.R., J. Maina, R. Moothien-Pillay and A.C. Baker  
2005. Effects of geography, taxa, water flow, and temperature variation on coral bleaching intensity in Mauritius. *Marine Ecology Progress Series* 298: 131-142.
- McClanahan, T.R., M. Ateweberhan, N.A.J. Graham, S.K. Wilson, C.R. Sebastian, M.M.M. Guillaume and H. Bruggemann  
2007. Western Indian Ocean coral communities: Bleaching responses and susceptibility to extinction. *Marine Ecology Progress Series* 337: 1-13.
- Mioche, D.  
1998. *Aspects du cycle de la matière sur les récifs frangeants de l'île de la Réunion (Océan Indien) : signature des dégradations et mécanismes de l'eutrophisation*. Doctorat de Spécialité en Écologie marine, Université Réunion, France, 292 pp.
- Mioche, D. and P. Cuet

2002. Community metabolism on the reef flats at Reunion (Indian Ocean): natural versus anthropogenic disturbance. *Proceedings of the 9<sup>th</sup> International Coral Reef Symposium* Bali, Indonesia 1: 431-436.
- Mioche, D., R. Troadec, N. Germain, C. Esbelin, C. Tourrand, M. Psaradellis, O. Naim and J. Coudray  
2002. Relations entre l'équilibre sédimentaire des plages, la structure des communautés et le budget des carbonates dans l'écosystème corallien de la Réunion. *The Journal of Nature (Reunion)* 14 (1): 35-43.
- Montaggioni, L.F.  
1978. *Recherche géologiques sur les complexes récifaux de l'archipel des Mascareignes (Océan Indien occidental)*. Doctorat ès Sciences, Université Aix-Marseille II, 2 vol.
- Montaggioni, L.F. and G.F. Faure  
1980. Les récifs coralliens des Mascareignes (Océan Indien). *Collection des Travaux du Centre Universitaire, Université de la Réunion*, 151 pp.
- Moothien-Pillay, R., H. Terashima and H. Kawasaki  
2002. The extent and intensity of the 1998 mass bleaching event on the reefs of Mauritius, Indian Ocean. *Galaxea* 4: 43-52.
- Naim, O.  
1989. *Les platiers récifaux de l'île de la Réunion. Géomorphologie et état de santé des peuplements benthiques*. Rapport Université Réunion (Laboratoire d'Ecologie Marine) - Agence Urbanisme Réunion, 151 pp.  
1993. Seasonal responses of a fringing reef community to eutrophication (Reunion Island, Western Indian Ocean). *Marine Ecology Progress Series* 99: 307-315.  
2002. *Acropora formosa* (Dana, 1846) (Anthozoaire, Scléactiniaire), une espèce de corail structurant les biocénoses de platiers récifaux (Ile de la Réunion, S.O. Océan Indien). *The Journal of Nature (Reunion)* 14 (3): 103-111.  
2006. The coral reef communities at *Saint-Gilles La Saline* in 1987 (Reunion Island, South West Indian Ocean). *The Journal of Nature (Reunion)* 18 (1): 13-31.
- Naim, O., P. Cuet and Y. Letourneur  
1997. Experimental shift in benthic community structure. *Proceedings of the 8<sup>th</sup> International Coral Reef Symposium*, Panama 2: 1873-1878.
- Naim, O., P. Cuet and V. Mangar  
2000. The Mascarene Islands. In: *Coral reefs of the Indian Ocean : their Ecology and Conservation* (McClanahan, T., Sheppard, C., Obura, D.O. eds), Oxford Univ. Press, Pp. 353-381.
- Naim, O., P. Chabanet, T. Done, C. Tourrand and Y. Letourneur  
2002. Regeneration of a reef flat ten years after the impact of the cyclone Firinga (Reunion, SW Indian Ocean). *Proceedings of the 9<sup>th</sup> International Coral Reef Symposium*, Bali, Indonesia 1: 547-552.
- Naim, O., C. Tourrand, E. Ballesteros, S. Semple, L. Bigot, B. Cauvin, P. Cuet and L.F. Montaggioni  
2013a. Fringing reefs of Reunion Island and eutrophication effects. Part 2: Long-term monitoring of primary producers. *Atoll Research Bulletin*, 597, 36 p.
- Naim, O., C. Tourrand, G.F. Faure, L. Bigot, B. Cauvin, S. Semple and L.F. Montaggioni  
2013b. Fringing reefs of Reunion Island and eutrophication effects. Part 3: Long-term monitoring of living corals. *Atoll Research Bulletin*, 598, 32 p.
- Obura, D.O.  
2005. Resilience and climate change. Lessons from coral reefs and bleaching in the Western Indian Ocean. *Estuarine, Coastal & Shelf Science* 63: 353-372.
- Obura, D.O., J. Tamelander and O. Lindén, eds  
2008. *Coral Reef Degradation in the Indian Ocean (CORDIO)*. Status report. 2 vol., 494 pp.
- Pichon, M.  
1971. Comparative studies of the main features of some coral reefs of Madagascar, La Réunion and Mauritius. *Symposium of the Zoological Society London* 28 : 185-216.
- Quod, J.P.

1999. Consequences of the 1998 coral bleaching event for the islands of the western Indian Ocean. *In: Coral Reef Degradation in the Indian Ocean. Status report and project presentations* (Lindén, O. Sporrang, N., eds). SIDA /SAREC Mar. Sci. Progr., Sweden, Pp. 53-59.
- Sammarco, P.W.  
1982. Echinoid grazing as a structuring force in coral communities: whole reef manipulations. *Journal of Experimental Marine Biology & Ecology* 61 (1): 31-55.
- Scopélitis, J., S. Andrefouët, S. Phinn, P. Chabanet, O. Naim, C. Tourrand and T. Done  
2009. Changes of coral communities over 35 years: integration *in situ* and remote-sensing data on Saint-Leu reef (La Réunion, Indian Ocean). *Estuarine, Coastal & Shelf Science* 30: 1-11
- Semple, S.  
1997. The use of temporal variations in algal cover and biomass as a tool for the quantification of the degree of eutrophication of a fringing coral reef in Reunion Island, S.W. Indian Ocean. *Oceanologica Acta* 20 (6): 851-861.
- Sheppard, C.R.C.  
2000. Coral reefs of the Western Indian Ocean: an overview. In : *In: Coral reefs of the Indian Ocean : their Ecology and Conservation* (McClanahan, T., Sheppard, C., Obura, D.O. eds), Oxford Univ. Press, Pp. 3-38.
- Shinn, E.A.  
1976. Coral reef recovery in Florida and the Persian Gulf. *Environmental Geology* 1: 241–254.
- Souter, D. and O. Lindén, eds  
2005. *Coral Reef Degradation in the Indian Ocean (CORDIO) Status report* 2 vol., 288 pp.
- Spalding, M.D., C. Ravilious and E. Green  
2001. *World atlas of coral reefs*. University of California, 424 pp.
- Stoddart, D.R.  
1973. Coral reefs of the Indian Ocean. *In: Biology and Geology of coral reefs* (Jones, O.A., Endean, R., eds), London Academic press: Pp. 51-91.
- Tessier, E., L. Bigot, C. Cadet, B. Cauvin, P. Chabanet, C. Conand, J.-B. Nicet and J.-P. Quod  
2008. Les récifs coralliens de la Réunion en 2007: état des lieux et réseau de suivi. *Revue d'Ecologie (Terre & Vie)* 63: 85-102.
- Turner, J. and R. Klaus  
2005. Coral reefs of the Mascarenes, Western Indian Ocean. *Philosophical Transactions of the Royal Society of London A* 363: 229-250.
- Turner, J., E. Hardman, R. Klaus, I. Fagoonee, D. Daby, R. Baghooli and S. Persands  
2000. The reefs of Mauritius. *In: Coral Reef Degradation in the Indian Ocean* (Souter, D., Obura, D., and Linden, O. (eds)). CORDIO/SAREC, Pp. 94-107.
- Turquet, J.L., L. Bigot, C. Cambert, C. Conand, R. Garnier, E. Mirault and J.-P. Quod  
2002. Bleaching on *Acropora* dominated reefs in Reunion. *In: Coral Degradation in the Indian Ocean (CORDIO). Status report 2002* (Linden O., Souter D., Wilhemsoson D., Obura, D., eds.), Pp. 179-184.
- Wild, C., O. Hoegh-Guldberg, M.S. Naumann, M.F. Colombo-Pallotta, M. Ateweberhan, W.K. Fitt, R. Iglesias-Prieto, C. Palmer and J. Bythell  
2011. Climate change impedes scleractinian corals as primary reef ecosystem engineers. *Marine and Freshwater Research* 62: 205–215.
- Wilkinson, C., ed.  
2008. *Status of coral reefs of the world: 2008*. Global Monitoring Network and Reef and Rainforest Research Centre, Townsville, Australia, 296 pp.