



HAL
open science

Continuous-time limit of dynamic games with incomplete information and a more informed player.

Fabien Gensbittel

► **To cite this version:**

Fabien Gensbittel. Continuous-time limit of dynamic games with incomplete information and a more informed player.. 2015. hal-00910970v2

HAL Id: hal-00910970

<https://hal.science/hal-00910970v2>

Preprint submitted on 1 Jan 2015 (v2), last revised 11 Sep 2015 (v3)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Continuous-time limit of dynamic games with incomplete information and a more informed player.

Fabien Gensbittel*

January 1, 2015

Abstract

We study a two-player, zero-sum, dynamic game with incomplete information where one of the players is more informed than his opponent. We analyze the limit value as the players play more and more frequently. The more informed player observes the realization of a Markov process (X, Y) on which the payoffs depend, while the less informed player only observes Y and his opponent's actions. We show the existence of a limit value as the time span between two consecutive stages goes to zero. This value is characterized through an auxiliary optimization problem and as the unique viscosity solution of a second order Hamilton-Jacobi equation with convexity constraints.

Acknowledgments : The author gratefully acknowledges the support of the Agence Nationale de la Recherche, under grant ANR JEUDY, ANR-10-BLAN 0112.

1 Introduction.

This paper contributes to the literature on zero-sum dynamic games with incomplete information, by considering the case where one player is always more informed than his opponent.

A key feature appearing in recent contributions to the field of zero-sum dynamic games is the interplay between discrete-time and continuous-time dynamic models, as in Cardaliaguet-Laraki-Sorin [10], Neyman [27] or Cardaliaguet-Rainer-Rosenberg-Vieille [9], where the authors consider sequences of discrete-time dynamic games in which the players play more and more frequently. Such an analysis is related to the study of a sequence of discretizations in time of a given continuous-time dynamic game. In the present work, we adopt this method in order to study a continuous-time zero-sum dynamic game where one player is always more informed than his opponent and where the state variable evolves according to an exogenous Markov process. Precisely, we consider a model with two payoff-relevant variables $(X_t, Y_t)_{t \geq 0}$ which are evolving over time: X is a Markov chain with finite state space and Y is a diffusion process whose drift parameter depend on the current value of X . The process X is privately observed by the more informed player (say player 1) while Y is publicly observed, allowing the less informed player (player 2) to learn information about the variable X during the game. We analyze the sequence of discrete-time games indexed by $n \geq 1$ with incomplete information and perfect observation of actions, where stages occur at times $\frac{q}{n}$ for $q \geq 0$. At each stage, player 1 observes a pair of signals $(X_{\frac{q}{n}}, Y_{\frac{q}{n}})$ while player 2 only observes $Y_{\frac{q}{n}}$. The stage payoff

*Toulouse School of Economics (GREMAQ, Université Toulouse 1 Capitole), Manufacture des Tabacs, MF213, 21, Allée de Brienne 31015 Toulouse Cedex 6. E-mail:fabien.gensbittel@tse-fr.eu

function is assumed to depend on actions of both players and on $(X_{\frac{q}{n}}, Y_{\frac{q}{n}})$. The global payoff is a discounted sum of the stage payoffs with discount factor $\lambda_n = \int_0^{1/n} r e^{-rt} dt = 1 - e^{-r/n}$, where $r > 0$ is a given continuous-time discount rate. We assume that the stage payoffs are not observed and we study the limit value of these games as the players play more and more frequently.

We provide two characterizations for the limit value of these games as n goes to infinity. The first one is a probabilistic representation formula where the optimization variable is the set of admissible belief processes for the less informed player. Such a formula already appears in Sorin [28] as an illustration of the classical *Cav(u)* theorem of Aumann and Maschler [1]. A similar discrete-time formula was introduced by De Meyer in [12] in order to obtain a continuous-time limit value in a class of financial games and this approach led to several extensions in continuous-time models (see Cardaliaguet-Rainer [7, 8], Grün [19, 20], Gensbittel [16, 17], and more recently Cardaliaguet-Rainer-Rosenberg-Vieille [9] and Gensbittel-Grün [18]). This representation formula is important as it provides a characterization of optimal processes of revelation (martingales of posteriors induced by optimal strategies).

The second one is a variational characterization, the limit value is shown to be the unique viscosity solution of a second-order Hamilton-Jacobi equation with convexity constraints as introduced by Cardaliaguet [4, 5] and generalized in Cardaliaguet-Rainer [6], Grün [19], Cardaliaguet-Rainer-Rosenberg-Vieille [9] and Gensbittel-Grün [18].

2 Main results.

Notation 2.1. For any topological space E , $\Delta(E)$ denotes the set of Borel probability distributions on E endowed with the weak topology and the associated Borel σ -algebra. δ_x denotes the Dirac measure on $x \in E$. Finite sets are endowed with the discrete topology and Cartesian products with the product topology. $\mathbb{D}([0, \infty), E)$ denotes the set of càdlàg trajectories taking values in E , endowed with the topology of convergence in Lebesgue measure. The notations $\langle \cdot, \cdot \rangle$ and $|\cdot|$ stand for the canonical scalar product and the associated norm in \mathbb{R}^n .

Let us at first describe the continuous-time game we will approximate. This description is incomplete as we do not define strategies in continuous-time. Rather, we define below strategies in the different time-discretizations of this game. The notion of value for this game will therefore be the limit value along a sequence of discretizations when the mesh of the corresponding partitions goes to zero.

We assume that $(X_t)_{t \in [0, \infty)}$ is a continuous-time homogeneous Markov chain with finite state space K , infinitesimal generator $R = (R_{k,k'})_{k,k' \in K}$ and initial law $p \in \Delta(K)$. We identify $\Delta(K)$ with the canonical simplex of \mathbb{R}^K , i.e.:

$$\Delta(K) = \{p \in \mathbb{R}^K \mid \forall k \in K, p(k) \geq 0, \sum_{k \in K} p(k) = 1\}.$$

Then, define the real-valued process $(Y_t)_{t \in [0, \infty)}$ as the unique solution of the following stochastic differential equation (SDE)

$$\forall t \geq 0, Y_t = y + \int_0^t b(X_s, Y_s) ds + \int_0^t \sigma(Y_s) dW_t, \quad (2.1)$$

where $(W_t)_{t \in [0, \infty)}$ is a standard Brownian motion independent of X and $y \in \mathbb{R}$ is a given initial condition. The process Y may be seen as some noisy observation of the process X .

We assume that the functions b and σ in (2.1) are bounded and Lipschitz, and that there exists $\epsilon > 0$ such that for all $y \in \mathbb{R}$, $\sigma(y) \geq \epsilon$. The state process $Z := (X, Y)$ with values in $K \times \mathbb{R}$ is a well defined Feller Markov process, with semi-group of transition probabilities denoted $(P_t)_{t \geq 0}$.

Let I, J denote finite action sets for the two players (players 1 and 2), $g : (K \times \mathbb{R}) \times I \times J \rightarrow \mathbb{R}$ a bounded payoff function which is Lipschitz with respect to the second variable, and $r > 0$ a fixed discount rate.

We consider the following (heuristic) zero-sum game, played on the time interval $[0, \infty)$:

- Player 1 observes the trajectory of $Z = (X, Y)$.
- Player 2 observes only the trajectory of Y .
- They play the game $G(p, y)$ with total expected payoff for player 1:

$$\mathbb{E}\left[\int_0^{+\infty} r e^{-rt} g(X_t, Y_t, i_t, j_t) dt\right],$$

where i_t (resp. j_t) denote the action of player 1 at time t (resp. of player 2).

- Actions are observed during the game (and potentially convey relevant information).

We aim at studying the value function of this game and how information is used by the more informed player when playing optimally. In order to achieve this goal, we introduce a sequence of time-discretizations of the game. For simplicity, and without loss of generality, let us consider the uniform partition of $[0, +\infty)$ of mesh $1/n$. The corresponding discrete-time game, denoted $G_n(p, y)$ proceeds as follows:

- The variable $Z_{\frac{q}{n}} = (X_{\frac{q}{n}}, Y_{\frac{q}{n}})$ is observed by player 1 before stage q for $q \geq 0$.
- The variable $Y_{\frac{q}{n}}$ is observed by player 2 before stage q for $q \geq 0$.
- At each stage, both players choose simultaneously a pair of actions $(i_q, j_q) \in I \times J$.
- Chosen actions are observed after each stage.
- Stage payoff of player 1 equals $g(Z_{\frac{q}{n}}, i_q, j_q)$ (realized stage payoffs are not observed).
- The total expected payoff of player 1 is

$$\mathbb{E}\left[\lambda_n \sum_{q \geq 0} (1 - \lambda_n)^q g(Z_{\frac{q}{n}}, i_q, j_q)\right],$$

with $\lambda_n = 1 - e^{-r/n}$

Remark 2.2. When σ is constant and b depends only on X , the observation of player 2 correspond to a normally distributed random variable with mean $\int_{\frac{q}{n}}^{\frac{q+1}{n}} b(X_s) ds$ and variance $\frac{\sigma^2}{n}$. It may therefore be interpreted as a noisy observation of X .

The description of the game is common knowledge and we consider the game played in behavior strategies: at round q , player 1 and player 2 select simultaneously and independently an action $i_q \in I$ for player 1 and $j_q \in J$ for player 2 using some lotteries depending on their past observations.

Formally, a behavior strategy σ for player 1 is a sequence $(\sigma_q)_{q \geq 0}$ of transition probabilities:

$$\sigma_q : ((K \times \mathbb{R}) \times I \times J)^q \times (K \times \mathbb{R}) \rightarrow \Delta(I),$$

where $\sigma_q(Z_0, i_0, j_0, \dots, Z_{\frac{q-1}{n}}, i_{q-1}, j_{q-1}, Z_{\frac{q}{n}})$ denotes the lottery used to select the action i_q played at round q by player 1 when past actions played during the game are $(i_0, j_0, \dots, i_{q-1}, j_{q-1})$ and the sequence of observations of player 1 is $(Z_0, \dots, Z_{\frac{q}{n}})$. Let Σ denote the set of behavior strategies for player 1. Similarly, a behavior strategy τ for player 2 is a sequence $(\tau_q)_{q \geq 0}$ of transition probabilities depending on his past observations

$$\tau_q : (\mathbb{R} \times I \times J)^q \times \mathbb{R} \rightarrow \Delta(J).$$

Let \mathcal{T} denote the set of behavior strategies for player 2.

Let $\mathbb{P}_{(n,p,y,\sigma,\tau)} \in \Delta(\mathbb{D}([0, \infty), K \times \mathbb{R}) \times (I \times J)^{\mathbb{N}})$ denote the probability on the set of trajectories of Z and actions induced by the strategies σ, τ . The payoff function in $G_n(p, y)$ is defined by

$$\gamma_n(\nu, \sigma, \tau) := \mathbb{E}_{\mathbb{P}_{(n,p,y,\sigma,\tau)}} \left[\lambda_n \sum_{q \geq 0} (1 - \lambda_n)^q g(Z_{\frac{q}{n}}, i_q, j_q) \right].$$

The (lower) value of this game is defined by

$$V_n(p, y) := \sup_{\sigma \in \Sigma} \inf_{\tau \in \mathcal{T}} \gamma_n(p, y, \sigma, \tau).$$

Remark 2.3. V_n is actually the value of the game, meaning that the sup and inf in the above definition commute. However, we omit the proof of this result, which is based, e.g., on dynamic programming, as everything that will be said remains true for the lower value (i.e. sup inf) defined above.

We also need to consider the value function u of the non-revealing one-stage game $\Gamma(p, y)$, which is a finite game with payoff g in which player 1 cannot use his private information. Precisely,

$$u(p, y) := \sup_{\sigma \in \Delta(I)} \inf_{\tau \in \Delta(J)} \sum_{i \in I} \sum_{j \in J} \sum_{k \in K} p(k) \sigma(i) \tau(j) g(k, y, i, j),$$

and the value exists (i.e. the sup and inf commute in the above formula) as it is a finite game. It follows from standard arguments that u is Lipschitz in (p, y) .

The main results proved in sections 3 and 4 are two different characterizations for the limit of the sequence of value functions V_n .

Let us now introduce some notations.

Notation 2.4.

- The natural filtration \mathcal{F}^A of a process $(A_t)_{t \in [0, \infty)}$ is defined by $\mathcal{F}_t^A = \sigma(A_s, s \leq t)$. The associated right-continuous filtration is denoted $\mathcal{F}^{A,+}$ with $\mathcal{F}_t^{A,+} := \bigcap_{s > t} \mathcal{F}_s^A$.

- For any topological space E , $\mathbb{D}([0, \infty), E)$ denotes the set of E -valued càdlàg trajectories.
- For all $(p, y) \in \Delta(K) \times \mathbb{R}$, $\mathbb{P}_{p,y} \in \Delta(\mathbb{D}([0, +\infty), K \times \mathbb{R}))$ denotes the law of the process $Z = (X, Y)$ with initial law $p \otimes \delta_y$.

Our first main result is the following probabilistic characterization.

Theorem 2.5. For all $(p, y) \in \Delta(K) \times \mathbb{R}$,

$$V_n(p, y) \xrightarrow{n \rightarrow \infty} V(p, y) := \max_{(Z_t, \pi_t)_{t \geq 0} \in \mathcal{B}(p, y)} \mathbb{E} \left[\int_0^{+\infty} r e^{-rt} u(\pi_t, Y_t) dt \right], \quad (2.2)$$

where $\mathcal{B}(p, y) \subset \Delta(\mathbb{D}([0, \infty), (K \times \mathbb{R}) \times \Delta(K)))$ denotes the set of laws of càdlàg processes $(Z_t, \pi_t)_{t \in [0, \infty)}$ such that:

- $(Z_t)_{t \geq 0}$ has law $\mathbb{P}_{p,y}$ and is an $\mathcal{F}^{(Z, \pi)}$ -Markov process.
- For all $t \geq 0$, for all $k \in K$, $\pi_t(k) = \mathbb{P}(X_t = k | \mathcal{F}_t^{(\pi, Y)})$.

Let us comment briefly this result. We generalize here the idea that the problem the informed player is facing can be decomposed into two parts: at first he may decide how information will be used during the whole game, and then maximize his payoff under this constraint. To apply this method of decomposition, we need to identify precisely the set $\mathcal{B}(p, y)$ of achievable processes of posterior beliefs on X of the less informed player. The filtration $\mathcal{F}^{(\pi, Y)}$ represents information of player 2, which observes the process Y (a lower bound on information). The condition that Z is $\mathcal{F}^{(Z, \pi)}$ -Markov reflects the fact that player 2 cannot learn any information on the process X which is not known by player 1 (an upper bound on information) and the second condition simply says that π represents the process of beliefs of player 2 on X_t . Maximizers of the right-hand side of equation (2.2) represent optimal processes of revelation for the informed player and induce asymptotically optimal strategies for the informed player in the sequence of discretized games (see the proof of Theorem 2.5).

We now turn to the second characterization. Define $b(y) := b(k, y)_{k \in K} \in \mathbb{R}^K$ and for all $k \in K$ and $t \geq 0$, define the optional projection¹:

$$\chi_t(k) := \mathbb{P}(X_t = k | \mathcal{F}_t^{Y, +}).$$

Using Theorem 9.1 in [24] (see also the Note p.360 about the Markov property), the process $\psi := (\chi, Y)$ with values in $\mathbb{R}^K \times \mathbb{R}$ is a diffusion process satisfying the following stochastic differential equation:

$$\forall t \geq 0, \psi_t = \psi_0 + \int_0^t c(\psi_t) dt + \int_0^t \kappa(\psi_t) d\bar{W}_t, \quad (2.3)$$

where \bar{W} is a standard $\mathcal{F}^{Y, +}$ -Brownian motion and the vectors $c(p, y)$ and $\kappa(p, y)$ in $\mathbb{R}^{K+1} = \mathbb{R}^K \times \mathbb{R}$ are defined by

$$c(p, y) := (\mathbb{T}R p, \langle p, b(y) \rangle),$$

$$\kappa(p, y) := \left(\left(\frac{p_k}{\sigma(y)} (b(k, y) - \langle b(y), p \rangle) \right)_{k \in K}, \sigma(y) \right),$$

¹In all the proofs, we consider only natural or right-continuous filtrations, but we adopt the same convention as in [21] and do not complete the filtrations to avoid complex or ambiguous notations. Note that optional projections of càdlàg processes are well-defined and have almost surely càdlàg paths (see appendix 1 in [13]).

where ${}^{\mathbb{T}}R$ denotes the transpose of the matrix R and probabilities are seen as column vectors. We deduce from standard properties of diffusion processes that for any function $f \in C^2(\mathbb{R}^{K+1})$ with polynomial growth (say) and for all $0 \leq s \leq t$:

$$\mathbb{E}[f(\psi_t)|\mathcal{F}_s^\psi] = f(\psi_s) + \mathbb{E}\left[\int_s^t A(f)(\psi_u)du|\mathcal{F}_s^\psi\right]$$

where $A(f)$ is the differential operator defined by (using the notation $z = (p, y)$)

$$Af(z) = \langle Df(z), c(z) \rangle + \frac{1}{2} \langle \kappa(z), D^2 f(z) \kappa(z) \rangle.$$

In order to state our second main result, we need to define precisely the notion of weak solution we will use. Let $p \in \Delta(K)$, we define the tangent space at p by

$$T_{\Delta(K)}(p) := \{x \in \mathbb{R}^K \mid \exists \varepsilon > 0, p + \varepsilon x, p - \varepsilon x \in \Delta(K)\}.$$

Let \mathcal{S}^m denote the set of symmetric matrices of size m . Let $S \in \mathcal{S}^K$ and $p \in \Delta(K)$, we define

$$\lambda_{\max}(p, S) := \max \left\{ \frac{\langle x, Sx \rangle}{\langle x, x \rangle} \mid x \in T_{\Delta(K)}(p) \setminus \{0\} \right\}$$

and by convention $\lambda_{\max}(p, X) = -\infty$ whenever $T_{\Delta(K)}(p) = \{0\}$.

Theorem 2.6. *V is the unique continuous viscosity solution of*

$$\min\{rV + H(z, DV(z), D^2V(z)); -\lambda_{\max}(p, D_p^2V(z))\} = 0 \quad (2.4)$$

where for all $(z, \xi, S) \in (\Delta(K) \times \mathbb{R}) \times \mathbb{R}^{K+1} \times \mathcal{S}^{K+1}$:

$$H(z, \xi, S) := -\langle \xi, c(z) \rangle - \frac{1}{2} \langle \kappa(z), S \kappa(z) \rangle - ru(z),$$

and where DV, D^2V denote the gradient and the Hessian matrix of V and $D_p^2V(z)$ the Hessian matrix of the function V with respect to the variable p .

Let us recall the definitions of sub and super-solutions.

Definition 2.7. *We say that a bounded lower semi-continuous function f is a (viscosity) supersolution of the equation (2.4) on $\Delta(K) \times \mathbb{R}$ if for any test function ϕ, C^2 in a neighborhood of $\Delta(K) \times \mathbb{R}$ (in $\mathbb{R}^K \times \mathbb{R}$) such that $\phi \leq f$ on $\Delta(K) \times \mathbb{R}$ with equality in $(p, y) \in \Delta(K) \times \mathbb{R}$, we have*

$$\lambda_{\max}(p, D_p^2\phi(p, y)) \leq 0 \text{ and } r\phi(p, y) - A(\phi)(p, y) - ru(p, y) \geq 0.$$

We say that a bounded upper semi-continuous function f is a (viscosity) subsolution of the equation (2.4) on $\Delta(K) \times \mathbb{R}$ if for any test function ϕ, C^2 in a neighborhood of $\Delta(K) \times \mathbb{R}$ (in $\mathbb{R}^K \times \mathbb{R}$) such that $\phi \geq f$ on $\Delta(K) \times \mathbb{R}$ with equality in $(p, y) \in \Delta(K) \times \mathbb{R}$, we have

$$\lambda_{\max}(p, D_p^2\phi(p, y)) < 0 \Rightarrow r\phi(p, y) - A(\phi)(p, y) - ru(p, y) \leq 0.$$

The proof of Theorem 2.6 is based on theorem 2.5 and on dynamic programming.

2.1 Possible extensions and open problems.

We list below miscellaneous remarks.

- In comparison to [9], in the statement of Theorem 2.5, we maximize over a set of joint distributions (Z, π) rather than on the set of induced distributions for (π, Y) , which are the only relevant variables for the computation of the objective functional. The latter set of distributions is exactly the set of joint laws of càdlàg processes (π, Y) such that for all bounded continuous function ϕ on $\Delta(K) \times \mathbb{R}$ which are convex with respect to the first variable, we have:

$$\forall 0 \leq s \leq t, \mathbb{E}[\phi(\pi_t, Y_t) | \mathcal{F}_s^{(\pi, Y)}] \geq Q_{t-s}(\phi)(\pi_s, Y_s),$$

where Q is the semi-group of the diffusion process ψ . We do not prove this claim but it follows quite easily from Strassen's Theorem and the same techniques used in Lemma 4 in [9] and Lemma 5.11 in [18]. However, such a proof would not be constructive (due to Strassen's theorem) and therefore, we do not think that this result would be more interesting stated this way. Indeed, in order to construct asymptotically optimal strategies following the proof of Theorem 2.5, player 1 has to compute the joint law of (Z, π) anyway (precisely the conditional law of π given Z at times q/n for $q \geq 0$).

- One may generalize all the present results for the lower value functions to the case of infinite actions spaces I, J (even if the value u does not exist) by adapting the method developed in [16]. Note that the proof of the same kind of results for the upper value functions may rely on different tools as shown in [16], and that the extension of these results in the present model remains an open question.
- It can be shown directly (with classical arguments) that the functions V_n and V are continuous. However, this does not simplify nor shorten the proofs.
- It is reasonable to think that Theorem 2.5 can be extended to the case of a more general Feller processes (X, Y) , at least for diffusions with smooth coefficients. However, such an extension leads to the following open question: is it possible to write an Hamilton-Jacobi equation in the case of a diffusion process $Z = (X, Y)$ taking values in $\mathbb{R}^m \times \mathbb{R}^p$? Note that such an equation would be stated in an infinite dimensional space of probability measures.
- It would be interesting to try to find explicit solutions for simple examples with two states for X and with simple payoff functions and simple diffusion parameters for Y . Such an analysis and the comparison with the examples studied in [9] is left for future research.

3 Proof of Theorem 2.5

Recall the definition of conditional independence.

Definition 3.1. *Let $(\Omega, \mathcal{A}, \mathbb{P})$ a probability space and $\mathcal{F}, \mathcal{G}, \mathcal{H}$ three sub σ -fields of \mathcal{A} . We say that \mathcal{F} and \mathcal{G} are conditionally independent given \mathcal{H} if*

$$\forall F \in \mathcal{F}, \forall G \in \mathcal{G}, \mathbb{P}(F \cap G | \mathcal{H}) = \mathbb{P}(F | \mathcal{H})\mathbb{P}(G | \mathcal{H}).$$

This relation is denoted $\mathcal{F} \amalg_{\mathcal{H}} \mathcal{G}$ and the definition extends to random variables by considering the σ -fields they generate.

The next definition is related to the characterization of the Markov property in terms of conditional independence and will be useful in the sequel.

Definition 3.2. *Given two random processes $(A_q, B_q)_{q \geq 0}$ (with values in some Polish spaces) defined on $(\Omega, \mathcal{A}, \mathbb{P})$. We say that $(A_q)_{q \geq 0}$ is non-anticipative with respect to $(B_q)_{q \geq 0}$ if*

$$\forall q \geq 0, (A_0, \dots, A_q) \prod_{B_0, \dots, B_q} (B_m)_{m \geq 0}.$$

The next result is a classical property of conditional independence and its proof is postponed to the appendix.

Lemma 3.3. *Given two random processes $(A_q, B_q)_{q \geq 0}$ (with values in some Polish spaces), the process $(A_q)_{q \geq 0}$ is non-anticipative with respect to $(B_q)_{q \geq 0}$ if and only if there exists (on a possibly enlarged probability space) a sequence of independent random variables $(\xi_q)_{q \geq 0}$ uniformly distributed on $[0, 1]$ and independent of $(B_q)_{q \geq 0}$, and a sequence of measurable functions f_q (defined on appropriate spaces) such that for all $q \geq 0$*

$$A_q = f_q(B_p, \xi_m, m \leq q).$$

The proof of Theorem 2.5 is divided in two steps and relies on the technical Lemma 3.8, whose proof is postponed to the next subsection.

Step 1: We prove that $\liminf V_n \geq V$.

Let $\sigma^*(p, y)$ and $\tau^*(p, y)$ be measurable selections of optimal strategies for player 1 and 2 respectively, in the game $\Gamma(p, y)$ with value $u(p, y)$.

We start with a continuous-time process $(Z_t, \pi_t)_{t \geq 0}$ in $\mathcal{B}(p, y)$. We consider the discrete-time process $(Z_{\frac{q}{n}}, \pi_{\frac{q}{n}})_{q \geq 0}$. Using the Markov property at times $\frac{q}{n}$, we deduce that $(\pi_{\frac{q}{n}})_{q \geq 0}$ is non-anticipative with respect to $(Z_{\frac{q}{n}})_{q \geq 0}$. We now construct a strategy $\bar{\sigma}$ in $G_n(p, y)$ depending on the process (Z, π) . Using the conditional independence property (see Lemma 3.3), there exists a sequence $(\xi_q)_{q \geq 0}$ of independent random variables uniformly distributed on $[0, 1]$ and independent from $(Z_{\frac{q}{n}})_{q \geq 0}$, and a sequence of measurable functions $(f_q)_{q \geq 0}$ such that

$$\pi_{\frac{q}{n}} = f_q((Z_{\frac{m}{n}}, \xi_m)_{m \leq q}) \text{ for all } q \geq 0.$$

We define player 1's strategy σ as follows:

$$\bar{\sigma}_q(Z_0, \dots, Z_{\frac{q}{n}}, \xi_0, \dots, \xi_q) := \sigma^*(\pi_{\frac{q}{n}}, Y_{\frac{q}{n}}).$$

This does not define formally a behavior strategy but these transition probabilities induce a joint law for $(Z_{\frac{q}{n}}, i_q)_{q \geq 0}$ which can always be disintegrated in a behavior strategy (that does not depend on player 2's actions) since the induced process $(i_q)_{q \geq 0}$ is by construction non-anticipative with respect to $(Z_{\frac{q}{n}})_{q \geq 0}$ (using again Lemma 3.3). By taking the conditional expectation given $(Y_{\frac{\ell}{n}}, \xi_\ell, i_\ell, j_\ell)_{\ell=0, \dots, q}$, the payoff at stage q against any strategy τ is such that:

$$\mathbb{E}_{n,p,\bar{\sigma},\tau}[g(X_{\frac{q}{n}}, Y_{\frac{q}{n}}, i_q, j_q)] = \mathbb{E}_{n,p,\bar{\sigma},\tau}\left[\sum_{k \in K} \pi_{\frac{q}{n}}(k)g(k, Y_{\frac{q}{n}}, i_q, j_q)\right] \geq \mathbb{E}_{n,p,\bar{\sigma},\tau}[u(\pi_{\frac{q}{n}}, Y_{\frac{q}{n}})].$$

Therefore, $\bar{\sigma}$ is such that

$$V_n(p, y) \geq \inf_{\tau} \gamma_n(p, y, \bar{\sigma}, \tau) \geq \sum_{q \geq 0} \lambda_n (1 - \lambda_n)^q \mathbb{E}[u(\pi_{\frac{q}{n}}, Y_{\frac{q}{n}})].$$

Define $(\tilde{\pi}^n, \tilde{Z}^n)$ as the piecewise-constant process equal to (Z, π) at times $\frac{q}{n}$ for $q \geq 0$. Then $(\tilde{\pi}^n, \tilde{Z}^n)$ converges in probability to (Z, π) (see e.g. Lemma VI.6.37 in [21]) and therefore

$$\sum_{q \geq 0} \lambda_n (1 - \lambda_n)^q \mathbb{E}[u(\pi_{\frac{q}{n}}, Y_{\frac{q}{n}})] = \mathbb{E} \left[\int_0^\infty r e^{-rt} u(\tilde{\pi}_t^n, \tilde{Y}_t^n) dt \right] \xrightarrow{n \rightarrow \infty} \mathbb{E} \left[\int_0^\infty r e^{-rt} u(\pi_t, Y_t) dt \right]$$

As $(Z, \pi) \in \mathcal{B}(p, y)$ was chosen arbitrarily, we deduce that:

$$\liminf_{n \rightarrow \infty} V_n(p, y) \geq V(p, y)$$

Step 2: We prove that $\limsup V_n \leq V$.

Let us fix (p, y) and let ε_n a positive sequence going to zero. For all $n \geq 1$, let σ^n be an ε_n -optimal behavior strategy for player 1 in $G_n(p, y)$. We will construct a strategy τ^n for player 2 by induction such that for all $q \geq 0$ the expected payoff at round q is not greater than

$$\mathbb{E}_{n,p,y,\sigma^n,\tau^n}[u(\hat{p}_q, Y_{\frac{q}{n}}) + C|p_q - \hat{p}_q|_1], \quad (3.1)$$

for some constant C independent of n , where $|\cdot|_1$ denotes the ℓ_1 -norm and where for all $q \geq 0$, \hat{p}_q and p_q denote respectively the conditional laws of $X_{\frac{q}{n}}$ given the information of player 2 before and after playing round q . Precisely, for all $k \in K$:

$$\begin{aligned} \hat{p}_q(k) &:= \mathbb{P}_{(n,p,y,\sigma^n,\tau^n)}(X_{\frac{q}{n}} = k \mid Y_0, i_0, j_0, \dots, Y_{\frac{q-1}{n}}, i_{q-1}, j_{q-1}, Y_{\frac{q}{n}}), \\ p_q(k) &:= \mathbb{P}_{(n,p,y,\sigma^n,\tau^n)}(X_{\frac{q}{n}} = k \mid Y_0, i_0, j_0, \dots, Y_{\frac{q}{n}}, i_q, j_q). \end{aligned}$$

Note that the computation of \hat{p}_q does not depend on τ_q^n . We can therefore define by induction $\tau_q^n := \tau^*(\hat{p}_q, Y_{\frac{q}{n}})$. Then, inequality (3.1) follows directly from Lemmas V.2.5 and V.2.6 in [25]. We now suppress the indices $(n, p, y, \sigma^n, \tau^n)$ from the probabilities and expectations. Using that u is Lipschitz with respect to p , we have

$$\mathbb{E}[u(\hat{p}_q, Y_{\frac{q}{n}}) + C|p_q - \hat{p}_q|_1] \leq \mathbb{E}[u(p_q, Y_{\frac{q}{n}}) + 2C|p_q - \hat{p}_q|_1]$$

Define also:

$$\tilde{p}_{q+1} := \mathbb{P}(X_{\frac{q+1}{n}} = k \mid Y_0, i_0, j_0, \dots, Y_{\frac{q}{n}}, i_q, j_q) = (e^{\frac{1}{n} \text{TR}} p_q)(k).$$

Note that for all $q \geq 0$, the sequence $(\tilde{p}_{q+1}, \hat{p}_{q+1}, p_{q+1})$ is a martingale so that using Jensen's inequality.

$$\mathbb{E}[(\tilde{p}_{q+1})^2] \leq \mathbb{E}[(\hat{p}_{q+1})^2].$$

On the other hand, using the previous equality, we can choose the constant C so that almost surely

$$\forall q \geq 0, \quad |\tilde{p}_{q+1} - p_q| \leq \frac{C}{n}.$$

Mimicking the proof of [9], we have

$$\begin{aligned} \mathbb{E} \left[\sum_{q \geq 0} \lambda_n (1 - \lambda_n)^q |p_q - \hat{p}_q|_1 \right] &= \sum_{k \in K} \sum_{q \geq 0} \lambda_n (1 - \lambda_n)^q \mathbb{E}[|p_q(k) - \hat{p}_q(k)|] \\ &\leq \sum_{k \in K} \left(\sum_{q \geq 0} \lambda_n (1 - \lambda_n)^q \mathbb{E}[|p_q(k) - \hat{p}_q(k)|^2] \right)^{1/2} \\ &= \sum_{k \in K} \left(\sum_{q \geq 0} \lambda_n (1 - \lambda_n)^q \mathbb{E}[(p_q(k))^2 - (\hat{p}_q(k))^2] \right)^{1/2} \end{aligned}$$

which is also equal to

$$\sum_{k \in K} \left(\sum_{q \geq 0} \lambda_n (1 - \lambda_n)^q \mathbb{E}[(p_q(k))^2 - (\tilde{p}_{q+1}(k))^2 + (\tilde{p}_{q+1}(k))^2 - (\hat{p}_{q+1}(k))^2 + (\hat{p}_{q+1}(k))^2 - (\hat{p}_q(k))^2] \right)^{1/2}$$

and therefore is bounded from above by

$$\sum_{k \in K} \left(\sum_{q \geq 0} \lambda_n (1 - \lambda_n)^q \mathbb{E}[(\hat{p}_{q+1}(k))^2 - (\hat{p}_q(k))^2] + \frac{2C}{n} \right)^{1/2} \leq N(\lambda_n + \frac{2C}{n})^{1/2}$$

We proved that:

$$V_n(p, y) \leq \sum_{q \geq 0} \lambda_n (1 - \lambda_n)^q \mathbb{E}[u(\pi_{\frac{q}{n}}, Y_{\frac{q}{n}})] + N(\lambda_n + \frac{2C}{n})^{1/2} + \varepsilon_n$$

In order to conclude the proof, we consider the continuous-time process $(\tilde{Z}^n, \tilde{\pi}^n)$ which is piecewise-constant and equal to $(Z_{\frac{q}{n}}, p_q)$ at times q/n . Let us at first extract a subsequence of $V_n(p, y)$ which converges to $\limsup V_n(p, y)$. Then, using Lemma 3.8, there exists a further subsequence of $(\tilde{Z}^n, \tilde{\pi}^n)$ which converges in law to some process (Z, π) in $\mathcal{B}(p, y)$. We have therefore along this subsequence

$$\sum_{q \geq 0} \lambda_n (1 - \lambda_n)^q \mathbb{E}[u(\pi_{\frac{q}{n}}, Y_{\frac{q}{n}})] = \mathbb{E}[\int_0^\infty re^{-rt} u(\tilde{\pi}_t^n, \tilde{Y}_t^n) dt] \longrightarrow \mathbb{E}[\int_0^\infty re^{-rt} u(\pi_t, Y_t) dt],$$

so that

$$\limsup_{n \rightarrow \infty} V_n(p, y) \leq \mathbb{E}[\int_0^\infty re^{-rt} u(\pi_t, Y_t) dt] \leq V(p, y).$$

3.1 A technical Lemma

In reference to the paper of Meyer and Zheng [26], we will denote MZ the following topology on the set of càdlàg paths.

Notation 3.4. For a separable metric space (E, d) , the MZ -topology on the set $\mathbb{D}([0, \infty), E)$ of càdlàg functions is the topology of convergence in measure when $[0, \infty)$ is endowed with the measure $e^{-x} dx$. The associated weak topology over the set $\Delta(\mathbb{D}([0, \infty), E))$ when $\mathbb{D}([0, \infty), E)$ is endowed with the MZ -topology will be denoted $\mathcal{L}(MZ)$.

Remark 3.5. In contrast to the Skorokhod topology (*Sk* hereafter), if $E = F \times F'$ is a product of separable metric spaces, the MZ topology is a product topology, i.e. (as topological spaces)

$$(\mathbb{D}([0, \infty), F \times F'), MZ) = (\mathbb{D}([0, \infty), F), MZ) \times (\mathbb{D}([0, \infty), F'), MZ).$$

The following remark will be used in the proofs.

Remark 3.6. If E is a Polish space, the space $(\mathbb{D}([0, \infty), E), MZ)$ is a separable metric space which is not topologically complete. However, its Borel σ -algebra is the same as the one generated by the *Sk* topology and its topology is weaker than the *Sk* topology for which the space is Polish, implying that all the probability measures are MZ -tight. Therefore, all the results about disintegration and measurable selection usually stated for Polish spaces and which depend only on the Borel structure apply to this space.

Recall that the transition probabilities of Z are denoted $(P_t)_{t \geq 0}$, i.e. for any bounded measurable function ϕ on $K \times \mathbb{R}$, we have

$$P_t(\phi)(z) := \mathbb{E}_z[\phi(Z_t)] = \int_{\mathcal{Z}} \phi dP_t(z),$$

and that P is a Feller semi-group implying that $(z, t) \rightarrow P_t(\phi)(z)$ is continuous for any bounded continuous function ϕ .

Notation 3.7. Given a process $(Z_t)_{t \in [0, \infty)}$ of law $\mathbb{P}_{p, y}$, we define the process $(\widehat{Z}_t^n)_{t \in [0, \infty)} \in \mathbb{D}([0, \infty), K \times \mathbb{R})$ by

$$\forall t \geq 0, \widehat{Z}_t^n := Z_{\lfloor \frac{nt}{n} \rfloor}$$

where $\lfloor a \rfloor$ denotes the greatest integer lower or equal to a .

Lemma 3.8. Let (p, y) be given, and let us consider a sequence of càdlàg processes (Z^n, π^n) that are piecewise constant on the partition $\{\lfloor \frac{q}{n}, \frac{q+1}{n} \rfloor\}_{q \geq 0}$ and such that

- Z^n has the same law as \widehat{Z}^n (see the above notation).
- $(\pi_{\frac{q}{n}})_{q \geq 0}$ is non-anticipative with respect to $(Z_{\frac{q}{n}})_{q \geq 0}$.
- For all $t \geq 0$, for all $k \in K$, $\pi_t^n(k) = \mathbb{P}(X_t^n = k | \mathcal{F}_t^{(\pi^n, Y^n)})$.

Then, the sequence (Z^n, π^n) admits an $\mathcal{L}(MZ)$ -convergent subsequence and all the limit points belong to $\mathcal{B}(p, y)$.

Proof. Let \mathbb{Q}_n denote a sequence of laws of processes $(Z_t^n, \pi_t^n)_{t \in [0, \infty)}$. It follows from Proposition VI.6.37 in [21] that Z^n $\mathcal{L}(MZ)$ -converges to Z of law $\mathbb{P}_{p, y}$. On the other hand, Theorem 4 in [26] together with a diagonal extraction implies that the set of possible laws for $(Z_t^n, \pi_t^n)_{t \geq 0}$ is MZ -relatively sequentially compact, and we may extract some convergent subsequence²

Let us now prove that the limit belongs to $\mathcal{B}(p, y)$. Assume (without loss of generality) that the sequence of processes $(Z_t^n, \pi_t^n)_{t \geq 0}$ $\mathcal{L}(MZ)$ -converges to $(Z_t, \pi_t)_{t \geq 0}$. Note at first that the law of $(Z_t)_{t \geq 0}$ is $\mathbb{P}_{p, y}$ since the projection of the trajectories on the first coordinate is continuous (see Remark 3.5).

Using Skorokhod's representation Theorem for separable metric spaces (see Theorem 11.7.31 in [14]), we can assume that the processes are defined on the same probability space and that $(Z_t^n, \pi_t^n)_{t \geq 0} \xrightarrow{MZ} (Z_t, \pi_t)_{t \geq 0}$ almost surely. Up to extracting a subsequence, we can also assume that there exists a subset I of full measure in $[0, \infty)$ (i.e. $\int_I e^{-x} dx = 1$) such that for all $t \in I$, $(Z_t^n, \pi_t^n) \rightarrow (Z_t, \pi_t)$ almost surely.

We now prove that for all $t \geq 0$ and all $k \in K$

$$\pi_t(k) = \mathbb{P}(X_t = k | \mathcal{F}_t^{(Y, \pi)}).$$

For any $t \in I$, any finite family (t_1, \dots, t_r) in $I \cap [0, t]$ and any bounded continuous function ϕ defined on $(\mathbb{R} \times \Delta(K))^r$, we have

$$\mathbb{E}[(\pi_t^n(k) - \mathbf{1}_{X_t^n=k})\phi(Y_{t_1}^n, \pi_{t_1}^n, \dots, Y_{t_r}^n, \pi_{t_r}^n)] = 0.$$

²Precisely, for all $T > 0$ we may first apply this result to each coordinate of the processes $(Z_{t \wedge T}^n, \pi_{t \wedge T}^n)_{t \geq 0}$. Then, since convergent sequences are tight (see Theorem 11.5.3 in [14] and remark 3.6), we apply Lemma A.4 to deduce that the set of laws $\{\mathbb{Q}_n, n \geq 1\}$ is tight. Applying the direct part of Prohorov's theorem, which is valid for separable metric spaces, we may extract some convergent subsequence.

It follows by bounded convergence that

$$\mathbb{E}[(\pi_t - \mathbb{1}_{X_t=k})\phi(Y_{t_1}, \pi_{t_1}, \dots, Y_{t_r}, \pi_{t_r})] = 0.$$

We deduce that

$$\pi_t(k) = \mathbb{P}(X_t = k | \mathcal{F}_t^{(Y, \pi)}).$$

Given an arbitrary t , we take a decreasing sequence in I with limit t and applying Lemma A.7 (see appendix), we obtain:

$$\pi_t(k) = \mathbb{P}(X_t = k | \mathcal{F}_t^{(Y, \pi, +)}),$$

which implies the result using the tower property of conditional expectations.

It remains to prove the Markov property. Let $t_1 \leq \dots \leq t_r \leq s \leq t$ in I , and ϕ, ϕ' some bounded continuous functions defined on $(\mathbb{R} \times \Delta(K))^r$ and $K \times \mathbb{R}$, we claim that

$$\mathbb{E}[\phi'(Z_t)\phi(Y_{t_1}, \pi_{t_1}, \dots, Y_{t_r}, \pi_{t_r})] = \mathbb{E}[P_{t-s}(\phi')(Z_s)\phi(Y_{t_1}, \pi_{t_1}, \dots, Y_{t_r}, \pi_{t_r})].$$

Indeed, for all n , we have

$$\mathbb{E}[\phi'(Z_t^n)\phi(Y_{t_1}^n, \pi_{t_1}^n, \dots, Y_{t_r}^n, \pi_{t_r}^n)] = \mathbb{E}[P_{\lfloor nt \rfloor - \lfloor ns \rfloor}(\phi')(Z_s^n)\phi(Y_{t_1}^n, \pi_{t_1}^n, \dots, Y_{t_r}^n, \pi_{t_r}^n)],$$

and the conclusion follows by bounded convergence. The property extends to arbitrary $t_1 \leq \dots \leq t_r \leq s \leq t$ by taking decreasing sequences in I and we conclude as above that Z is an $\mathcal{F}^{(Z, \pi)}$ Markov process. \square

Let us end this section with a second technical lemma whose proof is similar to Lemma 3.8.

Lemma 3.9. *The set-valued map $(p, y) \rightarrow (\mathcal{B}(p, y), \mathcal{L}(MZ))$ has a closed graph with compact values.*

Proof. Since Z is a Feller process, the map $(p, y) \rightarrow \mathbb{P}_{p, y}$ is $\mathcal{L}(Sk)$ -continuous (hence $\mathcal{L}(MZ)$ -continuous, see e.g. [15]). We omit the rest of the proof as it follows exactly from the same arguments as Lemma 3.8 with obvious modifications. \square

4 The variational characterization

We state at first some properties of the function V .

Proposition 4.1. *V is upper-continuous and for all $y \in \mathbb{R}$, $p \rightarrow V(p, y)$ is concave on $\Delta(K)$.*

Proof. That V is upper semi-continuous follows directly from Lemma 3.9.

Concavity follows from the same method as the well-known splitting Lemma (see e.g. Chapter V.1 in [25]). Given $y \in \mathbb{R}$, $p_1, p_2 \in \Delta(K)$ and $\lambda \in [0, 1]$, $\mathbb{P}_1 \in C(p_1, y)$ and $\mathbb{P}_2 \in C(p_2, y)$, let us construct $\mathbb{P}_\lambda \in C(\lambda p_1 + (1 - \lambda)p_2, y)$ as follows. Assume that (Z^1, π^1) and (Z^2, π^2) are independent and of respective laws \mathbb{P}_1 and \mathbb{P}_2 . Let ξ be a random variable independent of (Z^1, π^1) and (Z^2, π^2) and such that $\mathbb{P}(\xi = 1) = \lambda$ and $\mathbb{P}(\xi = 2) = 1 - \lambda$. Define (Z, π) as the process equal to (Z^i, π^i) on $\{\xi = i\}$. It follows easily by conditioning on ξ that

$$\mathbb{E}\left[\int_0^\infty r e^{-rt} u(\pi_t, Y_t) dt\right] = \lambda \mathbb{E}\left[\int_0^\infty r e^{-rt} u(\pi_t^1, Y_t^1) dt\right] + (1 - \lambda) \mathbb{E}\left[\int_0^\infty r e^{-rt} u(\pi_t^2, Y_t^2) dt\right].$$

If we assume that (Z, π) has a law $\mathbb{P}_\lambda \in C(\lambda p_1 + (1 - \lambda)p_2, y)$, then for any $\varepsilon > 0$, we can choose \mathbb{P}_1 and \mathbb{P}_2 as ε -optimal probabilities so that

$$\begin{aligned} V(\lambda p_1 + (1 - \lambda)p_2, y) &\geq \mathbb{E}\left[\int_0^\infty re^{-rt}u(\pi_t, Y_t)dt\right] \\ &= \lambda\mathbb{E}\left[\int_0^\infty re^{-rt}u(\pi_t^1, Y_t^1)dt\right] + (1 - \lambda)\mathbb{E}\left[\int_0^\infty re^{-rt}u(\pi_t^2, Y_t^2)dt\right] \\ &\geq \lambda V(p_1, y) + (1 - \lambda)V(p_2, y) - \varepsilon, \end{aligned}$$

and this proves that V is concave with respect to p as ε can be chosen arbitrarily small.

In order to conclude, it remains therefore to prove that (Z, π) has a law $\mathbb{P}_\lambda \in C(\lambda p_1 + (1 - \lambda)p_2, y)$. Note at first that $(Z_t)_{t \geq 0}$ has law $\mathbb{P}_{\lambda p_1 + (1 - \lambda)p_2, y}$ by construction. Moreover, $\mathcal{F}_t^{\pi, Y}$ is included in $\sigma(\xi) \vee \mathcal{F}_t^{\pi^1, Y^1} \vee \mathcal{F}_t^{\pi^2, Y^2}$. Using independence (see Lemma A.2), we have therefore for all $k \in K$ and all $t \geq 0$:

$$\begin{aligned} \mathbb{P}(X_t = k | \mathcal{F}_t^{Y^1, \pi^1}, \mathcal{F}_t^{Y^2, \pi^2}, \xi) &= \mathbb{P}(X_t^1 = k | \mathcal{F}_t^{Y^1, \pi^1}, \mathcal{F}_t^{Y^2, \pi^2}, \xi) \mathbb{1}_{\xi=1} + \mathbb{P}(X_t^2 | \mathcal{F}_t^{Y^1, \pi^1}, \mathcal{F}_t^{Y^2, \pi^2}, \xi) \mathbb{1}_{\xi=2} \\ &= \mathbb{P}(X_t^1 = k | \mathcal{F}_t^{Y^1, \pi^1}) \mathbb{1}_{\xi=1} + \mathbb{P}(X_t^2 = k | \mathcal{F}_t^{Y^2, \pi^2}) \mathbb{1}_{\xi=2} = \pi_t^1(k) \mathbb{1}_{\xi=1} + \pi_t^2(k) \mathbb{1}_{\xi=2} = \pi_t(k), \end{aligned}$$

and using the tower property of conditional expectations, we deduce that

$$\pi_t(k) = \mathbb{P}(X_t = k | \mathcal{F}_t^{(\pi, Y)}).$$

To prove the Markov property, let $s \geq t$ and ϕ some bounded continuous function on $\Delta(K) \times \mathbb{R}$. As above, we have:

$$\begin{aligned} \mathbb{E}[\phi(Z_s) | \mathcal{F}_t^{Z^1, \pi^1}, \mathcal{F}_t^{Z^2, \pi^2}, \xi] &= \sum_i \mathbb{1}_{\xi=i} \mathbb{E}[\phi(Z_s^i) | \mathcal{F}_t^{Z^1, \pi^1}, \mathcal{F}_t^{Z^2, \pi^2}, \xi] \\ &= \sum_i \mathbb{1}_{\xi=i} \mathbb{E}[\phi(Z_s^i) | \mathcal{F}_t^{Z^i}] \\ &= \sum_i \mathbb{1}_{\xi=i} P_{s-t}(\phi)(Z_t^i) = P_{s-t}(\phi)(Z_t). \end{aligned}$$

The conclusion follows by using the tower property of conditional expectation with the intermediate σ -field $\mathcal{F}_t^{Z, \pi}$. \square

4.1 Dynamic programming.

Notation 4.2. *In the following, we will use the notation $\mathbb{E}_{p, y}$ to denote the expectation associated to the diffusion process ψ starting at time 0 with initial position $\psi_0 = (p, y)$.*

We now state a dynamic programming principle which will be the key element for the proof of Theorem 2.6.

Proposition 4.3. *For all $(p, y) \in \Delta(K) \times \mathbb{R}$, for all $h \geq 0$, we have*

$$V(p, y) = \max_{(\pi, Y) \in \mathcal{B}(p, y)} \mathbb{E}\left[\int_0^h re^{-rt}u(\pi_t, Y_t)dt + e^{-rh}V(\pi_h, Y_h)\right]. \quad (4.1)$$

As a consequence,

$$V(p, y) \geq \mathbb{E}_{p, y} \left[\int_0^h r e^{-rt} u(\psi_t) dt + e^{-rh} V(\psi_h) \right]. \quad (4.2)$$

Moreover, if (π, Y) is an optimal process for $V(p, y)$, then for all $h \geq 0$:

$$V(p, y) = \mathbb{E} \left[\int_0^h r e^{-rt} u(\pi_t, Y_t) dt + e^{-rh} V(\pi_h, Y_h) \right]. \quad (4.3)$$

Proof. We prove at first that the maximum is reached in the right-hand side of (4.1). Let us define the MZ -topology on the set $\mathbb{D}([0, h], K \times \mathbb{R} \times \Delta(K))$ as the convergence in Lebesgue measure of the trajectories together with the convergence of the value of the process at time h . Note that this topology coincides (up to an identification) with the induced topology on the subset of $\mathbb{D}([0, \infty), K \times \mathbb{R} \times \Delta(K))$ made by trajectories that are constant on $[h, \infty)$. Using this identification and adapting the arguments of Lemma 3.8, the set of laws of the restrictions of the processes $(Z, \pi) \in \mathcal{B}(p, y)$ to the time interval $[0, h]$ is $\mathcal{L}(MZ)$ -sequentially relatively compact in $\Delta(\mathbb{D}([0, h], K \times \mathbb{R} \times \Delta(K)))$. The existence of a maximum follows since the map

$$\mathbb{P} \in \Delta(\mathbb{D}([0, h], K \times \mathbb{R} \times \Delta(K))) \longrightarrow \mathbb{E}_{\mathbb{P}} \left[\int_0^h r e^{-rt} u(\pi_t, Y_t) dt + e^{-rh} V(\pi_t, Y_t) \right],$$

is $\mathcal{L}(MZ)$ upper-semi-continuous.

We now prove (4.1). We begin with a measurable selection argument.

The function $\mathbb{P} \in \mathcal{B}(p, y) \rightarrow J(\mathbb{P}) := \mathbb{E}[\int_0^\infty r e^{-rt} u(\pi_t, Y_t) dt]$ is $\mathcal{L}(MZ)$ -continuous, and the set-valued map $(p, y) \rightarrow \mathcal{B}(p, y)$ is $\mathcal{L}(MZ)$ upper-semi-continuous. We deduce the subset O of the space $\Delta(K) \times \mathbb{R} \times \Delta(\mathbb{D}([0, \infty), (K \times \mathbb{R}) \times \Delta(K)))$ defined by

$$O := \{(p, y, \mathbb{P}) \mid \mathbb{P} \in \mathcal{B}(p, y), J(\mathbb{P}) \geq V(p, y)\}$$

is Borel-measurable (see Remark 3.6). Moreover, Lemma 3.8 implies that for any (p, y) , there exists some \mathbb{P} such that $(p, y, \mathbb{P}) \in O$. It follows therefore from Von Neumann's selection Theorem (see e.g. Proposition 7.49 in [2]) that there exists an optimal universally-measurable selection ϕ from $\Delta(K) \times \mathbb{R}$ to $\mathcal{B}(p, y)$ such that for all $(p, y) \in \Delta(K) \times \mathbb{R}$, $\phi(p, y) \in O$.

Let $(Z, \pi) \in \mathcal{B}(p, y)$ and $h \geq 0$ and let μ_h denote the joint law of (π_h, Y_h) . By construction, ϕ is μ_h -almost surely equal to a Borel map $\tilde{\phi}$. Using Lemma A.6, we can construct a process $(\tilde{\pi}_s)_{s \geq h}$ (on some extension of the probability space) such that the conditional law of $(Z_{h+s}, \tilde{\pi}_{h+s})_{s \geq 0}$ given $\mathcal{F}_h^{(Y, \pi)}$ is precisely $\tilde{\phi}(\pi_h, Y_h)$.

Let us consider the process $(Z, \hat{\pi})$ where $\hat{\pi}$ is equal to π on $[0, h)$ and to $\tilde{\pi}$ on $[h, \infty)$. Using the preceding construction, if we assume that the process $(Z, \hat{\pi})$ has a law in $\mathcal{B}(p, y)$, we deduce that:

$$\begin{aligned} V(p, y) &\geq \mathbb{E} \left[\int_0^\infty r e^{-rt} u(\hat{\pi}_t, Y_t) dt \right] \\ &= \mathbb{E} \left[\int_0^h r e^{-rt} u(\pi_t, Y_t) dt \right] + e^{-rh} \mathbb{E} \left[\mathbb{E} \left[\int_0^\infty r e^{-rt} u(\tilde{\pi}_{h+t}, Y_{h+t}) dt \mid \mathcal{F}_h^{(Y, \pi)} \right] \right] \\ &\geq \mathbb{E} \left[\int_0^h r e^{-rt} u(\pi_t, Y_t) dt \right] + e^{-rh} \mathbb{E} [V(\pi_h, Y_h)] \end{aligned}$$

which would prove that

$$V(p, y) \geq \max_{(Z, \pi) \in \mathcal{B}(p, y)} \mathbb{E} \left[\int_0^h r e^{-rt} u(\pi_t, Y_t) dt + e^{-rh} V(\pi_h, Y_h) \right]. \quad (4.4)$$

To conclude the proof of (4.4), we now check that the process $(Z, \hat{\pi})$ has a law in $\mathcal{B}(p, y)$.

At first, note that $(Z_t)_{t \geq 0}$ is a Markov process with initial law $p \otimes \delta_y$ by construction. Let us prove that for all $t \geq 0$, $\hat{\pi}_t(k) = \mathbb{P}(X_t = k | \mathcal{F}_t^{(Y, \hat{\pi})})$. The result is obvious by construction for $t < h$. For $t \geq h$, let us consider two finite families (t_1, \dots, t_m) in $[h, t]$ and (t'_1, \dots, t'_ℓ) in $[0, h)$ and two bounded continuous function ϕ, ϕ' defined on $(\Delta(K) \times \mathbb{R})^m$ and $(\Delta(K) \times \mathbb{R})^\ell$. Then:

$$\begin{aligned} & \mathbb{E}[\mathbb{1}_{X_t=k} \phi(\hat{\pi}_{t_1}, Y_{t_1}, \dots, \hat{\pi}_{t_m}, Y_{t_m}) \phi'(\hat{\pi}_{t'_1}, Y_{t'_1}, \dots, \hat{\pi}_{t'_\ell}, Y_{t'_\ell})] \\ &= \mathbb{E}[\mathbb{E}[\mathbb{1}_{X_t=k} \phi(\tilde{\pi}_{t_1}, Y_{t_1}, \dots, \tilde{\pi}_{t_m}, Y_{t_m}) | \mathcal{F}_h^{(\pi, Y)}] \phi'(\pi_{t'_1}, Y_{t'_1}, \dots, \pi_{t'_\ell}, Y_{t'_\ell})] \\ &= \mathbb{E}[\tilde{\pi}_t(k) \phi(\tilde{\pi}_{t_1}, Y_{t_1}, \dots, \tilde{\pi}_{t_m}, Y_{t_m}) \phi'(\pi_{t'_1}, Y_{t'_1}, \dots, \pi_{t'_\ell}, Y_{t'_\ell})] \\ &= \mathbb{E}[\hat{\pi}_t(k) \phi(\hat{\pi}_{t_1}, Y_{t_1}, \dots, \hat{\pi}_{t_m}, Y_{t_m}) \phi'(\hat{\pi}_{t'_1}, Y_{t'_1}, \dots, \hat{\pi}_{t'_\ell}, Y_{t'_\ell})]. \end{aligned}$$

This property extends to bounded measurable functions of any finite family (t_i) in $[0, t]$ by monotone class and we deduce that $\hat{\pi}_t(k) = \mathbb{P}(X_t = k | \mathcal{F}_t^{(Y, \hat{\pi})})$.

We now prove the Markov property. We consider only the case $s \geq t \geq h$, as the other cases can be studied by similar arguments. Given two finite families (t_1, \dots, t_m) in $[h, s]$ and (t'_1, \dots, t'_ℓ) in $[0, h)$ and three bounded continuous functions g, ϕ, ϕ' defined on $\Delta(K) \times \mathbb{R}$, $(\Delta(K) \times \mathbb{R})^m$ and $(\Delta(K) \times \mathbb{R})^\ell$ respectively, we deduce as above that

$$\begin{aligned} & \mathbb{E}[g(Z_t) \phi(\hat{\pi}_{t_1}, Y_{t_1}, \dots, \hat{\pi}_{t_m}, Y_{t_m}) \phi'(\hat{\pi}_{t'_1}, Y_{t'_1}, \dots, \hat{\pi}_{t'_\ell}, Y_{t'_\ell})] \\ &= \mathbb{E}[\mathbb{E}[g(Z_t) \phi(\tilde{\pi}_{t_1}, Y_{t_1}, \dots, \tilde{\pi}_{t_m}, Y_{t_m}) | \mathcal{F}_h^{(\pi, Y)}] \phi'(\pi_{t'_1}, Y_{t'_1}, \dots, \pi_{t'_\ell}, Y_{t'_\ell})] \\ &= \mathbb{E}[P_{t-s} g(Z_s) \phi(\hat{\pi}_{t_1}, Y_{t_1}, \dots, \hat{\pi}_{t_m}, Y_{t_m}) \phi'(\hat{\pi}_{t'_1}, Y_{t'_1}, \dots, \hat{\pi}_{t'_\ell}, Y_{t'_\ell})]. \end{aligned}$$

This property extends to bounded measurable functions of any finite family (t_i) in $[0, s]$ by monotone class and we deduce that $P_{t-s} g(Z_s) = \mathbb{E}[g(Z_t) | \mathcal{F}_t^{(Y, \hat{\pi})}]$. This concludes the proof of (4.4).

In order to conclude the proof of (4.1), we now prove the reverse inequality.

Let (Z, π) be an admissible process and $h > 0$. We check easily that the conditional law of $(Z_{h+s}, \pi_{h+s})_{s \geq 0}$ given $\mathcal{F}_h^{(\pi, Y)}$ belongs almost surely to $\mathcal{B}(\pi_h, Y_h)$. It follows that

$$\begin{aligned} \mathbb{E} \left[\int_0^\infty r e^{-rt} u(\pi_t, Y_t) dt \right] &= \mathbb{E} \left[\int_0^h r e^{-rt} u(\pi_t, Y_t) dt \right] + e^{-rh} \mathbb{E} \left[\mathbb{E} \left[\int_0^\infty r e^{-rt} u(\pi_{h+t}, Y_{h+t}) dt | \mathcal{F}_h^{(\pi, Y)} \right] \right] \\ &\leq \mathbb{E} \left[\int_0^h r e^{-rt} u(\pi_t, Y_t) dt \right] + e^{-rh} \mathbb{E} [V(\pi_h, Y_h)]. \end{aligned}$$

The conclusion follows by taking the supremum over all admissible processes (Z, π) .

The inequality (4.2) follows directly from (4.1). Precisely, given a process Z with initial law $p \otimes \delta_y$, define π by $\pi_t(k) := \chi_t(k) = \mathbb{P}[X_t = k | \mathcal{F}_t^{(Y, +)}]$ (optional projection). As explained before, (Z, π) has a law in $\mathcal{B}(p, y)$ and is a diffusion process of semi-group Q .

We finally prove (4.3). If $(Z, \pi) \in \mathcal{B}(p, y)$ is an optimal process (existence of a maximum follows from Theorem 4 in [26], see also the footnote in Lemma 3.8), then using the same

arguments as above, we have for all $h \geq 0$:

$$V(p, y) = \mathbb{E}\left[\int_0^\infty r e^{-rt} u(\pi_t, Y_t) dt\right] \leq \mathbb{E}\left[\int_0^h r e^{-rt} u(\pi_t, Y_t) dt\right] + e^{-rh} \mathbb{E}[V(\pi_h, Y_h)],$$

and the conclusion follows from (4.1). \square

4.2 Proof of Theorem 2.6.

Proof of theorem 2.6. The proof is divided in two parts showing respectively that the lower semicontinuous envelope V_* of V is subsolution and that V is supersolution of (2.4). Uniqueness and continuity will follow from the comparison result (Theorem A.9) whose proof is postponed to the appendix.

part 1: We prove that the lower semicontinuous envelope of V , denoted V_* , is a supersolution of (2.4).

Let ϕ be any smooth test function such that $\phi \leq V_*$ with equality in $(p, y) \in \Delta(K) \times \mathbb{R}$. As V_* is bounded, we may assume without loss of generality that ϕ is bounded. Consider a sequence $(p_n, y_n) \rightarrow (p, y)$ such that $V(p_n, y_n) \rightarrow V_*(p, y)$. From (4.4), we deduce that

$$V(p_n, y_n) - e^{-rh} \mathbb{E}_{p_n, y_n}[\phi(\psi_h)] - \mathbb{E}_{p_n, y_n}\left[\int_0^h r e^{-rs} u(\psi_s) ds\right] \geq 0.$$

Letting $n \rightarrow \infty$, we obtain that (recall that ψ is a Feller process):

$$\phi(p, y) - e^{-rh} \mathbb{E}_{p, y}[\phi(\psi_h)] - \mathbb{E}_{p, y}\left[\int_0^h r e^{-rs} u(\psi_s) ds\right] \geq 0.$$

Applying Itô's formula, we have

$$\mathbb{E}_{p, y}[\phi(\psi_h)] = \phi(p, y) + \mathbb{E}_{p, y}\left[\int_0^h A(\phi)(\psi_s) ds\right].$$

Dividing by h and letting then $h \rightarrow 0$, it follows from usual arguments that

$$r\phi(p, y) - A(\phi)(p, y) - ru(p, y) \geq 0. \quad (4.5)$$

Let us prove that V_* is concave with respect to p . Let $y \in \mathbb{R}$ and $p = \lambda p_1 + (1 - \lambda)p_2$ for some $p, p_1, p_2 \in \Delta(K)$ and $\lambda \in [0, 1]$. Let (p^n, y^n) a sequence converging to (p, y) such that $V(p^n, y^n) \rightarrow V_*(p, y)$. Then, there exists $p_1^n, p_2^n \in \Delta(K)$ such that $p^n = \lambda p_1^n + (1 - \lambda)p_2^n$ and $(p_1^n, p_2^n) \rightarrow (p_1, p_2)$ (it is for example a consequence of Lemma 8.2 in [23]). It follows that

$$V(p^n, y^n) \geq \lambda V(p_1^n, y^n) + (1 - \lambda)V(p_2^n, y^n).$$

By letting $n \rightarrow \infty$ and using the definition of V_* , we deduce that

$$V_*(p, y) \geq \lambda V_*(p_1, y) + (1 - \lambda)V_*(p_2, y),$$

which proves that V_* is concave. We deduce that $\lambda_{\max}(p, D^2 \phi_p(p, y)) \leq 0$, and together with (4.5) this concludes the proof of the supersolution property.

part 2: We prove that V is subsolution of (2.4).

Let ϕ be smooth test function such that $\phi \geq V$ with equality at $\bar{z} = (\bar{p}, \bar{y})$. We have to prove that if $\lambda_{\max}(\bar{p}, D_{\bar{p}}^2\phi(\bar{z})) < 0$, then $rV(\bar{z}) - A(\phi)(\bar{z}) - ru(\bar{z}) \leq 0$.

Using Proposition 4.3, let $(Z, \pi) \in \mathcal{B}(\bar{z})$ be an optimal process, so that for all $h \geq 0$, we have

$$V(\bar{z}) = \mathbb{E}\left[\int_0^h re^{-rt}u(\pi_s, Y_s)ds + e^{-rh}V(\pi_h, Y_h)\right]. \quad (4.6)$$

Since $\lambda_{\max}(\bar{p}, D_{\bar{p}}^2\phi(\bar{z})) < 0$ (see e.g. the proof of Theorem 3.3. in [5]), there exists $\delta > 0$ such that for all $z = (p, \bar{y})$ with $p \in \Delta(K)$ such that $p - \bar{p} \in T_{\Delta(K)}(\bar{p})$, we have:

$$V(z) \leq V(\bar{z}) + \langle D_p\phi(\bar{z}), p - \bar{p} \rangle - \delta|p - \bar{p}|^2.$$

As $\mathbb{E}[\pi_0] = \bar{p}$, the variable π_0 belongs almost surely to the smallest face of $\Delta(K)$ containing \bar{p} so that $\pi_0 - \bar{p} \in T_{\Delta(K)}(\bar{p})$. On the other hand, $Y_0 = \bar{y}$ so that (4.6) with $h = 0$ implies

$$V(\bar{z}) = \mathbb{E}[V(\pi_0, \bar{y})] \leq V(\bar{z}) - \delta\mathbb{E}[|\pi_0 - \bar{p}|^2].$$

We deduce that $\pi_0 = \bar{p}$ almost surely.

Recall the definition of the process χ as an optional projection:

$$\forall k \in K, \forall s \geq 0, \chi_s(k) = \mathbb{P}(X_s = k | \mathcal{F}_s^{Y, +}).$$

Lemma A.7 implies that $\pi_s(k) = \mathbb{P}(X_s = k | \mathcal{F}_s^{(\pi, Y), +})$, and we deduce that $\mathbb{E}[\pi_s | \mathcal{F}_s^{Y, +}] = \chi_s$ using the tower property of conditional expectations. Coming back to (4.6), Jensen's inequality implies:

$$V(\bar{z}) = \mathbb{E}\left[\int_0^h re^{-rt}u(\pi_s, Y_s)ds + e^{-rh}V(\pi_h, Y_h)\right] \leq \mathbb{E}\left[\int_0^h re^{-rt}u(\pi_s, Y_s)ds + e^{-rh}V(\chi_h, Y_h)\right].$$

Since $V \leq \phi$, we obtain

$$V(\bar{z}) = \phi(\bar{z}) \leq \mathbb{E}\left[\int_0^h re^{-rt}u(\pi_s, Y_s)ds + e^{-rh}\phi(\chi_h, Y_h)\right].$$

Dividing the above inequality by h , and letting h go to zero, it follows from the usual arguments (using that $\pi_s \rightarrow \pi_0$ when $s \rightarrow 0$, and Itô's formula) that:

$$rV(\bar{z}) - A(\phi)(\bar{z}) - ru(\bar{z}) \leq 0.$$

□

A Technical Proofs and auxiliary tools.

A.1 Proofs of Lemma 3.3

Let us now recall some properties of conditional independence. As we will manipulate conditional laws, we introduce a specific notation in order to shorten statements and proofs.

Notation A.1. Let E be a Polish space and A be an E -valued random variable defined on some probability space $(\Omega, \mathcal{A}, \mathbb{P})$.

- $\llbracket A \rrbracket$ denotes the law of A .

- Given a σ -field $\mathcal{F} \subset \mathcal{A}$, $\llbracket A \mid \mathcal{F} \rrbracket$ denotes a version of the conditional law of A given \mathcal{F} , hence an \mathcal{F} -measurable random variable with values in $\Delta(E)$ (see e.g. [2] Proposition 7.26 for this last point).

Lemma A.2. *Let E be a Polish space and A, B be E -valued random variables defined on $(\Omega, \mathcal{A}, \mathbb{P})$. Let $\mathcal{F} \subset \mathcal{G}$ be two sub σ -fields of \mathcal{A} .*

- If B is independent of (A, \mathcal{F}) , then $\llbracket A \mid \mathcal{F} \rrbracket = \llbracket A \mid \mathcal{F}, B \rrbracket$.
- If $\llbracket A \mid \mathcal{F} \rrbracket = \llbracket A \mid \mathcal{G} \rrbracket$, then for any σ -field \mathcal{H} such that $\mathcal{F} \subset \mathcal{H} \subset \mathcal{G}$, we have

$$\llbracket A \mid \mathcal{F} \rrbracket = \llbracket A \mid \mathcal{H} \rrbracket = \llbracket A \mid \mathcal{G} \rrbracket$$

Lemma A.3.

- Let A, B, C be three random variables (with values in some Polish space) defined on the same probability space. A is independent of B conditionally on C if and only if $\llbracket B \mid C \rrbracket = \llbracket B \mid C, A \rrbracket$.
- $A \amalg_C B$ if and only if there exists (on a possibly enlarged probability space) a random variable ξ uniform on $[0, 1]$ independent of (A, C) , and a measurable function f such that $B = f(C, \xi)$.

Proof. See Proposition 5.6 and 5.13 in [22]. □

Proof of Lemma 3.3. The “if” part is obvious. Let us prove the “only if” part. For $q = 0$, this is just Lemma A.3. However, we need to be more precise on how to construct this variable. We assume that there exists a family of independent variables $(\zeta_0, \dots, \zeta_n)$ uniformly distributed on $[0, 1]$ and independent of $(A_0, B_0, \dots, A_n, B_n)$. Then, the variable ξ_0 given by Lemma A.3 can be constructed as a function of (A_0, B_0, ζ_0) (see the proof of Proposition 5.13 in [22]). Let us now proceed by induction and assume the above property is true for $p \leq q$ and that ξ_p is measurable with respect to $(A_0, B_0, \zeta_0, \dots, A_p, B_p, \zeta_p)$. Since

$$(A_0, \dots, A_{q+1}) \amalg_{(B_0, \dots, B_{q+1})} (B_0, \dots, B_n),$$

we have

$$\llbracket B_0, \dots, B_n \mid B_0, \dots, B_{q+1}, A_0, \dots, A_{q+1} \rrbracket = \llbracket B_0, \dots, B_n \mid B_0, \dots, B_{q+1} \rrbracket.$$

Using the second point of Lemma A.2, we deduce that

$$\llbracket B_0, \dots, B_n \mid B_0, \dots, B_{q+1}, A_{q+1} \rrbracket = \llbracket B_0, \dots, B_n \mid B_0, \dots, B_{q+1} \rrbracket.$$

Using now the induction hypothesis and the first point of Lemma A.2, we also have

$$\begin{aligned} \llbracket B_0, \dots, B_n \mid B_0, \dots, B_{q+1}, \xi_0, \dots, \xi_q \rrbracket &= \llbracket B_0, \dots, B_n \mid B_0, \dots, B_{q+1} \rrbracket, \\ \llbracket B_0, \dots, B_n \mid B_0, \dots, B_{q+1}, \xi_0, \dots, \xi_q, A_{q+1} \rrbracket &= \llbracket B_0, \dots, B_n \mid B_0, \dots, B_{q+1}, A_{q+1} \rrbracket. \end{aligned}$$

Finally, we deduce that $A_{q+1} \amalg_{(\xi_0, \dots, \xi_q, B_0, \dots, B_{q+1})} (B_0, \dots, B_n)$ and the result follows then by applying Lemma A.3. □

A.2 Auxiliary Tools

The following lemma is classical.

Lemma A.4. *Let E, E' be two separable metric spaces and A, A' two tight (resp. closed, convex) subsets of $\Delta(E)$ and $\Delta(E')$. Then the set $\mathcal{P}(A, A')$ of probabilities on $E \times E'$ having marginals in the sets A and A' is itself tight (resp. closed, convex).*

Proof. Let us prove the tightness property. Let $\mu \in A$, $\nu \in A'$ and $\pi \in \mathcal{P}(\mu, \nu)$. By assumption, for any $\varepsilon > 0$ there is a compact K_ε of E , independent of the choice of μ in A , such that $\mu(E/K_\varepsilon) \leq \varepsilon$, and a compact K'_ε , independent of the choice of ν in A' such that $\nu(E'/K'_\varepsilon) \leq \varepsilon$. Then for any pair of random variables (U, V) of law π :

$$\mathbb{P}[(U, V) \notin K_\varepsilon \times L_\varepsilon] \leq \mathbb{P}[U \notin K_\varepsilon] + \mathbb{P}[V \notin L_\varepsilon] \leq 2\varepsilon$$

The closed and convex properties follow directly from the continuity and linearity of the application mapping π to its marginals. \square

The following theorem is well-known and allows to construct variables with prescribed conditional laws.

Theorem A.5. *(Blackwell-Dubins [3])*

Let E be a polish space with $\Delta(E)$ the set of Borel probabilities on E , and $([0, 1], \mathcal{B}([0, 1]), \lambda)$ the unit interval equipped with Lebesgue's measure. There exists a measurable mapping

$$\Phi : \Delta(E) \times [0, 1] \longrightarrow E$$

such that for all $\mu \in \Delta(E)$, the law of $\Phi(\mu, U)$ is μ where U is the canonical element in $[0, 1]$.

In the proof of Proposition 4.3, we use indirectly this result together with the a disintegration theorem. Precisely:

Lemma A.6. *Let E, F be Polish spaces, $(\Omega, \mathcal{A}, \mathbb{P})$ be some probability space, Y be an E -valued random variable defined on Ω , and \mathcal{F} a sub- σ -field of \mathcal{A} . Assume that f is an \mathcal{F} -measurable map from Ω to $\Delta(E \times F)$ such that the marginal $f_1(\omega) \in \Delta(E)$ of $f(\omega)$ on the first coordinate is a version of the conditional law of Y given \mathcal{F} . Then, (up to enlarging the probability space, there exists a random variable Z such that $f(\omega)$ is a version of the conditional law of (Y, Z) given \mathcal{F} .*

Proof. Up to enlarging the probability space, we may assume that there exists some random variable U uniformly distributed on $[0, 1]$ and independent of (Y, \mathcal{F}) . One can define using Theorem A.5 a variable $(\tilde{Y}, \tilde{Z}) = \Phi(f(\omega), U)$ having the property that $f_1(\omega)$ is a version of the conditional law of \tilde{Y} given \mathcal{F} . Let $g(\omega, \tilde{Y})$ be a version of the conditional law of \tilde{Z} given (\mathcal{F}, \tilde{Y}) , it follows easily that $Z = \Phi(g(\omega, \tilde{Y}), U)$ fulfills the required properties. \square

The next Lemma is a generalized martingale backward convergence theorem directly adapted from the corresponding result for classical forward martingales that can be found in chapter III of [25].

Lemma A.7. *Let X_n be an uniformly bounded sequence of real-valued random variables defined on some probability space $(\Omega, \mathcal{A}, \mathbb{P})$. Let $(\mathcal{F}_n)_{n \geq 0}$ be a decreasing sequence of sub σ -fields of \mathcal{A} . Assume that X_n converges almost surely to some variable X , then $\mathbb{E}[X_n | \mathcal{F}_n]$ converges almost surely to $\mathbb{E}[X | \bigcap_{n \geq 0} \mathcal{F}_n]$.*

Proof. Define $X_n^+ = \sup_{m \geq n} X_m$ and $Y_n^+ = \mathbb{E}[X_n^+ | \mathcal{F}_n]$. The sequence X_n^+ is non-increasing with limit X and we have

$$Y_{n+1}^+ = \mathbb{E}[X_{n+1}^+ | \mathcal{F}_{n+1}] \leq \mathbb{E}[X_n^+ | \mathcal{F}_{n+1}] = \mathbb{E}[Y_n^+ | \mathcal{F}_{n+1}].$$

Y_n^+ is therefore a backward sub-martingale and converges almost surely to some variable Y^+ (see e.g. Theorem 30 p.24 in [13]) which is $\bigcap_{n \geq 0} \mathcal{F}_n$ -measurable. Similarly, define $X_n^- = \inf_{m \geq n} X_m$, and $Y_n^- = \mathbb{E}[X_n^- | \mathcal{F}_n]$. Then Y_n^- is a backward supermartingale which converges almost surely to Y^- . To conclude, note that

$$Y_n^- \leq \mathbb{E}[X | \mathcal{F}_n] \leq Y_n^+,$$

and that $\mathbb{E}[Y_n^+ - Y_n^-] = \mathbb{E}[X_n^+ - X_n^-] \rightarrow 0$ by bounded convergence. Since $\mathbb{E}[X | \mathcal{F}_n]$ converges almost surely to $\mathbb{E}[X | \bigcap_{n \geq 0} \mathcal{F}_n]$, we deduce that $\mathbb{E}[X_n | \mathcal{F}_n]$ converges almost surely to $\mathbb{E}[X | \bigcap_{n \geq 0} \mathcal{F}_n]$ as $Y_n^- \leq \mathbb{E}[X_n | \mathcal{F}_n] \leq Y_n^+$. \square

A.3 comparison

In this section we adapt the comparison principle given in [9] for super solutions and sub solutions of (2.4).

Remark A.8. *Note that the process χ takes values in $\Delta(K)$, and that our assumptions on b and σ imply that the functions c and κ are Lipschitz continuous and bounded on $\Delta(K) \times \mathbb{R}$. In the following, we will assume without loss of generality that the functions c and κ are bounded and Lipschitz on the whole space \mathbb{R}^{K+1} (the explicit formula cannot be used directly since the resulting functions would be unbounded and only locally Lipschitz). Similarly, we assume that the function u is bounded and Lipschitz on the whole space \mathbb{R}^{K+1} .*

With our assumptions on c and κ , it is well known (see e.g. [11], p.19) that there exists a constant C (depending on the Lipschitz constants of c, κ, u) such that for any $\eta > 0$, $z, z' \in \Delta(K) \times \mathbb{R}$, $\xi \in \mathbb{R}^{K+1}$ and symmetric matrices $S, S' \in \mathcal{S}^{K+1}$ with

$$\begin{pmatrix} S & 0 \\ 0 & S' \end{pmatrix} \leq \eta \begin{pmatrix} I & -I \\ -I & I \end{pmatrix},$$

we have

$$\begin{aligned} |u(z) - u(z')| &\leq C|z - z'| \\ |\langle b(z), \xi \rangle - \langle b(z'), \xi \rangle| &\leq C|\xi||z - z'|, \\ -\frac{1}{2}\langle \kappa(z'), -S'\kappa(z') \rangle &\leq -\frac{1}{2}\langle \kappa(z), S\kappa(z) \rangle + C\eta|z - z'|^2. \end{aligned}$$

Let us state the comparison principle.

Theorem A.9. *Let w_1 be a subsolution and w_2 be a supersolution of (2.4), then $w_1 \leq w_2$.*

The rest of this subsection is devoted to the proof of this result. Let w_1 be a subsolution and w_2 be a supersolution of (2.4) (recall that w_1, w_2 are bounded functions). Our aim is to show that $w_1 \leq w_2$. We argue by contradiction, and assume that

$$M := \sup_{z \in \Delta(K) \times \mathbb{R}} \{w_1(z) - w_2(z)\} > 0. \quad (\text{A.1})$$

Because of the lack of compactness, let $\beta > 0$ and $g(y) := \sqrt{(1+y^2)}$. Define

$$M' := \sup_{z \in \Delta(K) \times \mathbb{R}} \{w_1(z) - w_2(z) - 2\beta g(y)\}.$$

We will choose β sufficiently small so that $M' > \frac{2C_1\beta}{r} > 0$ with $C_1 = \|\kappa\|_\infty^2 + \|b\|_\infty$.

We first regularize the maps w_1 and w_2 by quadratic sup and inf-convolution respectively. This technique is classical (see [11] for details), for $\delta > 0$ and $z \in \mathbb{R}^{K+1}$ we define:

$$w_1^\delta(z) := \max_{z' \in \Delta(K) \times \mathbb{R}} \left\{ w_1(z') - \frac{1}{2\delta} |z - z'|^2 \right\}$$

and

$$w_{2,\delta}(z) := \min_{z' \in \Delta(K) \times \mathbb{R}} \left\{ w_2(z') + \frac{1}{2\delta} |z - z'|^2 \right\}.$$

Note that w_1^δ and $w_{2,\delta}$ are defined on the whole space \mathbb{R}^{K+1} and that w_1^δ is semiconvex while $w_{2,\delta}$ is semiconcave. Moreover, we have the following growth property (uniformly in y)

$$\lim_{|p| \rightarrow +\infty} |p|^{-1} w_1^\delta(p, y) = -\infty, \quad \lim_{|p| \rightarrow +\infty} |p|^{-1} w_{2,\delta}(p, y) = +\infty.$$

Define (with $z^i = (p^i, y^i)$):

$$M_\delta := \sup_{z^1, z^2 \in \mathbb{R}^{K+1}} \left\{ w_1^\delta(z^1) - w_{2,\delta}(z^2) - \beta(g(y^1) + g(y^2)) - \frac{1}{2\delta} |z^1 - z^2|^2 \right\}. \quad (\text{A.2})$$

We have the classical result.

Lemma A.10. *For any $\delta > 0$, the problem (A.2) has at least one maximum point. If (z_δ^1, z_δ^2) is such a maximum point and if $(z_\delta^1)' \in \Delta(K) \times \mathbb{R}$ and $(z_\delta^2)'' \in \Delta(K) \times \mathbb{R}$ are such that*

$$w_1^\delta(z_\delta^1) = w_1((z_\delta^1)'_\delta) - \frac{1}{2\delta} |z_\delta^1 - (z_\delta^1)'_\delta|^2 \quad \text{and} \quad w_{2,\delta}(z_\delta^2) = w_2((z_\delta^2)''_\delta) + \frac{1}{2\delta} |z_\delta^2 - (z_\delta^2)''_\delta|^2 \quad (\text{A.3})$$

then, as $\delta \rightarrow 0$, $M_\delta \rightarrow M'$ while

$$\frac{|z_\delta^1 - z_\delta^2|^2}{2\delta} + \frac{|z_\delta^1 - (z_\delta^1)'_\delta|^2}{2\delta} + \frac{|z_\delta^2 - (z_\delta^2)''_\delta|^2}{2\delta} \rightarrow 0.$$

We first prove that the regularized sub/supersolutions are sub/supersolutions of slightly modified equations.

Lemma A.11. *Assume that w_1^δ has a second order Taylor expansion at a point z . Then*

$$\min\{rw_1(z) + H(z', Dw_1^\delta(z), D^2w_1^\delta(z)); -\lambda_{\max}(p', D_p^2w_1^\delta(z))\} \leq 0, \quad (\text{A.4})$$

where $z' \in \Delta(K) \times \mathbb{R}$ is such that $w_1^\delta(z) = w_1(z') - \frac{1}{2\delta} |z - z'|^2$.

Similarly, if $w_{2,\delta}$ has a second order Taylor expansion at a point z , then

$$rw_2(z) + H(z'', Dw_{2,\delta}(z), D^2w_{2,\delta}(z)) \geq 0, \quad (\text{A.5})$$

where $z'' \in \Delta(K) \times \mathbb{R}$ is such that $w_{2,\delta}(z) = w_2(z'') + \frac{1}{2\delta} |z - z''|^2$.

Proof. We do the proof for w_1^δ , the second part being similar. Assume that w_1^δ has a second order Taylor expansion at a point \bar{z} and set, for $\gamma > 0$ small,

$$\phi_\gamma(z) := \langle Dw_1^\delta(\bar{z}), z - \bar{z} \rangle + \frac{1}{2} \langle z - \bar{z}, D^2 w_1^\delta(\bar{z})(z - \bar{z}) \rangle + \frac{\gamma}{2} |z - \bar{z}|^2.$$

Let \bar{z}' denote a point in $\Delta(K) \times \mathbb{R}$ such that $w_1^\delta(\bar{z}) = w_1(\bar{z}') - \frac{1}{2\delta} |\bar{z} - \bar{z}'|^2$. Then $w_1^\delta - \phi_\gamma$ has a maximum at \bar{z} , which implies, by definition of w_1^δ , that

$$w_1(z') - \frac{1}{2\delta} |z' - z|^2 \leq \phi_\gamma(z) - \phi_\gamma(\bar{z}) + w_1^\delta(\bar{z}) \quad \forall z \in \mathbb{R}^{K+1}, \forall z' \in \Delta(K) \times \mathbb{R},$$

with an equality for $(z, z') = (\bar{z}, \bar{z}')$. If we choose $z = z' - \bar{z}' + \bar{z}$ in the above formula, we obtain:

$$w_1(z') \leq \phi_\gamma(z' - \bar{z}' + \bar{z}) + \frac{1}{2\delta} |\bar{z}' - \bar{z}|^2 - \phi_\gamma(\bar{z}) + w_1^\delta(\bar{z}) \quad \forall z' \in \Delta(K) \times \mathbb{R},$$

with an equality at $z' = \bar{z}'$. As w_1 is a subsolution, we obtain therefore, using the right-hand side of the above inequality as a test function,

$$\min \{rw_1(\bar{z}') + H(\bar{z}', D\phi_\gamma(\bar{z}), D^2\phi_\gamma(\bar{z})); -\lambda_{\max}(\bar{z}', D_p^2\phi_\gamma(\bar{z}))\} \leq 0. \quad (\text{A.6})$$

By construction, we have $D\phi_\gamma(\bar{z}) = Dw_1^\delta(\bar{z})$, $D^2\phi_\gamma(\bar{z}) = D^2w_1^\delta(\bar{z}) + \gamma I$ and $w_1(\bar{z}') \geq w_1^\delta(\bar{z})$. The conclusion follows therefore by letting $\gamma \rightarrow 0$. \square

In order to use inequality (A.4), we have to produce points at which w_1^δ is strictly concave with respect to the first variable. For this reason, as in [9], we introduce a additional penalization. For $\sigma > 0$ and $z^i = (p^i, y^i) \in \mathbb{R}^{K+1}$, we consider

$$M_{\delta,\sigma} := \sup_{(z^1, z^2) \in (\mathbb{R}^{K+1})^2} \left\{ w_1^\delta(z^1) - w_{2,\delta}(z^2) - \beta(g(y^1) + g(y^2)) + \sigma g(|p^1|) - \frac{1}{2\delta} |z^1 - z^2|^2 \right\}.$$

One easily checks that there exists a maximizer (\hat{z}^1, \hat{z}^2) to the above problem. In order to use Jensen's Lemma (Lemma A.3 in [11]), we also need this maximum to be strict. For this we modify the penalization: we set for $i = 1, 2$:

$$\xi_1(p^1) = g(|p^1|) - \sigma g(|p^1 - \hat{p}^1|), \quad \zeta_i(y^i) = -\beta g(y^i) - \sigma g(y^i - \hat{y}^i).$$

We choose $\sigma > 0$ sufficiently small so that ξ_1 has a positive second order derivative. By definition,

$$M_{\delta,\sigma} = \sup_{(z^1, z^2) \in (\mathbb{R}^{K+1})^2} \left\{ w_1^\delta(z^1) - w_{2,\delta}(z^2) + \zeta_1(y^1) + \zeta_2(y^2) + \sigma \xi(|p^1|) - \frac{1}{2\delta} |z^1 - z^2|^2 \right\},$$

and the above problem has a strict maximum at (\hat{z}^1, \hat{z}^2) . As the map $(z^1, z^2) \rightarrow w_1^\delta(z^1) - w_{2,\delta}(z^2) + \zeta_1(y^1) + \zeta_2(y^2) + \sigma \xi_1(p^1) - \frac{1}{2\delta} |z^1 - z^2|^2$ is semiconcave, Jensen's Lemma (together with Alexandrov theorem) states that, for any $\varepsilon > 0$, there is vector $a_\varepsilon \in (\mathbb{R}^{K+1})^2$ with $|a_\varepsilon| \leq \varepsilon$, such that the problem

$$M_{\delta,\sigma,\varepsilon} := \sup_{z^1, z^2 \in (\mathbb{R}^{K+1})^2} \left\{ w_1^\delta(z^1) - w_{2,\delta}(z^2) + \zeta_1(y^1) + \zeta_2(y^2) + \sigma \xi(|p^1|) - \frac{1}{2\delta} |z^1 - z^2|^2 + \langle a_\varepsilon, (z^1, z^2) \rangle \right\},$$

has a maximum point $(z_{\delta,\sigma,\varepsilon}^1, z_{\delta,\sigma,\varepsilon}^2) \in (\mathbb{R}^{K+1})^2$ at which the maps w_1^δ and $w_{2,\delta}$ have a second order Taylor expansion. From Lemma A.11, we have

$$\min \{rw_1(z_{\delta,\sigma,\varepsilon}^1) + H((z^1)'_{\delta,\sigma,\varepsilon}, Dw_1^\delta(z_{\delta,\sigma,\varepsilon}^1), D^2w_1^\delta(z_{\delta,\sigma,\varepsilon}^1)) ; -\lambda_{\max}((z^1)'_{\delta,\sigma,\varepsilon}, D_p^2w_1^\delta(z_{\delta,\sigma,\varepsilon}^1))\} \leq 0, \quad (\text{A.7})$$

and

$$rw_2(z_{\delta,\sigma,\varepsilon}^2) + H((z^2)''_{\delta,\sigma,\varepsilon}, Dw_{2,\delta}(z_{\delta,\sigma,\varepsilon}^2), D^2w_{2,\delta}(z_{\delta,\sigma,\varepsilon}^2)) \geq 0, \quad (\text{A.8})$$

where $(z^1)'_{\delta,\sigma,\varepsilon}$ and $(z^2)''_{\delta,\sigma,\varepsilon}$ are points in $\Delta(K) \times \mathbb{R}$ at which one has

$$w_1^\delta(z_{\delta,\sigma,\varepsilon}^1) = w_1((z^1)'_{\delta,\sigma,\varepsilon}) - \frac{1}{2\delta} |z_{\delta,\sigma,\varepsilon}^1 - (z^1)'_{\delta,\sigma,\varepsilon}|^2 \text{ and } w_{2,\delta}(z_{\delta,\sigma,\varepsilon}^2) = w_2((z^2)''_{\delta,\sigma,\varepsilon}) + \frac{1}{2\delta} |z_{\delta,\sigma,\varepsilon}^2 - (z^2)''_{\delta,\sigma,\varepsilon}|^2.$$

Using the properties of inf and sup-convolutions, we have:

$$Dw_1^\delta(z_{\delta,\sigma,\varepsilon}^1) = -\frac{1}{\delta} (z_{\delta,\sigma,\varepsilon}^1 - (z^1)'_{\delta,\sigma,\varepsilon}) \text{ and } Dw_{2,\delta}(z_{\delta,\sigma,\varepsilon}^2) = \frac{1}{\delta} (z_{\delta,\sigma,\varepsilon}^2 - (z^2)''_{\delta,\sigma,\varepsilon}). \quad (\text{A.9})$$

By definition of $M_{\delta,\sigma,\varepsilon}$ we have for all $(z^1, z^2) \in (\mathbb{R}^{K+1})^2$:

$$w_1^\delta(z^1) - w_{2,\delta}(z^2) + \zeta_1(y^1) + \zeta_2(y^2) + \sigma\xi_1(p_1) \leq M_{\delta,\sigma,\varepsilon} + \frac{1}{2\delta} |z^1 - z^2|^2 - \langle a_\varepsilon, (z^1, z^2) \rangle,$$

with an equality at $(z_{\delta,\sigma,\varepsilon}^1, z_{\delta,\sigma,\varepsilon}^2)$. Hence

$$Dw_1^\delta(z_{\delta,\sigma,\varepsilon}^1) + \begin{pmatrix} \sigma D\xi_1(p_{\delta,\sigma,\varepsilon}^1) \\ -\beta g'(y_{\delta,\sigma,\varepsilon}^1) - \sigma g'(y_{\delta,\sigma,\varepsilon}^1 - \hat{y}^1) \end{pmatrix} = \frac{1}{\delta} (z_{\delta,\sigma,\varepsilon}^1 - z_{\delta,\sigma,\varepsilon}^2) - a_\varepsilon^1 \quad (\text{A.10})$$

$$-Dw_{2,\delta}(z_{\delta,\sigma,\varepsilon}^2) + \begin{pmatrix} 0 \\ -\beta g'(y_{\delta,\sigma,\varepsilon}^2) - \sigma g'(y_{\delta,\sigma,\varepsilon}^2 - \hat{y}^1) \end{pmatrix} = \frac{1}{\delta} (z_{\delta,\sigma,\varepsilon}^2 - z_{\delta,\sigma,\varepsilon}^1) - a_\varepsilon^2 \quad (\text{A.11})$$

while

$$\begin{pmatrix} S & 0 \\ 0 & S' \end{pmatrix} \leq \frac{1}{\delta} \begin{pmatrix} I & -I \\ -I & I \end{pmatrix} \quad (\text{A.12})$$

with

$$S := D^2w_1^\delta(z_{\delta,\sigma,\varepsilon}^1) + \begin{pmatrix} \sigma D^2\xi_1(p_{\delta,\sigma,\varepsilon}^1) & 0 \\ 0 & -\beta g''(y_{\delta,\sigma,\varepsilon}^1) - \sigma g''(y_{\delta,\sigma,\varepsilon}^1 - \hat{y}^1) \end{pmatrix}$$

$$S' := -D^2w_{2,\delta}(z_{\delta,\sigma,\varepsilon}^2) + \begin{pmatrix} 0 & 0 \\ 0 & -\beta g''(y_{\delta,\sigma,\varepsilon}^2) - \sigma g''(y_{\delta,\sigma,\varepsilon}^2 - \hat{y}^1) \end{pmatrix}$$

This implies that $S \leq -S'$ (see [11] p.19) and therefore

$$D_p^2w_1^\delta(z_{\delta,\sigma,\varepsilon}^1) \leq D^2w_{2,\delta}(z_{\delta,\sigma,\varepsilon}^2) - D^2\xi_1(p_{\delta,\sigma,\varepsilon}^1). \quad (\text{A.13})$$

We now check that $\lambda_{\max}(((p^1)'_{\delta,\sigma,\varepsilon}), D_p^2w_1^\delta(z_{\delta,\sigma,\varepsilon}^1)) < 0$. Using the definition of $w_{2,\delta}$, for all $p \in \mathbb{R}^K$ and $p'' \in \Delta(K)$,

$$w_{2,\delta}(p, y_{\delta,\sigma,\varepsilon}^2) \leq w_2(p'', (y^2)''_{\delta,\sigma,\varepsilon}) + \frac{1}{2\delta} (|p - p''|^2 + |y_{\delta,\sigma,\varepsilon}^2 - (y^2)''_{\delta,\sigma,\varepsilon}|^2),$$

with an equality at $(p, p'') = (p_{\delta,\sigma,\varepsilon}^2, (p^2)''_{\delta,\sigma,\varepsilon})$. If $m \in T_{\Delta(K)}((p^2)''_{\delta,\sigma,\varepsilon})$ with $|m|$ small enough, taking $p := p_{\delta,\sigma,\varepsilon}^2 + m$ and $p'' = (p^2)''_{\delta,\sigma,\varepsilon} + m$ gives

$$w_{2,\delta}(p_{\delta,\sigma,\varepsilon}^2 + m, y_{\delta,\sigma,\varepsilon}^2) \leq w_2((p^2)''_{\delta,\sigma,\varepsilon} + m, (y^2)''_{\delta,\sigma,\varepsilon}) + \frac{1}{2\delta} (|p_{\delta,\sigma,\varepsilon}^2 - (p^2)''_{\delta,\sigma,\varepsilon}|^2 + |y_{\delta,\sigma,\varepsilon}^2 - (y^2)''_{\delta,\sigma,\varepsilon}|^2), \quad (\text{A.14})$$

with equality for $m = 0$. As w_2 is concave with respect to the first variable (see e.g. Lemma 3.2 in [5]), the above inequality implies that $\lambda_{\max}((p^2)''_{\delta,\sigma,\varepsilon}, D_p^2 w_{2,\delta}(z_{\delta,\sigma,\varepsilon}^2)) \leq 0$. In view of (A.13) we get therefore

$$\lambda_{\max}((p^1)'_{\delta,\sigma,\varepsilon}, D_p^2 w_1^\delta(z_{\delta,\sigma,\varepsilon}^1)) \leq -\sigma \lambda_{\min}((p^1)'_{\delta,\sigma,\varepsilon}, D^2 \xi_1(p_{\delta,\sigma,\varepsilon}^1)) < 0,$$

because $D^2 \xi_1 > 0$ by construction. So (A.7) becomes

$$r w_1(z_{\delta,\sigma,\varepsilon}^1) + H((z^1)'_{\delta,\sigma,\varepsilon}, D w_1^\delta(z_{\delta,\sigma,\varepsilon}^1), D^2 w_1^\delta(z_{\delta,\sigma,\varepsilon}^1)) \leq 0 \quad (\text{A.15})$$

We compute the difference of the two inequalities (A.15) and (A.8) above:

$$\begin{aligned} r(w_1^\delta(z_{\delta,\sigma,\varepsilon}^1) - w_{2,\delta}(z_{\delta,\sigma,\varepsilon}^2)) + H((z^1)'_{\delta,\sigma,\varepsilon}, D w_1^\delta(z_{\delta,\sigma,\varepsilon}^1), D^2 w_1^\delta(z_{\delta,\sigma,\varepsilon}^1)) \\ - H((z^2)''_{\delta,\sigma,\varepsilon}, D w_{2,\delta}(z_{\delta,\sigma,\varepsilon}^2), D^2 w_{2,\delta}(z_{\delta,\sigma,\varepsilon}^2)) \leq 0, \end{aligned}$$

where, in view of (A.9) and the definitions of S, S' (and using that g', g'' and $|D\xi_1|, |D^2\xi_1|$ are bounded by 1)

$$H((z^1)'_{\delta,\sigma,\varepsilon}, D w_1^\delta(z_{\delta,\sigma,\varepsilon}^1), D^2 w_1^\delta(z_{\delta,\sigma,\varepsilon}^1)) \geq H((z^1)'_{\delta,\sigma,\varepsilon}, \frac{1}{\delta}(z_{\delta,\sigma,\varepsilon}^1 - z_{\delta,\sigma,\varepsilon}^2), S) - C_1(\beta + \varepsilon + \sigma),$$

$$H((z^2)''_{\delta,\sigma,\varepsilon}, D w_{2,\delta}(z_{\delta,\sigma,\varepsilon}^2), D^2 w_{2,\delta}(z_{\delta,\sigma,\varepsilon}^2)) \leq H((z^2)''_{\delta,\sigma,\varepsilon}, \frac{1}{\delta}(z_{\delta,\sigma,\varepsilon}^1 - z_{\delta,\sigma,\varepsilon}^2), -S') + C_1(\beta + \varepsilon + \sigma),$$

Next, we have:

$$|u((z^1)'_{\delta,\sigma,\varepsilon}) - u((z^2)''_{\delta,\sigma,\varepsilon})| \leq C |(z^1)'_{\delta,\sigma,\varepsilon} - (z^2)''_{\delta,\sigma,\varepsilon}|,$$

$$|\langle b((z^1)'_{\delta,\sigma,\varepsilon}), \frac{1}{\delta}(z_{\delta,\sigma,\varepsilon}^1 - z_{\delta,\sigma,\varepsilon}^2) \rangle - \langle b((z^2)''_{\delta,\sigma,\varepsilon}), \frac{1}{\delta}(z_{\delta,\sigma,\varepsilon}^1 - z_{\delta,\sigma,\varepsilon}^2) \rangle| \leq \frac{C}{\delta} |(z^1)'_{\delta,\sigma,\varepsilon} - (z^2)''_{\delta,\sigma,\varepsilon}| |z_{\delta,\sigma,\varepsilon}^1 - z_{\delta,\sigma,\varepsilon}^2|,$$

$$-\frac{1}{2} \langle \kappa((z^2)''_{\delta,\sigma,\varepsilon}), -S' \kappa((z^2)''_{\delta,\sigma,\varepsilon}) \rangle \leq -\frac{1}{2} \langle \kappa((z^1)'_{\delta,\sigma,\varepsilon}), S \kappa((z^1)'_{\delta,\sigma,\varepsilon}) \rangle + \frac{C}{\delta} |(z^1)'_{\delta,\sigma,\varepsilon} - (z^2)''_{\delta,\sigma,\varepsilon}|^2.$$

We deduce that:

$$\begin{aligned} r(w_1^\delta(p_{\delta,\sigma,\varepsilon}) - w_{2,\delta}(p_{\delta,\sigma,\varepsilon})) \leq C \left(\frac{1}{\delta} |(z^1)'_{\delta,\sigma,\varepsilon} - (z^2)''_{\delta,\sigma,\varepsilon}|^2 + |(z^1)'_{\delta,\sigma,\varepsilon} - (z^2)''_{\delta,\sigma,\varepsilon}| \right) \\ + \frac{C}{\delta} |(z^1)'_{\delta,\sigma,\varepsilon} - (z^2)''_{\delta,\sigma,\varepsilon}| |z_{\delta,\sigma,\varepsilon}^1 - z_{\delta,\sigma,\varepsilon}^2| + 2C_1(\beta + \varepsilon + \sigma). \end{aligned}$$

As σ and ε tend to 0, the $z_{\delta,\sigma,\varepsilon}^1, z_{\delta,\sigma,\varepsilon}^2, (z^1)'_{\delta,\sigma,\varepsilon}$ and $(z^2)''_{\delta,\sigma,\varepsilon}$ converges (up to a subsequence) to $z_\delta^1, z_\delta^2, (z^1)'_\delta$ and $(z^2)''_\delta$, where (z_δ^1, z_δ^2) is a maximum in (A.2) and where $(z^1)'_\delta$ and $(z^2)''_\delta$ satisfy (A.3). The above inequality together with the definition of M_δ implies:

$$\begin{aligned} r M_\delta \leq r(w_1^\delta(z_\delta^1) - w_{2,\delta}(z_\delta^2)) \leq C \left(\frac{1}{\delta} |(z^1)'_\delta - (z^2)''_\delta|^2 + |(z^1)'_\delta - (z^2)''_\delta| \right) \\ + \frac{C}{\delta} |(z^1)'_\delta - (z^2)''_\delta| |z_\delta^1 - z_\delta^2| + 2C_1\beta. \end{aligned}$$

We finally let $\delta \rightarrow 0$: in view of Lemma A.10 the above inequality yields to $r M' = \lim_{\delta \rightarrow 0} r M_\delta \leq 2C_1\beta$, which contradicts our initial assumption. Therefore $w_1 \leq w_2$ and the proof is complete.

References

- [1] R. J. Aumann and M. Maschler, *Repeated Games with Incomplete Information*, with the collaboration of R. Stearns, Cambridge, MA: MIT Press, 1995.
- [2] D.P. Bertsekas and S.E. Shreve, *Stochastic optimal control: The discrete time case*, Academic Press, 1978.
- [3] D. Blackwell and L.E. Dubins, *An extension of Skorohod's almost sure representation theorem*, Proceedings of the American Mathematical Society, 1983, **89**, 691–692.
- [4] P. Cardaliaguet, *Differential Games with Asymmetric Information*, SIAM J. Control Optim., 2007, **46**, pp. 816–838.
- [5] P. Cardaliaguet, *A double obstacle problem arising in differential game theory*, Journal of Mathematical Analysis and Applications, 2009, **360**, pp. 95–107.
- [6] P. Cardaliaguet and C. Rainer, *Stochastic Differential Games with Asymmetric Information*, Applied Mathematics and Optimization, 2009, **59**, pp. 1–36.
- [7] P. Cardaliaguet and C. Rainer, *On a Continuous-Time Game with Incomplete Information*, Math. of Oper. Res., 2009, **34**, pp. 769–794.
- [8] P. Cardaliaguet and C. Rainer, *Games with incomplete information in continuous time and for continuous types*, Dynamic Games and Applications, 2012, **2**, pp. 206–227.
- [9] P. Cardaliaguet, C. Rainer, D. rosenberg, N. Vieille, *Markov games with frequent actions and incomplete information*, 2013, Preprint, [Arxiv-1307.3365](https://arxiv.org/abs/1307.3365)
- [10] P. Cardaliaguet, R. Laraki and S. Sorin, *A Continuous Time Approach for the Asymptotic Value in Two-Person Zero-Sum Repeated Games*, SIAM J. Control Optim., 2012, **50**, pp. 1573–1596.
- [11] M.G. Crandall, H. Ishii, and P. L. Lions, *User's guide to viscosity solutions of second order partial differential equations*, Bulletin of the American Mathematical Society, 1992, **27**, pp.1–67.
- [12] B. De Meyer, *Price dynamics on a stock market with asymmetric information*, Games and Econ. Behav., 2010, **69**, pp. 42–71.
- [13] C. Dellacherie, J.-P. Meyer, *Probabilities and potential*, B. North-Holland Mathematics Studies, 1982.
- [14] R. M. Dudley, *Real analysis and probability*, Cambridge University Press, 2002.
- [15] S.N. Ethier and T.G. Kurtz, *Markov processes: characterization and Convergence*, John Wiley & Sons Inc., 1986.
- [16] F. Gensbittel, *Extensions of the Cav(u) theorem for repeated games with one-sided information*, to appear in Math of Oper. Res.
- [17] F. Gensbittel, *Covariance control problems over martingale arising from game theory*, SIAM J. Control Optim., 2013, **51**, 1152–1185.

- [18] F. Gensbittel and C. Grün, *Zero-sum stopping games with asymmetric information*, Preprint, [Arxiv-1412.1412](https://arxiv.org/abs/1412.1412).
- [19] C. Grün, *On Dynkin games with incomplete information*, SIAM J. Control Optim., 2013, **51**, pp. 4039–4065.
- [20] C. Grün, *A BSDE approach to stochastic differential games with incomplete information*, Stochastic Processes and their Applications, 2012, **122**, pp. 1917–1946.
- [21] J. Jacod and A.N. Shiryaev, *Limit theorems for stochastic processes, Second Edition*, Springer-Verlag, 2003.
- [22] O. Kallenberg, *Foundations of modern probability*, Springer Verlag, 2002.
- [23] R. Laraki, *On the Regularity of the Convexification Operator on a Compact Set*, Journal of Convex Analysis, 2004, **11**, 209-234.
- [24] R.S. Lipster and A.N. Shiryaev, *Statistics of random Processes. I: General theory. Second edition.*, Springer-Verlag, 2001.
- [25] J. F. Mertens, S. Sorin and S. Zamir, *Repeated Games*, CORE Discussion Papers 9420, 9421 and 9422, Université Catholique De Louvain, Belgium, 1994.
- [26] P.A. Meyer and W.A. Zheng, *Tightness criteria for laws of semimartingales*, Annales de l’Institut Henri Poincaré (B) Probability and Statistics, 1984, **20**, pp. 353–372.
- [27] A. Neyman, *Stochastic Games with Short-Stage Duration*, Dynamic Games and Applications, 2013, **3**, pp. 236–278.
- [28] S. Sorin, *A First Course on Zero-Sum Repeated Games*, Springer, 2002.
- [29] D.W. Stroock and S.R.S. Varadhan, *Multidimensional diffusion processes*, Springer, 1979.