

Chains of causalities of environmental impacts

Robert Joumard, Santiago Mancebo Quintana, Gerassimos Arapis, Tomasz Zacharz

▶ To cite this version:

Robert Joumard, Santiago Mancebo Quintana, Gerassimos Arapis, Tomasz Zacharz. Chains of causalities of environmental impacts. Towards the definition of a measurable environmentally sustainable transport, Feb 2008, Norway. pp.1-12. hal-00916735

HAL Id: hal-00916735

https://hal.science/hal-00916735

Submitted on 10 Dec 2013

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Chains of causalities of environmental impacts

Robert Journard¹, Santiago Mancebo Quintana², Gerassimos Arapis³ & Tomasz Zacharz⁴

Abstract

In order to prepare an encompassing ex-ante assessment of the impacts of the transport system on the environment by building impact indicators, the pressure-state-impact concept is widen into the concept of chain of causalities or process. A process is defined by an output of the transport system, a homogeneous series of physical, chemical, biological, psychological relationships between elements, and a final target. All the known today impacts of the transport system on the environment are described as a list of 43 chains of causalities. The knowledge of the relationships will be used to design indicators or to know what does and not does measure a given indicator.

Key-words: environmental impact, classification, process, chain of causalities, transport.

1. Introduction

To build tools for assessing the impacts on the environment of a transport system or sub-system asks for a definition of the impacts on the environment, defined by final targets and modifications of target. For instance, the final targets of the traffic safety are mainly the humans with death and injuries. To assess each impact, the best way should be to measure the impact itself, by counting or evaluating for instance the number of people injured or dead because the traffic system. But such counting can't be made only *ex-post* and does not give any indication on the causes of the impact, because the impact cannot be linked by a one-to-one relationship with the accidents: the accidents are not the only causality of human death and injuries: local air pollution, greenhouse effect, hazards, among others cause death and injuries. The account of death and injuries due to the accident needs to take into account the process of accident. It is especially easy in the field of traffic safety, much more complex for most of the impacts.

For an evaluation *ex-ante* or for looking for the causes of an *ex-post* evaluation, a clear and precise relationship has to be established with the transport system. Each process, each chain of causalities from the source to each final impact on the environment has to be described in detail: in terms of sources, intermediate and final targets, mechanisms between intermediate sources and intermediate targets. Such description allows us also to express clearly what a potential indicator measures and does not measure, and on which scientific mechanisms an indicator should be based. For instance the global warming potential evaluates the global temperature increase and not really the final impacts of greenhouse effect as sea level increase, the amount of fauna, flora and human habitat destruction, the food chain changes... The knowledge of the physical mechanism of the climate and temperature modifications as a function of greenhouse gas emissions allows to build the shape of the indicator 'global warming potential'.

At the same time, the description of the chains of causalities allows us to define quite precisely the term 'environment': What are the impacts on the environment? What are their characteristics or

¹ Inrets, France, journard@inrets.fr

² Univ. Politécnica de Madrid, Spain

³ Agricultural Univ. of Athens, Greece

⁴ Komag, Poland

typical features?

The most common presentation of the environment, especially by economists, considers it as a resource used by the humans for producing economic goods. This resource is an ecosystem, i.e. the association between a physicochemical and abiotic (the biotope) environment and a living community characteristic of the latter (the biocenosis), including fossil resources. This resource is destroyed but can be renewed at a given extend: the environmental issue is a question of resource flow and capacity of the biosphere to support the effects of the human activities (carrying capacity): It calls the 7th principle of the Rio declaration (UNCED, 1992): "...to conserve, protect and restore [...] the integrity of the Earth's ecosystem [...] the pressures their societies place on the global environment". The pressure-state-impact (PSI) system from OECD seems well applicable to this meaning with a pressure representing a flow.

In parallel, the environment is often understood as the quality of our physical environment or the quality of life: a calm area with pure air and pure water, a beautiful landscape... It calls the first principle of the Rio declaration: "Human beings [...] are entitled to a healthy and productive life in harmony with nature". It is here often difficult to consider only flows or pressures.

These both meanings of the environment correspond roughly to the external and internal territory sustainability by Wackernagel and Rees (1999): the internal sustainability consists in protecting its direct environment and living area, but the external sustainability consists in protecting the world.

According to USEPA (1996)	OECD (2002)	COST 350 (2002)	Goger (2006), Goger & Joumard (2007)	Journard & Nicolas (2007)	chain
All impacts from transport	All impacts from transport	All impacts from transport (red: global; black: local/regional)	Impacts from transport due to pollutant emissions (grey: out of the scope)	All impacts from transport (red: irreversible; black: reversible)	Corresponding chain in Table 4
	use of natural resources	non-renewable energy consumpt. use of material resources		non-renewable resource use, including energy	21
	climate change	climate change	Greenhouse effect	greenhouse effect	42
		ozone depletion	Ozone depletion		15, 16
pollutant emissions	air quality	air pollution	Direct restricted health effects Direct ecotoxicity	local air quality	3
			Photochemical pollution	regional air quality (smog)	17, 18, 19, 20
highway and airport runoff toxic release sewage dumping	water quality	water pollution	Eutrophication	water quality / uses and régime	5 24, 25, 26 8
release of deicing compounds		soil pollution			24, 25, 26
waste					23
direct waste from vehicles		waste production			34

noise	noise	noise nuisance/vibratio n		light and noise	41 29, 30, 31
				nuisances	14
		traffic accidents		traffic safety	22
roadkill, wildlife collisions					35
Habitat disruption and land take by	severance	barrier effects / land fragmentation		landuse	6, 7
infrastructure		land uptake			9, 11
		soil erosion			
		hydrologic/hydra ulic risks			27
			Sensitive		37
	visual impacts		pollution	visual qualities of	38, 39
				landscape/townsc ape	40
	historical / effe archaeological / aesth		Degradation of common man-made heritage	man-made	2, 12, 27
	nature conservation	cultural heritage	Degradation of historic man-made heritage	heritage	12
			Ţ	biodiversity and	9, 10, 13
	loss of biodiversity		(Direct ecotoxicity)	protected areas	
	acidification		Acidification		1, 2
hazardeous material incident				technological and natural hazards	28
introduction of non-native species					32
habitat disruption by wakes / anchors					33
					36
					43

Table 1: Correspondence between the environmental impacts as listed by 5 references, and the list of processes proposed in this paper.

2. Precise list of environmental impacts

Such definitions are much too global and rough to be useful for describing the environmental issue or the impact on the environment of a human activity as the transport system, and for designing environmental impact indicators. An exhaustive list of the chains of causalities is necessary to present a full picture. But the definition of the environmental or ecological impacts is neither clear nor precise in the literature. When lists of environmental impacts are given, they are often heterogeneous, merging impacts and sources: See some examples Table 1. For instance USEPA (1996) lists mainly the pressures or the first consequences of the transport system on the environment rather than environmental impacts (although designed as impacts). The use of natural resources, the hydrologic and hydraulic risks, the traffic safety and the final impacts as sensitive

pollution are missing; A contrario rarely mentioned impacts are listed as introduction of non-native species, habitat disruption by wakes or anchors, direct waste from vehicles, roadkill and wildlife collisions. The other references examined merge impacts on the environment as climate change or visual effects, and intermediate states of the environment as local air quality, water quality. Goger (2006) and Goger & Joumard (2007) give the most precise list but only due to atmospheric pollutant emissions: In this field, impacts are distinguished when they are due to different chains of causalities, taking into account the fact that, as stated in Wäger (2006), the impact categories shall together enable an encompassing assessment of relevant impacts, which are known today (completeness), but at the same time should have the least overlap as possible (independence).

3. The concept of chain of causalities

It is the reason why we prefer to enlarge the PSI picture to the concept of process or chain of causalities between a cause and a final impact, with possibly a succession of couples cause-impact. A good example is the greenhouse effect with the greenhouse gases (GHG) as a first cause, which by physical phenomenon increases the earth temperature, which modifies the global and local climates, with impacts on the agriculture, sea level, with impacts on all the biocenosis including the humans. If an initial pressure can be easily detected (GHG emissions), they are afterwards a lot of intermediate states and impacts. Another advantage of the concept of process or chain of causalities is to be much wider than a physical flow problem: any process can be taken into account, as cultural, psychological, psycho-physical, or biological effect.

A chain of causalities can be described through:

- The element(s) of the transport system, which is at the begin of the process, taking into account the life cycle approach, ie. considering all the activities involved. 3 main subsystems are involved (infrastructure, fuel, and vehicle), and for each of them 5 types of activities (production, existence, use, maintenance, and destruction). All together there are 13 subsystems-activities: See Table 2. The 13 subsystems can be simplified into 4, as coloured in Table 2 and used in Table 4, by considering the 3 main subsystems but extracting the traffic, i.e. the use of the infrastructure, final energy and vehicle.




Table 2: Typology of the main transport subsystems involved in the environmental impacts. Colours correspond to wider subsystems as used in Table 4.

- The final targets: Goger (2006) and Goger & Joumard (2007) consider 3 targets (nature, humans, man-made heritage) and a pseudo-target, the earth. In addition the Eco-indicator approach (Brand et al., 1998; Goedkoop & Spriemsma, 2000) includes three types of endpoint damages: resources, ecosystem quality, and human health. The 2 first are subdivisions of the target nature. The (human) health is defined by World Health Organisation (WHO, 1946) as "a state of complete physical, mental and social well-being and not merely the absence of disease or infirmity". Therefore it is useful to distinguish health in a restricted meaning (absence of disease or infirmity) and the complement so-called human well-being, because the processes are often

very different. Finally we get the target structure presented Table 3.

	Pseudo-target								
	Resources	Earth:							
	Ecosystems:	Covers all the							
Nature	Nature understood as ecosystems, i.e. the association	targets: the three							
Nature	between a physicochemical and abiotic (the biotope)	previous targets							
	environment and a living community characteristic of	(ecosystems,							
	the latter (the biocenosis)	humans and man							
Humans:	Human health:	made heritage)							
Humankind which we extract	In a restricted meaning	and physical							
from nature and focus on its	Human wall baing	environments							
health as defined by the WHO	Human well-being	such as the							
	Man-made heritage:								
With a distinction is a	With a distinction is made between common and historic buildings								

Table 3: Structure of the targets of the impacts on the environment.

- The in-between elements, i.e. the chain of causalities between the transport system and the final targets, to be described in detail. To design impact indicators, it is important to know the scientific milieu able to understand the process, and therefore to give the scientific disciplines involved. We propose a first and simple science structure: physics, chemistry, biology, psychology/sociology. It is important also to know if the process is linear or not, and if the transport system characteristics are major or minor explanation parameters, in order to know how these characteristics can be used for indicator building. Finally the reversibility is a major parameter from the sustainability point of view; The distance and time scales indicate who is concerned, if it is a local/global, shot/medium/long term impact.

It disaggregates the different impacts found in the literature in order to understand the complexity of the processes involved, to identify the related sciences and to estimate the order of magnitude of the impact in space and time.

4. Typology of chains of causalities

According to this structure, a typology of the chains of causalities of the environmental impacts of the transport system is proposed Table 4. 29 chains are distinguished, and 43 when taking into account differentiation in the last steps of the process corresponding to different final targets. The chains are independent and encompass all the relevant impacts found in the literature.

The description of the chains could be more detailed, by dividing a chain into two or more chains, if it is considered as not homogeneous in terms of process or targets. In addition some chains can be missing.

A contrario, the aggregation of impacts is possible when the knowledge necessary to build impact indicators is similar and if the main characteristics of the chain are similar. As, to be practical, the number of categories should amount to a not too high number, and considering the importance of each impact and the availability of indicators, some impacts could be merged, or minor chains be deleted. Because it is important to give the possibility to further users to perform such simplifications, the chain structure has to be as detailed as possible: It is easier to merge and delete than to add processes.

	Sou	rces										Tar	get	
Iinfrastructure	Energy	Vehicle	Traffic	First step of the chain (pressure)	Identificat ion	reversibilit y, distance and time scale from the source	scientific disciplines invo	Chain of causalities (states and processes) and <i>final impact</i> (main scientific disciplines involved: P: Physics; C: Chemistry: B: Biology; PS: Psychology / Sociology)				Health	Human well-being	Man-made heritage Earth
	*						(incl. secondary effect of photochemical pollution) Dispersion in the	dispersion in air, water and soil (P), ecotoxicity on fauna and flora (B). Decrease of ecosystem health, loss of biodiversity.	1	,	E S			
	*	*	Emissions	Acidificati on	Mm, year		deposition on surfaces (P), chemical reactions with materials (C). Loss of man-made heritage (PS), destruction of archaeological, classical or historic remains (P), loss of cultural legacy (PS).	2]	M	
	*	*	*	Emission of particles	Direct	km, day	Dispersion in the atmosphere and water (P),	direct ecotoxicity on fauna and flora (B).	3		E S			
			*	and air pollutants	toxicity	, ,	sometimes dispersion in food (P),	direct restricted health effects (B).	4			Н		
	*		* *	Emissions NOx	Eutrophic ation	10 km, year		nere and water (P), increase of plant oxia of fauna and flora (B).	5		E S			
*					Habitat	practically	S	(B). Loss of ecosystem health, loss of iodiversity.	6		E S			
*			*	Land take	fragmenta tion	irreversible , km, year	Cutting of the human habita	Cutting of the human habitat, reduction of living areas of people (B, PS)					H W B	
* * *				Land take	Hydraulic changes	km, year	Hydraulic changes, modification of fauna, mainly, and flora habitat (P, B).				E S			

* * *	*	*		Land take by infrastructur e building	Land take	practically irreversible , km, year	Waterproofing of areas, low waterproofing of areas, modification of our	cerease of ecosystems (P, B). Loss of iodiversity. coss of natural and wildlife protected areas. loss of available land for humans, tdoor recreation areas (PS).	9 10 11	R ?	E S E S	H W B	
								cal, classical or historic remains (P), <i>ultural legacy</i> (PS).	12				M
	*			Agriculture for biofuels	Biofuel agricultur e	km, year	Transformation of natural areas, disappearance of fauna and flora (B).			R ?	E S		
*			*	Emission of light	Light pollution	Mm, min		uminosity of the open space (P), ehaviour (B), effects on biota health.	14		E S		
				Emission of	Ozone		Dispersion in the atmosphere (P), chemical	ecotoxicity on fauna and flora (B).	15		E S		
		* halogen compounds	alogen depletion	earth, year	reaction (C) depletion of ozone layer, increase of UV on the earth (P),	health effects (B).	16		Н				
								health effects (B).	17		Н		
								loss of agriculture productivity (B).	18	R			
								ecotoxicity on fauna and flora (B).	19		E S		
*:	*	* *	Emission of NOx, NMVOC, CO.	Photoche mical pollution	Mm, day	Dispersion in the atmosphere (P), chemical reaction (C) and therefore increase of photochemical pollutants as ozone,	deposition on surfaces (P), chemical reactions with materials (C), loss of man-made heritage (PS), destruction of archaeological, classical or historic remains (P), loss of cultural legacy (PS).	20				M	
								Secondary effects: - greenhouse gas (see greenhouse effect) - acidification (see acidification)	-		(E S)		(M)
*	*	*	*	Non- renewable	Non- renewable	irreversible , Mm, 100	Decrease of metals, fossil	fuels availability for the future (P).	21	R			

			*	resource use	resource use	years						
			* *	Accidents	Traffic Safety	partially irreversible , m, -	Human death, injuries (B).	22		Н		
*	*	*		Waste disposal	Non- recyclable waste	Partially irrevers. (nuclear waste), all	Includes the nuclear waste. Dissemination in the nature (P), impacts on health and ecosystems (B).	23	E S	Н	H W B	
				Emission of			ecosystem health (B).	24	E S			
*			*	gazeous, liquid or	Soil and water pollution	100 km,	Dispersion in the soil and health effects (B).	25		Н		
				solid pollutants			year	water (P), recreational areas forbidden (PS).	26			H W B
* *				Risk of floods	Hydraulic risk	km, year	Destruction of natural and human habitat (P).	27	E S	Н		M
	*		*	Risk for industrial safety	Technolog ical hazards	km to earth, day to century	Industrial accidents, included of nuclear power plants. Dispersion in the atmosphere, soil and water (P), biological impacts on humans and biota (B).	28	E S	Н		
			*				Diffusion (P), disappearance of calm areas (PS).	29			H W B	
*			*	Emission of noise	Noise	km, hour	Diffusion in air, absorption or reflection by surfaces (P), annoyance for people (PS), health effects (B).	30		Н	H W B	
							Diffusion in air, absorption or reflection by surfaces (P), <i>ecosystem health</i> (B).	31	E S			
			*	Introduction of non- native species	Introducti on of non- native species	Earth, irreversible	Small individuals, seeds disperse and survive (B), modification of biocenosis. <i>Loss of biodiversity</i> .	32	E S			
			*	Emission of wakes	Habitat disruption	km	Microhabitat changes. Loss of biodiversity, loss of ecosystem health.	33	E S			

					by wakes / anchors								
			*	Emission of waste	Direct waste from vehicles	100 m, year	Waste thrown directly from the vehicles, accumulation. <i>Annoyance</i> (PS), especially if the landscape is of high quality.	34			H W B		
			*	Biota collision	Biota collision	partially irreversible , m, -	Fauna collision from small insects to big mammals or fish, damage by anchors. <i>Loss of biodiversity</i> (B).	35	E S				
			*	Risk of fire	Fire risk	10 km, year	Fire ignition by sparks, matches or accidents. <i>Destruction of natural and human habitat</i> (P).	36	E S	Н	H W B		
	*		* *	Emission of VOC	Odours	100 m, hour	Dispersion in the atmosphere (P) at short distance, sensitive pollution perceived by smell (PS).	37			H W B		
	*		*	Emission of PM	Soiling	100 m, year	Dispersion in the atmosphere (P) at short distance, deposition on surfaces (P), chemical reactions with materials (C), sensitive pollution perceived by the sight (PS).	38			H W B		
	*		*	Emission of PM and atmospheric pollutants	Visibility	100 m, day	Dispersion in the atmosphere (P) at mid distance, chemical reaction in air (C), sensitive pollution perceived by the sight (PS).	39			H W B		
* * *				Land use	Visual qualities of landscape/ townscape	practically irreversible , km, year	Infrastructure presence, <i>annoyance</i> (PS), especially if the landscape is of high quality.	40			H W B		
			*	Emission of vibration	Vibration	100 m, hour	Heavy traffic (HDV, trains) vibrations, mass diffusion, destruction of earth houses (P), annoyance (PS).	41			H W B	M	
*	*	*	* *	Emission of air pollutants	Greenhou se effect	irreversible , earth, century	Dispersion in the atmosphere (P), sometimes chemical reaction (C) and therefore creation of secondary pollutants, increase of the greenhouse effect (P), climate change (P), sea level increase (P), destruction or modification of habitat for fauna, flora and humans (P), change in food chain (B), economic losses (PS)	42					Е

	*	*	* * *	Emission of aerosols	Dimming	100 km and earth, day to month	Dispersion in the atmosphere (P), physical reactions (P) and sometimes chemical reactions (C), regional dimming, regional temperature decrease, global climate changes, <i>destruction or modification of habitat for fauna, flora and humans</i> (P), <i>change in food chain</i> (B), <i>economic losses</i> (PS)	43				E	3
--	---	---	-------	----------------------	---------	---	---	----	--	--	--	---	---

Table 4: Proposed list of the main chains of causalities of environmental impacts with some characteristics.

5. Conclusion

To describe the environmental impacts of an activity as transport through a complete list of independent chains of causalities allows us firstly to give a precise definition of the term 'environment'. In the literature, the differences in the impacts considered translate often the research area of the author, and, when the work is more global, the local perception of the environmental or ecological issue. For instance the loss of visibility above the cities, due to air pollution, is always cited in North America, but never in Europe, although the physical situations are similar. It is especially important to define the term environment, when today the environmental issue is taken into account by most of the transport specialists without precise knowledge of this field: In this case the environmental issue is very often reduced to greenhouse gases or to few well known impacts, or are reduced unconsciously to impacts for which simple to use assessment tools are available.

According to COST 356 (Journard, 2008), an indicator of environmentally sustainable transport is a variable, based on measurements, representing potential or actual impacts on the environment, or factors that may cause such impacts, due to transport systems, flows or policies, as accurately as possible and necessary. The precise description of the environmental processes constitutes then a powerful tool for indicator assessment, similar to that done by USEPA (1996). A priori, it can be stated that the more to the right the indicator is, the more precise the final impact is. It is mainly a tool to define what precisely an indicator does represent: Does it represent the final impact, or an intermediate one? How accurately is the process translated into the indicator function? Which relevant impacts are not taken into account by existing indicators? Isn't it possible double counting?

When the aim is to design new indicators of environmentally sustainable transport, the knowledge of the process indicates which scientists should be asked about the best way to represent the impact. It is also a comprehensive basis to study the social perception of the environmental issue by survey, whom outputs can be used to balance the quality of local air, of regional air, noise, greenhouse effect... according to the focus placed on each of these impacts, as made for the Personal Security Index designed by the Canadian Council on Social Development: See Tsoukalas & MacKenzie, 2003).

References

- Brand G., Braunschweig A., Scheidegger A. & Schwank O. (1998): Weighting in Ecobalances with the ecoscarcity method Ecofactors 1997. BUWAL (SAFEL) Environment Series, No. 297, Bern. www.e2mc.com/BUWAL297%20english.pdf
- COST 350 (2002): Integrated Assessment of Environmental Impact of Traffic and Transport Infrastructure, executive summary of final deliverable of Workin Group 1. 7 p. /www.rws.nl/rws/dww/home/cost350/
- Goedkoop M. & Spriemsma R. (2000): *The Eco-Indicator 99: A damage oriented method for Life Cycle Impact Assessment.* Methodology report. 2nd Edition, 214 p. www.pre.nl/eco-indicator99/ei99-reports.htm
- Goger T. (2006): Un indicateur d'impact environnemental global des polluants atmosphériques émis par les transports. Thèse, Insa Lyon, 28 nov. 2006, et rapport Inrets, n°LTE 0633, Bron, France, 283 p.
- Goger T. & Journard R. (2007): A method of building an aggregated indicator of air-pollution impacts. 3rd int. conf. *Sustainable development 2007*, 25-27 April 2007, Algarve, Portugal.
- Journard R. (2008): Definitions of indicator within the COST action 356 EST. Seminar COST 356 EST "Towards the definition of a measurable environmentally sustainable transport", 20 February 2008, Oslo, Norway.

- Joumard R. & Nicolas J.P. (2007): Méthodologie d'évaluation de projets de transport dans le cadre du développement durable. 12^e coll. int. *Évaluation environnementale et transports*: concepts, outils et méthodes, 18-22 juin 2007, Genève, Suisse.
- OECD (2002): *Impact of Transport Infrastructure Investment on Regional Development*. http://www.cemt.org/pub/pubpdf/JTRC/02RTRinvestE.pdf
- Tsoukalas S. & MacKenzie A. (2003): Personal Security Index 2003: A reflection of how Canadians feel five years later.CCSD, Ottawa, 87 p. www.ccsd.ca/pubs/2003/psi/
- UNCED (1992): Rio Declaration on Environment and Development. 3 p. www.unep.org/Documents.Multilingual/Default.asp?DocumentID=78&ArticleID=1163
- USEPA (1996): Indicators of environmental impacts of transportation Highway, Rail, aviation and maritime transport. USEPA report, 230-R-96-009, Washington, USA, 268 p.
- Wackernagel M. & Rees W. (1999): Notre empreinte écologique, Ecosociété, Paris, 207 p.
- Wäger P. (2006): Life Cycle Assessment a short overview. COST356 WG1 meeting, Madrid, 18-19 October.
- WHO (1946): Preamble to the Constitution of the World Health Organization as adopted by the International Health Conference, New York, 19-22 June, 1946; signed on 22 July 1946 by the representatives of 61 States (Official Records of the World Health Organization, no. 2, p. 100) and entered into force on 7 April 1948.