



# Local and global proper infiniteness for continuous $C(X)$ -algebras

Etienne Blanchard

## ► To cite this version:

Etienne Blanchard. Local and global proper infiniteness for continuous  $C(X)$ -algebras. 2013. hal-00922786v5

**HAL Id: hal-00922786**

**<https://hal.science/hal-00922786v5>**

Preprint submitted on 17 Jan 2014 (v5), last revised 19 Feb 2014 (v9)

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# LOCAL AND GLOBAL PROPER INFINITENESS FOR CONTINUOUS $C(X)$ -ALGEBRAS

ETIENNE BLANCHARD

**ABSTRACT.** All unital continuous  $C^*$ -bundles with properly infinite fibres are properly infinite  $C^*$ -algebras if and only if the full unital free product  $\mathcal{T}_2 *_\mathbb{C} \mathcal{T}_2$  of two copies of the Cuntz extensions  $\mathcal{T}_2$  generated by two isometries with orthogonal ranges is a  $K_1$ -injective  $C^*$ -algebra ([BRR08, Theorem 5.5], [Blan10, Proposition 4.2]). We show in this article that there is a state  $\psi_n$  with faithful GNS representation on the Cuntz extension  $\mathcal{T}_n$  such that the reduced unital free product  $(\mathcal{T}_n, \psi_n) *_\mathbb{C} (\mathcal{T}_n, \psi_n)$  is  $K_1$ -injective for all  $n \geq 3$ .

## 1. INTRODUCTION

The classification programme of nuclear  $C^*$ -algebras through K-theoretical invariants launched by G. Elliott ([Ell94]) led A. Toms and W. Winter to introduce the strong self-absorption property for simple unital  $C^*$ -algebras ([TW07]). This notion is a pretty rigid one: Any separable unital continuous  $C(X)$ -algebra  $A$  the fibres of which are isomorphic to the same strongly self-absorbing  $C^*$ -algebra  $D$  is a trivial  $C(X)$ -algebra provided the compact Hausdorff base space  $X$  has finite topological dimension. (The strongly self-absorbing  $C^*$ -algebra  $D$  tensorially absorbs the Jiang-Su algebra  $\mathcal{Z}$  ([Win09]). Hence, the  $C^*$ -algebra  $D$  is  $K_1$ -injective ([Ror04]) and the  $C(X)$ -algebra  $A$  satisfies  $A \cong D \otimes C(X)$  ([DW08]).) But I. Hirshberg, M. Rørdam and W. Winter have built a non-trivial unital continuous  $C^*$ -bundle over the infinite dimensional compact product  $\prod_{n=0}^\infty S^2$  such that all its fibres are isomorphic to the strongly self-absorbing UHF algebra of type  $2^\infty$  ([HRW07, Example 4.7]). More recently, M. Dădărlat has constructed in [Dad09, §3] for all pair  $(\Gamma_0, \Gamma_1)$  of discrete countable torsion groups a unital separable continuous  $C(X)$ -algebra  $A$  such that:

- the base space  $X$  is the compact Hilbert cube  $X = \mathfrak{X}$  of infinite dimension,
- all the fibres  $A_x$  ( $x \in \mathfrak{X}$ ) are isomorphic to the strongly self-absorbing Cuntz  $C^*$ -algebra  $\mathcal{O}_2$  generated by two isometries  $s_1, s_2$  satisfying  $1_{\mathcal{O}_2} = s_1 s_1^* + s_2 s_2^*$ ,
- $K_i(A) \cong C(Y_0, \Gamma_i)$  for  $i = 0, 1$ , where  $Y_0 \subset [0, 1]$  is the canonical Cantor set.

These K-theoretical conditions imply that the  $C(\mathfrak{X})$ -algebra  $A$  is not a trivial one. But this argument does not anymore work when the stringy self-absorbing algebra  $D$  is the Cuntz algebra  $\mathcal{O}_\infty$  ([Cun77]), in so far as  $K_0(\mathcal{O}_\infty) = \mathbb{Z}$  is a torsion free group.

In this note, we attack this trivialization problem for unital continuous  $C(X)$ -algebras with fibres  $\mathcal{O}_\infty$  in a different way: All unital continuous  $C(X)$ -algebras with properly infinite fibres are properly infinite  $C^*$ -algebras if and only if the full unital

---

2010 *Mathematics Subject Classification.* Primary: 46L80; Secondary: 46L06, 46L35.

*Key words and phrases.*  $C^*$ -algebra, Classification, Proper Infiniteness.

free product  $\mathbf{A} = \mathcal{T}_2 *_\mathbb{C} \mathcal{T}_2$  of two distinct copies of the unital extension  $\mathcal{T}_2$  of the Cuntz algebras  $\mathcal{O}_2$  by the compact operators ([Cun77]) is a  $K_1$ -injective  $C^*$ -algebra, *i.e.* the canonical map from  $\mathcal{U}(\mathbf{A})/\mathcal{U}^0(\mathbf{A})$  to  $K_1(\mathbf{A})$  is injective (Corollary 4.3). That free product  $\mathbf{A}$  unittally embeds in the free product  $\mathcal{T}_2 *_\mathbb{C} \mathcal{T}_3$  ([ADEL04]) and works by M. Rieffel ([Rief83], [Rief87]), K. Dykema, U. Haagerup, M. Rørdam ([DHR97], [Roh09]) enable us to prove that at least some reduced quotient  $\mathfrak{B}$  of  $\mathcal{T}_2 *_\mathbb{C} \mathcal{T}_3$  is a unital  $K_1$ -injective  $C^*$ -algebra which contains the algebraic unital free product  $\mathcal{T}_2 \otimes_\mathbb{C} \mathcal{T}_3$  of the two Cuntz extensions  $\mathcal{T}_2$  and  $\mathcal{T}_3$  (Proposition 5.2).

## 2. NOTATIONS

We present in this section the main notations which are used in this article.

**Definition 2.1.** ([Dix69], [Kas88], [Blan97]) *Let  $X$  be a compact Hausdorff space and let  $C(X)$  be the  $C^*$ -algebra of continuous function on  $X$ .*

- *A unital  $C(X)$ -algebra is a unital  $C^*$ -algebra  $A$  endowed with a unital morphism of  $C^*$ -algebra from  $C(X)$  to the centre of  $A$ .*
- *For all closed subset  $F \subset X$  and all  $a \in A$ , one denotes by  $a|_F$  the image of  $a$  in the quotient  $A|_F := A/C_0(X \setminus F) \cdot A$ . If  $x \in X$  is a point in  $X$ , one calls fibre at  $x$  the quotient  $A_x := A|_{\{x\}}$  and one write  $a_x$  for  $a|_{\{x\}}$ .*
- *The  $C(X)$ -algebra  $A$  is said to be continuous if the upper semicontinuous map  $x \in X \mapsto \|a_x\| \in \mathbb{R}_+$  is continuous for all  $a \in A$ .*

**Definition 2.2.** ([Pim95]) *Let  $X$  be a compact Hausdorff space.*

a) *The full Fock Hilbert  $C(X)$ -module  $\mathcal{F}(E)$  of a Hilbert  $C(X)$ -module  $E$  is the direct sum of Hilbert  $C(X)$ -module*

$$\mathcal{F}(E) := \bigoplus_{m \in \mathbb{N}} E^{(\otimes_{C(X)} m)}, \quad (2.1)$$

where  $E^{(\otimes_{C(X)} m)} := \begin{cases} E \otimes_{C(X)} \dots \otimes_{C(X)} E & (m \text{ terms}) \text{ if } m \geq 1, \\ C(X) & \text{if } m = 0. \end{cases}$

b) *The Pimsner-Toeplitz  $C(X)$ -algebra  $\mathcal{T}(E)$  of a full Hilbert  $C(X)$ -module  $E$ , *i.e.* with non-zero fibres, is the unital subalgebra of the  $C(X)$ -algebra  $\mathcal{L}_{C(X)}(\mathcal{F}(E))$  of adjointable  $C(X)$ -linear operator acting on  $\mathcal{F}(E)$  generated by the creation operators  $\ell(\zeta)$  ( $\zeta \in E$ ), where:*

$$\begin{aligned} - \ell(\zeta) (f \cdot \hat{1}_{C(X)}) &:= f \cdot \zeta = \zeta \cdot f & \text{for } f \in C(X) & \text{and} \\ - \ell(\zeta) (\zeta_1 \otimes \dots \otimes \zeta_k) &:= \zeta \otimes \zeta_1 \otimes \dots \otimes \zeta_k & \text{for } \zeta_1, \dots, \zeta_k \in E & \text{if } k \geq 1. \end{aligned} \quad (2.2)$$

**Remarks 2.3.** a) ([Cun81], [BRR08]) For all integer  $n \geq 2$ , the  $C^*$ -algebra  $\mathcal{T}_n := \mathcal{T}(\mathbb{C}^n)$  is the universal unital  $C^*$ -algebra generated by  $n$  isometries  $s_1, \dots, s_n$  satisfying the relation

$$s_1 s_1^* + \dots + s_n s_n^* \leq 1. \quad (2.3)$$

b) A unital  $C^*$ -algebra  $A$  is *properly infinite* if and only if one the following equivalent conditions holds ([Cun77], [Ror03, Proposition 2.1]):

- $A$  contains two isometries with mutually orthogonal range projections,

- $A$  contains a unital copy of the simple  $C^*$ -algebra  $\mathcal{O}_\infty$ .

### 3. LOCAL PROPER INFINITENESS

Let  $(C^*(\mathbb{Z}), \Delta)$  be the unital Hopf  $C^*$ -algebra ([Wor95]) generated by a unitary  $\mathbf{u}$  with spectrum  $\mathbb{T} := \{z \in \mathbb{C}; z^*z = 1\}$  (often written  $S^1$ ) and with coproduct  $\Delta(\mathbf{u}) = \mathbf{u} \otimes \mathbf{u}$ .

If  $X$  is a compact Hausdorff space and  $E$  is a separable Hilbert  $C(X)$ -module with non-zero fibres, there is only one coaction  $\alpha_E$  of  $(C^*(\mathbb{Z}), \Delta)$  on the Pimsner-Toeplitz  $C(X)$ -algebra  $\mathcal{T}(E)$  such that  $\alpha_E(\ell(\zeta)) = \ell(\zeta) \otimes \mathbf{u}$  for all  $\zeta \in E$ , *i.e.*

$$\begin{aligned} \mathcal{T}(E) &\rightarrow \mathcal{T}(E) \otimes C^*(\mathbb{Z}) &= C(\mathbb{T}, \mathcal{T}(E)) \\ \ell(\zeta) &\mapsto \ell(\zeta) \otimes \mathbf{u} &= (z \mapsto \ell(z\zeta)) \end{aligned} \quad (3.1)$$

Equation (3.1) implies that the fixed point  $C(X)$ -subalgebra  $\mathcal{T}_{C(X)}(E)^{\alpha_E}$  under this coaction is the closed linear span  $\mathcal{T}_{C(X)}(E)^{\alpha_E} = [C(X).1 + \sum_{k \geq 1} \ell(E)^k \cdot (\ell(E)^k)^*]$ .

Besides, the following local proper infiniteness property holds.

**Proposition 3.1.** ([Blac04]) *Let  $X$  be a compact Hausdorff space and let  $E$  be a separable Hilbert  $C(X)$ -module with infinite dimensional fibres.*

*If  $x$  is any point in  $X$ , then there is a closed neighbourhood  $F \subset X$  of  $x$  such that the restriction  $\mathcal{T}(E)|_F$  of the Pimsner-Toeplitz  $C(X)$ -algebra  $\mathcal{T}(E)$  is properly infinite, *i.e.* there is a unital  $*$ -homomorphism  $\mathcal{O}_\infty \rightarrow \mathcal{T}(E)|_F$ .*

*Proof.* Let  $\zeta \in E$  be a norm 1 section satisfying  $\|\zeta_y\| = 1$  for every point  $y$  in a closed neighbourhood  $F \subset X$  of the point  $x$ . Denote by  $\pi_F$  the quotient  $*$ -homomorphism  $C(X) \rightarrow C(X)/C_0(X \setminus F) \cong C(F)$ .

- (1) There is an isomorphism of Hilbert  $C(F)$ -module  $E|_F \cong C(F) \cdot \zeta \otimes_{\pi_F} 1 \oplus E'|_F$ , where  $E'$  is the orthocomplement of  $\zeta$  in the Hilbert  $C(X)$ -module  $E$ .
- (2) There is an isomorphism of  $C(F)$ -algebra

$$\mathcal{T}(E)|_F = C^* \langle \mathcal{T}(E)^{\alpha_E}|_F, \ell(\zeta)|_F \rangle \cong (\mathcal{T}(E)^{\alpha_E})|_F \rtimes \mathbb{N},$$

where the action of the semigroup  $\mathbb{N}$  on the  $C(F)$ -algebra  $\mathcal{T}(E)^{\alpha_E}|_F$  is given by  $a \mapsto \ell(\zeta)|_F \cdot a \cdot \ell(\zeta)|_F^*$ .

- (3) The restriction  $\mathcal{T}(E)|_F$  is properly infinite by [Blan13, Lemma 6.1].

□

*Remarks 3.2.* a) We can also prove Proposition 3.1 thanks to the semiprojectivity of the Cuntz  $C^*$ -algebra  $\mathcal{O}_\infty$  ([Blac04, Theorem 3.2]).

b) The restriction  $\mathcal{T}(E)|_F$  is properly infinite if the closed subset  $F \subset X$  is perfect (*i.e.* without any isolated point) and there is a section  $\zeta \in E$  with  $\|\zeta_x\| \geq 1$  for all points  $x \in F$ .

#### 4. GLOBAL PROPER INFINITENESS

Proposition 3.1 of the present paper, Proposition 2.5 of [BRR08] and section 6 of [Blan13] entail the following:

**Proposition 4.1.** *Let  $X$  be a second countable perfect compact Hausdorff space and let  $A$  be a separable unital continuous  $C(X)$ -algebra with properly infinite fibres.*

1) *There exist:*

- (a) *a finite integer  $n \geq 1$ ,*
- (b) *a covering  $X = \overset{\circ}{F}_1 \cup \dots \cup \overset{\circ}{F}_n$  by the interiors of closed balls  $F_1, \dots, F_n$ ,*
- (c) *unital embeddings of  $C^*$ -algebra  $\sigma_k : \mathcal{O}_\infty \rightarrow A|_{F_k}$  ( $1 \leq k \leq n$ ).*

2) *The tensor product  $M_p(\mathbb{C}) \otimes A$  is properly infinite for all large enough integers  $p$ .*

3) *For all  $i, j$  in  $\{1, \dots, n\}$ , there is a unitary  $u_{i,j} \in \mathcal{U}(A|_{F_i \cap F_j})$  such that*

$$\sigma_i(s_m)|_{F_i \cap F_j} = u_{i,j} \cdot \sigma_j(s_m)|_{F_i \cap F_j} \quad \text{for all } m \in \mathbb{N}.$$

*Proof.* 1) For all point  $x \in X$ , the semiprojectivity of the  $C^*$ -subalgebra  $\mathcal{O}_\infty \hookrightarrow A_x$  ([Blac04, Theorem 3.2]) implies that there are a closed neighbourhood  $F \subset X$  of the point  $x$  and a  $C(F)$ -linear unital embedding  $\mathcal{O}_\infty \otimes C(F) \hookrightarrow A|_F$ . The compactness of the topological space  $X$  enables to conclude.

2) derives from Proposition 3.1, [BRR08, Proposition 2.7] and [Ror97, Proposition 2.1].

3) One sets  $u_{i,j} := \sum_{m \in \mathbb{N}} \sigma_i(s_m)|_{F_i \cap F_j} \cdot \sigma_j(s_m)^*|_{F_i \cap F_j}$  ([Blan13, Proposition 6.3]).  $\square$

*Remark 4.2.* Assertion (2) alone does not imply that the  $C^*$ -algebra  $A$  is properly infinite: There exists a unital  $C^*$ -algebra  $A$  which is not properly infinite, but such that  $M_2(\mathbb{C}) \otimes A$  is properly infinite ([Ror03, Proposition 4.5]).

Note the following link with the  $K_1$ -injectivity for properly infinite  $C^*$ -algebras.

**Corollary 4.3.** *Let  $j_0, j_1$  denote the two canonical unital embeddings of the  $C^*$ -algebra  $\mathcal{T}_2$  in the full unital free product  $\mathbf{A} := \mathcal{T}_2 *_\mathbb{C} \mathcal{T}_2$  and let  $u \in \mathcal{U}(\mathcal{T}_2 *_\mathbb{C} \mathcal{T}_2)$  be a  $K_1$ -trivial unitary such that  $j_1(s_1) = u \cdot j_0(s_1)$  ([BRR08, Lemma 2.4]).*

1) *The following conditions are equivalent:*

- (a) *The full unital free product  $\mathcal{T}_2 *_\mathbb{C} \mathcal{T}_2$  is  $K_1$ -injective.*
- (b) *The unitary  $u \in \mathcal{U}(\mathcal{T}_2 *_\mathbb{C} \mathcal{T}_2)$  belongs to the connected component  $\mathcal{U}^0(\mathcal{T}_2 *_\mathbb{C} \mathcal{T}_2)$ .*
- (c) *Every separable unital continuous  $C(X)$ -algebra  $A$  with properly infinite fibres is a properly infinite  $C^*$ -algebra.*

2) *If the  $C^*$ -algebra  $\mathcal{T}_2 *_\mathbb{C} \mathcal{T}_2$  is  $K_1$ -injective, then the Pimsner-Toeplitz algebra  $\mathcal{T}(E)$  of any Hilbert  $C(X)$ -module  $E$  with fibres of dimension at least 2 is properly infinite and there is an isomorphism of  $C(X)$ -algebra  $\mathcal{T}(E) \cong \mathcal{T}(E)^{\alpha_E} \rtimes \mathbb{N}$ .*

*Proof.* 1) (a) $\Rightarrow$ (b) A unital  $C^*$ -algebra  $A$  is  $K_1$ -injective if and only if every unitary  $v \in \mathcal{U}(A)$  is homotopic to  $1_A$  in the group  $\mathcal{U}(A)$ . Hence, (b) is a special case of (a).

(b) $\Rightarrow$ (c) Take a finite covering  $X = \overset{o}{F}_1 \cup \dots \cup \overset{o}{F}_n$  of the compact space  $X$  such that there are unital embeddings  $\sigma_k : \mathcal{T}_2 \rightarrow A|_{F_k}$  ( $1 \leq k \leq n$ ). Set  $G_k := F_1 \cup \dots \cup F_k \subset X$  for all  $1 \leq k \leq n$  and let us construct by induction isometries  $w_k \in A|_{G_k}$  such that the two projections  $w_k w_k^*$  and  $1_{|G_k} - w_k w_k^*$  are properly infinite and full in  $A|_{G_k}$ :

- If  $k = 1$ , the isometry  $w_1 := \sigma_1(s_1)$  has the requested properties.
- If  $1 \leq k \leq n-1$  and the isometry  $w_k \in A|_{G_k}$  is already constructed, then Lemma 2.4 of [BRR08] implies that there exist a unital homomorphism of  $C^*$ -algebra  $\pi_k : \mathcal{T}_2 *_C \mathcal{T}_2 \rightarrow A|_{G_k \cap F_{k+1}}$  and a  $K_1$ -trivial unitary  $u_{k+1} \in \mathcal{U}(A|_{G_k \cap F_{k+1}})$  satisfying:

$$\begin{aligned} & - \pi_k(j_0(s_1)) = w_k|_{G_k \cap F_{k+1}} \text{ and} \\ & - \pi_k(j_1(s_1)) = \sigma_{k+1}(s_1)|_{G_k \cap F_{k+1}} = u_{k+1} \cdot w_k|_{G_k \cap F_{k+1}}. \end{aligned}$$

If condition (b) holds, the unitary  $u_{k+1}$  can even be taken in  $\mathcal{U}^0(A|_{G_k \cap F_{k+1}})$ , so that it admits a unitary lifting  $z_{k+1} \in \mathcal{U}^0(A|_{F_{k+1}})$  by Lemma 2.1.7 of [LLR00]. Let  $w_{k+1} \in A|_{G_{k+1}}$  be the only isometry satisfying the two constraints:

$$w_{k+1}|_{G_k} = w_k \quad \text{and} \quad w_{k+1}|_{F_{k+1}} = (z_{k+1})^* \cdot \sigma_{k+1}(s_1).$$

It verifies that the two projections  $w_{k+1} w_{k+1}^*$  and  $1_{|G_{k+1}} - w_{k+1} w_{k+1}^*$  are properly infinite and full in  $A|_{G_{k+1}}$ .

The proper infiniteness of the projection  $w_n w_n^* \in A|_{G_n} = A$  implies that the unit  $1_A = w_n^* \cdot w_n w_n^* \cdot w_n$  is also properly infinite in  $A$ , *i.e.*  $A$  is a properly infinite  $C^*$ -algebra.

(c) $\Rightarrow$ (a) The  $C^*$ -algebra  $\mathcal{D} := \{f \in C([0, 1], \mathcal{T}_2 *_C \mathcal{T}_2); f(0) \in j_0(\mathcal{T}_2) \text{ and } f(1) \in j_1(\mathcal{T}_2)\}$  is a unital continuous  $C([0, 1])$ -algebra the fibres of which are all properly infinite. Thus, condition (c) implies that  $\mathcal{D}$  is a properly infinite  $C^*$ -algebra, a statement equivalent to the  $K_1$ -injectivity of the full free product  $\mathcal{T}_2 *_C \mathcal{T}_2$  ([Blan10, Proposition 4.2]).

2) All the fibres of the  $C(X)$ -algebra  $\mathcal{T}(E)$  are simple and properly infinite. Thus, the  $C^*$ -algebra  $\mathcal{T}(E)$  is properly infinite as soon as condition (c) of assertion 1) holds. One can then construct by induction isometries  $w_k \in \mathcal{T}(E)|_{G_k}$  ( $1 \leq k \leq n$ ) such that:

- the projections  $w_k w_k^*$  and  $1_{|G_k} - w_k w_k^*$  are properly infinite and full in  $A|_{G_k}$ ,
- there is an isomorphism of  $C(G_k)$ -algebra  $\mathcal{T}(E)|_{G_k} \cong \mathcal{T}(E)^{\alpha_E}|_{G_k} \rtimes_{ad w_k} \mathbb{N}$ .

□

*Remarks 4.4.* a) The isometry  $w_n \in \mathcal{T}(E)$  built in assertion 2) cannot belong to the subspace  $\ell(E) \subset \mathcal{T}(E)$  in general. Indeed, if  $\mathfrak{X}$  is the compact Hilbert cube of infinite topological dimension and  $\mathcal{E}$  is the non-trivial Hilbert  $C(\mathfrak{X})$ -module constructed by J. Dixmier and A. Douady ([DD63], [BK04a, Proposition 3.6]), then there exist by Proposition 3.1 finitely many sections  $\zeta_1, \dots, \zeta_n$  in  $\mathcal{E}$  such that

$$\mathcal{T}(\mathcal{E}) = C^* \langle \mathcal{T}(\mathcal{E})^{\alpha_{\mathcal{E}}}, \ell(\zeta_1), \dots, \ell(\zeta_n) \rangle.$$

But there is no  $\zeta \in \sum_{1 \leq k \leq n} C(X) \cdot \zeta_k \subset \mathcal{E}$  such that  $\mathcal{T}(\mathcal{E}) = C^* \langle \mathcal{T}(\mathcal{E})^{\alpha_{\mathcal{E}}}, \ell(\zeta) \rangle$  since:

- any section  $\zeta \in \mathcal{E}$  satisfies  $\zeta_x = 0$  for at least one point  $x \in \mathfrak{X}$  ([DD63, 14], [BK04a, Proposition 3.6]) and
- $C^* \langle \mathcal{T}(\mathcal{E})^{\alpha_{\mathcal{E}}}, \ell(\zeta)_x \rangle \cong \mathcal{T}(\mathcal{E}_x)^{\alpha_{\mathcal{E}_x}} \subsetneq \mathcal{T}(\mathcal{E}_x) \cong \mathcal{O}_{\infty}$ .

b) If  $\mathfrak{A}$  denotes the universal UHF-algebra with  $K_1(\mathfrak{A}) = \mathbb{Q}$ , there is a non-trivial unital continuous  $C(\mathfrak{X})$ -algebra  $D$  with  $D_x \cong \mathcal{O}_\infty \otimes \mathfrak{A}$  for all  $x \in \mathfrak{X}$  ([Dad09]). But is there a non-trivial unital continuous  $C(\mathfrak{X})$ -algebra with fibres isomorphic to  $\mathcal{O}_\infty$ ? Note that  $K_0(\mathcal{O}_\infty \otimes \mathfrak{A}) = \mathbb{Q} \neq \mathbb{Z} = K_0(\mathcal{O}_\infty)$ .

## 5. A $K_1$ -INJECTIVE QUOTIENT

The universal Cuntz extension  $\mathcal{T}_2$  generated by two isometries with orthogonal ranges unittally embeds in the Cuntz extension  $\mathcal{T}_3$  ([Cun81, Lemma 3.1]). Accordingly, the full unital free product  $\mathcal{T}_2 *_\mathbb{C} \mathcal{T}_2$  unittally embeds in the free product  $\mathcal{T}_2 *_\mathbb{C} \mathcal{T}_3$  ([ADEL04, Proposition 2.4]). We show in this section that at least some quotient of  $\mathcal{T}_2 *_\mathbb{C} \mathcal{T}_3$  is a simple  $K_1$ -injective  $C^*$ -algebra which contains the algebraic unital free product  $\mathcal{T}_2 \otimes_\mathbb{C} \mathcal{T}_3$ .

Let  $\ell^2(\mathbb{N})$  be the Hilbert space of all sequences  $\xi = (\xi_k) \in \mathbb{C}^\mathbb{N}$  which satisfy  $\|\xi\|^2 = \sum_k |\xi_k|^2 < \infty$  and take an orthonormal basis  $\{e_0, e_1, e_2, \dots\}$  on  $\ell^2(\mathbb{N})$ . Define for all integer  $n \geq 2$ :

- the  $n$  isometries  $s_{n,1}, \dots, s_{n,n}$  in  $\mathbb{B}(\ell^2(\mathbb{N}))$  satisfying  $s_{n,i} \cdot e_k = e_{nk+i}$  ( $k \in \mathbb{N}$ ),
- the Cuntz extension  $\mathcal{T}_n := C^*(s_{n,1}, \dots, s_{n,n})$  ([Cun77]),
- the state  $\psi_n(a) = \frac{1}{n} \sum_{i=1}^n \langle e_i, a \cdot e_i \rangle$  on the  $C^*$ -algebra  $\mathcal{T}_n \subset \mathbb{B}(\ell^2(\mathbb{N}))$ ,
- the unitary  $u_n := s_{n,1} s_{n,2}^* + \dots + s_{n,n} s_{n,1}^* + p \in \mathcal{T}_n \subset \mathbb{B}(\ell^2(\mathbb{N}))$ , where  $p$  is the rank 1 projection  $p = 1 - \sum_i s_{n,i} s_{n,i}^* = \theta_{e_0, e_0}$  in  $\mathbb{B}(\ell^2(\mathbb{N}))$ .

**Lemma 5.1.** 1)  $\psi_n(a) = \psi_n(u_n^* a u_n)$  for any operator  $a \in \mathcal{T}_n$  and  $\psi_n((u_n)^j) = 0$  for any integer  $j \in \{1, \dots, n-1\}$ ,

2) The state  $\psi_n$  on  $\mathcal{T}_n$  has a faithful GNS representation.

3) The restriction of the state  $\psi_n$  to the  $C^*$ -subalgebra  $C^*(u_n) \subset \mathcal{T}_n$  is faithful.

*Proof.* 1) The unitary  $u_n$  satisfies  $(u_n)^n = 1$  and  $(u_n)^j \cdot e_1 = e_{n-j+1}$  for all  $1 \leq j \leq n$ . Thus, there exists an isomorphism  $\alpha_n : C^*(u_n) \rightarrow C^*(\mathbb{Z}/n\mathbb{Z})$  and

$$\begin{aligned} \psi_n(a) &= \frac{1}{n} \sum_{i=1}^n \langle (u_n)^i \cdot e_1, a (u_n)^i \cdot e_1 \rangle \\ &= \frac{1}{n} \sum_{i=1}^n \langle (u_n)^{i+1} \cdot e_1, a (u_n)^{i+1} \cdot e_1 \rangle = \psi_n(u_n^* a u_n). \end{aligned}$$

We also have  $\psi_n((u_n)^j) = \langle e_1, (u_n)^j \cdot e_1 \rangle = \langle e_1, e_{n-j+1} \rangle = 0$  for all  $j \in \{1, \dots, n-1\}$ .

2) The  $C^*$ -algebra  $\mathcal{T}_n$  is contained in  $\mathbb{B}(\ell^2(\mathbb{N}))$  and the Hilbert space  $\ell^2(\mathbb{N})$  is the closure of  $\mathcal{T}_n \cdot e_0$ . Thus, if  $a \in \mathcal{T}_n \setminus \{0\}$  is non-zero, there exists an element  $b \in \mathcal{T}_n$  such that

$$\begin{aligned} 0 < \|ab \cdot e_0\|^2 &= \|ab s_{n,1}^* \cdot e_1\|^2 = \langle e_1, s_{n,1} b^* \cdot a^* a \cdot b s_{n,1}^* e_1 \rangle \\ &= \sum_{i=1}^n \langle e_i, s_{n,1} b^* \cdot a^* a \cdot b s_{n,1}^* e_i \rangle \\ &= n \cdot \psi_n(s_{n,1} b^* \cdot a^* a \cdot b s_{n,1}^*). \end{aligned}$$

3) The composition  $\psi_n \circ \alpha_n^{-1}$  equals the only Haar state  $h_n$  on the finite dimensional Hopf  $C^*$ -algebra  $(C^*(\mathbb{Z}/n\mathbb{Z}), \Delta)$ , which is faithful on  $C^*(\mathbb{Z}/n\mathbb{Z})$ .  $\square$

As a consequence, the following holds:

**Proposition 5.2.** *Let  $(\mathfrak{B}, \psi) := (\mathcal{T}_2, \psi_2) *_\mathbb{C} (\mathcal{T}_3, \psi_3)$  be the reduced unital free product defined by D. Voiculescu ([Voi85]).*

- 1) *The  $C^*$ -algebra  $\mathfrak{B}$  contains the algebraic unital free product generated by  $\mathcal{T}_2$  and  $\mathcal{T}_3$*
- 2) *The  $C^*$ -algebra  $\mathfrak{B}$  is simple and has stable rank 1.*
- 3) *The  $C^*$ -algebra  $\mathfrak{B}$  is  $K_1$ -injective.*

*Proof.* 1) This embedding result is essentially contained in Proposition 4.2 of [DS01]. But we include here the main steps of that proof for completeness. Define:

- the tensor product  $C^*$ -algebra  $B := \mathcal{T}_2 \otimes \mathcal{T}_3$ ,
- the state  $\phi := \psi_2 \otimes \psi_3$  on the  $C^*$ -algebra  $B$ ,
- the full countably generated Hilbert  $B$ -bimodule  $E := L^2(B, \phi) \otimes B$ ,
- the full Fock Hilbert  $B$ -bimodule  $\mathcal{F}(E) = B \oplus E \oplus (E \otimes E) \oplus \dots$ ,
- the Pimsner-Toeplitz  $C^*$ -algebra  $\mathcal{T}(E)$  generated in  $\mathcal{L}(\mathcal{F}(E))$  by the *creation* operators  $\ell(\xi)$ ,  $\xi \in E$  ([Pim95]).
- the conditional expectation  $\mathfrak{E} : \mathcal{T}(E) \rightarrow B$  given by compression with the orthogonal projection from  $\mathcal{F}_B(E)$  onto the first summand  $B \subset \mathcal{F}(E)$  ([DS01]).

Take a unitary  $v$  in the  $C^*$ -subalgebra  $C^*(\ell(\Lambda_\phi 1 \otimes 1)) \subset \mathcal{T}(E)$  such that  $\mathfrak{E}(v^j) = 0$  for any non-zero integer  $j$  and let  $\pi_2 : \mathcal{T}_2 \rightarrow \mathcal{T}(E)$ ,  $\pi_3 : \mathcal{T}_3 \rightarrow \mathcal{T}(E)$  be the two unital  $*$ -morphisms  $\pi_2(a) = v^2(1 \otimes a \otimes 1_{\mathcal{T}_3})v^{-2}$  and  $\pi_3(a) = v^3(1 \otimes 1_{\mathcal{T}_2} \otimes a)v^{-3}$ . Then, the algebraic amalgamated free product  $\mathcal{T}_2 *_\mathbb{C} \mathcal{T}_3$  studied by B. Blackadar ([Blac78]) and G. Pedersen ([Ped94]) is (isomorphic to) the unital  $*$ -algebra generated by  $\pi_2(\mathcal{T}_2)$  and  $\pi_3(\mathcal{T}_3)$  in  $\mathcal{T}(E)$ , whereas Voiculescu's reduced unital free product  $(\mathcal{T}_2, \psi_2) *_\mathbb{C} (\mathcal{T}_3, \psi_3)$  ([Voi85]) is (isomorphic to) its closure (see *e.g.* [Blan09, Theorem 4.1]).

2) The previous Lemma and [BD01, Theorem 1.3] imply that there is a unital embedding  $(C^*(\mathbb{Z}/2\mathbb{Z}), h_2) *_\mathbb{C} (C^*(\mathbb{Z}/3\mathbb{Z}), h_3) \hookrightarrow (\mathfrak{B}, \psi) = (\mathcal{T}_2, \psi_2) *_\mathbb{C} (\mathcal{T}_3, \psi_3)$ , if  $h_2$  and  $h_3$  denote the Haar states on the compact groups  $(C^*(\mathbb{Z}/2\mathbb{Z}), \Delta_2)$  and  $(C^*(\mathbb{Z}/3\mathbb{Z}), \Delta_3)$ . Hence, the reduced free product  $\mathfrak{B}$  is simple ([Av82, Proposition 3.1+its corollary]) and this unital  $C^*$ -algebra has stable rank 1 ([DHR97, Theorem 3.8]), *i.e.* the open subset of invertible elements is dense in  $\mathfrak{B}$ .

3) With the notations of [Rief83], the set  $Lg_1(\mathfrak{B})$  of invertible elements is dense in the stable rank 1  $C^*$ -algebra  $\mathfrak{B}$ , so that this  $C^*$ -algebra  $\mathfrak{B}$  is  $K_1$ -injective by [Rief83, Proposition 1.6] and [Rief87, Theorem 2.10]. (A self-contained proof is also available in Theorem 3.2.11 of [Roh09].)  $\square$

*Remark 5.3.* The same proof shows that the reduced free product  $(\mathcal{T}_{n_1}, \psi_{n_1}) *_\mathbb{C} (\mathcal{T}_{n_2}, \psi_{n_2})$  is simple and  $K_1$ -injective for all integers  $n_1 \geq 2$  and  $n_2 \geq 3$ .

**Question 5.4.** *Is the full free product  $\mathcal{T}_2 *_\mathbb{C} \mathcal{T}_3$  also  $K_1$ -injective?*

*Note that the full free products  $\mathcal{T}_2 *_\mathbb{C} \mathcal{T}_2$  and  $\mathcal{T}_2 *_\mathbb{C} \mathcal{T}_3$  unitaly embed in the  $C^*$ -algebras  $\mathcal{O}_3 *_\mathbb{C} \mathcal{O}_3$  and  $\mathcal{O}_3 *_\mathbb{C} \mathcal{O}_4$  ([ADEL04]).*



## REFERENCES

- [ADEL04] S. Armstrong, K. Dykema, R. Exel, H. Li, *On embeddings of full amalgamated free product of  $C^*$ -algebras*, Proc. Amer. Math. Soc. **132** (2004), 2019–2030.
- [Av82] D. Avitzour, *Free products of  $C^*$ -algebras*, Trans. Amer. Math. Soc. **271** (1982), 423–465.
- [Blac78] B. Blackadar, *Weak expectation and nuclear  $C^*$ -algebras*, Indiana Univ. Math. J. **27** (1978), 1021–1026.
- [Blac04] B. Blackadar, *Semiprojectivity in simple  $C^*$ -algebras*, Adv. Stud. Pure Math. **38** (2004), 1–17.
- [Blan97] E. Blanchard, *Subtriviality of continuous fields of nuclear  $C^*$ -algebras*, with an appendix by E. Kirchberg, J. Reine Angew. Math. **489** (1997), 133–149.
- [Blan09] E. Blanchard, *Amalgamated free products of  $C^*$ -bundles*, Proc. Edinburgh Math. Soc. **52** (2009), 23–36.
- [Blan10] E. Blanchard,  *$K_1$ -injectivity for properly infinite  $C^*$ -algebras*, Clay Math. Proc. **11** (2010), 49–54.
- [Blan13] E. Blanchard, *Continuous fields with fibres  $\mathcal{O}_\infty$* , to appear in Math. Scand.
- [BD01] E. Blanchard, K. Dykema, *Embeddings of reduced free products of operator algebras*, Pacific J. Math. **199** (2001), 1–19.
- [BK04a] E. Blanchard, E. Kirchberg, *Global Glimm halving for  $C^*$ -bundles*, J. Op. Th. **52** (2004), 385–420.
- [BK04b] E. Blanchard, E. Kirchberg, *Non-simple purely infinite  $C^*$ -algebras: the Hausdorff case*, J. Funct. Anal. **207** (2004), 461–513.
- [BRR08] E. Blanchard, R. Rohde, M. Rørdam, *Properly infinite  $C(X)$ -algebras and  $K_1$ -injectivity*, J. Noncommut. Geom. **2** (2008), 263–282.
- [Cun77] J. Cuntz, *Simple  $C^*$ -Algebras Generated by Isometries*, Commun. Math. Phys. **57** (1977), 173–185.
- [Cun81] J. Cuntz,  *$K$ -theory for certain  $C^*$ -algebras*, Ann. of Math. **113** (1981), 181–197.
- [DD63] J. Dixmier, A. Douady, *Champs continus d’espaces hilbertiens et de  $C^*$ -algèbres*, Bull. Soc. Math. France **91** (1963), 227–284.
- [Dix69] J. Dixmier, *Les  $C^*$ -algèbres et leurs représentations*, Gauthiers-Villars Paris (1969).
- [DS01] K. Dykema, D. Shlyakhtenko, *Exactness of Pimsner-Toeplitz  $C^*$ -algebras* Proc. Edinb. Math. Soc. **44** (2001), 425–444.
- [DHR97] K. Dykema, U. Haagerup, M. Rørdam, *The stable rank of some free product  $C^*$ -algebras*, Duke Math. J. **90** (1997), 95–121 [Correction in DMJ **94** (1998), 213].
- [Dad09] M. Dădărlat, *Fiberwise  $KK$ -equivalence of continuous fields of  $C^*$ -algebras*, J. K-Theory **3** (2009), 205–219.
- [DW08] M. Dădărlat, W. Winter, *Trivialization of  $C(X)$ -algebras with strongly self-absorbing fibres*, Bull. Soc. Math. France **136** (2008), 575–606.
- [Ell94] G. Elliott, *The classification problem for amenable  $C^*$ -algebras*, Proc. Internat. Congress of Mathematicians (Zurich, Switzerland, 1994), Birkhauser Verlag, Basel (1995), 922–932.
- [HRW07] I. Hirshberg, M. Rørdam, W. Winter,  *$C_0(X)$ -algebras, stability and strongly self-absorbing  $C^*$ -algebras*, Math. Ann. **339** (2007), 695–732.
- [HR98] J. Hjelmborg, M. Rørdam, *On stability of  $C^*$ -algebras*, J. Funct. Anal. **155** (1998), 153–170.
- [Kas88] G.G. Kasparov, *Equivariant  $KK$ -theory and the Novikov conjecture*, Invent. Math. **91** (1988), 147–201.
- [KR02] E. Kirchberg, M. Rørdam, *Infinite non-simple  $C^*$ -algebras: absorbing the Cuntz algebra  $\mathcal{O}_\infty$* , Advances in Math **167** (2002), 195–264.
- [LLR00] F. Larsen, N. J. Laustsen, M. Rørdam, *An Introduction to  $K$ -theory for  $C^*$ -algebras*, London Mathematical Society Student Texts **49** (2000) CUP, Cambridge.
- [Ped94] G.K. Pedersen, *Pullback and pushout constructions in  $C^*$ -algebra theory*, J. Funct. Anal. **167** (1999), 243–344.
- [Pim95] M. V. Pimsner, *A class of  $C^*$ -algebras generalizing both Cuntz-Krieger algebras and crossed products by  $\mathbb{Z}$* , Free probability theory, Fields Inst. Commun. **12** (1997), 189–212.

- [Rief83] M. Rieffel, *Dimension and stable rank in the  $K$ -theory of  $C^*$ -algebras*, Proc. London Math. Soc. **46** (1983), 301-333.
- [Rief87] M. Rieffel, *The homotopy groups of the unitary groups of noncommutative tori*, J. Operator Theory **17** (1987), 237-254.
- [Roh09] R. Rohde,  *$K_1$ -injectivity of  $C^*$ -algebras*, Ph.D. thesis at Odense (2009)
- [Ror97] M. Rørdam, *Stability of  $C^*$ -algebras is not a stable property.*, Doc. Math. J. **2** (1997), 375–386.
- [Ror03] M. Rørdam, *A simple  $C^*$ -algebra with a finite and an infinite projection*, Acta Math. **191** (2003), 109142.
- [Ror04] M. Rørdam, *The stable and the real ranks of  $\mathcal{Z}$ -absorbing  $C^*$ -algebras*, Internat. J. Math. **15** (2004), 1065–1084.
- [TW07] A. Toms, W. Winter, *Strongly self-absorbing  $C^*$ -algebras*, Trans. Amer. Math. Soc. **359** (2007), 3999-4029.
- [Voi85] D. Voiculescu, *Symmetries of some reduced free product  $C^*$ -algebras*, Operator Algebras and Their Connections with Topology and Ergodic Theory (Buşteni, 1983), Lecture Notes in Math. **112** (1985), 556–588.
- [Win09] W. Winter, *Strongly self-absorbing  $C^*$ -algebras are  $\mathcal{Z}$ -stable*, J. Noncommut. Geom. **5** (2011), 253-264.
- [Wor95] S. L. Woronowicz, *Compact quantum groups*, Symétries quantiques (Les Houches, 1995), North-Holland, Amsterdam (1998), 845-884.

Etienne.Blanchard@math.jussieu.fr

IMJ, Bâtiment Sophie Germain, Case 7012, F-75013 Paris