HAL open science

Subtriviality of continuous fields of nuclear C*-algebras, with an appendix by E. Kirchberg

Etienne Blanchard

▶ To cite this version:

Etienne Blanchard. Subtriviality of continuous fields of nuclear C*-algebras, with an appendix by E. Kirchberg. Journal für die reine und angewandte Mathematik, 1997, 498, pp.133—149. hal-00922873

HAL Id: hal-00922873

https://hal.science/hal-00922873

Submitted on 31 Dec 2013

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Subtriviality of continuous fields of nuclear C*-algebras

By Etienne Blanchard

Last update 15/08/04

Abstract

We extend in this paper the characterisation of a separable nuclear C*-algebra given by Kirchberg proving that given a unital separable continuous field of nuclear C*-algebras A over a compact metrizable space X, the $C(X)$-algebra A is isomorphic to a unital $C(X)$-subalgebra of the trivial continuous field $\mathcal{O}_2 \otimes C(X)$, image of $\mathcal{O}_2 \otimes C(X)$ by a norm one projection.

AMS classification: 46L05.

0 Introduction

In order to study deformations in the C*-algebraic framework, Dixmier introduced the notion of continuous field of C*-algebras over a locally compact space ([7]). In the same way as there is a faithful representation in a Hilbert space for any C*-algebra thanks to the Gelfand–Naimark–Segal construction, a separable continuous field of C*-algebras A over a compact metrizable space X always admits a continuous field of faithful representations π in a Hilbert $C(X)$-module, i.e. there exists a family of representations $\{\pi_x,\ x \in X\}$, in a separable Hilbert space H which factorize through a faithful representation of the fibre A_x such that for each $a \in A$, the map $x \mapsto \pi_x(a)$ is strongly continuous ([4, théorème 3.3]).

In a work on tensor products over $C(X)$ of continuous fields of C*-algebras over X ([16]), Kirchberg and Wassermann raised the question of whether the continuous field of C*-algebras A could be subtrivialized, i.e. whether one could find a continuous field of faithful representations π such that the map $x \mapsto \pi_x(a) \in L(H)$ is actually norm continuous for all a in A. In this case, given any C*-algebra B, the minimal tensor product $A \otimes B$ is a $C(X)$-subalgebra of the trivial continuous field $[L(H) \otimes B] \otimes C(X)$ and is therefore a continuous field with fibres $(A \otimes B)_x = A_x \otimes B$. They proved that a non-exact continuous field with exact fibres cannot be subtrivialized and they constructed such examples.

The non-trivial example of the continuous field of rotation algebras over the unit circle $\mathbb{T}$ had already been studied by Haagerup and Rørdam in [10]. More precisely, they constructed continuous functions u, v from $\mathbb{T}$ to the unitary group $U(H)$ of the infinite-dimensional separable Hilbert space H satisfying the commutation relation $u_t v_t = t v_t u_t$

for all $t \in \mathbb{T}$ and the uniform continuity condition $\max\{\|u_t - u_{t'}\|, \|v_t - v_{t'}\|\} < C'|t - t'|^{1/2}$ where C' is a computable constant.

Our purpose in the present paper is to show that the subtrivialization is always possible in the nuclear separable case through a generalisation of the following theorem of Kirchberg using $\mathcal{R}KK$-theory arguments:

Theorem 0.1 *([15]) A unital separable* C*-*algebra A is exact if and only if it is isomorphic to a* C*-*subalgebra of $\mathcal{O}_2$. Moreover the* C*-*algebra A is nuclear if and only if A is isomorphic to a* C*-*subalgebra of $\mathcal{O}_2$ containing the unit $1_{\mathcal{O}_2}$ of $\mathcal{O}_2$, image of $\mathcal{O}_2$ by a unital completely positive projection.*

As a matter of fact, we get an equivalent characterisation of nuclear separable continuous fields of C*-algebras (theorem 3.2) which is made possible thanks to $C(X)$-linear homotopy invariance of the bifunctor $\mathcal{R}KK(X; -, -)$ (theorem 1.6) and $C(X)$-linear Weyl-von Neumann absorption results (proposition 2.5). This also enables us to have a better understanding of the characterisation of separable continuous fields of nuclear C*-algebras given by Bauval in [2].

In an added appendix, the corresponding characterisation of exact separable continuous fields of C*-algebras as $C(X)$-subalgebras of $\mathcal{O}_2 \otimes C(X)$ given by Eberhard Kirchberg is described (theorem A.1).

I would like to thank E. Kirchberg for his enlightenment on the exact case. I also want to express my gratitude to C. Anantharaman-Delaroche and J. Cuntz for fruitful discussions.

1 Preliminaries

1.1 $C(X)$-algebras

Let X be a compact Hausdorff space and $C(X)$ be the C*-algebra of continuous functions on X with complex values. We start by recalling the following definition.

Definition 1.1 *([13]) A $C(X)$-algebra is a* C*-*algebra A endowed with a unital morphism from $C(X)$ in the centre of the multiplier algebra $M(A)$ of A.*

Remark: We do not assume that $C(X)$ embeds into $M(A)$. For instance, there is a natural structure of $C([0,2])$-algebra on the C*-algebra $C([0,1])$.

For $x \in X$, define the kernel $C_x(X)$ of the evaluation map $ev_x : C(X) \to \mathbb{C}$ at x; denote by A_x the quotient of a $C(X)$-algebra A by the closed ideal $C_x(X)A$ and by a_x the image of an element $a \in A$ in the fibre A_x. Then the function

$$x \mapsto \|a_x\| = \inf\{\|[1 - f + f(x)]a\|, f \in C(X)\}$$

is upper semi-continuous for any $a \in A$ and the $C(X)$-algebra A is said to be a continuous field of C*-algebras over X if the function $x \mapsto \|a_x\|$ is actually continuous for every $a \in A$ ([7]).

Examples *1.* If A is a $C(X)$-algebra and D is a C*-algebra, the spatial tensor product $B = A \otimes D$ is naturally endowed with a structure of $C(X)$-algebra through the map $f \in C(X) \mapsto f \otimes 1_{M(D)} \in M(A \otimes D)$. In particular, if $A = C(X)$, the tensor product B is a trivial continuous field over X with constant fibre $B_x \simeq D$
2. Given a $C(X)$-algebra A, define the unital $C(X)$-algebra $\mathcal{A}$ generated by A and $u[C(X)]$ in $M[A \oplus C(X)]$ where $u(g)(a \oplus f) = ga \oplus gf$ for $a \in A$ and $f, g \in C(X)$. It defines a continuous field of C*-algebras over X if and only if the $C(X)$-algebra A is continuous ([4, proposition 3.2]).

Remark: If A is a separable continuous field of non-zero C*-algebras (not necessarily unital) over the compact Hausdorff space X, the positive cone $C(X)_+$ and so the C*-algebra $C(X)$ are separable. Hence, the topological space X is metrizable.

Definition 1.2 *([4, 5]) Given a continuous field of* C*-*algebras A over the compact Hausdorff space X, a continuous field of representations of a $C(X)$-algebra D in the multiplier C*-algebra $M(A)$ of A is a $C(X)$-linear morphism $\pi : D \to M(A)$, i.e. for each $x \in X$, the induced representation π_x of D in $M(A_x)$ factorizes through the fibre D_x.*

If the $C(X)$-algebra D admits a continuous field of faithful representations π in the $C(X)$-algebra $M(A)$ where A is a continuous field over X, i.e. the induced representation of the fibre D_x in $M(A_x)$ is faithful for every point $x \in X$, the function

$$x \mapsto \|\pi_x(d)\| = \sup\{\|(\pi(d)a)_x\|, a \in A \,\text{such that}\, \|a\| \leq 1\}$$

is lower semi-continuous for all $d \in D$ and the $C(X)$-algebra D is therefore continuous.

In particular a separable $C(X)$-algebra D is continuous if and only if there exists a Hilbert $C(X)$-module $\mathcal{E}$ such that D admits a continuous field of faithful representations in the multiplier algebra $M(\mathcal{K}(\mathcal{E})) = \mathcal{L}(\mathcal{E})$ of the continuous field over X of compact operators $\mathcal{K}(\mathcal{E})$ acting on $\mathcal{E}$ ([4, théorème 3.3]).

Let us also mention the characterisation of separable continuous fields of nuclear C*-algebras over a compact metrizable space X given by Bauval in [2] using a natural $C(X)$-linear version of nuclearity introduced by Kasparov and Skandalis in [14]§6.2 : a $C(X)$-linear completely positive σ from a $C(X)$-algebra A into a $C(X)$-algebra B is said to be $C(X)$-nuclear if and only if given any compact set F in A and any strictly positive real number ε, there exist an integer k and $C(X)$-linear completely positive contractions $T : A \to M_k(\mathbb{C}) \otimes C(X)$ and $S : M_k(\mathbb{C}) \otimes C(X) \to B$ such that for all $a \in F$, one has the inequality

$$\|\sigma(a) - (S \circ T)(a)\| < \varepsilon.$$

One can then state the following results. The first assertion is a simple $C(X)$-linear reformulation of the Choi-Effros theorem and the second one is due to Bauval.

Proposition 1.3 *Let X be a compact metrizable space and A be a separable $C(X)$-algebra.*

1. *([14]§6.2) Given a $C(X)$-algebra B and a closed ideal $J \subset B$, any contractive $C(X)$-nuclear map $A \to B/J$ admits a contractive $C(X)$-linear completely positive lift $A \to B$.*

2. *([2, théorème 7.2]) The $C(X)$-algebra A is a continuous fields of nuclear C*-algebras over X if and only if the identity map $id_A : A \to A$ is $C(X)$-nuclear.*

Remark: In assertion *1.*, the ideal $J = (C(X)B)J = C(X)J$ is a $C(X)$-algebra.

1.2 $C(X)$-extensions

Given a compact Hausdorff space X, we introduce a natural $C(X)$-linear version of the semi-group $Ext(-,-)$ defined by Kasparov ([12, 13]).

Call a morphism of $C(X)$-algebras a $*$-homomorphism between $C(X)$-algebras which is $C(X)$-linear.

Definition 1.4 *A $C(X)$-extension of a $C(X)$-algebra A by a $C(X)$-algebra B is a short exact sequence*

$$0 \to B \to D \xrightarrow{\pi} A \to 0$$

in the category of $C(X)$-algebras. The $C(X)$-extension is said to be trivial if the map π admits a cross section $s : A \to D$ which is a morphism of $C(X)$-algebras.

As in the C*-algebraic case a $C(X)$-extension $0 \to B \to D \to A \to 0$ of A by B defines unambiguously an homomorphism from D to the multiplier algebra $M(B)$ of B, which gives a morphism of $C(X)$-algebras $\sigma : A \to M(B)/B$ (called the Busby invariant of the extension) and the $C(X)$-extension is trivial if and only if the map σ lifts to a morphism of $C(X)$-algebras $A \to M(B)$. Conversely, given a morphism of $C(X)$-algebras $\sigma : A \to M(B)/B$, setting $D = \{(a, m) \in A \times M(B), \sigma(a) = q(m)\}$ where q is the quotient map $M(B) \to M(B)/B$, one has a $C(X)$-extension $0 \to B \to D \to A \to 0$ (see [12]§7).

Remark: A $C(X)$-extension $0 \to B \to D \to A \to 0$ induces for every $x \in X$ a C*-extension $0 \to B_x \to D_x \to A_x \to 0$.

In order to define the sum of two $C(X)$-extensions, recall that the Cuntz algebra $\mathcal{O}_2$ is the unital C*-algebra generated by two orthogonal isometries s_1 and s_2 subject to the relation $1 = s_1 s_1^* + s_2 s_2^*$ ([6]). Then if $\mathcal{K}$ is the C*-algebra of compact operators on the infinite-dimensional separable Hilbert space, one defines the sum of two $C(X)$-extensions σ_1 and σ_2 of the $C(X)$-algebra A by the stable $C(X)$-algebra $\mathcal{K} \otimes B$ through the choice of a unital copy of $\mathcal{O}_2$ in the multiplier algebra $M(\mathcal{K})$ of $\mathcal{K}$ to be the $C(X)$-extension

$$\sigma_1 \oplus \sigma_2 : a \mapsto q(s_1 \otimes 1)\sigma_1(a)q(s_1^* \otimes 1) + q(s_2 \otimes 1)\sigma_2(a)q(s_2^* \otimes 1) \in M(\mathcal{K} \otimes B)/(\mathcal{K} \otimes B),$$

where q is the quotient map $M(\mathcal{K} \otimes B) \to M(\mathcal{K} \otimes B)/(\mathcal{K} \otimes B)$.

Definition 1.5 *Given a compact Hausdorff space X and two $C(X)$-algebras A and B, $Ext(X;A,B)$ is the semi-group of $C(X)$-extensions of A by $\mathcal{K}\otimes B$ divided by the equivalence relation $\sim$ where $\sigma_1\sim\sigma_2$ if there exist a unitary $U\in M(\mathcal{K}\otimes B)$ of image $q(U)$ in the quotient $M(\mathcal{K}\otimes B)/(\mathcal{K}\otimes B)$ and two trivial $C(X)$-extensions π_1 and π_2 such that for all $a\in A$,*

$$(\sigma_2\oplus\pi_2)(a)=q(U)^*(\sigma_1\oplus\pi_1)(a)q(U)\ \textit{(in } M(\mathcal{K}\otimes B)/(\mathcal{K}\otimes B)\textit{)}.$$

Let $Ext(X;A,B)^{-1}$ be the group of invertible elements of $Ext(X;A,B)$, i.e. the group of classes of $C(X)$-extension σ such that there exists a $C(X)$-extension θ with $\sigma\oplus\theta$ trivial. One can generalise Kasparov's theorem of homotopy invariance of the group $Ext(A,B)^{-1}$ to the framework of $C(X)$-algebras as follows.

Theorem 1.6 *([12]) Assume that A is a separable $C(X)$-algebra and that B is a σ-unital $C(X)$-algebra. Then the group $Ext(X;A,B)^{-1}$ is isomorphic to the group $\mathcal{R}KK^1(X;A,B)$ and is therefore $C(X)$-linear homotopy invariant in both entries A and B.*

Proof : Let us first make the following observation. Given a $C(X)$-algebra B and a Hilbert B-module $\mathcal{E}$, denote by $\mathcal{L}(\mathcal{E})$ the set of bounded B-linear operators on $\mathcal{E}$ which admit an adjoint ([11]). Then any operator $T\in\mathcal{L}(\mathcal{E})$ is B-linear and so $C(X)$-linear. This argument provides a natural extension of the Stinespring-Kasparov theorem ([12]) to the framework of $C(X)$-algebras. Consequently, if A is a separable $C(X)$-algebra and B is a σ-unital $C(X)$-algebra, the class of a $C(X)$-extension $\sigma:A\to M(\mathcal{K}\otimes B)/(\mathcal{K}\otimes B)$ is invertible in $Ext(X;A,B)$ if and only if there is a $C(X)$-linear completely positive contractive lift $A\to M(\mathcal{K}\otimes B)$.

Let $\mathcal{RE}(X;A,B)$ be the set of Kasparov $C(X)$–A,B-bimodules ([13, definition 2.19]), i.e. the set of Kasparov A,B bimodules $(\mathcal{E},F)$ such that the representation $A\to\mathcal{L}(\mathcal{E})$ is a $C(X)$-representation. Call a $C(X)$-linear operator homotopy an element $\{(\mathcal{E},F_t),0\le t\le 1\}\in\mathcal{RE}(X;A,B\otimes C([0,1]))$ such that $t\mapsto F_t$ is norm continuous and define on $\mathcal{RE}(X;A,B)$ the equivalence relation corresponding to the one defined by Skandalis in [18, definition 2]. The constructions given by Kasparov in [12, section 7] imply that, if the $C(X)$-algebra B is σ-unital, the group of equivalence classes $\mathcal{R}\widetilde{KK}(X;A,B\otimes\mathcal{C}_1)$ is isomorphic to $Ext(X;A,B)^{-1}$, where $\mathcal{C}_1$ is the first (graded) Clifford algebra.

On the other hand, given two graded $C(X)$-algebras A and B with A separable, the different steps of the demonstration of [18, theorem 19] provide us with an isomorphism between the two groups $\mathcal{R}\widetilde{KK}(X;A,B)$ and $\mathcal{R}KK(X;A,B)$ since proposition 2.21 of [13] defines an intersection product in $\mathcal{R}\widetilde{KK}$-theory and lemma 18 of [18] gives us the equality

$$(ev_0\otimes id_{C(X)})^*(1_{C(X)})=(ev_1\otimes id_{C(X)})^*(1_{C(X)})\text{ in }\mathcal{R}\widetilde{K}K(X;C([0,1])\otimes C(X),C(X)),$$

where $1_{C(X)}$ is the Kasparov $C(X),C(X)$-bimodule $(C(X),0)$ and $ev_t:C([0,1])\to\mathbb{C}$ is the evaluation map at $t\in[0,1]$. $\square$

Remarks: *1.* Kuiper's theorem implies that the law of addition on the abelian group $Ext(X;A,B)^{-1}$ is independent of the choice of the unital copy of $\mathcal{O}_2$ in $M(\mathcal{K})$.

2. If A is a separable nuclear continuous field of C*-algebras over X and B is a $C(X)$-algebra, every $C(X)$-linear morphism from A to the quotient $M(\mathcal{K}\otimes B)/(\mathcal{K}\otimes B)$ is $C(X)$-nuclear and therefore admits a $C(X)$-linear completely positive lifting $A\to M(\mathcal{K}\otimes B)$ thanks to proposition 1.3. Accordingly one has the equality

$$Ext(X;A,B)^{-1}=Ext(X;A,B).$$

2 An absorption result

In this section we prove a continuous generalisation of a statement contained in [15] which will enable us to get a $C(X)$-linear Weyl-von Neumann type result (proposition 2.5). Let us start with the following definition of Cuntz.

Definition 2.1 *([6]) A simple* C*-*algebra A distinct from $\mathbb{C}$ is said to be purely infinite if and only if for any non-zero $a,b\in A$, there exist elements $x,y\in A$ such that $a=xby$.*

Then, we can state a proposition from Kirchberg's classification work, based on Glimm's lemma ([7], § 11.2). A sketch of proof can also be found in [1, proposition 5.1].

Proposition 2.2 *([15]) Let A be a purely infinite simple* C*-*algebra and D be a separable* C*-*subalgebra of $M(A)$. Assume that $V:D\to A$ is a nuclear contraction.*

Then there exists a sequence (a_n) of elements in A of norm less than 1 such that $V(d)=\lim_{n\to\infty}a_n^*da_n$ for all $d\in D$.

Remark: A simple ring has by definition exactly two distinct two sided ideals and is therefore non-zero.

Corollary 2.3 *Let A be a continuous field of purely infinite simple* C*-*algebras over a compact Hausdorff space X and assume that D is a separable $C(X)$-subalgebra of the multiplier algebra $M(A)$ such that there is a unital $C(X)$-embedding of the $C(X)$-algebra $\mathcal{O}_\infty\otimes C(X)$ in the commutant D' of D in $M(A)$ and the identity map $id_D:D\to M(A)$ is a continuous field of faithful representations.*

If $V:D\to A$ is a $C(X)$-nuclear contraction, there exists a sequence (a_n) in the unit ball of A with the property that for all $d\in D$,

$$V(d)=\lim_{n\to\infty}a_n^*da_n.$$

Proof : If F is a compact generating set for D, it is enough to prove that given a strictly positive real number $\varepsilon>0$, there exists an element a in the unit ball of A such that $\|V(d)-a^*da\|<\varepsilon$ for all $d\in F$.

For $x\in X$, the fibre A_x is a purely infinite simple C*-algebra and the map $d\mapsto V(d)_x\in A_x$ factorizes through $D_x\simeq(id_D)_x(D)\subset M(A_x)$ since id_D is a continuous field of faithful representations. As a consequence, the previous proposition implies that we can find an element $g\in A$ with $\|g\|\leq 1$ satisfying for all $d\in F$ the inequality

$$\|\Big[V(d)-g^*dg\Big]_x\|<\varepsilon.$$

Thus, by upper semi-continuity and compactness, there exist a finite open covering $\{U_1, \ldots, U_n\}$ of the space X and elements $g_1, \ldots, g_n$ in the unit ball of A such that for all $d \in F$ and $x \in U_i$, $1 \leq i \leq n$,

$$\|\Big[V(d) - g_i^* d g_i\Big]_x\| < \varepsilon.$$

Choose n orthogonal isometries $w_1, \ldots, w_n$ in the C*-algebra $\mathcal{O}_\infty \otimes 1_{C(X)} \subset D'$ and let $\{\phi_i\}$ be a partition of the unit $1_{C(X)}$ subordinate to the covering $\{U_i\}$ of X. The element $a = \sum_i \phi_i^{1/2} w_i g_i \in A$ verifies:

1. $a^* a = \sum_{i,j} \sqrt{\phi_i \phi_j}\, g_i^* w_i^* w_j g_j = \sum_i \phi_i g_i^* g_i \leq 1_{M(A)}$,
2. for $d \in F$ and $x \in X$, $\|\Big[V(d) - a^* d a\Big]_x\| \leq \sum_i \phi_i(x) \|\Big[V(d) - g_i^* d g_i\Big]_x\| < \varepsilon$. $\square$

Let us mention the following technical corollary which will be needed in theorem 3.2.

Corollary 2.4 *If $p \in \mathcal{O}_2 \otimes C(X)$ is a projection such that for all points $x \in X$, p_x is non-zero, then there exists an isometry $u \in \mathcal{O}_2 \otimes C(X)$ such that $p = uu^*$.*

Proof : Let $\mathcal{D}_2 = \lim_{n\to\infty} {\mathcal{O}_2}^{\otimes n}$ be the infinite tensor product of $\mathcal{O}_2$.

Given a projection $q \in \mathcal{D}_2 \otimes C(X)$ such that $\|q_x\| = 1$ for all $x \in X$, we first show that there exists an element $v \in \mathcal{D}_2 \otimes C(X)$ satisfying $1_{\mathcal{D}_2 \otimes C(X)} = v^* q v$. Namely, by density of the algebraic tensor product

$$\Big[\bigcup_n {\mathcal{O}_2}^{\otimes n}\Big] \odot C(X) = \bigcup_n \Big[{\mathcal{O}_2}^{\otimes n} \odot C(X)\Big]$$

in the C*-algebra $\mathcal{D}_2 \otimes C(X)$ and functional calculus one can find an integer $n > 0$ and a projection $r \in {\mathcal{O}_2}^{\otimes n} \otimes C(X) \subset \mathcal{D}_2 \otimes C(X)$ such that $\|q - r\| < 1$, which implies that $r = s^* q s$ for some element $s \in \mathcal{D}_2 \otimes C(X)$. Take then a faithful state φ on ${\mathcal{O}_2}^{\otimes n}$ and consider the $C(X)$-linear completely positive map

$$V : [{\mathcal{O}_2}^{\otimes n} \otimes 1_{\mathcal{O}_2}] \otimes C(X) \to {\mathcal{O}_2}^{\otimes n+1} \otimes C(X)$$

defined by the formula $V(d) = (\varphi \otimes id_{C(X)})(d) 1_{{\mathcal{O}_2}^{\otimes n+1} \otimes C(X)}$ for $d \in [{\mathcal{O}_2}^{\otimes n} \otimes 1_{\mathcal{O}_2}] \otimes C(X) \simeq {\mathcal{O}_2}^{\otimes n} \otimes C(X)$. According to corollary 2.3, there exists an element $t \in {\mathcal{O}_2}^{\otimes n+1} \otimes C(X)$ such that

$$1_{\mathcal{D}_2 \otimes C(X)} = 1_{{\mathcal{O}_2}^{\otimes n+1} \otimes C(X)} = t^* r t = (st)^* q (st).$$

Consider now the set $\mathcal{P}$ of projections p in $\mathcal{O}_2 \otimes C(X)$ such that $p_x \neq 0$ for all points $x \in X$. If p belongs to $\mathcal{P}$, there exists an isometry $v \in \mathcal{O}_2 \otimes C(X)$ such that $p \geq vv^*$ since the K-trivial purely infinite separable unital nuclear C*-algebra $\mathcal{D}_2$ satisfying the U.C.T. is isomorphic to $\mathcal{O}_2$ ([15]). As a consequence, if t is the isometry $t = v(s_1 \otimes 1)v^*$, the projection $r = tt^*$ (Murray-von Neumann equivalent to $1_{\mathcal{O}_2 \otimes C(X)}$) verifies

$$p - r \geq r' = v(s_2 s_2^* \otimes 1)v^* \in \mathcal{P}.$$

The non-empty set $\mathcal{P}$ therefore satisfies the conditions (π_1)–(π_4) defined by Cuntz in [6]. But the C*-algebra $\mathcal{O}_2 \otimes C(X)$ is K_0-triviality thanks to [6, theorem 2.3] and the theorem 1.4 of [6] enables us to conclude. □

One now deduces from corollary 2.3 the following absorption results ([21, 12, 15]):

Proposition 2.5 *Let A be a σ-unital continuous field of purely infinite simple nuclear C*-algebras over a compact Hausdorff space X and let $\mathcal{K}$ be the C*-algebra of compact operators on the separable Hilbert space H. Denote by q the quotient map $M(\mathcal{K}\otimes A) \to M(\mathcal{K}\otimes A)/(\mathcal{K}\otimes A)$.*

1. *Assume that D is a unital separable $C(X)$-subalgebra of the multiplier algebra $M(\mathcal{K}\otimes A)$ with same unit such that there is a unital $C(X)$-embedding of the $C(X)$-algebra $\mathcal{O}_\infty \otimes C(X)$ in the commutant of D in $M(\mathcal{K}\otimes A)$ and the identity map id_D is a continuous field of faithful representations of D in $M(\mathcal{K}\otimes A)$.*

   (a) *If V is a unital $C(X)$-linear completely positive map from D in $M(\mathcal{K}\otimes A)$ which is zero on the intersection $D\cap(\mathcal{K}\otimes A)$, there exists a sequence of isometries s_n in $M(\mathcal{K}\otimes A)$ such that for every $d\in D$,*
   $$V(d) - s_n^* d s_n \in \mathcal{K}\otimes A \text{ and } V(d) = \lim_n s_n^* d s_n.$$

   (b) *If π is a unital morphism of $C(X)$-algebras from D into $M(\mathcal{K}\otimes A)$ which is zero on the intersection $D\cap(\mathcal{K}\otimes A)$, there exists a sequence of unitaries u_n in $M(\mathcal{K}\otimes A)$ such that for every $d\in D$,*
   $$(d\oplus\pi(d)) - u_n^* d u_n \in \mathcal{K}\otimes A \text{ and } (d\oplus\pi(d)) = \lim_n u_n^* d u_n.$$

   (c) *Let B be a $C(X)$-algebra and assume that the quotient $D/(D\cap(\mathcal{K}\otimes A))$ is isomorphic to the $C(X)$-algebra $\mathcal{B}$, where $\mathcal{B}$ is the unital $C(X)$-algebra generated by $C(X)$ and B in $M[B\oplus C(X)]$ ([4, définition 2.7]).*

   Then, if $\pi : B \to M(\mathcal{K}\otimes A)$ is a $C(X)$-linear homomorphism, there exists a unitary $U\in M(\mathcal{K}\otimes A)$ such that for all $b\in B\subset M(\mathcal{K}\otimes A)/(\mathcal{K}\otimes A)$,
   $$b\oplus(q\circ\pi)(b) = q(U)^*\, b\, q(U).$$

2. *Assume that the continuous field A is separable and let D be a separable $C(X)$-subalgebra of $M(A)$ containing A such that the identity representation $D\to M(A)$ is a continuous field of faithful representations and there is a unital $C(X)$-embedding of the $C(X)$-algebra $\mathcal{O}_\infty\otimes C(X)$ in the commutant of D in $M(A)$. Define the quotient $C(X)$-algebra $B = D/A$.*

   If $\pi : \mathcal{K}\otimes B \to M(\mathcal{K}\otimes A)$ is a morphism of $C(X)$-algebras, there exists a unitary $U\in M(\mathcal{K}\otimes A)$ such that for all $b\in\mathcal{K}\otimes B\subset M(\mathcal{K}\otimes A)/(\mathcal{K}\otimes A)$,
   $$b\oplus(q\circ\pi)(b) = q(U)^*\, b\, q(U).$$

Proof : *1.* It derives from corollary 2.3 by the same method as the one developed by Kasparov in [11, theorem 5 and 6]. Nevertheless, for the convenience of the reader we describe the different steps of the demonstration.
1.a) Let F be a compact generating set for D containing the unit $1_{M(\mathcal{K}\otimes A)}$. Then given a real number $\varepsilon > 0$, it is enough to find an element $a \in M(\mathcal{K} \otimes A)$ such that $V(d) - a^*da \in \mathcal{K} \otimes A$ and $\|V(d) - a^*da\| < 3\varepsilon$ for all $d \in F$.

Let $\{e_n\}$ be an increasing, positive, quasicentral, countable approximate unit in the ideal $\mathcal{K} \otimes A$ of the C*-algebra generated by $\mathcal{K} \otimes A + V(D)$. If we set $f_0 = (e_0)^{1/2}$ and $f_k = (e_k - e_{k-1})^{1/2}$ for $k \geq 1$, we can then assume, passing to a subsequence of (e_n) if necessary, that $\|V(d)f_k - f_kV(d)\| < 2^{-k}\varepsilon$ for all $k \in \mathbb{N}$ and $d \in F$. This implies that the series $\sum_k [V(d)f_k - f_kV(d)]f_k$ is convergent in $\mathcal{K} \otimes A$ and its norm is less than ε. Furthermore, the series $\sum_k [f_kV(d)f_k]$ is strictly convergent in $M(\mathcal{K} \otimes A)$ for all $d \in F$ since $\sum_k f_k^2$ is strictly convergent to 1.

Notice now that the maps $V_k(d) = f_kV(d)f_k$ are all $C(X)$-nuclear since the separable continuous field $\mathcal{K} \otimes A$ is nuclear. The corollary 2.3 therefore enables us to choose by induction $a_k \in \mathcal{K} \otimes A$ satisfying the following conditions:

1. $\forall d \in F \quad \|V_k(d) - a_k^*da_k\| < 2^{-k}\varepsilon$,
2. $\forall d \in F, \forall l < k \quad \|a_l^*da_k\| < 2^{-l-k}\varepsilon$,
3. $\sum_k a_k$ is strictly convergent toward an element $a \in M(A)$.

One then checks as in [11, theorem 5] that the limit a satisfies the desired properties.

1.b) Take a compact generating F for D containing $1_{M(\mathcal{K}\otimes A)}$ and consider the homomorphism $\pi' = 1 \otimes \pi : D \to M(\mathcal{K} \otimes (\mathcal{K} \otimes A)) \simeq M(\mathcal{K} \otimes A)$. Given $\delta > 0$, one can find, thank to the previous assertion, an isometry $s \in M(\mathcal{K} \otimes A)$ such that

$$s^*ds - \pi'(d) \in \mathcal{K} \otimes A \text{ and } \|s^*ds - \pi'(d)\| < \delta \text{ for all } d \in K^*K.$$

As a consequence, if we fix $\varepsilon > 0$, the choice of δ small enough gives us the inequality $\|pd - dp\| < \varepsilon$, and so $\|d - [pdp + p^\perp dp^\perp]\| < 2\varepsilon$ for all $d \in F$, where $p = ss^*$ and $p^\perp = 1 - p$.

Define the unital map $\Theta : D \to M(p^\perp(\mathcal{K} \otimes A)p^\perp)$ by the formula $\Theta(d) = p^\perp dp^\perp$. According to the stabilisation theorem of Kasparov ([11, theorem 2]), one can construct a unitary $w \in M(\mathcal{K} \otimes A)$ verifying for all $d \in F$ the inequality

$$\|d - w^*[\pi'(d) \oplus \Theta(d)]w\| < 3\varepsilon.$$

To finish the demonstration, notice that the two homomorphisms π' and $\pi' \oplus \pi$ are unitarily equivalent.

1.c) Consider the unital extension $\widetilde{\pi}$ of π to $\mathcal{B}$. Then, the morphism $\widetilde{\pi} \circ q : D \to M(\mathcal{K} \otimes A)$ reduces the demonstration to the previous assertion.

2. The identity representation of the unital $C(X)$-algebra $\mathcal{D} = (\mathcal{K} \otimes D) + C(X) \subset M(\mathcal{K} \otimes A)$ is clearly a continuous field of faithful representations since the unital $C(X)$-representation $C(X) \to M(A)$ is a continuous field of faithful representations. Extend

the map $\pi : \mathcal{K} \otimes B = (\mathcal{K} \otimes D)/(\mathcal{K} \otimes A) \to M(\mathcal{K} \otimes A)$ to a unital morphism of $C(X)$-algebras $\tilde{\pi} : \mathcal{D}/(\mathcal{K} \otimes A) \to M(\mathcal{K} \otimes A)$. Applying assertion *1.b)* to the unital homomorphism $d \mapsto (\tilde{\pi} \circ q)(d)$ from the $C(X)$-subalgebra $\mathcal{D} \subset M(\mathcal{K} \otimes A)$ to the multiplier algebra $M(\mathcal{K} \otimes A)$ now leads to the desired conclusion. □

3 The subtriviality

Given a separable continuous field of nuclear C*-algebras A over X, the strategy to prove the subtriviality of the $C(X)$-algebra A will be the same as the one developed by Kirchberg in [15] to prove theorem 0.1 whose main ideas of demonstration are also explained in [1, Théorème 6.1]. We associate to A a $C(X)$-extension by an hereditary C*-subalgebra of the trivial continuous field $\mathcal{O}_2 \otimes C(X)$ (proposition 3.1) and then prove that after stabilisation, this $C(X)$-extension splits by $\mathcal{R}KK$-theory arguments (theorem 3.2).

3.1 Let us construct the fundamental $C(X)$-extension associated to an exact separable continuous field of C*-algebras.

Proposition 3.1 *Given a compact Hausdorff space X and a non-zero separable unital exact $C(X)$-algebra A, there exist a unital $C(X)$-subalgebra F of $\mathcal{O}_2 \otimes C(X)$ with same unit and an hereditary subalgebra I of $\mathcal{O}_2 \otimes C(X)$ such that I is an ideal in F and the $C(X)$-algebra A is isomorphic to the quotient $C(X)$-algebra F/I.*

Furthermore, if the topological space X is perfect (i.e. without any isolated point) and the $C(X)$-algebra A is continuous, the canonical map $F \to M(I)$ is a continuous field of faithful representations.

Proof : Thanks to the characterisation of separable exact C*-algebras obtained by Kirchberg (theorem 0.1), one may assume that the C*-algebra A is a C*-subalgebra of $\mathcal{O}_2$ containing the unit of $\mathcal{O}_2$.

Let $G \subset \mathcal{O}_2 \otimes C(X)$ be the trivial continuous field $A \otimes C(X)$ over X. Then the kernel of the $C(X)$-linear morphism $\pi : G \to A$ defined by $\pi(a \otimes f) = fa$ is the ideal $J = C_\Delta(X \times X)G$ where $C_\Delta(X \times X)$ is the ideal in $C(X \times X)$ of functions which are zero on the diagonal. Indeed suppose that $T \in G$ verifies $\pi(T) = 0$. Then given $\varepsilon > 0$, take a finite number of elements $a_i \in A$, $f_i \in C(X)$ such that $\|T - \sum_i a_i \otimes f_i\| < \varepsilon$; one has $\|T - \sum_i (1 \otimes f_i - f_i \otimes 1)(a_i \otimes 1)\| < \varepsilon + \|\pi(\sum_i a_i \otimes f_i)\| < 2\varepsilon$.

Define then the hereditary subalgebra $I = J[\mathcal{O}_2 \otimes C(X)]J$ in $\mathcal{O}_2 \otimes C(X)$ generated by J. It is a $C(X)$-algebra since it is closed by Cohen theorem (see e.g. [4, proposition 1.8]) and the product $(1_{\mathcal{O}_2} \otimes f)(bc) = b(1_{\mathcal{O}_2} \otimes f)c$ belongs to I for all $f \in C(X)$ and $b, c \in I$. If we set $F = I + G$, the intersection $G \cap I$ is reduced by construction to the subalgebra J, and so we have a $C(X)$-extension

$$0 \to I \to F \to A \to 0.$$

Assume now that the space X is perfect and that the $C(X)$-algebra A is continuous. We need to prove that the map $F_x \to M(I_x)$ is injective for each $x \in X$. Let $a \in G$ and $b \in I$ be two elements such that the sum $d = a + b \in F$ verifies for a given point $x \in X$ the equality

$$d_x I_x + I_x d_x = 0 \text{ (in } [\mathcal{O}_2 \otimes C(X)]_x \simeq \mathcal{O}_2 \otimes \mathbb{C}).$$

To end the proof, we have to show that d_x is zero. For every $f \in C_\Delta(X \times X)$, one has $(bf)_x = -(af)_x \in J_x$, whence $b_x \in J_x$ and so $d_x \in G_x$. But the representation of $G_x \simeq A$ in $M(J_x) \simeq M(C_x(X)A)$ is injective since X is perfect and A is continuous, from which we deduce that $d_x = 0$. $\square$

Remark: With the previous notations, if the $C(X)$-algebra A is nuclear and ψ is a unital completely positive projection from $\mathcal{O}_2$ onto A, the map $\pi \circ (\psi \otimes id_{C(X)})$ is a unital $C(X)$-linear completely positive map from $\mathcal{O}_2 \otimes C(X)$ onto the $C(X)$-subalgebra A which is zero on the nuclear hereditary $C(X)$-subalgebra I.

3.2 We can now state the main theorem:

Theorem 3.2 *Let X be a compact metrizable space and A be a unital separable $C(X)$-algebra with a unital embedding of the $C(X)$-algebra $C(X)$ in A.*

The following assertions are equivalent:

1. *A is a continuous field of nuclear* C*-*algebras over X;*
2. *there exist a unital monomorphism of $C(X)$-algebras $\alpha : A \hookrightarrow \mathcal{O}_2 \otimes C(X)$ and a unital $C(X)$-linear completely positive map $E : \mathcal{O}_2 \otimes C(X) \to A$ such that $E \circ \alpha = id_A$.*

Proof : *2⇒1* By assumption the identity map $id_A = E \circ id_{\mathcal{O}_2 \otimes C(X)} \circ \alpha : A \to A$ is nuclear since the C*-algebra $\mathcal{O}_2 \otimes C(X)$ is nuclear and so the C*-algebra A is nuclear. Besides the $C(X)$-algebra A is isomorphic to the $C(X)$-subalgebra $\alpha(A)$ of the continuous field $\mathcal{O}_2 \otimes C(X)$ and is therefore continuous.

1⇒2 • Let us first deal with the case where the space X is perfect.

Given a unital nuclear separable continuous fields A over X which is unitaly embedded in the C*-algebra $\mathcal{O}_2$, consider the $C(X)$-extension

$$0 \to I \to F \xrightarrow{\pi} A \to 0$$

constructed in proposition 3.1 and take the associated $C(X)$-extension

$$0 \to \mathcal{K} \otimes I \otimes \mathcal{O}_2 \to D = (\mathcal{K} \otimes F \otimes 1_{\mathcal{O}_2}) + (\mathcal{K} \otimes I \otimes \mathcal{O}_2) \to \mathcal{K} \otimes A \to 0.$$

The $C(X)$-nuclear quotient map $\sigma = \sigma \circ id_{\mathcal{K} \otimes A}$ from the separable nuclear continuous field $\mathcal{K} \otimes A$ to the quotient $D/(\mathcal{K} \otimes I \otimes \mathcal{O}_2) \subset M(\mathcal{K} \otimes I \otimes \mathcal{O}_2)/(\mathcal{K} \otimes I \otimes \mathcal{O}_2)$ admits a $C(X)$-linear completely positive lifting $\mathcal{K} \otimes A \to D$ ($\subset \mathcal{K} \otimes [\mathcal{O}_2 \otimes C(X)] \otimes \mathcal{O}_2$) thanks to proposition 1.3. This means that the class of σ is invertible in $Ext(X; \mathcal{K} \otimes A, I \otimes \mathcal{O}_2)$ (see the second remark following theorem 1.6).

But the group $Ext(X; \mathcal{K} \otimes A, I \otimes \mathcal{O}_2)^{-1}$ is $C(X)$-linear homotopy invariant (theorem 1.6), hence zero since the endomorphism $\phi_2(a) = s_1 a s_1^* + s_2 a s_2^*$ of $\mathcal{O}_2$ is homotopic to the identity map $id_{\mathcal{O}_2}$ ([6, proposition 2.2]) and so $[\theta] = 2[\theta]$ in $Ext(X; \mathcal{K} \otimes A, I \otimes \mathcal{O}_2)^{-1}$ for any invertible extension θ of $\mathcal{K} \otimes A$ by $I \otimes \mathcal{O}_2$. As a consequence, the $C(X)$-extension defined by σ is stably trivial. Furthermore, the identity representation of

$D \subset M(\mathcal{K} \otimes I \otimes \mathcal{O}_2)$ is a continuous field of faithful representations (proposition 3.1) and the assertion *2.* of proposition 2.5 implies that the quotient morphism $(id_{\mathcal{K}} \otimes \pi \otimes id_{\mathcal{O}_2})$ from D to $\mathcal{K} \otimes A$ admits a cross section α which is a morphism of $C(X)$-algebras.

This monomorphism α is going to enable us to conclude by standard arguments, using theorem 0.1 and the result of Elliott ([9]) that the C*-algebra $\mathcal{O}_2$ is isomorphic to $\mathcal{O}_2 \otimes \mathcal{O}_2$.

Choose a non-zero minimal projection e_{11} in the C*-algebra $\mathcal{K}$ of compact operators that we embed in $\mathcal{O}_2$ and let φ be a state on $\mathcal{O}_2$ such that $\varphi(e_{11}) = 1$. If we take a unital completely positive projection ψ of $\mathcal{O}_2$ onto the nuclear C*-subalgebra $A \subset \mathcal{O}_2$ (theorem 0.1), the composed map

$$E = (\varphi \otimes id_A) \circ \Big(id_{\mathcal{O}_2} \otimes [\pi \circ (\psi \otimes id_{C(X)})] \otimes \varphi\Big)$$

is a unital $C(X)$-linear completely positive map from $\mathcal{O}_2 \otimes [\mathcal{O}_2 \otimes C(X)] \otimes \mathcal{O}_2$ onto A. Take also an isometry $u \in \mathcal{O}_2 \otimes C(X)$ such that $\alpha(e_{11} \otimes 1_A) = uu^*$ (corollary 2.4) and define the unital $C(X)$-algebra morphism

$$\beta : A \longrightarrow \mathcal{O}_2 \otimes [\mathcal{O}_2 \otimes C(X)] \otimes \mathcal{O}_2 \simeq \mathcal{O}_2 \otimes C(X)$$

by the formula $\beta(a) = u^*\alpha(e_{11} \otimes a)u$. If $\widetilde{E} : \mathcal{O}_2 \otimes C(X) \to A$ is the completely positive unital map $d \mapsto E(udu^*)$, one gets for all $a \in A$ the equality

$$(\widetilde{E} \circ \beta)(a) = (E \circ \alpha)(e_{11} \otimes a) = a$$

• Let us now come back to the general case of a compact space X.

Define the continuous field $B = A \otimes C([0,1])$ over the perfect compact space $Y = X \times [0,1]$. According to the previous discussion, there exist a unital completely positive map $\widetilde{E} : \mathcal{O}_2 \otimes C(Y) \to B$ and a $C(X) \otimes C([0,1])$-linear monomorphism $\widetilde{\alpha} : B \to \mathcal{O}_2 \otimes C(Y)$ such that $\widetilde{E} \circ \widetilde{\alpha} = id_B$. If $ev_1 : C([0,1]) \to \mathbb{C}$ is the evaluation map at $x = 1 \in [0,1]$, define the two maps $E : \mathcal{O}_2 \otimes C(X) \to A$ and $\alpha : A \to \mathcal{O}_2 \otimes C(X)$ by

$$E(d) = (id_A \otimes ev_1) \circ \widetilde{E}(d \otimes 1_{C([0,1])}) \text{ and } \alpha(a) = (id_{\mathcal{O}_2 \otimes C(X)} \otimes ev_1) \circ \widetilde{\alpha}(a \otimes 1_{C([0,1])}).$$

Then E is a unital $C(X)$-linear completely positive map, α is a unital $C(X)$-linear monomorphism and one has the identity $E \circ \alpha = id_A$. $\square$

Remark: Assume that X is a locally compact metrizable space and that the $C_0(X)$-algebra A is a nuclear continuous field of C*-algebras over X, where $C_0(X)$ denotes the algebra of continuous functions on X vanishing at infinity. If $\widetilde{X}$ is the Alexandroff compactification of X, the unital $C(\widetilde{X})$-algebra $\mathcal{A}$ generated by A and $C(\widetilde{X})$ in the multiplier algebra $M[A \oplus C(\widetilde{X})]$ is a separable unital continuous field of C*-algebras over $\widetilde{X}$ ([4, proposition 3.2]). By theorem 3.2, there exists therefore a $C(\widetilde{X})$-linear monomorphism $\alpha : \mathcal{A} \hookrightarrow \mathcal{O}_2 \otimes C(\widetilde{X})$ and the $C_0(X)$-algebra A is isomorphic to the $C_0(X)$-subalgebra $\alpha(A)$ of $\mathcal{O}_2 \otimes C_0(X)$.

4 Concluding remarks

4.1 A $C(X)$-subalgebra of $\mathcal{O}_2 \otimes C(X)$ is by construction exact and continuous. Conversely, if A is a non-zero exact separable unital continuous field of C*-algebras over a perfect metrizable compact space X, one has by proposition 3.1 a $C(X)$-extension

$$0 \to I \to F \to A \to 0$$

where F is a $C(X)$-subalgebra of $\mathcal{O}_2 \otimes C(X)$. If the identity map $A \to A = F/I$ admits a $C(X)$-linear completely positive lifting $A \to F$, the same method as the one used in theorem 3.2 will imply that the exact continuous field A is isomorphic to a $C(X)$-subalgebra of the trivial continuous field $\mathcal{O}_2 \otimes C(X)$.

It is therefore interesting to know whether this map admits a $C(X)$-linear completely positive lifting in the not discrete case.

4.2 Let us have a look at one of the technical problems involved, the Hahn-Banach type extension property in the continuous field framework for finite type $C(X)$-submodules.

Let A be a separable unital continuous field of C*-algebras over a compact metrizable space X and let D be a finitely generated $C(X)$-submodule which is an operator system. Assume that $\phi : D \to C(X)$ is a $C(X)$-linear unital completely positive map. Then for $x \in X$, there exists, thanks to [4, proposition 3.13], a continuous field of states Φ^x on A, i.e. a $C(X)$-linear unital positive map from A to $C(X)$, such that for all $d \in D$,

$$\Phi^x(d)(x) = \phi(d)(x).$$

As a consequence, given $\varepsilon > 0$ and a finite subset $\mathcal{F}$ of D, one can build by continuity and compactness a continuous field of states Φ on A such that

$$\max\{\|\Phi(d) - \phi(d)\|, d \in \mathcal{F}\} < \varepsilon.$$

But one cannot find in general any continuous field of states on A whose restriction to D is ϕ. Indeed, consider the $C(\widetilde{\mathbb{N}})$-algebra $A = \mathbb{C}^2 \otimes C(\widetilde{\mathbb{N}})$ where $\widetilde{\mathbb{N}} = \mathbb{N} \cup \{\infty\}$ is the Alexandroff compactification of the space $\mathbb{N}$ of positive integers. Define the positive element $a \in C_\infty(\widetilde{\mathbb{N}})A \subset A$ by the formulas

$$a_n = \begin{cases} (\frac{1}{n+1}, 0) & \text{if } n \text{ even} \\ (0, \frac{1}{n+1}) & \text{if } n \text{ odd} \end{cases}$$

and let ϕ be the $C(\widetilde{\mathbb{N}})$-linear unital completely positive map with values in $C(\widetilde{\mathbb{N}})$ defined on the $C(\widetilde{\mathbb{N}})$-submodule generated by the two $C(\widetilde{\mathbb{N}})$-linearly independent elements 1_A and a through the formula

$$\phi(a)(n) = \tfrac{1}{n+1} \text{ if } n < \infty \text{ and } \phi(a)(\infty) = 0.$$

Suppose that the continuous field of states Φ is a $C(\widetilde{\mathbb{N}})$-linear extension of ϕ to A. Then as A. Bauval already noticed it, one has the contradiction

$$\begin{aligned} 1 = \Phi(1_A)(\infty) &= \Phi\left((1,0) \otimes 1\right)(\infty) + \Phi\left((0,1) \otimes 1\right)(\infty) \\ &= \lim_{n\to\infty} \Phi\left((1,0) \otimes 1\right)(2n+1) + \lim_{n\to\infty} \Phi\left((0,1) \otimes 1\right)(2n) \\ &= 0 + 0 = 0. \end{aligned}$$

Appendix *by Eberhard Kirchberg*

(Humboldt Universität zu Berlin)

In this appendix, we solve in proposition A.3 the lifting question raised in paragraph 4.1 through a continuous generalisation of joint work of E.G. Effros and U. Haagerup on lifting problems for C*-algebras ([8], see also [22]). This result enables us to state the following characterisation of separable exact continuous fields of C*-algebras:

Theorem A.1 *Let X be a compact metrizable space and A be a (unital) separable continuous field of* C*-*algebras over X.*

Then the C*-*algebra A is exact if and only if there exists a (unital) monomorphism of $C(X)$-algebras $A \hookrightarrow \mathcal{O}_2 \otimes C(X)$.*

Let us start with a technical $C(X)$-linear version of Auerbach's theorem ([17, proposition 1.c.3]) for a continuous field of C*-algebras A over X which gives us local bases over $C(X)$ with continuous coordinate maps for particular free $C(X)$-submodules of finite type in A.

Define a $C(X)$–operator system in A to be a $C(X)$-submodule which is an operator system.

Lemma A.2 *([8, lemma 2.4]) Let A be a separable unital continuous field of* C*-*algebras over a compact metrizable space X, $E \subset A$ be a $C(X)$–operator system and assume that there exists an integer $n \in \mathbb{N}^*$ such that for all $x \in X$, the dimension* $\dim E_x$ *of the operator system $E_x \subset A_x$ equals n. Then the following holds.*

Given any point $x \in X$, there exist an open neighbourhood $\mathcal{U}$ of x in X, self-adjoint $C(X)$-linear contractions $\varphi_i : A \to C(X)$ and self-adjoint elements $f_i \in E$ with $\|f_i\| \leq 2$ for $1 \leq i \leq n$ such that

$$\forall a \in C_0(\mathcal{U})E,\ a = \textstyle\sum_i \varphi_i(a) f_i.$$

Furthermore, there exists a continuous field of states $\Psi : A \to C(X)$ such that the restriction of the map $2n\Psi - id_A$ to the operator system E is completely positive.

Proof : Let us fix a point $x \in X$. Then there exist, thanks to Auerbach's theorem, a normal basis $\{r_1, \ldots, r_n\}$ of the fibre E_x where each r_i is self adjoint and norm one hermitian functionals $\phi_j : A_x \to \mathbb{C}$, $1 \leq j \leq n$, such that $\phi_j(r_i) = \delta_{i,j}$.

Consider the polar decomposition $\phi_j = \phi_j^+ - \phi_j^-$ where ϕ_j^+ and ϕ_j^- are positive functionals such that $1 = \|\phi_j\| = \|\phi_j^+\| + \|\phi_j^-\|$. By [4] lemme 3.12, there exist $C(X)$-linear positive maps φ_j^+ and $\varphi_j^- : A \to C(X)$ which extend the functionals ϕ_j^+ and ϕ_j^- on the fibre A_x to the $C(X)$-algebra A with the property that $\varphi_j^+(1) = \|\phi_j^+\|$ and $\varphi_j^-(1) = \|\phi_j^-\|$. Take also n norm 1 self-adjoint elements $e_i \in E$ satisfying the equality $(e_i)_x = r_i$ and define the matrix $T = [\varphi_j(e_i)]_{i,j} \in M_n(\mathbb{R}) \otimes C(X)$.

One has by construction $T_x = 1_{M_n(\mathbb{R})}$; the set $\mathcal{U}_1 \subset X$ of points $y \in X$ for which the spectrum of $T_y \in M_n(\mathbb{R})$ is included in the open set $\{z \in \mathbb{C}, |z| > 1/2\}$ is therefore an open neighbourhood of x in X ([4, proposition 2.4 *b)*]). If η is a continuous function on X with values in $[0,1]$ which is 0 outside $\mathcal{U}_1$ and 1 on an open neighbourhood $\mathcal{U}$ of the point $x \in X$, the self-adjoint elements $f_1, \ldots, f_n$ of norm less than 2 are then well defined in $C_0(\mathcal{U}_1)E$ by the formula

$$T\begin{pmatrix} f_1 \\ \vdots \\ f_n \end{pmatrix} = \eta \begin{pmatrix} e_1 \\ \vdots \\ e_n \end{pmatrix}$$

and satisfy the relation $\varphi_j(f_i)(y) = \delta_{i,j}$ for each $y \in \mathcal{U}$ since the matrix T_y is invertible, whence the desired equality for every $a \in C_0(\mathcal{U})E$.

Keeping the same fixed point x, define now the continuous field of states $\Phi = \frac{1}{n}\sum_i(\varphi_i^+ + \varphi_i^-)$. Then one gets for all $a \in C_0(\mathcal{U})E$ the equality:

$$(2n\Phi - id_A)(a) = \sum_{1\leq i\leq n} \Big[\varphi_i^+(a)(2-f_i) + \varphi_i^-(a)(2+f_i)\Big].$$

The restriction of the map $(2n\Phi - id_A)$ to $C_0(\mathcal{U})E$ is therefore completely positive and an appropriate partition of the unit $1_{C(X)}$ enables us to conclude. □

Noticing that a $C(X)$-linear map $\sigma : A \to B$ between $C(X)$-algebras is completely positive if and only if each induced map $\sigma_x : A_x \to B_x$ is completely positive (see for instance [4, proposition 2.9]), the lemma A.2 allows us to state a continuous version of theorem 2.5 of [8]. Replacing then the continuous field A by $A \oplus M_{2^\infty}(\mathbb{C}) \otimes C(X)$ (where $M_{2^\infty}(\mathbb{C}) = \overline{\lim_{n\to\infty} M_{2^n}(\mathbb{C})}$) and working with finitely generated $C(X)$–operator systems $E_k \subset A \oplus \bigcup_n M_{2^n}(\mathbb{C}) \otimes C(X)$ for which the function $x \mapsto \dim(E_k)_x$ is continuous, one derives the following desired $C(X)$-linear completely positive lifting result.

Proposition A.3 *([8, theorem 3.4]) Suppose that A and B are two unital separable exact continuous fields of* C*-*algebras over a compact space X with $A = B/J$ for some nuclear ideal J in B.*

Then there exists a $C(X)$-linear unital completely positive lifting $A \to B$ of id_A.

Proof : Let us define the two continuous fields $\mathcal{A} = A \oplus M_{2^\infty}(\mathbb{C}) \otimes C(X)$ and $\mathcal{B} = B \oplus M_{2^\infty}(\mathbb{C}) \otimes C(X)$. It is clearly enough to find a $C(X)$-linear unital completely positive cross section θ of the quotient morphism $\mathcal{B} \to \mathcal{A}$ (by [4, theorem 3.3]).

Consider a dense sequence $\{a_k\}$ in the self-adjoint part of $\mathcal{A}$ where each a_k belongs to the dense subalgebra $A \oplus \bigcup_n M_{2^n}(\mathbb{C}) \otimes C(X)$ of $\mathcal{A}$ and $a_1 = 1$. Let us show that we may assume inductively that $C(X)$–operator system E_n generated by the a_k, $1 \leq k \leq n$, satisfies the equality $\dim(E_n)_x = n$ for every $n \in \mathbb{N}^*$ and every $x \in X$. The inductive step is the following. Given $n \geq 2$, there exists by construction an integer l such that $E_n \subset A \oplus M_{2^l}(\mathbb{C}) \otimes C(X)$. Set $a'_n = a_n + 2^{-n-1} d_l \otimes 1_{C(X)}$ where

$$d_l = 1_{M_{2^l}(\mathbb{C})} \otimes \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix} \in M_{2^{l+1}}(\mathbb{C}) \subset M_{2^\infty}(\mathbb{C}).$$

Then the $C(X)$-module $E'_n = E_{n-1} + C(X)a'_n$ verifies for each $x \in X$ the equality $\dim(E'_n)_x = \dim(E_{n-1})_x + 1$.

Using proposition 1.3, one can now finish the proof by the same method as the one developed by E.G. Effros and U. Haagerup in [8].3 (see also [22, theorem 6.10]). □

References

[1] *C. Anantharaman-Delaroche*, Classification des C*-algèbres purement infinies [d'après E. Kirchberg], Séminaire Bourbaki **805** (1995).

[2] *A. Bauval*, $\mathcal{R}KK(X)$-nucléarité (d'après G. Skandalis), *to appear in* K-theory.

[3] *B. Blackadar*, K-theory for Operator Algebras, M.S.R.I. Publications **5**, Springer Verlag, New York (1986).

[4] *E. Blanchard*, Déformations de C*-algèbres de Hopf, Bull. Soc. Math. France **124** (1996), 141–215.

[5] *E. Blanchard*, Tensor products of $C(X)$-algebras over $C(X)$, Astérisque **232** (1995), 81–92.

[6] *J. Cuntz*, K-theory for certain C*-algebras, Ann. of Math. **113** (1981), 181–197.

[7] *J. Dixmier*, Les C*-algèbres et leurs représentations, Gauthiers-Villars Paris (1969).

[8] *E.G. Effros and U. Haagerup*, Lifting problems and local reflexivity for C*-algebras, Duke Math. J. **52** (1985), 103–128.

[9] *G. Elliott*, On the classification of C*-algebras of real rank zero, J. Reine angew. Math. **443** (1993), 179–219.

[10] *U. Haagerup and M. Rørdam*, Perturbations of the rotation C*-algebras and of the Heisenberg commutation relation, Duke Math. J. **77** (1995), 627–656.

[11] *G.G. Kasparov*, Hilbert C^*-modules: theorems of Stinespring and Voiculescu, J. Operator Theory **4** (1980), 133–150.

[12] *G.G. Kasparov*, The operator K-functor and extensions of C*-algebras, Math. U.S.S.R. Izv. **16** (1981), 513–572. Translated from Izv. Acad. Nauk S.S.S.R., Ser. Math. **44** (1980), 571–636.

[13] *G.G. Kasparov*, Equivariant KK-theory and the Novikov conjecture, Invent. Math. **91** (1988), 147–201.

[14] *G.G. Kasparov and G. Skandalis*, Groups acting on buildings, Operator K-theory and Novikov conjecture, K-theory **4** (1991), 303–337.

[15] *E. Kirchberg*, The classification of purely infinite C*-algebras using Kasparov's theory, preliminary version (3rd draft), Humboldt Universität zu Berlin (1994).

[16] *E. Kirchberg and S. Wassermann*, Operations on continuous bundles of C*-algebras, Math. Annalen **303** (1995), 677–697.

[17] *J. Lindenstrauss and L. Tzafriri*, Classical Banach Spaces I, Ergebnisse Math., Springer Verlag, 1977.

[18] *G. Skandalis*, Some remarks on Kasparov theory, J. Funct. Anal. **56** (1984), 337–347.

[19] *G. Skandalis*, Kasparov's bivariant K-theory and applications, Expo. Math. **9** (1991), 193–250.

[20] *G. Skandalis*, Une notion de nucléarité en K-théorie (d'après J. Cuntz), K-theory **1** (1988), 549–573.

[21] *D. Voiculescu*, A non-commutative Weyl-von Neumann theorem, Rev. Roum. Math. Pures et Appl. **21** (1976), 97–113.

[22] *S. Wassermann*, Exact C*-algebras and Related Topics, Lecture Notes Series 19, GARC, Seoul National University, 1994.

Inst. Math. Luminy, CNRS–Luminy case 930, F–13288 Marseille CEDEX 9
E-mail: E.Blanchard@iml.univ-mrs.fr