



**Mathematical renormalization in quantum
electrodynamics via noncommutative generating series**
Gérard Henry Edmond Duchamp, Vincel Hoang Ngoc Minh, Quoc Hoan Ngo,
Karol A. Penson, Pierre Simonnet

► **To cite this version:**

Gérard Henry Edmond Duchamp, Vincel Hoang Ngoc Minh, Quoc Hoan Ngo, Karol A. Penson, Pierre Simonnet. Mathematical renormalization in quantum electrodynamics via noncommutative generating series. 2014. hal-00927641v1

HAL Id: hal-00927641
<https://hal.science/hal-00927641v1>

Preprint submitted on 13 Jan 2014 (v1), last revised 23 Feb 2017 (v2)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Mathematical renormalization in quantum electrodynamics via noncommutative generating series¹

G.H.E. Duchamp^{5,2}, V. Hoang Ngoc Minh^{1,2},
Q.H. Ngô², K. Penson³, P. Simonnet⁴.

¹Université Lille II, 1, Place Déliot, 59024 Lille, France.

²LIPN - UMR 7030, CNRS, 93430 Villetaneuse, France.

³Université Paris VI - LPTMC, 4, Place Jussieu, 75252 Paris Cedex 05, France.

⁴Université de Corse, Avenue Jean Nicoli, BP 52, 20250 Corte, France.

⁵Université Paris XVIII, 93430 Villetaneuse, France.

Abstract

In this work, we focus on the approach by noncommutative formal power series to study the combinatorial aspects of the renormalization at the singularities in $\{0, 1, +\infty\}$ of the solutions of nonlinear differential equations involved in quantum electrodynamics.

Contents

1	Introduction	2
2	Background	6
2.1	Combinatorics on shuffle and stuffle Hopf algebras	6
2.1.1	Schützenberger’s monoidal factorization	6
2.1.2	Extended Schützenberger’s monoidal factorization	9
2.2	Indiscernability over a class of formal power series	15
2.2.1	Residual calculus and representative series	15
2.2.2	Continuity and indiscernability	17
2.3	Polylogarithms and harmonic sums	18
2.3.1	Structures of polylogarithms and harmonic sums	18
2.3.2	Results <i>à la Abel</i> for noncommutative generating and global regularizations	20
3	Polysystem and differential realization	25
3.1	Polysystem and convergence criterion	25
3.1.1	Serial estimates from above	25
3.1.2	Upper bounds <i>à la Cauchy</i>	28
3.2	Polysystem and nonlinear differential equation	30
3.2.1	Nonlinear differential equation (with three singularities)	30
3.2.2	Asymptotic behavior of the successive differentiation of solution via extended Fliess fundamental formula	31

¹Version du 13-01-2014 13:52

3.3	Differential realization	33
3.3.1	Differential realization	33
3.3.2	Fliess' local realization theorem	35

1 Introduction

During the last century, the functional expansions were common in physics as well as in engineering and have been developped, by Tomonaga, Schwinger and Feynman [16], to represent the nonlinear dynamical systems in quantum electrodynamics. The main difficulty of this approach is the divergence of these expansions at the singularity 0 or at $+\infty$ (see [2]) and leads to the problems of *regularization* and *renormalization* which can be solved by combinatorial techniques : Feynman diagrams [18] and their derivatives [14, 23], noncommutative formal power series [19, 20], trees [10],

Recently, following the previous vein, and basing on

- the shuffle and quasi-shuffle Hopf algebras [5], the combinatorics on noncommutative formal power series were intensively amplified for the asymptotic analysis of nonlinear dynamical systems with three regular singularities in² $\{0, 1, +\infty\}$ [39, 40],
- the monodromy and the Galois differential groups of the Knizhnik-Zamolodchikov differential equation KZ_3 [38, 40], *i.e.* the following noncommutative evolution equation³

$$\frac{dG(z)}{dz} = \left(\frac{x_0}{z} + \frac{x_1}{1-z} \right) G(z), \quad (1)$$

the Schützenberger's monoidal factorization facilitates mainly the renormalization and the computation of the associators⁴ via the universal one, *i.e.* the associator Φ_{KZ} of Drinfel'd.

²Any differential equation with three singularities in $\{a, b, c\}$, via homographic transformation $(z-a)(c-b)(z-b)^{-1}(c-a)^{-1}$ can be changed into a differential equation with three singularities in $\{0, 1, +\infty\}$.

³Here, x_0 and x_1 are noncommutative variables. More precisely (with $i^2 = -1$),

$$x_0 := \frac{t_{1,2}}{2i\pi} \quad \text{and} \quad x_1 := -\frac{t_{2,3}}{2i\pi},$$

where $t_{1,2}$ and $t_{2,3}$ belong to $\mathcal{T}_3 = \{t_{1,2}, t_{1,3}, t_{2,3}\}$ satisfying the infinitesimal 3-braid relations

$$[t_{1,3}, t_{1,2} + t_{2,3}] = [t_{2,3}, t_{1,2} + t_{1,3}] = 0.$$

⁴They were introduced in quantum field theory by Drinfel'd and it plays an important role for the still open problem of the effective determination of the polynomial invariants of knots and links via Kontsevich's integral (see [6, 23]) and Φ_{KZ} , was obtained firstly, in [23], with explicit coefficients which are polyzetas and regularized polyzetas (see [40, 41] for the computation of the other associators involving *only* convergent polyzetas as local coordinates, and for three algorithmical process to regularize the divergent polyzetas).

In fact, these associators are noncommutative formal power series on two variables and regularize the Chen generating series of the differential forms admitting singularities at 0 or at 1 along the integration paths on the universal covering $\widehat{\mathbb{C} - \{0, 1\}}$ of \mathbb{C} without points 0 and 1. Their coefficients which are, up to multiple of power of $2i\pi$, polynomial on polyzetas, *i.e.* the following real numbers⁵ (for $r \geq 1, s_1 \geq 2$ and $s_2, \dots, s_r \geq 1$) [42, 23, 51]

$$\zeta(s_1, \dots, s_r) = \sum_{n_1 > \dots > n_r > 0} \frac{1}{n_1^{s_1} \dots n_r^{s_r}} \quad (2)$$

and it is reasonable to think that these numbers admit a structure of \mathbb{N} -graded, by weight, algebra over the rational numbers deduced from the combinatorial aspects of the shuffle and quasi-shuffle Hopf algebras [40, 41].

More precisely, for $s_1 \geq 2, s_2, \dots, s_r \geq 1$, the polyzeta $\zeta(s_1, \dots, s_r)$ can be obtained as the limit of

- the polylogarithm $\text{Li}_{s_1, \dots, s_r}(z)$, for $z \rightarrow 1$ (recalled in Section 2.3.1)

$$\text{Li}_{s_1, \dots, s_r}(z) = \sum_{n_1 > \dots > n_r > 0} \frac{z^{n_1}}{n_1^{s_1} \dots n_r^{s_r}} \quad (3)$$

- the harmonic sum $\text{H}_{s_1, \dots, s_r}(N)$, for $N \rightarrow +\infty$ (recalled in Section 2.3.1)

$$\text{H}_{s_1, \dots, s_r}(N) = \sum_{n_1 > \dots > n_r > 0}^N \frac{1}{n_1^{s_1} \dots n_r^{s_r}}. \quad (4)$$

Then, by a theorem of Abel, one has

$$\zeta(s_1, \dots, s_r) = \lim_{z \rightarrow 1} \text{Li}_{s_1, \dots, s_r}(z) = \lim_{N \rightarrow +\infty} \text{H}_{s_1, \dots, s_r}(N). \quad (5)$$

Since the algebras of polylogarithms and of harmonic sums are isomorphic respectively to the shuffle algebra $(\mathbb{Q}\langle X \rangle, \sqcup)$ and the quasi-shuffle algebra $(\mathbb{Q}\langle Y \rangle, \boxplus)$ both admitting the Lyndon words $\mathcal{Lyn}X$ over $X = \{x_0, x_1\}$ and $\mathcal{Lyn}Y$ over $Y = \{y_i\}_{i \geq 1}$, as transcendence bases (recalled in Section 2.1) then, by using

- the correspondence between the composition in \mathbb{N}_+ with the words in Y^* and in X^*x_1 , *i.e.*

$$(s_1, \dots, s_r) \longleftrightarrow y_{s_1} \dots y_{s_r} \longleftrightarrow x_0^{s_1-1} x_1 \dots x_0^{s_r-1} x_1, \quad (6)$$

- the non commutative generating series in the factorized forms, *i.e.*

$$\text{L} = \prod_{l \in \mathcal{Lyn}X} \exp(\text{Li}_{S_l} P_l) \quad \text{and} \quad \text{H} = \prod_{l \in \mathcal{Lyn}Y} \exp(\text{H}_{\Sigma_l} \Pi_l), \quad (7)$$

$$\text{Z}_{\sqcup} = \prod_{\substack{l \in \mathcal{Lyn}X \\ l \neq x_0, x_1}} \exp(\zeta(S_l) P_l) \quad \text{and} \quad \text{Z}_{\boxplus} = \prod_{\substack{l \in \mathcal{Lyn}Y \\ l \neq y_1}} \exp(\zeta(\Sigma_l) \Pi_l), \quad (8)$$

⁵The sum $s_1 + \dots + s_r$ is the *weight* of $\zeta(s_1, \dots, s_r)$ which equals to the weight of the composition (s_1, \dots, s_r) .

where $\{S_l\}_{l \in \mathcal{L}_{yn}X}$ and $\{\Sigma_l\}_{l \in \mathcal{L}_{yn}Y}$ are transcendence bases of $(\mathbb{Q}\langle X \rangle, \sqcup)$ and $(\mathbb{Q}\langle Y \rangle, \sqcup)$, $\{P_l\}_{l \in \mathcal{L}_{yn}X}$ and $\{\Pi_l\}_{l \in \mathcal{L}_{yn}Y}$ are bases of the Lie algebras of the primitive elements of the bi-algebras $\mathcal{H}_{\sqcup} = (\mathbb{Q}\langle X \rangle, \text{conc}, \Delta_{\sqcup}, 1_{X^*})$ and $\mathcal{H}_{\sqcup} = (\mathbb{Q}\langle Y \rangle, \text{conc}, \Delta_{\sqcup}, 1_{Y^*})$ in duality respectively with $\{S_l\}_{l \in \mathcal{L}_{yn}X}$ and $\{\Sigma_l\}_{l \in \mathcal{L}_{yn}Y}$,

we obtain the Abel like theorem [39, 40, 41], *i.e.* there exist two noncommutative formal series over Y , Z_1 and Z_2 with constant terms, such that :

$$\lim_{z \rightarrow 1} \exp\left(y_1 \log \frac{1}{1-z}\right) \pi_Y L(z) = Z_1, \quad (9)$$

$$\lim_{N \rightarrow \infty} \exp\left(\sum_{k \geq 1} H_{y_k}(N) \frac{(-y_1)^k}{k}\right) H(N) = Z_2. \quad (10)$$

Moreover, Z_1, Z_2 are equal and stand for the noncommutative generating series of the convergent polyzetas $\{\zeta(w)\}_{w \in Y^* - y_1 Y^*}$, or equivalently $\{\zeta(w)\}_{w \in x_0 X^* x_1}$:

$$Z_1 = Z_2 = \pi_Y Z_{\sqcup}, \quad (11)$$

where π_Y denotes the projection over Y^* .

This Abel theorem allows to explicit the counter-terms eliminating the divergence of

- the polylogarithms $\{Li_w(z)\}_{w \in x_1 X^*}$ for $z \rightarrow 1$,
- the harmonic sums $\{H_w(N)\}_{w \in y_1 Y^*}$ for $N \rightarrow \infty$

and leads naturally to an equation connecting these two algebraic structures

$$\prod_{\substack{l \in \mathcal{L}_{yn}Y \\ l \neq y_1}}^{\searrow} \exp(\zeta(\Sigma_l) \Pi_l) = \exp\left(\sum_{k \geq 2} \zeta(k) \frac{(-y_1)^k}{k}\right) \pi_Y \prod_{\substack{l \in \mathcal{L}_{yn}X \\ l \neq x_0, x_1}}^{\searrow} \exp(\zeta(S_l) P_l). \quad (12)$$

This identity (12) enables, in particular,

- to calculate the Euler-Mac Laurin constants and the Hadamard finite parties associated to the divergent polyzetas $\{\zeta(w)\}_{w \in y_1 Y^*}$,
- to describe *completely*, via identification of local coordinates in infinite dimension, the kernel of ζ by its *algebraic* generators, and thus this kernel is generated by homogenous polynomials.

In this paper, we will focus on the approach by noncommutative formal power series, adapted from [19, 20], and explaining the results of [39, 40, 41] allowing to study the combinatorial aspects of the renormalization at the singularities in $\{0, 1, +\infty\}$ of the solutions of linear differential equations (see Example 1 bellow) as well as the solutions of nonlinear differential equations (see Examples 2 and 3 bellow) described in Section 3.2 and involved in quantum electrodynamics.

Example 1 (Hypergeometric equation). *Let t_0, t_1, t_2 be parameters and*

$$z(1-z)\ddot{y}(z) + [t_2 - (t_0 + t_1 + 1)z]\dot{y}(z) - t_0 t_1 y(z) = 0.$$

Let $q_1(z) = y(z)$ and $q_2(z) = z(1-z)\dot{y}(z)$. One has

$$\begin{pmatrix} \dot{q}_1 \\ \dot{q}_2 \end{pmatrix} = \left(\frac{M_0}{z} + \frac{M_1}{1-z} \right) \begin{pmatrix} q_1 \\ q_2 \end{pmatrix},$$

where M_0 and M_1 are the following matrices

$$M_0 = - \begin{pmatrix} 0 & 0 \\ t_0 t_1 & t_2 \end{pmatrix} \quad \text{and} \quad M_1 = - \begin{pmatrix} 0 & 1 \\ 0 & t_2 - t_0 - t_1 \end{pmatrix}.$$

Or equivalently,

$$\begin{aligned} \dot{q}(z) &= A_0(q) \frac{1}{z} + A_1(q) \frac{1}{1-z}, \\ y(z) &= q(z), \end{aligned}$$

where A_0 and A_1 are the following parameterized linear vector fields

$$A_0(q) = -(t_0 t_1 q_1 + t_2 q_2) \frac{\partial}{\partial q_2} \quad \text{and} \quad A_1(q) = -q_1 \frac{\partial}{\partial q_1} - (t_2 - t_0 - t_1) q_2 \frac{\partial}{\partial q_2}.$$

Example 2 (Harmonic oscillator). *Let k_1, k_2 be parameters and*

$$\ddot{y}(z) + k_1 y(z) + k_2 y^2(z) = u_1(z).$$

which can be represented by the following state equations

$$\begin{aligned} \dot{q}(z) &= A_0(q) + A_1(q) u_1(z), \\ y(z) &= q(z), \end{aligned}$$

where A_0 and A_1 are the following vector fields

$$A_0(q) = -(k_1 q + k_2 q^2) \frac{\partial}{\partial q} \quad \text{and} \quad A_1(q) = \frac{\partial}{\partial q}.$$

Example 3 (Duffing's equation). *Let a, b, c be parameters and*

$$\ddot{y}(z) + a\dot{y}(z) + by(z) + cy^3(z) = u_1(z).$$

which can be represented by the following state equations

$$\begin{aligned} \dot{q}(z) &= A_0(q) + A_1(q) u_1(z), \\ y(z) &= q(z), \end{aligned}$$

where A_0 and A_1 are the following vector fields

$$A_0(q) = -(aq_2 + b^2 q_1 + cq_1^3) \frac{\partial}{\partial q_2} + q_2 \frac{\partial}{\partial q_1} \quad \text{and} \quad A_1(q) = \frac{\partial}{\partial q_2}.$$

This approach by noncommutative formal power series is adequate for studying the algebraic combinatorial aspects of the asymptotic analysis, at the singularities in $\{0, 1, +\infty\}$, of the nonlinear dynamical systems described in Section 3.2 because

- the polylogarithms form the universal Picard-Vessiot of these nonlinear differential equations [38] and their algebra is isomorphic to the shuffle algebra admitting $\{Li_l\}_{l \in \mathcal{L}yn X}$ as transcendence basis,
- the harmonic sums generate the coefficients of the ordinary Taylor expansion of their solutions (when these expansion exist) [39] and their algebra is isomorphic to the shuffle algebra admitting $\{H_l\}_{l \in \mathcal{L}yn Y}$ as transcendence basis,
- the polyzetas do appear as the fundamental arithmetical constants involved in the computations of the monodromies [35, 32], the Kummer type functional equations [36, 32], the asymptotic expansions of the solutions of these equations [39, 40] and their algebra is freely generated by the polyzetas encoded by irreducible Lyndon words [40].

Hence, a lot of algorithms can be deduced from these facts and the more general studies will be proceed (see [5, 11] for example).

The organisation of this paper is following

- In Section 2, we will give algebraic and analytic foundations, *i.e.* the combinatorial Hopf algebra of shuffles and the indiscernability respectively, for polylogarithms and harmonic sums.
- These will be exploited systematically, in Section 3, to expand the solutions of nonlinear dynamical systems with singular inputs.

2 Background

2.1 Combinatorics on shuffle and stuffle Hopf algebras

2.1.1 Schützenberger’s monoidal factorization

Let X be a finite totally ordered alphabet. The free monoid and the set of Lyndon words, over X , are denoted respectively by X^* and $\mathcal{L}yn X$. The neutral element of X^* is denoted by 1_{X^*} . Let $\mathbb{Q}\langle X \rangle$ be equipped by the concatenation and the shuffle defined by

$$\forall w \in X^*, \quad w \sqcup 1_{X^*} = 1_{X^*} \sqcup w = w, \quad (13)$$

$$\forall x, y \in X, \forall u, v \in X^*, \quad xu \sqcup yv = x(u \sqcup yv) + y(xu \sqcup v), \quad (14)$$

or by their dual co-products, $\Delta_{\text{conc}}, \Delta_{\sqcup}$, defined by

$$\forall w \in X^*, \quad \Delta_{\text{conc}}(w) = \sum_{u, v \in X^*, uv=w} u \otimes v, \quad (15)$$

$$\forall x \in X, \quad \Delta_{\sqcup}(x) = x \otimes 1 + 1 \otimes x, \quad (16)$$

satisfying, for any $u, v, w \in X^*$,

$$\langle \Delta_{\text{conc}}(w) \mid u \otimes v \rangle = \langle w \mid uv \rangle, \quad (17)$$

$$\langle \Delta_{\sqcup}(w) \mid u \otimes v \rangle = \langle w \mid u \sqcup v \rangle. \quad (18)$$

With the co-unit and the antipode defined by

$$\forall P \in \mathbb{Q}\langle X \rangle, \quad \epsilon(P) = \langle P \mid 1_{X^*} \rangle, \quad (19)$$

$$\forall w = x_{i_1} \dots x_{i_r} \in X^*, \quad a_{\sqcup}(w) = (-1)^{-|w|} x_{i_r} \dots x_{i_1}, \quad (20)$$

one gets two mutually dual Hopf algebras

$$\mathcal{H}_{\sqcup} = (\mathbb{Q}\langle X \rangle, \text{conc}, 1_{X^*}, \Delta_{\sqcup}, \epsilon, a_{\sqcup}), \quad (21)$$

$$\mathcal{H}_{\sqcup}^{\vee} = (\mathbb{Q}\langle X \rangle, \sqcup, 1_{X^*}, \Delta_{\text{conc}}, \epsilon, a_{\sqcup}). \quad (22)$$

By the Cartier-Quillen-Milnor-Moore theorem, the connected graded positively, co-commutative Hopf algebra \mathcal{H} is isomorphic to the envelopping algebra of the Lie algebra of its primitive elements which is equal to $\mathcal{L}ie_{\mathbb{Q}}\langle X \rangle$. Hence, any graded⁶ suited⁷ basis of $\mathcal{L}ie_{\mathbb{Q}}\langle X \rangle$, can be completed, by the Poincaré-Birkhoff-Witt theorem, as a linear (graded) basis $\{b_w\}_{w \in X^*}$ for $\mathcal{U}(\mathcal{L}ie_{\mathbb{Q}}\langle X \rangle)$. One can then⁸ construct, by duality, a basis $\{\check{b}_w\}_{w \in X^*}$ of \mathcal{H}_{\sqcup} such that :

$$\forall u, v \in X^*, \quad \langle \check{b}_u \mid b_v \rangle = \delta_{u,v}, \quad (23)$$

and, for any $x \in X$, $\check{b}_x = x$ and for any $w = l_1^{i_1} \dots l_k^{i_k}$ with $l_1, \dots, l_k \in \mathcal{L}ynX$ and $l_1 > \dots > l_k$, one has

$$\check{b}_w = \frac{\check{b}_{l_1}^{i_1} \times \dots \times \check{b}_{l_k}^{i_k}}{i_1! \dots i_k!}. \quad (24)$$

For that, one can use a theorem of Viennot [50] to construct, by bijection with $\mathcal{L}ynX$, a basis for $\mathcal{L}ie_{\mathbb{Q}}\langle X \rangle$ and, by the PBW theorem, one completes then to a linear basis for $\mathcal{U}(\mathcal{L}ie_{\mathbb{Q}}\langle X \rangle)$.

On an other way, by a Radford's theorem [45], the elements of $\mathcal{L}ynX$ constitute also a transcendence basis for $(\mathbb{Q}\langle X \rangle, \sqcup)$ and it can be completed then to the linear basis $\{w\}_{w \in X^*}$ which is auto-dual :

$$\forall v, v \in X^*, \quad \langle u \mid v \rangle = \delta_{u,v}. \quad (25)$$

But the elements $l \in \mathcal{L}ynX - X$ are not primitive, for Δ_{\sqcup} , and then $\mathcal{L}ynX$ does not constitute a basis for $\mathcal{L}ie_{\mathbb{Q}}\langle X \rangle$. Denoting \mathcal{D}_X the diagonal series over X , one has

$$\mathcal{D}_X = \sum_{w \in X^*} w \otimes w = \sum_{l \in \mathcal{L}ynX} l \otimes l + \sum_{w \notin \mathcal{L}ynX} w \otimes w \quad (26)$$

⁶for the multidegree.

⁷i.e. a basis of polynomials indexed by their initial words which is a Lyndon word.

⁸because \mathcal{H} is graded in finite dimensions by the multidegree.

and \mathcal{D}_X can be also factorized as follows

$$\mathcal{D}_X = \prod_{l \in \mathcal{L}yn X}^{\searrow} \exp(l \otimes l). \quad (27)$$

Using the transcendence basis $\{l\}_{l \in \mathcal{L}yn X}$, Mélançon and Reutenauer give an other completion by putting [47]

$$S'_l = l, \quad \text{for } l \in \mathcal{L}yn X, \quad (28)$$

$$S'_w = \frac{l_1^{\sqcup i_1} \sqcup \dots \sqcup l_k^{\sqcup i_k}}{i_1! \dots i_k!}, \quad \text{for } w = l_1^{i_1} \dots l_k^{i_k}, l_1 > \dots > l_k, \quad (29)$$

$$= w + \sum_{v < w, |v|=|w|} c'_v v. \quad (30)$$

The linear basis $\{S'_w\}_{w \in X^*}$ admits the dual basis $\{P'_w\}_{w \in X^*}$ but it is not of PBW type although the elements of $\{P'_l\}_{l \in \mathcal{L}yn X}$ are primitive, for Δ_{\sqcup} [12].

Chen, Fox and Lyndon [8] constructed $\{P_w\}_{w \in X^*}$, so-called the PBW-Lyndon basis, for $\mathcal{U}(\mathcal{L}ie_{\mathbb{Q}}\langle X \rangle)$ as follows

$$P_x = x \quad \text{for } x \in X, \quad (31)$$

$$P_l = [P_s, P_r] \quad \text{for } l \in \mathcal{L}yn X, \text{ standard factorization of } l = (s, r), \quad (32)$$

$$P_w = P_{l_1}^{i_1} \dots P_{l_k}^{i_k} \quad \text{for } w = l_1^{i_1} \dots l_k^{i_k}, l_1 > \dots > l_k, l_1, \dots, l_k \in \mathcal{L}yn X. \quad (33)$$

Schützenberger constructed bases for $(\mathbb{Q}\langle X \rangle, \sqcup)$ defined by duality as follows :

$$\forall u, v \in X^*, \quad \langle S_u \mid P_v \rangle = \delta_{u,v}, \quad (34)$$

and obtained the transcendence basis $\{S_l\}_{l \in \mathcal{L}yn X}$ and the linear basis $\{S_w\}_{w \in X^*}$ as follows

$$S_l = x S_u, \quad \text{for } l = xu \in \mathcal{L}yn X, \quad (35)$$

$$S_w = \frac{S_{l_1}^{\sqcup i_1} \sqcup \dots \sqcup S_{l_k}^{\sqcup i_k}}{i_1! \dots i_k!} \quad \text{for } w = l_1^{i_1} \dots l_k^{i_k}, l_1 > \dots > l_k. \quad (36)$$

After that, Mélançon and Reutenauer [47] proved that⁹, for any $w \in X^*$,

$$P_w = w + \sum_{v > w, |v|=|w|} c_v v \quad \text{and} \quad S_w = w + \sum_{v < w, |v|=|w|} c_v v. \quad (37)$$

⁹ Recall that the duality preserves the (multi)-homogeneous degree and interchanges the triangularity of polynomials [47]. For that, one can construct the triangular matrices M and N admitting coefficients as the coefficients of the homogeneous of degree k triangular polynomials, $\{P_w\}_{w \in X^*}$ and $\{S_w\}_{w \in X^*}$ in the basis $\{w\}_{w \in X^*}$ respectively :

$$M_{u,v} = \langle P_u \mid v \rangle \quad \text{and} \quad N_{v,u} = \langle S_u \mid v \rangle.$$

The triangular matrices M and N are unipotent and satisfy the identity $N = ({}^t M)^{-1}$.

In other words, the elements of the bases $\{S_w\}_{w \in X^*}$ and $\{P_w\}_{w \in X^*}$ are lower and upper triangular respectively and they are of homogeneous by degree.

Thanks to the duality of the bases $\{P_w\}_{w \in X^*}$ and $\{S_w\}_{w \in X^*}$ and the basis $\{P_w\}_{w \in X^*}$ is of PBW type, one also has [47]

$$\mathcal{D}_X = \sum_{w \in X^*} S_w \otimes P_w \quad (38)$$

$$= \sum_{l \in \mathcal{Lyn} X} S_l \otimes P_l + \sum_{w \notin \mathcal{Lyn} X} S_w \otimes P_w \quad (39)$$

$$= \prod_{l \in \mathcal{Lyn} X}^{\searrow} \exp(S_l \otimes P_l). \quad (40)$$

Example 4 ([27]). Let $X = \{x_0, x_1\}$ with $x_0 < x_1$.

l	P_l	S_l
x_0	x_0	x_0
x_1	x_1	x_1
$x_0 x_1$	$[x_0, x_1]$	$x_0 x_1$
$x_0^2 x_1$	$[x_0, [x_0, x_1]]$	$x_0^2 x_1$
$x_0 x_1^2$	$[[x_0, x_1], x_1]$	$x_0 x_1^2$
$x_0^3 x_1$	$[x_0, [x_0, [x_0, x_1]]]$	$x_0^3 x_1$
$x_0^2 x_1^2$	$[x_0, [[x_0, x_1], x_1]]$	$x_0^2 x_1^2$
$x_0 x_1^3$	$[[[x_0, x_1], x_1], x_1]$	$x_0 x_1^3$
$x_0^4 x_1$	$[x_0, [x_0, [x_0, [x_0, x_1]]]]$	$x_0^4 x_1$
$x_0^3 x_1^2$	$[x_0, [x_0, [[x_0, x_1], x_1]]]$	$x_0^3 x_1^2$
$x_0^2 x_1 x_0 x_1$	$[[x_0, [x_0, x_1]], [x_0, x_1]]$	$2x_0^3 x_1^2 + x_0^2 x_1 x_0 x_1$
$x_0^2 x_1^3$	$[x_0, [[[x_0, x_1], x_1], x_1]]$	$x_0^2 x_1^3$
$x_0 x_1 x_0 x_1^2$	$[[x_0, x_1], [[x_0, x_1], x_1]]$	$3x_0^2 x_1^3 + x_0 x_1 x_0 x_1^2$
$x_0 x_1^4$	$[[[[x_0, x_1], x_1], x_1], x_1]$	$x_0 x_1^4$
$x_0^5 x_1$	$[x_0, [x_0, [x_0, [x_0, [x_0, x_1]]]]]$	$x_0^5 x_1$
$x_0^4 x_1^2$	$[x_0, [x_0, [x_0, [[x_0, x_1], x_1]]]]$	$x_0^4 x_1^2$
$x_0^3 x_1 x_0 x_1$	$[x_0, [[x_0, [x_0, x_1]], [x_0, x_1]]]$	$2x_0^4 x_1^2 + x_0^3 x_1 x_0 x_1$
$x_0^3 x_1^3$	$[x_0, [x_0, [[[x_0, x_1], x_1], x_1]]]$	$x_0^3 x_1^3$
$x_0^2 x_1 x_0 x_1^2$	$[x_0, [[x_0, x_1], [[x_0, x_1], x_1]]]$	$3x_0^3 x_1^3 + x_0^2 x_1 x_0 x_1^2$
$x_0^2 x_1^2 x_0 x_1$	$[[x_0, [[x_0, x_1], x_1]], [x_0, x_1]]$	$6x_0^3 x_1^3 + 3x_0^2 x_1 x_0 x_1^2 + x_0^2 x_1^2 x_0 x_1$
$x_0^2 x_1^4$	$[x_0, [[[[x_0, x_1], x_1], x_1], x_1]]$	$x_0^2 x_1^4$
$x_0 x_1 x_0 x_1^3$	$[[x_0, x_1], [[[x_0, x_1], x_1], x_1]]$	$4x_0^2 x_1^4 + x_0 x_1 x_0 x_1^3$
$x_0 x_1^5$	$[[[[[[x_0, x_1], x_1], x_1], x_1], x_1]]]$	$x_0 x_1^5$

2.1.2 Extended Schützenberger's monoidal factorization

Now, let $Y = \{y_i\}_{i \geq 1}$ be an infinite totally ordered alphabet¹⁰. The free monoid and the set of Lyndon words, over Y , are denoted respectively by Y^* and $\mathcal{Lyn} Y$. The neutral element of Y^* is denoted by 1_{Y^*} .

¹⁰by $y_1 > y_2 > y_3 > \dots$

Let $u = y_{i_1} \dots y_{i_k} \in Y^*$, the *length* and the *weight* of u are defined respectively as the numbers $|u| = k$ and $(u) = i_1 + \dots + i_k$.

Let us define the commutative product over $\mathbb{Q}\langle Y \rangle$, denoted by μ , as follows

$$\forall y_n, y_m \in Y, \quad \mu(y_n, y_m) = y_{n+m}, \quad (41)$$

or by its associated coproduct, Δ_+ , defined by

$$\forall y_n \in Y, \quad \Delta_+ y_n = \sum_{i=1}^{n-1} y_i \otimes y_{n-i} \quad (42)$$

satisfying,

$$\forall x, y, z \in Y, \quad \langle \Delta_+ x \mid y \otimes z \rangle = \langle x \mid \mu(y, z) \rangle. \quad (43)$$

Let $\mathbb{Q}\langle Y \rangle$ be equipped by

1. The concatenation (or by its associated coproduct, Δ_{conc}).
2. The *shuffle* product, *i.e.* the commutative product defined by [47]

$$\begin{aligned} \forall w \in Y^*, \quad w \sqcup 1_{Y^*} &= 1_{Y^*} \sqcup w = w, \\ \forall x, y \in Y, \forall u, v \in Y^*, \quad xu \sqcup yv &= x(u \sqcup yv) + y(xu \sqcup v) \end{aligned}$$

or by its associated coproduct, Δ_{\sqcup} , defined, on the letters, by

$$\forall y_k \in Y, \quad \Delta_{\sqcup} y_k = y_k \otimes 1 + 1 \otimes y_k$$

and extended by morphism. It satisfies

$$\forall u, v, w \in Y^*, \quad \langle \Delta_{\sqcup} w \mid u \otimes v \rangle = \langle w \mid u \sqcup v \rangle.$$

3. The *quasi-shuffle* product, *i.e.* the commutative product defined by [43], for any $w \in Y^*$,

$$w \boxplus 1_{Y^*} = 1_{Y^*} \boxplus w = w,$$

and, for any $y_i, y_j \in Y, \forall u, v \in Y^*$,

$$\begin{aligned} y_i u \boxplus y_j v &= y_j(y_i u \boxplus v) + y_i(u \boxplus y_j v) + \mu(y_i, y_j)(u \boxplus v), \\ &= y_j(y_i u \boxplus v) + y_i(u \boxplus y_j v) + y_{i+j}(u \boxplus v) \end{aligned}$$

or by its associated coproduct, Δ_{\boxplus} , defined, on the letters, by

$$\forall y_k \in Y, \quad \Delta_{\boxplus} y_k = \Delta_{\sqcup} y_k + \Delta_+ y_k$$

and extended by morphism. It satisfies

$$\forall u, v, w \in Y^*, \quad \langle \Delta_{\boxplus} w \mid u \otimes v \rangle = \langle w \mid u \boxplus v \rangle.$$

Note that Δ_{\boxplus} and Δ_{\sqcup} are morphisms for the concatenation (by definition) whereas Δ_+ is not a morphism for the product of $\mathbf{k}Y$ (for example $\Delta_+(y_1^2) = y_1 \otimes y_1$, whereas $\Delta_+(y_1)^2 = 0$).

Hence, with the counit \mathbf{e} defined by

$$\forall P \in \mathbb{Q}\langle Y \rangle, \quad \mathbf{e}(P) = \langle P \mid 1_{Y^*} \rangle, \quad (44)$$

one gets two pairs of mutually dual bialgebras

$$\mathcal{H}_{\sqcup} = (\mathbb{Q}\langle Y \rangle, \text{conc}, 1_{Y^*}, \Delta_{\sqcup}, \mathbf{e}) \quad \text{and} \quad \mathcal{H}_{\sqcup}^{\vee} = (\mathbb{Q}\langle Y \rangle, \sqcup, 1_{Y^*}, \Delta_{\text{conc}}, \mathbf{e}) \quad (45)$$

$$\mathcal{H}_{\sqcup} = (\mathbb{Q}\langle Y \rangle, \text{conc}, 1_{Y^*}, \Delta_{\sqcup}, \mathbf{e}) \quad \text{and} \quad \mathcal{H}_{\sqcup}^{\vee} = (\mathbb{Q}\langle Y \rangle, \sqcup, 1_{Y^*}, \Delta_{\text{conc}}, \mathbf{e}) \quad (46)$$

As in (26-27), one has

$$\mathcal{D}_Y = \sum_{w \in Y^*} w \otimes w \quad (47)$$

$$= \sum_{l \in \mathcal{L}yn Y} l \otimes l + \sum_{w \notin \mathcal{L}yn Y} w \otimes w \quad (48)$$

$$= \prod_{l \in \mathcal{L}yn Y}^{\times} \exp(l \otimes l). \quad (49)$$

By the CQMM theorem (see [5]), the connected \mathbb{N} -graded, co-commutative Hopf algebra \mathcal{H}_{\sqcup} is isomorphic to the enveloping algebra of the Lie algebra of its primitive elements which is equal to $\mathcal{L}ie_{\mathbb{Q}}\langle Y \rangle$:

$$\mathcal{H}_{\sqcup} \cong \mathcal{U}(\mathcal{L}ie_{\mathbb{Q}}\langle Y \rangle) \quad \text{and} \quad \mathcal{H}_{\sqcup}^{\vee} \cong \mathcal{U}(\mathcal{L}ie_{\mathbb{Q}}\langle Y \rangle)^{\vee}. \quad (50)$$

Hence, let us consider [8]

1. the PBW-Lyndon basis $\{p_w\}_{w \in Y^*}$ for $\mathcal{U}(\mathcal{L}ie_{\mathbb{Q}}\langle Y \rangle)$ constructed recursively

$$\begin{cases} p_y &= y & \text{for } y \in Y, \\ p_l &= [p_s, p_r] & \text{for } l \in \mathcal{L}yn Y, \text{ standard factorization of } l = (s, r), \\ p_w &= p_{l_1}^{i_1} \dots p_{l_k}^{i_k} & \text{for } w = l_1^{i_1} \dots l_k^{i_k}, l_1 > \dots > l_k, l_1 \dots, l_k \in \mathcal{L}yn Y, \end{cases}$$

2. and, by duality¹¹, the linear basis $\{s_w\}_{w \in Y^*}$ for $(\mathbb{Q}\langle Y \rangle, \sqcup, 1_{Y^*})$, *i.e.*

$$\forall u, v \in Y^*, \quad \langle p_u \mid s_v \rangle = \delta_{u,v}.$$

This basis can be computed recursively as follows [47]

$$\begin{cases} s_y &= y, & \text{for } y \in Y, \\ s_l &= y s_u, & \text{for } l = yu \in \mathcal{L}yn Y, \\ s_w &= \frac{s_{l_1}^{\sqcup i_1} \sqcup \dots \sqcup s_{l_k}^{\sqcup i_k}}{i_1! \dots i_k!} & \text{for } w = l_1^{i_1} \dots l_k^{i_k}, l_1 > \dots > l_k. \end{cases}$$

¹¹The dual family (i.e. the set of coordinate forms) of a basis lies in the algebraic dual which is here the space of noncommutative series, but as the enveloping algebra under consideration is graded in finite dimensions (here by the multidegree), these series are in fact (multihomogeneous) polynomials.

Hence, since $\{p_w\}_{w \in Y^*}$ is of PBW type then, as in (38)-(40), we get also

$$\mathcal{D}_Y = \sum_{w \in Y^*} s_w \otimes p_w \quad (51)$$

$$= \sum_{l \in \mathcal{L}yn Y} s_l \otimes p_l + \sum_{w \notin \mathcal{L}yn Y} s_w \otimes p_w \quad (52)$$

$$= \prod_{l \in \mathcal{L}yn Y} \exp(s_l \otimes p_l). \quad (53)$$

Similarly, by the CQMM theorem, the connected \mathbb{N} -graded, co-commutative Hopf algebra \mathcal{H}_{\boxplus} is isomorphic to the enveloping algebra of

$$\text{Prim}(\mathcal{H}_{\boxplus}) = \text{Im}(\pi_1) = \text{span}_{\mathbb{Q}}\{\pi_1(w) | w \in Y^*\}, \quad (54)$$

where, for any $w \in Y^*$, $\pi_1(w)$ is obtained as follows [40]

$$\pi_1(w) = w + \sum_{k \geq 2} \frac{(-1)^{k-1}}{k} \sum_{u_1, \dots, u_k \in Y^+} \langle w | u_1 \boxplus \dots \boxplus u_k \rangle u_1 \dots u_k. \quad (55)$$

Note that Equation (55) is equivalent to the following identity [40, 41]

$$w = \sum_{k \geq 0} \frac{1}{k!} \sum_{u_1, \dots, u_k \in Y^*} \langle w | u_1 \boxplus \dots \boxplus u_k \rangle \pi_1(u_1) \dots \pi_1(u_k). \quad (56)$$

In particular, for any $y_k \in Y$, we have successively [40, 41]

$$\pi_1(y_k) = y_k + \sum_{l \geq 2} \frac{(-1)^{l-1}}{l} \sum_{\substack{j_1, \dots, j_l \geq 1 \\ j_1 + \dots + j_l = k}} y_{j_1} \dots y_{j_l}, \quad (57)$$

$$y_n = \sum_{k \geq 1} \frac{1}{k!} \sum_{s'_1 + \dots + s'_k = n} \pi_1(y_{s'_1}) \dots \pi_1(y_{s'_k}). \quad (58)$$

Hence, by introducing the new alphabet $\bar{Y} = \{\bar{y}\}_{y \in Y} = \{\pi_1(y)\}_{y \in Y}$, one has

$$(\mathbb{Q}\langle \bar{Y} \rangle, \text{conc}, 1_{\bar{Y}^*}, \Delta_{\boxplus}) \cong (\mathbb{Q}\langle Y \rangle, \text{conc}, 1_{Y^*}, \Delta_{\boxplus}) \quad (59)$$

and

$$\mathcal{H}_{\boxplus} \cong \mathcal{U}(\mathcal{L}ie_{\mathbb{Q}}\langle \bar{Y} \rangle) \cong \mathcal{U}(\text{Prim}(\mathcal{H}_{\boxplus})), \quad (60)$$

$$\mathcal{H}_{\boxplus}^{\vee} \cong \mathcal{U}(\mathcal{L}ie_{\mathbb{Q}}\langle \bar{Y} \rangle)^{\vee} \cong \mathcal{U}(\text{Prim}(\mathcal{H}_{\boxplus}))^{\vee}. \quad (61)$$

By considering

1. the PBW-Lyndon basis $\{\Pi_w\}_{w \in Y^*}$ for $\mathcal{U}(\text{Prim}(\mathcal{H}_{\boxplus}))$ constructed recursively as follows [40]

$$\begin{cases} \Pi_y &= \pi_1(y) & \text{for } y \in Y, \\ \Pi_l &= [\Pi_s, \Pi_r] & \text{for } l \in \mathcal{L}yn Y, \text{ standard factorization of } l = (s, r), \\ \Pi_w &= \Pi_{l_1}^{i_1} \dots \Pi_{l_k}^{i_k} & \text{for } w = l_1^{i_1} \dots l_k^{i_k}, l_1 > \dots > l_k, l_1, \dots, l_k \in \mathcal{L}yn Y, \end{cases}$$

2. and, by duality, the linear basis $\{\Sigma_w\}_{w \in Y^*}$ for $(\mathbb{Q}\langle Y \rangle, \boxplus, 1_{Y^*})$, i.e.

$$\forall u, v \in Y^*, \quad \langle \Pi_u \mid \Sigma_v \rangle = \delta_{u,v}.$$

This basis can be computed recursively as follows [4, 40]

$$\begin{cases} \Sigma_y &= y & \text{for } y \in Y, \\ \Sigma_l &= \sum_{\substack{\{s'_1, \dots, s'_i\} \subset \{s_1, \dots, s_k\}, l_1 \geq \dots \geq l_n \in \mathcal{L}_{ynY} \\ (y_{s_1} \dots y_{s_k}) \stackrel{*}{=} (y_{s'_1} \dots y_{s'_n}, l_1, \dots, l_n)}} \frac{1}{i!} y_{s'_1} + \dots + s'_i \Sigma_{l_1 \dots l_n} & \text{for } l \in \mathcal{L}_{ynY} \\ & & l = y_{s_1} \dots y_{s_k}, \\ \Sigma_w &= \frac{1}{i_1! \dots i_k!} \Sigma_{l_1}^{\boxplus i_1} \boxplus \dots \boxplus \Sigma_{l_k}^{\boxplus i_k} & \text{for } l_1 > \dots > l_k \\ & & w = l_1^{i_1} \dots l_k^{i_k}, \end{cases}$$

Hence, we get the following extended Schützenberger's factorization of \mathcal{D}_Y [40, 41]

$$\mathcal{D}_Y = \prod_{l \in \mathcal{L}_{ynY}}^{\searrow} \exp(\Sigma_l \otimes \Pi_l). \quad (62)$$

Proposition 1 ([40]). *Let $\pi_Y : (\mathbb{Q} \oplus \mathbb{Q}\langle X \rangle x_1, \cdot) \longrightarrow (\mathbb{Q}\langle Y \rangle, \cdot)$ be the morphism mapping $x_0^{s_1-1} x_1 \dots x_0^{s_r-1} x_1$ to $y_{s_1} \dots y_{s_r}$ and π_X be its inverse. Its extension over $\mathbb{Q}\langle X \rangle$ verifying $\pi_Y(p) = 0$, for $p \in \mathbb{Q}\langle X \rangle x_0$, is still denoted by π_Y . Then*

1. *The polynomials $\{\pi_Y P_{\pi_X l}\}_{l \in \mathcal{L}_{ynY}}$ are homogeneous in weight, linearly independent and upper triangular :*

$$\pi_Y P_{\pi_X l} = \Pi_l + \sum_{v > l, (v)=(l)} p_v v.$$

2. *For any $w \in Y^*$, the following polynomial, homogeneous in weight,*

$$\pi_Y P_{\pi_X w} = \Pi_w + \sum_{v > w, (v)=(w)} c_v v$$

is of multi-degree (w) and $\{\pi_Y P_{\pi_X w}\}_{w \in Y^}$ forms a basis for $\mathbb{Q}\langle Y \rangle$.*

3. *Let $\{\Theta_w\}_{w \in Y^*}$ be the family of polynomials, homogeneous in weight, in duality with the family $\{\pi_Y P_{\pi_X w}\}_{w \in Y^*}$:*

$$\forall u, v \in Y^*, \quad \langle \pi_Y P_{\pi_X u} \mid \Theta_v \rangle = \delta_{u,v}.$$

Then, the family $\{\Theta_w\}_{w \in Y^}$ generates freely $(\mathbb{Q}\langle Y \rangle, \boxplus)$ and*

$$\forall w \in Y^*, \quad \Theta_w = \Sigma_w + \sum_{v < w, (v)=(w)} d_v v.$$

Remark 1. *The family $\{\Theta_l\}_{l \in \mathcal{L}_{ynY}}$ is not a transcendence basis of $(\mathbb{Q}\langle Y \rangle, \boxplus)$ because if $\{\Theta_l\}_{l \in \mathcal{L}_{ynY}}$ constitutes a transcendence basis of $(\mathbb{Q}\langle Y \rangle, \boxplus)$ then, for any $l \in \mathcal{L}_{ynY}$, $\pi_Y P_{\pi_X l}$ is primitive but it is false in general.*

Example 5 ([41]).

$$\begin{aligned}
\Pi_{y_4} &= y_4 - \frac{1}{2}y_1y_3 - \frac{1}{2}y_2y_2 - \frac{1}{2}y_3y_1 + \frac{1}{3}y_1^2y_2 + \frac{1}{3}y_1y_2y_1 + \frac{1}{3}y_2y_1^2 - \frac{1}{4}y_1^4, \\
\Pi_{y_3y_1} &= y_3y_1 - \frac{1}{2}y_2y_1^2 - y_1y_3 + \frac{1}{2}y_1^2y_2, \\
\Pi_{y_2y_2} &= y_2y_2 - \frac{1}{2}y_2y_1^2 - \frac{1}{2}y_1^2y_2 + \frac{1}{4}y_1^4, \\
\Pi_{y_2y_1^2} &= y_2y_1^2 - 2y_1y_2y_1 + y_1^2y_2, \\
\Pi_{y_1y_3} &= y_1y_3 - \frac{1}{2}y_1^2y_2 - \frac{1}{2}y_1y_2y_1 + \frac{1}{3}y_1^4, \\
\Pi_{y_1y_2y_1} &= y_1y_2y_1 - y_1^2y_2, \\
\Pi_{y_1^2y_2} &= y_1^2y_2 - \frac{1}{2}y_1^4, \\
\Pi_{y_1^4} &= y_1^4, \\
\Sigma_{y_4} &= y_4, \\
\Sigma_{y_3y_1} &= \frac{1}{2}y_4 + y_3y_1, \\
\Sigma_{y_2^2} &= \frac{1}{2}y_4 + y_2^2, \\
\Sigma_{y_2y_1^2} &= \frac{1}{6}y_4 + \frac{1}{2}y_3y_1 + \frac{1}{2}y_2y_2 + y_2y_1^2, \\
\Sigma_{y_1y_3} &= y_4 + y_3y_1 + y_1y_3, \\
\Sigma_{y_1y_2y_1} &= \frac{1}{2}y_4 + \frac{1}{2}y_3y_1 + y_2^2 + y_2y_1^2 + \frac{1}{2}y_1y_3 + y_1y_2y_1, \\
\Sigma_{y_1^2y_2} &= \frac{1}{2}y_4 + y_3y_1 + y_2^2 + y_2y_1^2 + y_1y_3 + y_1y_2y_1 + y_1^2y_2, \\
\Sigma_{y_1^4} &= \frac{1}{24}y_4 + \frac{1}{6}y_3y_1 + \frac{1}{4}y_2^2 + \frac{1}{2}y_2y_1^2 + \frac{1}{6}y_1y_3 + \frac{1}{2}y_1y_2y_1 + \frac{1}{2}y_1^2y_2 + y_1^4.
\end{aligned}$$

Example 6 ([41]). *Let us express $\{\pi_Y P_l\}_{l \in \mathcal{L}_{yn_{\leq 4}Y}}$ in the basis $\{\Pi_w\}_{w \in Y^*}$:*

l	$\pi_Y P_l$
x_0	0
x_1	Π_{y_1}
x_0x_1	$\Pi_{y_2} + \frac{\Pi_{y_1}^2}{2}$
$x_0^2x_1$	$\Pi_{y_3} + \frac{\Pi_{y_2y_1}}{2} + \Pi_{y_1y_2} + \frac{\Pi_{y_1^3}}{6}$
$x_0x_1^2$	$\Pi_{y_2y_1} - 2\Pi_{y_1y_2} - \frac{\Pi_{y_1^3}}{6}$
$x_0^3x_1$	$\Pi_{y_4} + \frac{\Pi_{y_3y_1}}{2} + \Pi_{y_1y_3} + \frac{\Pi_{y_2^2}}{2} + \frac{\Pi_{y_1^2y_2} + \Pi_{y_1}\Pi_{y_2}\Pi_{y_1} + \Pi_{y_2}\Pi_{y_1^2}}{6} + \frac{\Pi_{y_1^4}}{24}$
$x_0^2x_1^2$	$\Pi_{y_3}\Pi_{y_1} - \frac{\Pi_{y_2}\Pi_{y_1^2}}{2} + \frac{\Pi_{y_1}\Pi_{y_2}\Pi_{y_1}}{2} - \Pi_{y_1^2}\Pi_{y_2} - 2\Pi_{y_2^2} - \frac{\Pi_{y_1^4}}{3}$
$x_0x_1^3$	$\Pi_{y_2}\Pi_{y_1^2} - 3\Pi_{y_1}\Pi_{y_2}\Pi_{y_1} + 3\Pi_{y_1^2}\Pi_{y_2} + \frac{\Pi_{y_1^4}}{2}$

2.2 Indiscernability over a class of formal power series

2.2.1 Residual calculus and representative series

Definition 1. Let $S \in \mathbb{Q}\langle\langle X \rangle\rangle$ and let $P \in \mathbb{Q}\langle X \rangle$. The left (resp. right) residual of S by P is the formal power series $P \triangleleft S$ (resp. $S \triangleright P$) in $\mathbb{Q}\langle\langle X \rangle\rangle$ defined by:

$$\langle P \triangleleft S \mid w \rangle = \langle S \mid wP \rangle \quad (\text{resp.} \quad \langle S \triangleright P \mid w \rangle = \langle S \mid Pw \rangle).$$

We straightforwardly get, for any $P, Q \in \mathbb{Q}\langle X \rangle$:

$$P \triangleleft (Q \triangleleft S) = PQ \triangleleft S, \quad (S \triangleright P) \triangleright Q = S \triangleright PQ, \quad (P \triangleleft S) \triangleright Q = P \triangleleft (S \triangleright Q). \quad (63)$$

In case $x, y \in X$ and $w \in X^*$, we get¹²:

$$x \triangleleft (wy) = \delta_{x,y} w \quad \text{and} \quad xw \triangleright y = \delta_{x,y} w. \quad (64)$$

Lemma 1. (Reconstruction lemma) Let $S \in \mathbb{Q}\langle\langle X \rangle\rangle$. Then

$$S = \langle S \mid 1_{X^*} \rangle + \sum_{x \in X} x(S \triangleright x) = \langle S \mid 1_{X^*} \rangle + \sum_{x \in X} (x \triangleleft S)x.$$

Lemma 2. The residuals by a letter x is a derivation of $(\mathbb{Q}\langle\langle X \rangle\rangle, \sqcup)$:

$$x \triangleleft (u \sqcup v) = (x \triangleleft u) \sqcup v + u \sqcup (x \triangleleft v), \quad (u \sqcup v) \triangleright x = (u \triangleright x) \sqcup v + u \sqcup (v \triangleright x).$$

Proof. Use the recursive definitions of the shuffle product. \square

Consequently,

Lemma 3. For any Lie polynomial $Q \in \mathcal{L}ie_{\mathbb{Q}}\langle X \rangle$, the linear maps “ $Q \triangleleft$ ” and “ $\triangleright Q$ ” are derivations on $(\mathbb{Q}[\mathcal{L}yn X], \sqcup)$.

Proof. For any $l, l_1, l_2 \in \mathcal{L}yn X$, we have

$$\begin{aligned} \hat{l} \triangleleft (l_1 \sqcup l_2) &= l_1 \sqcup (\hat{l} \triangleleft l_2) + (\hat{l} \triangleleft l_1) \sqcup l_2 = l_1 \delta_{l_2, \hat{l}} + \delta_{l_1, \hat{l}} l_2, \\ (l_1 \sqcup l_2) \triangleright \hat{l} &= l_1 \sqcup (l_2 \triangleright \hat{l}) + (l_1 \triangleright \hat{l}) \sqcup l_2 = l_1 \delta_{l_2, \hat{l}} + \delta_{l_1, \hat{l}} l_2. \end{aligned}$$

\square

Lemma 4. For any Lyndon word $l \in \mathcal{L}yn X$ and \check{S}_l defined as in (??), one has

$$x_1 \triangleleft l = l \triangleright x_0 = 0 \quad \text{and} \quad x_1 \triangleleft \check{S}_l = \check{S}_l \triangleright x_0 = 0.$$

Proof. Since $x_1 \triangleleft$ and $\triangleright x_0$ are derivations and for any $l \in \mathcal{L}yn X - X$, the polynomial \check{S}_l belongs to $x_0 \mathbb{Q}\langle X \rangle x_1$ then the expected results follow. \square

Theorem 1. (On representative series) The following properties are equivalent for any series $S \in \mathbb{Q}\langle\langle X \rangle\rangle$:

¹²For any words u and $v \in X^*$, if $u = v$ then $\delta_{u,v} = 1$ else 0.

- (1) The left \mathbb{C} -module $Res_g(S) = \text{span}\{w \triangleleft S \mid w \in X^*\}$ is finite dimensional.
- (2) The right \mathbb{C} -module $Res_d(S) = \text{span}\{S \triangleright w \mid w \in X^*\}$ is finite dimensional.
- (3) There are matrices $\lambda \in \mathcal{M}_{1,n}(\mathbb{Q})$, $\eta \in \mathcal{M}_{n,1}(\mathbb{Q})$ and a representation of X^* in $\mathcal{M}_{n,n}$, such that

$$\begin{aligned} S &= \sum_{w \in X^*} [\lambda \mu(w) \eta] w, \\ &= \lambda \left(\prod_{l \in \text{Lyn} X}^{\rightarrow} e^{\mu(S_l) P_l} \right) \eta. \end{aligned}$$

A series that satisfies the items of this theorem will be called *representative series*. This concept can be found in [1, 44, 15]. The two first items are in [19, 24]. The third item can be deduced from [7, 13] for example and it was used to factorize first time, by Lyndon words, the output of bilinear and analytical dynamical systems respectively in [26, 27] and to study polylogarithms, hypergeometric functions and associated functions in [29, 31, 38]. The dimension of $Res_g(S)$ is equal to that of $Res_d(S)$, and to the minimal dimension of a representation satisfying the third point of Theorem 1. This rank is then equal to the rank of the Hankel matrix of S , that is the infinite matrix $(\langle S \mid uv \rangle)_{u,v \in X}$ indexed by $X^* \times X^*$ and is also called *Hankel rank* of S [19, 24]:

Definition 2. ([19, 24]) *The Hankel rank of a formal power series $S \in \mathbb{C}\langle\langle X \rangle\rangle$ is the dimension of the vector space*

$$\{S \triangleright \Pi \mid \Pi \in \mathbb{C}\langle X \rangle\}, \quad (\text{resp.} \quad \{\Pi \triangleleft S \mid \Pi \in \mathbb{C}\langle X \rangle\}).$$

The triplet (λ, μ, η) is called a *linear representation* of S . We define the minimal representation¹³ of S as being a representation of S of minimal dimension.

For any proper series S , the following power series is called “star of S ”

$$S^* = 1 + S + S^2 + \cdots + S^n + \dots \quad (65)$$

Definition 3. ([3, 49]) *A series S is called rational if it belongs to the closure in $\mathbb{Q}\langle\langle X \rangle\rangle$ of the noncommutative polynomial algebra by sum, product, and star operation of proper¹⁴ elements. The set of rational power series will be denoted by $\mathbb{Q}^{\text{rat}}\langle\langle X \rangle\rangle$.*

Lemma 5. *For any noncommutative rational series (resp. polynomial) R and for any polynomial P , the left and right residuals of R by P are rational (resp. polynomial).*

Theorem 2. (Schützenberger, [3, 49]) *Any noncommutative power series is representative if and only if it is rational.*

¹³It can be shown that all minimal representations are isomorphic (see [3]).

¹⁴A series S is said to be proper if $\langle S \mid \epsilon \rangle = 0$.

Example 7. Let $S = (x_0x_1)^*$ be the rational power series over $X = \{x_0, x_1\}$. Since

$$S = (x_0x_1)^* = 1_{X^*} + x_0x_1 + (x_0x_1)^2 + \dots$$

then

$$\begin{aligned} S \triangleright x_0 &= x_1 + x_1x_0x_1 + \dots, \\ S \triangleright x_1 &= 0, \\ S \triangleright x_0^2 &= 0, \\ S \triangleright x_0x_1 &= S. \end{aligned}$$

The Hankel rank of S is then 3.

2.2.2 Continuity and indiscernability

Definition 4. ([25, 39]) Let \mathcal{H} be a class of formal power series over X and let $S \in \mathbb{C}\langle\langle X \rangle\rangle$.

- (1) S is said to be continuous¹⁵ over \mathcal{H} if for any $\Phi \in \mathcal{H}$, the following sum, denoted by $\langle S \parallel \Phi \rangle$, is convergent in norm

$$\sum_{w \in X^*} \langle S \mid w \rangle \langle \Phi \mid w \rangle.$$

The set of continuous power series over \mathcal{H} will be denoted by $\mathbb{C}^{\text{cont}}\langle\langle X \rangle\rangle$.

- (2) S is said to be indiscernable¹⁶ over \mathcal{H} if and only if

$$\forall \Phi \in \mathcal{H}, \quad \langle S \parallel \Phi \rangle = 0.$$

Lemma 6. Let \mathcal{H} be a monoid containing $\{e^{tx}\}_{x \in X}^{t \in \mathbb{C}}$. Let $S \in \mathbb{C}^{\text{cont}}\langle\langle X \rangle\rangle$ be indiscernable over \mathcal{H} . Then for any $x \in X$, $x \triangleleft S$ and $S \triangleright x$ belong to $\mathbb{C}^{\text{cont}}\langle\langle X \rangle\rangle$ and they are indiscernable over \mathcal{H} .

Proof. Let us calculate $\langle x \triangleleft S \parallel \Phi \rangle = \langle S \parallel \Phi x \rangle$ and $\langle S \triangleright x \parallel \Phi \rangle = \langle S \parallel x \Phi \rangle$. Since

$$\lim_{t \rightarrow 0} \frac{e^{tx} - 1}{t} = x \quad \text{and} \quad \lim_{t \rightarrow 0} \frac{e^{tx} - 1}{t} = x$$

then, for any $\Phi \in \mathcal{H}$, by uniform convergence, one has

$$\begin{aligned} \langle S \parallel \Phi x \rangle &= \langle S \parallel \lim_{t \rightarrow 0} \Phi \frac{e^{tx} - 1}{t} \rangle = \lim_{t \rightarrow 0} \langle S \parallel \Phi \frac{e^{tx} - 1}{t} \rangle, \\ \langle S \parallel x \Phi \rangle &= \langle S \parallel \lim_{t \rightarrow 0} \frac{e^{tx} - 1}{t} \Phi \rangle = \lim_{t \rightarrow 0} \langle S \parallel \frac{e^{tx} - 1}{t} \Phi \rangle. \end{aligned}$$

¹⁵See [25, 39] for a convergence criterion and an example of continuous generating series.

¹⁶Here, we adapt this notion developed in [25] via the residual calculus.

Since S is indiscernable over \mathcal{H} then

$$\begin{aligned}\langle S \parallel \Phi x \rangle &= \lim_{t \rightarrow 0} \frac{1}{t} \langle S \parallel \Phi e^{tx} \rangle - \lim_{t \rightarrow 0} \frac{1}{t} \langle S \parallel \Phi \rangle = 0, \\ \langle S \parallel x \Phi \rangle &= \lim_{t \rightarrow 0} \frac{1}{t} \langle S \parallel e^{tx} \Phi \rangle - \lim_{t \rightarrow 0} \frac{1}{t} \langle S \parallel \Phi \rangle = 0.\end{aligned}$$

□

Proposition 2. *Let \mathcal{H} be a monoid containing $\{e^{tx}\}_{x \in X}^{t \in \mathbb{C}}$. The formal power series $S \in \mathbb{C}^{\text{cont}} \langle\langle X \rangle\rangle$ is indiscernable over \mathcal{H} if and only if $S = 0$.*

Proof. If $S = 0$ then it is immediate that S is indiscernable over \mathcal{H} . Conversely, if S is indiscernable over \mathcal{H} then by Lemma 6, for any word $w \in X^*$ and, by induction on the length of w , $w \triangleleft S$ is indiscernable over \mathcal{H} . In particular,

$$\langle w \triangleleft S \parallel \text{Id}_{\mathcal{H}} \rangle = \langle S \mid w \rangle = 0.$$

In other words, $S = 0$.

□

2.3 Polylogarithms and harmonic sums

2.3.1 Structures of polylogarithms and harmonic sums

Let Y be the infinite alphabet $\{y_i\}_{i \geq 1}$ equipped with the order $y_1 > y_2 > y_3 > \dots$ and let $\mathcal{Lyn}Y$ be the set of Lyndon words over Y . The length of $w = y_{s_1} \dots y_{s_r} \in Y^*$ is denoted by $|w|$ and its degree equals $s_1 + \dots + s_r$.

Let X be the finite alphabet $\{x_0, x_1\}$ equipped with the order $x_0 < x_1$ and

$$\mathcal{C} := \mathbb{C} \left[z, \frac{1}{z}, \frac{1}{1-z} \right] \quad \text{and} \quad \mathcal{G} := \left\{ z, \frac{1}{z}, \frac{z-1}{z}, \frac{z}{z-1}, \frac{1}{1-z}, 1-z \right\} \quad (66)$$

This ring \mathcal{C} is invariant under differentiation and under the homographic transformations belonging to the group \mathcal{G} whose elements commute the singularities $\{0, 1, +\infty\}$. The iterated integral over ω_0, ω_1 associated to the word $w = x_{i_1} \dots x_{i_k}$ over X^* (the monoid generated by X) and along the integration path $z_0 \rightsquigarrow z$ is the following multiple integral defined by

$$\int_{z_0 \rightsquigarrow z} \omega_{i_1} \dots \omega_{i_k} = \int_{z_0}^z \omega_{i_1}(t_1) \int_{z_0}^{t_1} \omega_{i_2}(t_2) \dots \int_{z_0}^{t_{r-1}} \omega_{i_r}(t_r), \quad (67)$$

where $t_1 \dots t_{r-1}$ is a subdivision of the path $z_0 \rightsquigarrow z$. In a shortened notation, we denote this integral by $\alpha_{z_0}^z(w)$ and¹⁷ $\alpha_{z_0}^z(1_{X^*}) = 1$. One can check that the polylogarithm $\text{Li}_{s_1, \dots, s_r}$ is also the value of the iterated integral over ω_0, ω_1 and along the integration path $0 \rightsquigarrow z$ [29, 31]:

$$\text{Li}_w(z) = \alpha_0^z(x_0^{s_1-1} x_1 \dots x_0^{s_r-1} x_1). \quad (68)$$

¹⁷Here, 1_{X^*} stands for the empty word over X .

Example 8.

$$\begin{aligned}
\alpha_0^z(x_0 x_1) &= \int_0^z \frac{ds}{s} \int_0^s \frac{dt}{1-t} \\
&= \int_0^z \frac{ds}{s} \int_0^s dt \sum_{k \geq 0} t^k \\
&= \sum_{k \geq 1} \int_0^z ds \frac{s^{k-1}}{k} \\
&= \sum_{k \geq 1} \frac{z^k}{k^2} \\
&= \text{Li}_2(z).
\end{aligned}$$

The definition of polylogarithms is extended over the words $w \in X^*$ by putting $\text{Li}_{x_0}(z) := \log(z)$. Thus, $\{\text{Li}_w\}_{w \in X^*}$, and then $\{P_w(z) := (1-z)^{-1} \text{Li}_w(z)\}_{w \in Y^*}$, are \mathbb{C} -linearly independent [35, 32]. Since, for any $v = y_{s_1} \dots y_{s_r} \in Y^*$, P_v is the ordinary generating function of the sequence $\{H_v(N)\}_{N \geq 0}$ [37]:

$$P_v(z) = \sum_{N \geq 0} H_v(N) z^N, \quad \text{with} \quad H_v(N) = \sum_{N \geq n_1 > \dots > n_r > 0} \frac{1}{n_1^{s_1} \dots n_r^{s_r}} \quad (69)$$

then, as a consequence of the classical isomorphism between convergent Taylor series and their associated sums, the harmonic sums $\{H_w\}_{w \in Y^*}$ are also \mathbb{C} -linearly independent. Firstly, $\ker P = \{0\}$ and $\ker H = \{0\}$, and secondly, P is a morphism for the Hadamard product:

$$P_u(z) \odot P_v(z) = \sum_{N \geq 0} H_u(N) H_v(N) z^N = \sum_{N \geq 0} H_{u \boxtimes v}(N) z^N = P_{u \boxtimes v}(z). \quad (70)$$

Proposition 3. ([37]) *Extended by linearity, the map*

$$\begin{aligned}
P : (\mathbb{C}\langle Y \rangle, \boxtimes) &\longrightarrow (\mathbb{C}\{P_w\}_{w \in Y^*}, \odot), \\
u &\longmapsto P_u
\end{aligned}$$

is an isomorphism of algebras. Moreover, the map

$$\begin{aligned}
H : (\mathbb{C}\langle Y \rangle, \boxtimes) &\longrightarrow (\mathbb{C}\{H_w\}_{w \in Y^*}, \cdot), \\
u &\longmapsto H_u = \{H_u(N)\}_{N \geq 0}
\end{aligned}$$

is an isomorphism of algebras.

Studying the equivalence between action of $\{(1-z)^l\}_{l \in \mathbb{Z}}$ over $\{P_w(z)\}_{w \in Y^*}$ and that of $\{N^k\}_{k \in \mathbb{Z}}$ over $\{H_w(N)\}_{w \in Y^*}$ (see [17]), we have

Theorem 3. ([39]) *The Hadamard \mathbb{C} -algebra of $\{P_w\}_{w \in Y^*}$ can be identified with that of $\{P_l\}_{l \in \mathcal{L}_{yn}Y}$. In the same way, the algebra of harmonic sums $\{H_w\}_{w \in Y^*}$ with polynomial coefficients can be identified with that of $\{H_l\}_{l \in \mathcal{L}_{yn}Y}$.*

2.3.2 Results à la Abel for noncommutative generating and global regularizations

Let L and H be the noncommutative generating series of respectively $\{Li_w\}_{w \in X^*}$ and $\{H_w(N)\}_{w \in Y^*}$ [35, 37]:

$$L = \sum_{w \in X^*} Li_w w \quad \text{and} \quad H = \sum_{w \in Y^*} H_w w. \quad (71)$$

Hence, L and H are image, by the tensor products $Li \otimes Id$ and $H \otimes Id$, of the diagonal series \mathcal{D}_X and \mathcal{D}_Y respectively. We get

Theorem 4 (Factorization of L and of H , [35, 32, 40]). *Let*

$$L_{\text{reg}} = \prod_{l \in \mathcal{L}yn X - X}^{\searrow} e^{Li_{S_l} P_l} \quad \text{and} \quad H_{\text{reg}}(N) = \prod_{l \in \mathcal{L}yn Y - \{y_1\}}^{\searrow} e^{H_{\Sigma_l(N)} \Sigma_l}.$$

Then $L(z) = e^{-x_1 \log(1-z)} L_{\text{reg}}(z) e^{x_0 \log z}$ and $H(N) = e^{H_{y_1(N)} y_1} H_{\text{reg}}(N)$.

For any $l \in \mathcal{L}yn X - X$ (resp. $\mathcal{L}yn Y - \{y_1\}$), the polynomial S_l (resp. Σ_l) is a finite combination of words in $x_0 X^* x_1$ (resp. $Y^* - y_1 Y^*$). Then we can state the following

Proposition 4 ([40]). *Let $Z_{\sqcup} := L_{\text{reg}}(1)$ and $Z_{\boxplus} := H_{\text{reg}}(\infty)$.*

Then Z_{\sqcup} and Z_{\boxplus} are group-like, for Δ_{\sqcup} and Δ_{\boxplus} respectively.

Proposition 5 (Successive differentiations of L , [38]). *Let $\partial_z = d/dz$ and let $\theta_0 = zd/dz$. For any $n \in \mathbb{N}$, we have*

$$\partial_z^n L(z) = D_n(z) L(z) \quad \text{and} \quad \theta_0^n L(z) = E_n(z) L(z),$$

where¹⁸

- the polynomials $D_n(z)$ and $E_n(z)$ in $\mathcal{C}\langle X \rangle$ are defined as follows

$$D_n(z) = \sum_{\text{wgt}(\mathbf{r})=n} \sum_{w \in X^{\deg(\mathbf{r})}} \prod_{i=1}^{\deg(\mathbf{r})} \binom{\sum_{j=1}^i r_j + j - 1}{r_i} \tau_{\mathbf{r}}(w),$$

$$E_n(z) = \sum_{\text{wgt}(\mathbf{r})=n} \sum_{w \in X^{\deg(\mathbf{r})}} \prod_{i=1}^{\deg(\mathbf{r})} \binom{\sum_{j=1}^i r_j + j - 1}{r_i} \rho_{\mathbf{r}}(w),$$

- for any $w = x_{i_1} \cdots x_{i_k}$ and $\mathbf{r} = (r_1, \dots, r_k)$ of degree $\deg(\mathbf{r}) = k$ and of weight $\text{wgt}(\mathbf{r}) = k + r_1 + \cdots + r_k$, the polynomials $\tau_{\mathbf{r}}(w) = \tau_{r_1}(x_{i_1}) \cdots \tau_{r_k}(x_{i_k})$ and $\rho_{\mathbf{r}}(w) = \rho_{r_1}(x_{i_1}) \cdots \rho_{r_k}(x_{i_k})$ are defined respectively by, for any $r \in \mathbb{N}$,

$$\tau_r(x_0) = \partial_z^r \frac{x_0}{z} = \frac{-r! x_0}{(-z)^{r+1}} \quad \text{and} \quad \tau_r(x_1) = \partial_z^r \frac{x_1}{1-z} = \frac{r! x_1}{(1-z)^{r+1}},$$

$$\rho_r(x_0) = \theta_0^r \frac{x_0}{z} = 0 \quad \text{and} \quad \rho_r(x_1) = \theta_0^r \frac{zx_1}{1-z} = Li_{-r}(z) x_1.$$

¹⁸Consider also $\theta_1 = (1-z)d/dz$. But $\theta_0 + \theta_1 = \partial_z$ and then $\theta_1^n L(z) = [D_n(z) - E_n(z)]L(z)$.

Example 9 (Coefficients of $\theta_0^n \text{Li}$). *Since, for any $w \in X^+$, one has*

$$\theta_0 \text{Li}_w = \begin{cases} \text{Li}_u & \text{if } w = x_0 u, \\ \text{Li}_0 \text{Li}_u & \text{if } w = x_1 u. \end{cases}$$

then one obtains, for example

- *For any $n \geq 1$ and for any $w \in X^*$, one has*

$$\theta_0^n \text{Li}_{x_0^n w}(z) = \text{Li}_w(z).$$

In particular,

$$\theta_0^n \text{Li}_{x_0^n}(z) = 1.$$

- *Letting $E(n, m)$ denote the Eulerian numbers :*

$$E(n, m) = \sum_{j=0}^m (-1)^j \binom{n+1}{j} (m+1-j)^n,$$

one has, for any $n \geq 0$,

$$\theta_0^{n+1} \text{Li}_{x_1}(z) = \frac{z}{(1-z)^{n+1}} \sum_{m=0}^n E(n, m) x^m =: \text{Li}_{-n}(z).$$

In particular,

$$\theta_0 \text{Li}_{x_1}(z) = \frac{z}{(1-z)} = \text{Li}_0(z).$$

- *For any $k \in \mathbb{N}^*$, one has*

$$\theta_0^k \text{Li}_{x_1^2}(z) = \sum_{j=0}^{k-1} \binom{k-1}{j} \text{Li}_{-j}(z) \text{Li}_{2+j-k}(z).$$

For example

$$\begin{aligned} \theta_0 \text{Li}_{x_1^2}(z) &= \text{Li}_0(z) \text{Li}_1(z), \\ \theta_0^2 \text{Li}_{x_1^2}(z) &= \text{Li}_{-1}(z) \text{Li}_1(z) + \text{Li}_0^2(z), \\ \theta_0^3 \text{Li}_{x_1^2}(z) &= \text{Li}_{-2}(z) \text{Li}_1(z) + 3 \text{Li}_{-1}(z) \text{Li}_0(z), \\ \theta_0^4 \text{Li}_{x_1^2}(z) &= \text{Li}_{-3}(z) \text{Li}_1(z) + 4 \text{Li}_{-2}(z) \text{Li}_0(z) + 3 \text{Li}_{-1}^2(z), \\ &\vdots \end{aligned}$$

The noncommutative generating series of polylogarithms [35, 32] satisfies the differential equation

$$dL = (x_0\omega_0 + x_1\omega_1)L \quad (72)$$

with boundary condition

$$L(z) \underset{z \rightarrow 0}{\sim} \exp(x_0 \log z) \quad \text{and} \quad L(z) \underset{z \rightarrow 1}{\sim} \exp(-x_1 \log(1-z)) Z_{\sqcup}. \quad (73)$$

This implies L is the exponential of a Lie series [35, 32]. Hence [38],

$$\log L(z) = \sum_{k \geq 1} \frac{(-1)^{k-1}}{k} \sum_{u_1, \dots, u_k \in X^+} \text{Li}_{u_1 \sqcup \dots \sqcup u_k}(z) u_1 \dots u_k \quad (74)$$

$$= \sum_{w \in X^*} \text{Li}_w(z) \pi_1(w). \quad (75)$$

Proposition 6 ([38]). *Let $G(z), H(z)$ be exponential solutions of (72). Then there exists a constant Lie series C such that $G(z) = H(z)e^C$.*

Theorem 5 ([38]). *Let $\text{Gal}_{\mathbb{C}}(DE)$ be the differential Galois group associated to the Drinfel'd equation. Then $\text{Gal}_{\mathbb{C}}(DE) = \{e^C \mid C \in \mathcal{L}ie_{\mathbb{C}}\langle\langle X \rangle\rangle\}$ and it contains the monodromy group defined as follows*

$$\mathcal{M}_0 L = L e^{2i\pi \mathbf{m}_0} \quad \text{and} \quad \mathcal{M}_1 L = L Z_{\sqcup}^{-1} e^{-2i\pi x_1} Z_{\sqcup} = L e^{2i\pi \mathbf{m}_1},$$

$$\text{where } \mathbf{m}_0 = x_0 \quad \text{and} \quad \mathbf{m}_1 = \prod_{l \in \text{Lyn} X - X}^{\searrow} e^{-\zeta(S_l) \text{ad}_{P_l}(-x_1)}.$$

Let us call $\text{Li}_{\mathcal{C}}$ the smallest algebra containing \mathcal{C} , closed under derivation and under integration with respect to ω_0 and ω_1 . It is the \mathcal{C} -module generated by the polylogarithms $\{\text{Li}_w\}_{w \in X^*}$.

Let $\Lambda(z) := \pi_Y L(z)$. Then [39]

$$\Lambda(z) \underset{z \rightarrow 1}{\sim} \exp\left(y_1 \log \frac{1}{1-z}\right) \pi_Y Z_{\sqcup}. \quad (76)$$

Since the coefficient of z^N in the ordinary Taylor expansion of $P_{y_1^k}(z)$ is $H_{y_1^k}(N)$ then let us put [39]

$$\text{Mono}(z) := e^{-(x_1+1) \log(1-z)} = \sum_{k \geq 0} P_{y_1^k}(z) y_1^k \quad (77)$$

$$\text{Const} := \sum_{k \geq 0} H_{y_1^k} y_1^k = \exp\left(-\sum_{k \geq 1} H_{y_k} \frac{(-y_1)^k}{k}\right). \quad (78)$$

We put also¹⁹ [39]

$$B(y_1) := \exp\left(\sum_{k \geq 1} \zeta(y_k) \frac{(-y_1)^k}{k}\right) \quad \text{and} \quad B'(y_1) := \exp(\gamma y_1) B(y_1). \quad (79)$$

¹⁹Here, the coefficient $\langle B(y_1) \mid y_1^k \rangle$ corresponds to the Euler–Mac Laurin constant associated to $\langle \text{Const}(N) \mid y_1^k \rangle$, i.e. the finite party of its asymptotic expansion in the scale of comparison $\{n^a \log^b(n)\}_{a \in \mathbb{Z}, b \in \mathbb{N}}$.

As a consequence of (76)-(78), one gets

Theorem 6 ([39]).

$$\pi_Y P(z) \underset{z \rightarrow 1}{\sim} \text{Mono}(z) \pi_Y Z_{\sqcup} \quad \text{and} \quad H(N) \underset{N \rightarrow \infty}{\sim} \text{Const}(N) \pi_Y Z_{\sqcup}.$$

Or equivalently,

$$\lim_{z \rightarrow 1} \exp\left(y_1 \log \frac{1}{1-z}\right) \Lambda(z) = \lim_{N \rightarrow \infty} \exp\left(\sum_{k \geq 1} H_{y_k}(N) \frac{(-y_1)^k}{k}\right) H(N) = \pi_Y Z_{\sqcup}.$$

Theorem 7 ([17]). *For any $g \in \mathcal{C}\{P_w\}_{w \in Y^*}$, there exist algorithmically computable coefficients $c_j \in \mathbb{C}, \alpha_j \in \mathbb{Z}, \beta_j \in \mathbb{N}$ such that*

$$g(z) \sim \sum_{j=0}^{+\infty} c_j (1-z)^{\alpha_j} \log^{\beta_j}(1-z) \quad \text{for } z \rightarrow 1,$$

and the coefficients $b_i \in \mathbb{C}, \eta_i \in \mathbb{Z}, \kappa_i \in \mathbb{N}$ such that

$$\langle g(z) \mid z^n \rangle \sim \sum_{i=0}^{+\infty} b_i n^{\eta_i} \log^{\kappa_i}(n) \quad \text{for } n \rightarrow \infty.$$

Theorem 7 means also that the $\{P_w\}_{w \in Y^*}$ admit a full singular expansion, at $z = 1$, and then their ordinary Taylor coefficients, $\{H_w(N)\}_{w \in Y^*}$ admit a full asymptotic expansion, for $N \rightarrow +\infty$. More precisely,

Corollary 1. *For any $w \in Y^*$ and for any $k \geq 1$, we have*

$$H_w(N) = \sum_{i=1}^{|w|} \alpha_i \log^i(N) + \gamma_w + \sum_{j=1}^k \sum_{i=0}^{|w|-1} \beta_{i,j} \frac{1}{N^j} \log^i(N) + O\left(\frac{1}{N^k}\right).$$

where the coefficients γ_w, α_i and $\beta_{i,j}$ belong to the $\mathbb{Q}[\gamma]$ -algebra generated by convergent polyzetas.

More generally, by Theorem 5, we get

Proposition 7. *For any commutative \mathbb{Q} -algebra A and for any Lie series $C \in \mathcal{L}ie_A\langle X \rangle$, let*

$$\bar{L} = Le^C, \quad \bar{\Lambda} = \pi_Y \bar{L}, \quad \bar{P}(z) = \frac{\bar{\Lambda}(z)}{1-z}.$$

Then

1. $\bar{Z}_{\sqcup} = Z_{\sqcup} e^C$ is group-like, for the co-product Δ_{\sqcup} ,
2. $\bar{L}(z) \underset{z \rightarrow 1}{\sim} \exp(-x_1 \log(1-z)) \bar{Z}_{\sqcup}$,
3. $\bar{P}(z) \underset{z \rightarrow 1}{\sim} \text{Mono}(z) \pi_Y \bar{Z}_{\sqcup}$,

$$4. \overline{H}(N) \xrightarrow[N \rightarrow \infty]{} \text{Const}(N) \pi_Y \overline{Z}_{\sqcup},$$

where, for any $w \in Y^*$ and $N \geq 0$, one defines the coefficient $\langle \overline{H}(N) \mid w \rangle$ of w in the power series $\overline{H}(N)$ as the coefficient $\langle \overline{P}_w(z) \mid z^N \rangle$ of z^N in the ordinary Taylor expansion of the polylogarithmic function $\overline{P}_w(z)$.

By Proposition 7, we get successively

Proposition 8 ([33]). *Let $\overline{\zeta}_{\sqcup}$ and $\overline{\zeta}_{\sqcup\sqcup}$ be the characters of respectively $(A\langle X \rangle, \sqcup)$ and $(A\langle Y \rangle, \sqcup\sqcup)$ satisfying*

$$\overline{\zeta}_{\sqcup}(x_0) = \overline{\zeta}_{\sqcup}(x_1) = 0 \quad \text{and} \quad \overline{\zeta}_{\sqcup\sqcup}(y_1) = 0.$$

Then

$$\begin{aligned} \sum_{w \in Y^*} \overline{\zeta}_{\sqcup}(w) w &= \overline{Z}_{\sqcup} = \prod_{l \in \mathcal{L}yn X - X}^{\searrow} \exp(\overline{\zeta}(S_l) P_l), \\ \sum_{w \in Y^*} \overline{\zeta}_{\sqcup\sqcup}(w) w &= \overline{Z}_{\sqcup\sqcup} = \prod_{l \in \mathcal{L}yn Y - \{y_1\}}^{\searrow} \exp(\overline{\zeta}(\Sigma_l) \Pi_l). \end{aligned}$$

Proposition 9. *Let $\{\overline{\gamma}_w\}_{w \in Y^*}$ be the Euler–Mac Laurin constants associated to $\{\overline{H}_w(N)\}_{w \in Y^*}$. Let \overline{Z}_γ be its noncommutative generating series. Then,*

1. *The following map realizes a character :*

$$\begin{aligned} \overline{\gamma}_\bullet &: (A\langle Y \rangle, \sqcup\sqcup) \longrightarrow (\mathbb{R}, \cdot), \\ w &\longmapsto \langle \overline{\gamma}_\bullet \mid w \rangle = \overline{\gamma}_w. \end{aligned}$$

2. *The noncommutative power series \overline{Z}_γ is group-like, for $\Delta_{\sqcup\sqcup}$.*

3. *There exists a group-like element $\overline{Z}_{\sqcup\sqcup}$, for the co-product $\Delta_{\sqcup\sqcup}$, such that*

$$\overline{Z}_\gamma = \sum_{w \in Y^*} \overline{\gamma}_w w = \exp(\gamma y_1) \overline{Z}_{\sqcup\sqcup}.$$

By Theorem 6, Propositions 7 and 9, we also get

Proposition 10. *For any $C \in \mathcal{L}ie_A\langle X \rangle$ such that $\overline{Z}_{\sqcup} = Z_{\sqcup} e^C$. Then*

$$\overline{Z}_\gamma = B(y_1) \pi_Y \overline{Z}_{\sqcup}, \quad \text{or equivalently by cancellation,} \quad \overline{Z}_{\sqcup\sqcup} = B'(y_1) \pi_Y \overline{Z}_{\sqcup},$$

where $B(y_1)$ and $B'(y_1)$ are given in (79).

By Propositions 7, the noncommutative generationg series \overline{Z}_{\sqcup} and $\overline{Z}_{\sqcup\sqcup}$ are group-like, for the co-product Δ_{\sqcup} and $\Delta_{\sqcup\sqcup}$ respectively, and we also have

$$\overline{Z}_{\sqcup} = \sum_{l \in \mathcal{L}yn X - X} \overline{\zeta}(S_l) P_l + \sum_{w \notin \mathcal{L}yn X - X} \overline{\zeta}_{\sqcup}(S_w) P_w, \quad (80)$$

$$\overline{Z}_{\sqcup\sqcup} = \sum_{l \in \mathcal{L}yn Y - \{y_1\}} \overline{\zeta}(\Sigma_l) \Pi_l + \sum_{w \notin \mathcal{L}yn Y - \{y_1\}} \overline{\zeta}_{\sqcup\sqcup}(\Sigma_w) \Pi_w. \quad (81)$$

Hence, by Proposition 10, we deduce in particular,

$$\sum_{l \in \mathcal{L}_{yn}Y - \{y_1\}} \bar{\zeta}(\Sigma_l) \Pi_l + \dots = B'(y_1) \sum_{l \in \mathcal{L}_{yn}X - X} \bar{\zeta}(S_l) \pi_Y P_l + \dots \quad (82)$$

By Proposition 1, the elements of the family $\{\pi_Y P_l\}_{l \in \mathcal{L}_{yn}X}$ are decomposable in the linear basis $\{\Pi_w\}_{w \in Y^*}$ of $\mathcal{U}(\text{Prim}(\mathcal{H}_{\mathbf{u}}))$. Thus, by identification of locale coordinates, *i.e.* the coefficients of $\{\Pi_l\}_{l \in \mathcal{L}_{yn}Y - \{y_1\}}$, we get the homogenous polynomial relations among convergent polyzetas [40].

3 Polysystem and differential realization

3.1 Polysystem and convergence criterion

3.1.1 Serial estimates from above

Here, generalizing a little, \mathbb{K} is supposed a \mathbb{C} -algebra and a complete normed vector space equipped with a norm denoted by $\|\cdot\|$.

For any $n \in \mathbb{N}$, $X^{\geq n}$ denotes the set of words over X of length greater than or equal to n . The set of formal power series (resp. polynomials) on X , is denoted by $\mathbb{K}\langle\langle X \rangle\rangle$ (resp. $\mathbb{K}\langle X \rangle$).

Definition 5. ([25, 39]) *Let ξ, χ be real positive functions over X^* . Let $S \in \mathbb{K}\langle\langle X \rangle\rangle$.*

(1) *S will be said ξ -exponentially bounded from above if it verifies*

$$\exists K \in \mathbb{R}_+, \exists n \in \mathbb{N}, \forall w \in X^{\geq n}, \quad \|\langle S \mid w \rangle\| \leq K \frac{\xi(w)}{|w|!}.$$

We denote by $\mathbb{K}^{\xi\text{-em}}\langle\langle X \rangle\rangle$ the set of formal power series in $\mathbb{K}\langle\langle X \rangle\rangle$ which are ξ -exponentially bounded from above.

(2) *S verifies the χ -growth condition if it satisfies*

$$\exists K \in \mathbb{R}_+, \exists n \in \mathbb{N}, \forall w \in X^{\geq n}, \quad \|\langle S \mid w \rangle\| \leq K \chi(w) |w|!.$$

We denote by $\mathbb{K}^{\chi\text{-gc}}\langle\langle X \rangle\rangle$ the set of formal power series in $\mathbb{K}\langle\langle X \rangle\rangle$ verifying the χ -growth condition.

Lemma 7. *We have*

$$R = \sum_{w \in X^*} |w|! w \Rightarrow \langle R^{\sqcup^2} \mid w \rangle = \sum_{\substack{u, v \in X^* \\ \text{supp}(u \sqcup v) \ni w}} |u|! |v|! \leq 2^{|w|} |w|!.$$

Proof. One has

$$\begin{aligned}
\sum_{\substack{u, v \in X^* \\ \text{supp}(u \sqcup v) \ni w}} |u|! |v|! &= \sum_{k=0}^{|w|} \sum_{\substack{|u|=k, |v|=|w|-k \\ \text{supp}(u \sqcup v) \ni w}} k! (|w| - k)! \\
&= \sum_{k=0}^{|w|} \binom{|w|}{k} k! (|w| - k)! \\
&= \sum_{k=0}^{|w|} |w|! = (1 + |w|) |w|!.
\end{aligned}$$

By induction on $|w|$, one has $1 + |w| \leq 2^{|w|}$. The expected result follows. \square

Proposition 11. *Let S_1 and S_2 verify the growth condition. Then $S_1 + S_2$ and $S_1 \sqcup S_2$ also verify the growth condition.*

Proof. The proof for $S_1 + S_2$ is immediate.

Next, since $\|\langle S_i \mid w \rangle\| \leq K_i \chi_i(w) |w|!$, for $i = 1$ or 2 and for $w \in X^*$, then²⁰

$$\begin{aligned}
\langle S_1 \sqcup S_2 \mid w \rangle &= \sum_{\text{supp}(u \sqcup v) \ni w} \langle S_1 \mid u \rangle \langle S_2 \mid v \rangle, \\
\Rightarrow \|\langle S_1 \sqcup S_2 \mid w \rangle\| &\leq K_1 K_2 \sum_{\substack{u, v \in X^* \\ \text{supp}(u \sqcup v) \ni w}} (\chi_1(u) |u|!) (\chi_2(v) |v|!).
\end{aligned}$$

Let $K = K_1 K_2$ and let χ be a real positive function over X^* such that

$$\forall w \in X^*, \quad \chi(w) = \max\{\chi_1(u) \chi_2(v) \mid u, v \in X^* \text{ and } \text{supp}(u \sqcup v) \ni w\}.$$

With the notations in Lemma 7, we get

$$\|\langle S_1 \sqcup S_2 \mid w \rangle\| \leq K \chi(w) \langle R^{\sqcup 2} \mid w \rangle.$$

Hence, $S_1 \sqcup S_2$ verifies the χ' -growth condition with $\chi'(w) = 2^{|w|} \chi(w)$. \square

Definition 6. ([25, 39]) *Let ξ be a real positive function defined over X^* , S will be said ξ -exponentially continuous if it is continuous over $\mathbb{K}^{\xi-\text{em}}\langle X \rangle$. The set of formal power series which are ξ -exponentially continuous is denoted by $\mathbb{K}^{\xi-\text{ec}}\langle X \rangle$.*

Lemma 8. [25, 39] *For any real positive function ξ defined over X^* , we have $\mathbb{K}\langle X \rangle \subset \mathbb{K}^{\xi-\text{ec}}\langle X \rangle$. Otherwise, for $\xi = 0$, we get $\mathbb{K}\langle X \rangle = \mathbb{K}^{0-\text{ec}}\langle X \rangle$. Hence, any polynomial is 0-exponentially continuous.*

Proposition 12. ([25, 39]) *Let ξ, χ be real positive functions over X^* and let $P \in \mathbb{K}\langle X \rangle$.*

²⁰ $\langle S_1 \sqcup S_2 \mid w \rangle$ is the coefficient of the word w in the power series $S_1 \sqcup S_2$.

(1) Let $S \in \mathbb{K}^{\xi-\text{em}}\langle\langle X \rangle\rangle$. The right residual of S by P belongs to $\mathbb{K}^{\xi-\text{em}}\langle\langle X \rangle\rangle$.

(2) Let $R \in \mathbb{K}^{\chi-\text{gc}}\langle\langle X \rangle\rangle$. The concatenation SR belongs to $\mathbb{K}^{\chi-\text{gc}}\langle\langle X \rangle\rangle$.

Proof. (1) Since $S \in \mathbb{K}^{\xi-\text{em}}\langle\langle X \rangle\rangle$ then

$$\exists K \in \mathbb{R}_+, \exists n \in \mathbb{N}, \forall w \in X^{\geq n}, \quad \|\langle S \mid w \rangle\| \leq K \frac{\xi(w)}{|w|!}.$$

If $u \in \text{supp}(P) := \{w \in X^* \mid \langle P \mid w \rangle \neq 0\}$ then, for any $w \in X^*$, one has $\langle S \triangleright u \mid w \rangle = \langle S \mid uw \rangle$ and $S \triangleright u$ belongs to $\mathbb{K}^{\xi-\text{em}}\langle\langle X \rangle\rangle$:

$$\exists K \in \mathbb{R}_+, \exists n \in \mathbb{N}, \forall w \in X^{\geq n}, \quad \|\langle S \triangleright u \mid w \rangle\| \leq [K\xi(u)] \frac{\xi(w)}{|w|!}.$$

It follows that $S \triangleright P$ is $\mathbb{K}^{\xi-\text{em}}\langle\langle X \rangle\rangle$ by taking $K_1 = K \max_{u \in \text{supp}(P)} \xi(u)$.

(2) Since $R \in \mathbb{K}^{\chi-\text{gc}}\langle\langle X \rangle\rangle$ then

$$\exists K \in \mathbb{R}_+, \exists n \in \mathbb{N}, \forall w \in X^{\geq n}, \quad \|\langle S \mid w \rangle\| \leq K \chi(w) |w|!.$$

Let $v \in \text{supp}(P)$ such that $v \neq \epsilon$. Since, for any $w \in X^*$, Rv belongs to $\mathbb{K}^{\chi-\text{gc}}\langle\langle X \rangle\rangle$ and one has $\langle Rv \mid w \rangle = \langle R \mid v \triangleleft w \rangle$:

$$\begin{aligned} \exists K \in \mathbb{R}_+, \exists n \in \mathbb{N}, \forall w \in X^{\geq n}, \quad \|\langle R \mid v \triangleleft w \rangle\| &\leq K \chi(v \triangleleft w) (|w| - |v|)! \\ &\leq K |w| \frac{\chi(w)}{\chi(v)}. \end{aligned}$$

Note if $v \triangleleft w = 0$ then $\langle Rv \mid w \rangle = 0$ and the previous conclusion holds. It follows that RP is $\mathbb{K}^{\chi-\text{gc}}\langle\langle X \rangle\rangle$ by taking $K_2 = K \min_{v \in \text{supp}(P)} \chi(v)^{-1}$. \square

Proposition 13. ([25, 39]) *Two real positive morphisms over X^* , ξ and χ are assumed to verify the condition*

$$\sum_{x \in X} \chi(x) \xi(x) < 1.$$

Then for any $F \in \mathbb{K}^{\chi-\text{gc}}\langle\langle X \rangle\rangle$, F is continuous over $\mathbb{K}^{\xi-\text{em}}\langle\langle X \rangle\rangle$.

Proof. If ξ, χ verify the upper bound condition then the following power series

$$\sum_{w \in X^*} \chi(w) \xi(w) = \left(\sum_{x \in X} \chi(x) \xi(x) \right)^*$$

is well defined. If $F \in \mathbb{K}^{\chi-\text{gc}}\langle\langle X \rangle\rangle$ and $C \in \mathbb{K}^{\xi-\text{em}}\langle\langle X \rangle\rangle$ then there exist $K_i \in \mathbb{R}_+$ and $n_i \in \mathbb{N}$ such that for any $w \in X^{\geq n_i}$, $i = 1, 2$, one has

$$\|\langle F \mid w \rangle\| \leq K_1 \chi(w) |w|! \quad \text{and} \quad \|\langle C \mid w \rangle\| \leq K_2 \frac{\xi(w)}{|w|!}.$$

Hence,

$$\begin{aligned} & \forall w \in X^*, |w| \geq \max\{n_1, n_2\}, \quad \|\langle F|w \rangle \langle C|w \rangle\| \leq K_1 K_2 \chi(w) \xi(w), \\ \Rightarrow & \sum_{w \in X^*} \|\langle F|w \rangle \langle C|w \rangle\| \leq K_1 K_2 \sum_{w \in X^*} \chi(w) \xi(w) = K_1 K_2 \left(\sum_{x \in X} \chi(x) \xi(x) \right)^*. \end{aligned}$$

□

3.1.2 Upper bounds à la Cauchy

Let q_1, \dots, q_n be commutative indeterminates over \mathbb{C} . The algebra of formal power series (resp. polynomials) over $\{q_1, \dots, q_n\}$ with coefficients in \mathbb{C} is denoted by $\mathbb{C}[[q_1, \dots, q_n]]$ (resp. $\mathbb{C}[q_1, \dots, q_n]$).

Definition 7. ([25, 39]) *Let*

$$f = \sum_{i_1, \dots, i_n \geq 0} f_{i_1, \dots, i_n} q_1^{i_1} \dots q_n^{i_n} \in \mathbb{C}[[q_1, \dots, q_n]].$$

We set

$$\begin{aligned} E(f) &:= \{\rho \in \mathbb{R}_+^n : \exists C_f \in \mathbb{R}_+ \text{ s.t. } \forall i_1, \dots, i_n \geq 0, |f_{i_1, \dots, i_n}| \rho_1^{i_1} \dots \rho_n^{i_n} \leq C_f\}. \\ \check{E}(f) &:= \text{the interior of } E(f) \text{ in } \mathbb{R}^n. \\ \text{CV}(f) &:= \{q \in \mathbb{C}^n : (|q_1|, \dots, |q_n|) \in \check{E}(f)\} : \text{the convergence domain of } f. \end{aligned}$$

The power series f is said to be convergent if $\text{CV}(f) \neq \emptyset$. Let \mathcal{U} be an open domain in \mathbb{C}^n and let $q \in \mathbb{C}^n$. The power series f is said to be convergent on q (resp. over \mathcal{U}) if $q \in \text{CV}(f)$ (resp. $\mathcal{U} \subset \text{CV}(f)$). We set

$$\mathbb{C}^{\text{cv}}[[q_1, \dots, q_n]] = \{f \in \mathbb{C}[[q_1, \dots, q_n]] : \text{CV}(f) \neq \emptyset\}.$$

Let $q \in \text{CV}(f)$. There exist the constants C_f, ρ and $\bar{\rho}$ such that

$$|q_1| < \bar{\rho} < \rho, \dots, |q_n| < \bar{\rho} < \rho \quad \text{and} \quad |f_{i_1, \dots, i_n}| \rho_1^{i_1} \dots \rho_n^{i_n} \leq C_f,$$

for $i_1, \dots, i_n \geq 0$. The convergence modulus of f at q is $(C_f, \rho, \bar{\rho})$.

Suppose that $\text{CV}(f) \neq \emptyset$ and let $q \in \text{CV}(f)$. If $(C_f, \rho, \bar{\rho})$ is a convergence modulus of f at q then $|f_{i_1, \dots, i_n} q_1^{i_1} \dots q_n^{i_n}| \leq C_f (\bar{\rho}_1/\rho_1)^{i_1} \dots (\bar{\rho}_n/\rho_n)^{i_n}$. Hence, at q , the power series f is majored termwise by

$$C_f \prod_{k=0}^m \left(1 - \frac{\bar{\rho}_k}{\rho_k}\right)^{-1}. \quad (83)$$

Hence, f is uniformly absolutely convergent in $\{q \in \mathbb{C}^n : |q_1| < \bar{\rho}, \dots, |q_n| < \bar{\rho}\}$ which is an open domain in \mathbb{C}^n . Thus, $\text{CV}(f)$ is an open domain in \mathbb{C}^n . Since the partial derivation $D_1^{j_1} \dots D_n^{j_n} f$ is estimated by

$$\|D_1^{j_1} \dots D_n^{j_n} f\| \leq C_f \frac{\partial^{j_1 + \dots + j_n}}{\partial \bar{\rho}^{j_1 + \dots + j_n}} \prod_{k=0}^m \left(1 - \frac{\bar{\rho}_k}{\rho_k}\right)^{-1}. \quad (84)$$

Proposition 14. ([25]) *We have $\text{CV}(f) \subset \text{CV}(D_1^{j_1} \dots D_n^{j_n} f)$.*

Let $f \in \mathbb{C}^{\text{cv}}[[q_1, \dots, q_n]]$. Let $\{A_i\}_{i=0,1}$ be a polysystem defined as follows

$$A_i(q) = \sum_{j=1}^n A_i^j(q) \frac{\partial}{\partial q_j}, \quad (85)$$

where for any $j = 1, \dots, n$, $A_i^j(q) \in \mathbb{C}^{\text{cv}}[[q_1, \dots, q_n]]$.

Lemma 9. ([22]) *For $i = 0, 1$ and $j = 1, \dots, n$, one has $A_i \circ q_j = A_i^j(q)$. Thus,*

$$\forall i = 0, 1, \quad A_i(q) = \sum_{j=1}^n (A_i \circ q_j) \frac{\partial}{\partial q_j}.$$

Let $(\rho, \bar{\rho}, C_f), \{(\rho, \bar{\rho}, C_i)\}_{i=0,1}$ be respectively the convergence modulus at

$$q \in \text{CV}(f) \bigcap_{\substack{i=0,1 \\ j=1, \dots, n}} \text{CV}(A_i^j) \quad (86)$$

of f and $\{A_i^j\}_{j=1, \dots, n}$. Let us consider the following monoid morphisms

$$\mathcal{A}(\epsilon) = \text{identity} \quad \text{and} \quad C(\epsilon) = 1, \quad (87)$$

$$\forall w = vx_i, x_i \in X, v \in X^*, \quad \mathcal{A}(w) = \mathcal{A}(v)A_i \quad \text{and} \quad C(w) = C(v)C_i. \quad (88)$$

Lemma 10. ([20]) *For any word w , $\mathcal{A}(w)$ is continuous over $\mathbb{C}^{\text{cv}}[[q_1, \dots, q_n]]$ and, for any $f, g \in \mathbb{C}^{\text{cv}}[[q_1, \dots, q_n]]$, one has*

$$\mathcal{A}(w) \circ (fg) = \sum_{u, v \in X^*} \langle u \sqcup v \mid w \rangle (\mathcal{A}(u) \circ f)(\mathcal{A}(v) \circ g).$$

These notations are extended, by linearity, to $\mathbb{K}\langle X \rangle$ and we will denote $\mathcal{A}(w) \circ f|_q$ the evaluation of $\mathcal{A}(w) \circ f$ at q .

Definition 8. ([20]) *Let $f \in \mathbb{C}^{\text{cv}}[[q_1, \dots, q_n]]$. The generating series of the polysystem $\{A_i\}_{i=0,1}$ and of the observation f is given by*

$$\sigma f := \sum_{w \in X^*} \mathcal{A}(w) \circ f w \in \mathbb{C}^{\text{cv}}[[q_1, \dots, q_n]]\langle\langle X \rangle\rangle.$$

Then the following generating series is called Fliess generating series of the polysystem $\{A_i\}_{i=0,1}$ and of the observation f at q :

$$\sigma f|_q := \sum_{w \in X^*} \mathcal{A}(w) \circ f|_q w \in \mathbb{C}\langle\langle X \rangle\rangle.$$

Lemma 11. ([20]) *Let $\{A_i\}_{i=0,1}$ be a polysystem. Then, the map*

$$\sigma : (\mathbb{C}^{\text{cv}}[[q_1, \dots, q_n]], \cdot) \longrightarrow (\mathbb{C}^{\text{cv}}[[q_1, \dots, q_n]]\langle\langle X \rangle\rangle, \sqcup),$$

is an algebra morphism, i.e. for $f, g \in \mathbb{C}^{\text{cv}}[[q_1, \dots, q_n]]$ and $\mu, \nu \in \mathbb{C}$, one has:

$$\sigma(\nu f + \mu h) = \nu \sigma f + \mu \sigma g \quad \text{and} \quad \sigma(fg) = \sigma f \sqcup \sigma g.$$

Lemma 12. ([22]) *Let $\{A_i\}_{i=0,1}$ be a polysystem and $f \in \mathbb{C}^{\text{cv}}[[q_1, \dots, q_n]]$. Then*

$$\begin{aligned} \forall x_i \in X, \quad \sigma(A_i \circ f) = x_i \triangleleft \sigma f &\in \mathbb{C}^{\text{cv}}[[q_1, \dots, q_n]]\langle\langle X \rangle\rangle \\ \forall w \in X^*, \quad \sigma(\mathcal{A}(w) \circ f) = w \triangleleft \sigma f &\in \mathbb{C}^{\text{cv}}[[q_1, \dots, q_n]]\langle\langle X \rangle\rangle. \end{aligned}$$

Lemma 13. ([25]) *Let $\tau = \min_{1 \leq k \leq n} \rho_k$ and $r = \max_{1 \leq k \leq n} \bar{\rho}_k / \rho_k$. We have*

$$\begin{aligned} \|\mathcal{A}(w) \circ f\| &\leq C_f \frac{(n+1)}{(1-r)^n} \frac{C(w) |w|!}{\binom{n+|w|-1}{|w|}} \left[\frac{n}{\tau(1-r)^{n+1}} \right]^{|w|} \\ &\leq C_f \frac{(n+1)}{(1-r)^n} C(w) \left[\frac{n}{\tau(1-r)^{n+1}} \right]^{|w|} |w|!. \end{aligned}$$

Theorem 8. ([25]) *Let $K = C_f(n+1)(1-r)^{-n}$ and let χ be the real positive function defined over X^* by*

$$\forall i = 0, 1, \quad \chi(x_i) = \frac{C_i n}{\tau(1-r)^{(n+1)}}.$$

Then the generating series σf of the polysystem $\{A_i\}_{i=0,1}$ and of the observation f satisfies the χ -growth condition.

It is the same for the Fliess generating series $\sigma f|_q$ of the polysystem $\{A_i\}_{i=0,1}$ and of the observation f at q .

3.2 Polysystem and nonlinear differential equation

3.2.1 Nonlinear differential equation (with three singularities)

Let us consider the singular inputs²¹ $u_0(z) := z^{-1}$ and $u_1(z) := (1-z)^{-1}$, and the following nonlinear dynamical system

$$\begin{cases} y(z) &= f(q(z)), \\ \dot{q}(z) &= A_0(q) u_0(z) + A_1(q) u_1(z), \\ q(z_0) &= q_0, \end{cases} \quad (89)$$

where the state $q = (q_1, \dots, q_n)$ belongs to the complex analytic manifold of dimension n , q_0 is the initial state, the observation f belongs to $\mathbb{C}^{\text{cv}}[[q_1, \dots, q_n]]$ and $\{A_i\}_{i=0,1}$ is the polysystem defined on (85).

Definition 9. ([27]) *The following power series is called transport operator of the polysystem $\{A_i\}_{i=0,1}$ and of the observation f*

$$\mathcal{T} := \sum_{w \in X^*} \alpha_{z_0}^z(w) \mathcal{A}(w).$$

²¹These singular inputs are not included in the studies of Fliess motivated, in particular, by the renormalization of $y(z)$ at $+\infty$ [20, 22].

By the factorization of the monoid by Lyndon words, we have [27]

$$\mathcal{T} = (\alpha_{z_0}^z \otimes \mathcal{A}) \left(\sum_{w \in X^*} w \otimes w \right) = \prod_{l \in \text{Lynd} X} \exp[\alpha_{z_0}^z(S_l) \mathcal{A}(P_l)]. \quad (90)$$

The Chen generating series along the path $z_0 \rightsquigarrow z$, associated to ω_0, ω_1 is

$$S_{z_0 \rightsquigarrow z} := \sum_{w \in X^*} \langle S \mid w \rangle w \quad \text{with} \quad \langle S \mid w \rangle = \alpha_{z_0}^z(w) \quad (91)$$

which solves the differential equation (72) with the initial condition $S_{z_0 \rightsquigarrow z_0} = 1$. Thus, $S_{z_0 \rightsquigarrow z}$ and $L(z)L(z_0)^{-1}$ satisfy the same differential equation taking the same value at z_0 and $S_{z_0 \rightsquigarrow z} = L(z)L(z_0)^{-1}$.

Any Chen generating series $S_{z_0 \rightsquigarrow z}$ is group like [46] and depends only on the homotopy class of $z_0 \rightsquigarrow z$ [9]. The product of $S_{z_1 \rightsquigarrow z_2}$ and $S_{z_0 \rightsquigarrow z_1}$ is $S_{z_0 \rightsquigarrow z_2} = S_{z_1 \rightsquigarrow z_2} S_{z_0 \rightsquigarrow z_1}$. Let $\varepsilon \in]0, 1[$ and $z_i = \varepsilon \exp(i\beta_i)$, for $i = 0$ or 1 . We set $\beta = \beta_1 - \beta_0$. Let $\Gamma_0(\varepsilon, \beta_0)$ (resp. $\Gamma_1(\varepsilon, \beta_1)$) be the path turning around 0 (resp. 1) in the positive direction from z_0 to z_1 . By induction on the length of w , one has

$$|\langle S_{\Gamma_i(\varepsilon, \beta)} \mid w \rangle| = (2\varepsilon)^{|w|_{x_i}} \frac{\beta^{|w|}}{|w|!}, \quad (92)$$

where $|w|$ denotes the length of w and $|w|_{x_i}$ denotes the number of occurrences of letter x_i in w , for $i = 0$ or 1 . For ε tends to 0^+ , these estimations yield $S_{\Gamma_i(\varepsilon, \beta)} = e^{i\beta x_i} + o(\varepsilon)$. In particular, if $\Gamma_0(\varepsilon)$ (resp. $\Gamma_1(\varepsilon)$) is a circular path of radius ε turning around 0 (resp. 1) in the positive direction, starting at $z = \varepsilon$ (resp. $1 - \varepsilon$), then, by the noncommutative residue theorem [35, 32], we get

$$S_{\Gamma_0(\varepsilon)} = e^{2i\pi x_0} + o(\varepsilon) \quad \text{and} \quad S_{\Gamma_1(\varepsilon)} = e^{-2i\pi x_1} + o(\varepsilon). \quad (93)$$

Finally, the asymptotic behaviors of L on (73) give [32, 35]

$$S_{\varepsilon \rightsquigarrow 1-\varepsilon} \underset{\varepsilon \rightarrow 0^+}{\sim} e^{-x_1 \log \varepsilon} Z_{\sqcup\sqcup} e^{-x_0 \log \varepsilon}. \quad (94)$$

In other terms, $Z_{\sqcup\sqcup}$ is the regularized Chen generating series $S_{\varepsilon \rightsquigarrow 1-\varepsilon}$ of differential forms ω_0 and ω_1 : $Z_{\sqcup\sqcup}$ is the noncommutative generating series of the finite parts of the coefficients of the Chen generating series $e^{x_1 \log \varepsilon} S_{\varepsilon \rightsquigarrow 1-\varepsilon} e^{x_0 \log \varepsilon}$, i.e. the concatenation of $e^{x_0 \log \varepsilon}$ and then $S_{\varepsilon \rightsquigarrow 1-\varepsilon}$ and finally, $e^{x_1 \log \varepsilon}$.

3.2.2 Asymptotic behavior of the successive differentiation of solution via extended Fliess fundamental formula

The Fliess' fundamental formula [20] can be then extended as follows

Theorem 9. ([39]) *We have*

$$y(z) = \mathcal{T} \circ f|_{q_0} = \sum_{w \in X^*} \langle S_{z_0 \rightsquigarrow z} \mid w \rangle \langle \mathcal{A}(w) \circ f|_{q_0} \mid w \rangle = \langle \sigma f|_{q_0} \parallel S_{z_0 \rightsquigarrow z} \rangle.$$

By the factorization of the Lie exponential series L , the expansions of the output y of nonlinear dynamical system with singular inputs follow

Corollary 2.

$$\begin{aligned}
y(z) &= \sum_{w \in X^*} g_w(z) \mathcal{A}(w) \circ f|_{q_0} \\
&= \sum_{k \geq 0} \sum_{n_1, \dots, n_k \geq 0} g_{x_0^{n_1} x_1 \dots x_0^{n_k} x_1}(z) \operatorname{ad}_{A_0}^{n_1} A_1 \dots \operatorname{ad}_{A_0}^{n_k} A_1 e^{\log z A_0} \circ f|_{q_0} \\
&= \prod_{l \in \mathcal{L}^{yn} X} \exp \left(g_{S_l}(z) \mathcal{A}(P_l) \circ f|_{q_0} \right) \\
&= \exp \left(\sum_{w \in X^*} g_w(z) \mathcal{A}(\pi_1(w)) \circ f|_{q_0} \right),
\end{aligned}$$

where, for any word w in X^* , g_w belongs to the polylogarithm algebra.

Since $S_{z_0 \rightsquigarrow z} = L(z)L(z_0)^{-1}$ and since $\sigma f|_{q_0}$ and $L(z_0)^{-1}$ are invariant by $\partial_z = d/dz$ and by $\theta_0 = zd/dz$ then we obtain the n -order differentiation of y :

$$\partial_z^n y(z) = \langle \sigma f|_{q_0} \parallel \partial^n S_{z_0 \rightsquigarrow z} \rangle = \langle \sigma f|_{q_0} \parallel \partial_z^n L(z)L(z_0)^{-1} \rangle \quad (95)$$

and the n -order differentiation of y with respect to θ_0 :

$$\theta_0^n y(z) = \langle \sigma f|_{q_0} \parallel \theta_0^n S_{z_0 \rightsquigarrow z} \rangle = \langle \sigma f|_{q_0} \parallel \theta_0^n L(z)L(z_0)^{-1} \rangle. \quad (96)$$

With the notations of Proposition 5, we get respectively

$$\partial_z^n y(z) = \langle \sigma f|_{q_0} \parallel [D_n(z)L(z)]L(z_0)^{-1} \rangle = \langle \sigma f|_{q_0} \triangleright D_n(z) \parallel L(z)L(z_0)^{-1} \rangle, \quad (97)$$

$$\theta_0^n y(z) = \langle \sigma f|_{q_0} \parallel [E_n(z)L(z)]L(z_0)^{-1} \rangle = \langle \sigma f|_{q_0} \triangleright E_n(z) \parallel L(z)L(z_0)^{-1} \rangle. \quad (98)$$

For $z_0 = \varepsilon \rightarrow 0^+$, the asymptotic behavior and the renormalization at $z = 1$ of $\partial_z^n y(z)$ and $\theta_0^n y(z)$ (or the asymptotic expansion and the renormalization of its Taylor coefficients at $+\infty$) are deduced from (94) and extend a little bit the results of [39] as follows

Corollary 3. *For any integer n , we have respectively²²*

²²We overget, by Note 18, the n -order differentiation of y w.r.t. θ_1 :

$$\theta_1^n y(z) = \langle \sigma f|_{q_0} \parallel \theta_1^n S_{z_0 \rightsquigarrow z} \rangle = \langle \sigma f|_{q_0} \parallel \theta_1^n L(z)L(z_0)^{-1} \rangle.$$

Hence,

$$\theta_1^n y(z) = \langle \sigma f|_{q_0} \parallel [D_n(z) - E_n(z)]L(z)L(z_0)^{-1} \rangle = \langle \sigma f|_{q_0} \triangleright [D_n(z) - E_n(z)] \parallel L(z)L(z_0)^{-1} \rangle$$

and then

$$\theta_1^n y(1) \underset{\varepsilon \rightarrow 0^+}{\widetilde{=}} \sum_{w \in X^*} \langle \mathcal{A}(w) \circ f|_{q_0} \mid w \rangle \langle [D_n(1 - \varepsilon) - E_n(1 - \varepsilon)] e^{-x_1 \log \varepsilon} Z_{\perp\perp} e^{-x_0 \log \varepsilon} \mid w \rangle.$$

The actions of $\theta_0 = u_0(z)^{-1}d/dz$ and $\theta_1 = u_1(z)^{-1}d/dz$ over $y(z)$ are equivalent to those of the residuals respectively by x_0 and x_1 over $\sigma f|_{q_0}$ and correspond to *functional* differentiations [21] while $\partial_z = d/dz$ is the ordinary differentiation and is equivalent to the residual by $x_0 + x_1$.

$$\begin{aligned}\partial_z^n y(1) & \widetilde{\varepsilon \rightarrow 0^+} \sum_{w \in X^*} \langle \mathcal{A}(w) \circ f|_{q_0} \mid w \rangle \langle D_n(1 - \varepsilon) e^{-x_1 \log \varepsilon} Z_{\sqcup} e^{-x_0 \log \varepsilon} \mid w \rangle, \\ \theta_0^n y(1) & \widetilde{\varepsilon \rightarrow 0^+} \sum_{w \in X^*} \langle \mathcal{A}(w) \circ f|_{q_0} \mid w \rangle \langle E_n(1 - \varepsilon) e^{-x_1 \log \varepsilon} Z_{\sqcup} e^{-x_0 \log \varepsilon} \mid w \rangle.\end{aligned}$$

Corollary 4. *The n -order differentiation of the output y of the system (89) is a \mathcal{C} -combination of the elements g belonging to the polylogarithm algebra.*

Moreover, if the ordinary Taylor expansions of $\partial_z^n y$ and $\theta_0^n y$ exist :

$$\partial_z^n y(z) = \sum_{k \geq 0} d_k z^k \quad \text{and} \quad \theta_0^n y(z) = \sum_{k \geq 0} t_k z^k$$

then the coefficients of these expansions belong to the algebra of harmonic sums and there exist algorithmically computable coefficients $a_i, a'_i \in \mathbb{Z}, b_i, b'_i \in \mathbb{N}$ and $c_i, c'_i \in \mathbb{Z}[\gamma]$ such that

$$d_k \widetilde{k \rightarrow \infty} \sum_{i \geq 0} c_i k^{a_i} \log^{b_i} k \quad \text{and} \quad t_k \widetilde{k \rightarrow \infty} \sum_{i \geq 0} c'_i k^{a'_i} \log^{b'_i} k.$$

3.3 Differential realization

3.3.1 Differential realization

Definition 10. *The Lie rank of a formal power series $S \in \mathbb{K}\langle\langle X \rangle\rangle$ is the dimension of the vector space generated by*

$$\{S \triangleright \Pi \mid \Pi \in \mathcal{Lie}_{\mathbb{K}}\langle X \rangle\}, \quad \text{or respectively by} \quad \{\Pi \triangleleft S \mid \Pi \in \mathcal{Lie}_{\mathbb{K}}\langle X \rangle\}.$$

Example 10. *Let $P = x_1 x_0 x_1 + x_0 x_1 x_0 x_1$ be a polynomial of degree 4 over $X = \{x_0, x_1\}$. A basis up to degree 4 of the Lie algebra $\mathcal{Lie}_{\mathbb{K}}\langle X \rangle$ is following (c.f. Example 4)*

$$\{P_{x_0}, P_{x_1}, P_{x_0 x_1}, P_{x_0^2 x_1}, P_{x_0 x_1^2}, P_{x_0^3 x_1}, P_{x_0 x_1^3}, P_{x_0^2 x_1^2}\}.$$

The Lie-Hankel matrix of P is following

	1_{X^*}	x_1	$x_0 x_1$	$x_1 x_0 x_1$
P_{x_0}	0	0	0	1
P_{x_1}	0	0	1	0
$P_{x_0 x_1}$	0	-1	1	0
$P_{x_0^2 x_1}$	0	-2	0	0
$P_{x_0 x_1^2}$	-2	0	0	0
$P_{x_0^2 x_1^2}$	-2	0	0	0

and its Lie-Hankel rank of P is then 4.

Definition 11. Let $S \in \mathbb{K}\langle\langle X \rangle\rangle$ and let us put

$$\begin{aligned}\text{Ann}(S) &:= \{\Pi \in \mathcal{L}ie_{\mathbb{K}}\langle X \rangle \mid S \triangleright \Pi = 0\}, \\ \text{Ann}^\perp(S) &:= \{Q \in (\mathbb{K}\langle\langle X \rangle\rangle, \sqcup) \mid Q \triangleright \text{Ann}(S) = 0\}.\end{aligned}$$

It is immediate that $\text{Ann}^\perp(S) \ni S$ and it follows that (see [22, 48]).

Lemma 14. Let $S \in \mathbb{K}\langle\langle X \rangle\rangle$. If S is of finite Lie rank, d , then the dimension of $\text{Ann}^\perp(S)$ equals d .

By Lemma 3, the residuals are derivations for shuffle product. Then,

Lemma 15. Let $S \in \mathbb{K}\langle\langle X \rangle\rangle$. Then:

- (1) For any Q_1 and $Q_2 \in \text{Ann}^\perp(S)$, one has $Q_1 \sqcup Q_2 \in \text{Ann}^\perp(S)$.
- (2) For any $P \in \mathbb{K}\langle X \rangle$ and $Q_1 \in \text{Ann}^\perp(S)$, one has $P \triangleleft Q_1 \in \text{Ann}^\perp(S)$.

Definition 12. ([22]) The formal power series $S \in \mathbb{K}\langle\langle X \rangle\rangle$ is differentially produced if there exist

- an integer d ,
- a power series $f \in \mathbb{K}[[\bar{q}_1, \dots, \bar{q}_d]]$,
- a homomorphism \mathcal{A} from X^* maps to the algebra of differential operators generated by

$$\mathcal{A}(x_i) = \sum_{j=1}^d A_i^j(\bar{q}_1, \dots, \bar{q}_d) \frac{\partial}{\partial \bar{q}_j},$$

where, for $j = 1, \dots, d$, $A_i^j(\bar{q}_1, \dots, \bar{q}_d)$ belongs to $\mathbb{K}[[\bar{q}_1, \dots, \bar{q}_d]]$ such that

$$\forall w \in X^*, \quad \langle S \mid w \rangle = \mathcal{A}(w) \circ f|_0.$$

The couple (\mathcal{A}, f) is called the differential representation of S of dimension d .

Proposition 15. ([48]) Let $S \in \mathbb{K}\langle\langle X \rangle\rangle$. If S is differentially produced then it verifies the growth condition and its Lie rank is finite.

Proof. Let (\mathcal{A}, f) be a differential representation of S of dimension d . Then, by the notations of Definition 8, we get

$$\sigma f|_0 = S = \sum_{w \in X^*} (\mathcal{A}(w) \circ f)|_0 w.$$

For any $j = 1, \dots, d$, we put

$$T_j = \sum_{w \in X^*} \frac{\partial(\mathcal{A}(w) \circ f)}{\partial \bar{q}_j} w.$$

Firstly, by Theorem 8, the generating series σf verifies the growth condition. Secondly, for any $\Pi \in \mathcal{L}ie_{\mathbb{K}}\langle X \rangle$ and for any $w \in X^*$, one has

$$\langle \sigma f \triangleright \Pi \mid w \rangle = \langle \sigma f \mid \Pi w \rangle = \mathcal{A}(\Pi w) \circ f = \mathcal{A}(\Pi) \circ (\mathcal{A}(w) \circ f).$$

Since $\mathcal{A}(\Pi)$ is a derivation over $\mathbb{K}[\bar{q}_1, \dots, \bar{q}_d]$:

$$\begin{aligned} \mathcal{A}(\Pi) &= \sum_{j=1}^d (\mathcal{A}(\Pi) \circ \bar{q}_j) \frac{\partial}{\partial \bar{q}_j}, \\ \Rightarrow \mathcal{A}(\Pi) \circ (\mathcal{A}(w) \circ f) &= \sum_{j=1}^d (\mathcal{A}(\Pi) \circ \bar{q}_j) \frac{\partial (\mathcal{A}(w) \circ f)}{\partial \bar{q}_j} \end{aligned}$$

then we deduce that

$$\begin{aligned} \forall w \in X^*, \quad \langle \sigma f \triangleright \Pi \mid w \rangle &= \sum_{j=1}^d (\mathcal{A}(\Pi) \circ \bar{q}_j) \langle T_j \mid w \rangle, \\ \iff \sigma f \triangleright \Pi &= \sum_{j=1}^d (\mathcal{A}(\Pi) \circ \bar{q}_j) T_j. \end{aligned}$$

That means $\sigma f \triangleright \Pi$ is \mathbb{K} -linear combination of $\{T_j\}_{j=1, \dots, d}$ and the dimension of the vector space $\text{span}\{\sigma f \triangleright \Pi \mid \Pi \in \mathcal{L}ie_{\mathbb{K}}\langle X \rangle\}$ is less than or equal to d . \square

3.3.2 Fliess' local realization theorem

Proposition 16. ([48]) *Let $S \in \mathbb{K}\langle\langle X \rangle\rangle$ be such that its Lie rank equals d . Then there exists a basis $S_1, \dots, S_d \in \mathbb{K}\langle\langle X \rangle\rangle$ of $(\text{Ann}^\perp(S), \sqcup) \cong (\mathbb{K}[S_1, \dots, S_d], \sqcup)$ such that the S_i 's are proper and for any $R \in \text{Ann}^\perp(S)$, one has*

$$R = \sum_{i_1, \dots, i_d \geq 0} \frac{r_{i_1, \dots, i_d}}{i_1! \dots i_d!} S_1^{\sqcup i_1} \sqcup \dots \sqcup S_d^{\sqcup i_d},$$

where the coefficients $\{r_{i_1, \dots, i_d}\}_{i_1, \dots, i_d \geq 0}$ belong to \mathbb{K} and $r_{0, \dots, 0} = \langle R \mid 1_{X^*} \rangle$.

Proof. By Lemma 14, such a basis exists. More precisely, since the Lie rank of S is d then there exist $P_1, \dots, P_d \in \mathcal{L}ie_{\mathbb{K}}\langle X \rangle$ such that $S \triangleright P_1, \dots, S \triangleright P_d \in (\mathbb{K}\langle\langle X \rangle\rangle, \sqcup)$ are \mathbb{K} -linearly independent. By duality, there exist $S_1, \dots, S_d \in (\mathbb{K}\langle\langle X \rangle\rangle, \sqcup)$ such that

$$\forall i, j = 1, \dots, d, \quad \langle S_i \mid P_j \rangle = \delta_{i,j}, \quad \text{and} \quad R = \prod_{i=1}^d \exp(S_i \mid P_i).$$

Expanding this product, one obtains, via PBW theorem, the expected expression for the coefficients $\{r_{i_1, \dots, i_d} = \langle R \mid P_1^{i_1} \dots P_d^{i_d} \rangle\}_{i_1, \dots, i_d \geq 0}$. Hence, $(\text{Ann}^\perp(S), \sqcup)$ is generated by S_1, \dots, S_d . \square

With the notations of Proposition 16, one has respectively.

Corollary 5. *If $S \in \mathbb{K}[S_1, \dots, S_d]$ then, for any $i = 0, 1$ and for any $j = 1, \dots, d$, one has $x_i \triangleleft S \in \text{Ann}^\perp(S) = \mathbb{K}[S_1, \dots, S_d]$.*

Corollary 6. *The power series S verifies the growth condition if and only if, for any $i = 1, \dots, d$, S_i also verifies the growth condition.*

Proof. Assume there exists $j \in [1, \dots, d]$ such that S_j does not verify the growth condition. Since $S \in \text{Ann}^\perp(S)$ then using the decomposition of S on S_1, \dots, S_d , one obtains a contradiction with the fact that S verifies the growth condition.

Conversely, using Proposition 11, we get the expected results. \square

Theorem 10. ([22]) *The formal power series $S \in \mathbb{K}\langle\langle X \rangle\rangle$ is differentially produced if and only if its Lie rank is finite and if it verifies the χ -growth condition.*

Proof. By Proposition 15, one gets a direct proof.

Conversely, since the Lie rank of S equals d then by Proposition 16, by putting $\sigma f|_0 = S$ and, for any $j = 1, \dots, d$, $\sigma \bar{q}_j = S_j$,

(1) we choose the observation f as follows

$$f(\bar{q}_1, \dots, \bar{q}_d) = \sum_{i_1, \dots, i_d \geq 0} \frac{r_{i_1, \dots, i_d}}{i_1! \dots i_d!} \bar{q}_1^{i_1} \dots \bar{q}_d^{i_d} \in \mathbb{K}[[\bar{q}_1, \dots, \bar{q}_d]]$$

such that

$$\sigma f|_0(\bar{q}_1, \dots, \bar{q}_d) = \sum_{i_1, \dots, i_d \geq 0} \frac{r_{i_1, \dots, i_d}}{i_1! \dots i_d!} (\sigma \bar{q}_1)^{\sqcup i_1} \sqcup \dots \sqcup (\sigma \bar{q}_d)^{\sqcup i_d},$$

(2) it follows that, for $i = 0, 1$ and for $j = 1, \dots, d$, the residual $x_i \triangleleft \sigma \bar{q}_j$ belongs to $\text{Ann}^\perp(\sigma f|_0)$ (see also Lemma 15),

(3) since σf verifies the χ -growth condition then, by Corollary 6, the generating series $\sigma \bar{q}_j$ and $x_i \triangleleft \sigma \bar{q}_j$ (for $i = 0, 1$ and for $j = 1, \dots, d$) verify also the growth condition. We then take (see Lemma 12)

$$\forall i = 0, 1, \quad \forall j = 1, \dots, d, \quad \sigma A_j^i(\bar{q}_1, \dots, \bar{q}_d) = x_i \triangleleft \sigma \bar{q}_j,$$

by expressing σA_j^i on the basis $\{\sigma \bar{q}_i\}_{i=1, \dots, d}$ of $\text{Ann}^\perp(\sigma f|_0)$,

(4) the homomorphism \mathcal{A} is then determined as follows

$$\forall i = 0, 1, \quad \mathcal{A}(x_i) = \sum_{j=0}^d A_j^i(\bar{q}_1, \dots, \bar{q}_d) \frac{\partial}{\partial \bar{q}_j},$$

where, by Lemma 9, one has $A_j^i(\bar{q}_1, \dots, \bar{q}_d) = \mathcal{A}(x_i) \circ \bar{q}_j$.

Thus, (\mathcal{A}, f) provides a differential representation²³ of dimension d of S . \square

²³In [22, 48], the reader can find the discussion on the *minimal* differential representation.

Moreover, one also has the following

Theorem 11. ([22]) *Let $S \in \mathbb{K}\langle\langle X \rangle\rangle$ be a differentially produced formal power series. Let (\mathcal{A}, f) and (\mathcal{A}', f') be two differential representations of dimension n of S . There exist a continuous and convergent automorphism h of \mathbb{K} such that*

$$\forall w \in X^*, \forall g \in \mathbb{K}, \quad h(\mathcal{A}(w) \circ g) = \mathcal{A}'(w) \circ (h(g)) \quad \text{and} \quad f' = h(f).$$

Since any rational power series verifies the growth condition and its Lie rank is less than or equal to its Hankel rank which is finite [22] then

Corollary 7. *Any rational power series and any polynomial over X with coefficients in \mathbb{K} are differentially produced.*

References

- [1] Abe, E.: Hopf algebra. Cambridge (1980)
- [2] Bender C.M., Brody D. C., Meister B.K.: Quantum field theory of partitions, J. Math. Phys. **40**, pp. 3239–3245.
- [3] Berstel, J., Reutenauer, C.: Rational series and their languages. Springer-Verlag (1988)
- [4] C. Bui, G. H. E. Duchamp, V. Hoang Ngoc Minh: Schützenberger’s factorization on the (completed) Hopf algebra of q –shuffle product (in preparation).
- [5] C. Bui, G. H. E. Duchamp, N. Hoang, V. Hoang Ngoc Minh, C. Tollu: Combinatorics of ϕ -deformed shuffle Hopf algebras.
- [6] Cartier, P. – *Développements récents sur les groupes de tresses. Applications à la topologie et à l’algèbre*, Sémin BOURBAKI, 42^{ème} 1989-1990, n°716.
- [7] Chari, R., Pressley, A.: A guide to quantum group. Cambridge (1994)
- [8] K.T. Chen, R.H. Fox, R.C. Lyndon: Free differential calculus, IV. The quotient groups of the lower central series, Ann. of Math. , 68 (1958) pp. 8195
- [9] Chen, K.T.: Iterated path integrals. Bull. Am. Math. Soc. **83**, 831-879 (1977)
- [10] Connes A., Kreimer D.: Hopf algebras, renormalization and noncommutative geometry, Comm. Math. Phys., 199 (1998), pp. 203242.
- [11] M. Deneufchâtel, G.H.E. Duchamp, Hoang Ngoc Minh, A.I. Solomon: *Independence of hyperlogarithms over function fields via algebraic combinatorics*, dans Lecture Notes in Computer Science (2011), Volume 6742/2011, 127-139.

- [12] M. Deneufchâtel, G.H. E. Duchamp, Hoang Ngoc Minh: Radford bases and Schützenberger’s Factorizations, arXiv:1111.6759 (November 2011)
- [13] Duchamp, G., Reutenauer, C.: Un critère de rationalité provenant de la géométrie noncommutative. *Invent. Math.* 613-622 (1997)
- [14] G.H.E. Duchamp, Hoang Ngoc Minh, A.I. Solomon, S. Goodenough: An interface between physics and number theory, in *Journal of Physics* (2011), 284(1), pp. 012-023.
- [15] Duchamp, G.H.E., Tollu, C.: Sweedler’s duals and Schützenberger’s calculus. “Conference on Combinatorics and Physics”, arXiv: 0712.0125v3
- [16] Dyson, F.J.: The radiation theories of Tomonaga, Schwinger and Feynman, *Physical Rev*, vol 75, (1949), pp. 486-502.
- [17] Costermans, C., Enjalbert, J.Y., Hoang Ngoc Minh: Algorithmic and combinatorial aspects of multiple harmonic sums. *Discrete Mathematics & Theoretical Computer Science Proceedings* (2005)
- [18] Feynman R.P., Hibbs A.R.: *Quantum Mechanics and Path Integrals*, McGraw-Hill, New York, (1965).
- [19] Fliess, M.: Matrices de Hankel. *J. Math. Pures Appl.* **53**, 197-222 (1974)
- [20] Fliess, M.: Fonctionnelles causales non linéaires et indéterminées non commutatives. *Bull. SMF* **109**, 3-40 (1981)
- [21] Fliess, M.: *Vers une notion de dérivation fonctionnelle causale*. *Annales de l’institut Henri Poincaré (C) Analyse non linéaire*, 3 no. 1 (1986), p. 67-76
- [22] Fliess, M.: *Réalisation locale des systèmes non linéaires, algèbres de Lie filtrées transitives et séries génératrices non commutatives*. *Inven. Math.* **71**(3), 521-537 (1983)
- [23] Lê, T.Q.T., Murakami, J.: Kontsevich’s integral for Kauffman polynomial. *Nagoya Math.*, 39-65 (1996)
- [24] Hespel, C.: Une étude des séries formelles noncommutatives pour l’Approximation et l’Identification des systèmes dynamiques, thèse docteur d’état, Université Lille 1 (1998)
- [25] Hoang Ngoc Minh: Contribution au développement d’outils informatiques pour résoudre des problèmes d’automatique non linéaire, Thèse, Lille, 1990
- [26] Hoang Ngoc Minh: Input/Output behaviour of nonlinear control systems: about exact and approximated computations. *IMACS-IFAC Symposium*, Lille, Mai 1991.

- [27] Hoang Ngoc Minh, Jacob, G., Oussous, N.: Input/Output Behaviour of Nonlinear Control Systems: Rational Approximations, Nilpotent structural Approximations. in *Analysis of controlled Dynamical Systems*, (B. Bonnard, B. Bride, J.P. Gauthier & I. Kupka eds.), Progress in Systems and Control Theory, Birkhäuser, 253-262 (1991)
- [28] Hoang Ngoc Minh: Summations of polylogarithms via evaluation transform. dans Math. & Comput. Simul. **1336**, 707-728 (1996)
- [29] Hoang Ngoc Minh: Fonctions de Dirichlet d'ordre n et de paramètre t . dans Discrete Math. **180**, 221-242 (1998)
- [30] Hoang Ngoc Minh: Calcul symbolique non commutatif: aspects combinatoires des fonctions spéciales et des nombres spéciaux. HDR, Lille 2000
- [31] Hoang Ngoc Minh, Jacob, G.: Symbolic integration of meromorphic differential systems via Dirichlet functions. Discrete Math. **210**, 87-116 (2000)
- [32] Hoang Ngoc Minh, Jacob, G., Oussous, N.E., Petitot, M.: Aspects combinatoires des polylogarithmes et des sommes d'Euler-Zagier. journal électronique du Séminaire Lotharingien de Combinatoire, B43e (2000)
- [33] Hoang Ngoc Minh, Jacob, G., Oussous, N.E., Petitot, M.: De l'algèbre des ζ de Riemann multivariées l'algèbre des ζ de Hurwitz multivariées. journal électronique du Séminaire Lotharingien de Combinatoire **44** (2001)
- [34] Hoang Ngoc Minh, Petitot, M.: Lyndon words, polylogarithmic functions and the Riemann ζ function. Discrete Math. **217**, 273-292 (2000)
- [35] Hoang Ngoc Minh, Petitot, M., Van der Hoeven, J.: Polylogarithms and Shuffle algebra. Proceedings of FPSAC'98 (1998)
- [36] Hoang Ngoc Minh, Petitot, M., Van der Hoeven, J.: L'algèbre des polylogarithmes par les séries génératrices. Proceedings of FPSAC'99 (1999)
- [37] Hoang Ngoc Minh: Finite polyzêtas, Poly-Bernoulli numbers, identities of polyzêtas and noncommutative rational power series. Proceedings of 4th International Conference on Words, 232-250 (2003)
- [38] Hoang Ngoc Minh: Differential Galois groups and noncommutative generating series of polylogarithms, in "Automata, Combinatorics and Geometry". 7th World Multi-conference on Systemics, Cybernetics and Informatics, Florida (2003)
- [39] Hoang Ngoc Minh: Algebraic combinatoric aspects of asymptotic analysis of nonlinear dynamical system with singular inputs. Acta Academiae Aboensis, Ser. B **67**(2), 117-126 (2007)
- [40] Hoang Ngoc Minh: On a conjecture by Pierre Cartier about a group of associators, in Acta Mathematica Vietnamica, Vol. 3, (2013).

- [41] Hoang Ngoc Minh.– *Structure of polyzetas and Lyndon words*, à paraître dans Vietnamese Mathematics Journal (2013), DOI 10.1007/10013-013-0047-x.
- [42] Hoffman, M.: The multiple harmonic series. Pac. J. Math. **152**(2), 275-290 (1992)
- [43] Hoffman, M.: The algebra of multiple harmonic series. J. Algebra (1997)
- [44] Hochschild, G.: The structure of Lie groups, Holden-Day (1965)
- [45] D.E. Radford: A natural ring basis for shuffle algebra and an application to group schemes, Journal of Algebra, 58, pp. 432-454, 1979.
- [46] Ree, R.: Lie elements and an algebra associated with shuffles. Ann. Math. **68**, 210-220 (1985)
- [47] Reutenauer, C.: Free Lie Algebras. London Math. Soc. Monographs (1993)
- [48] Reutenauer, C.: The local realisation of generating series of finite Lie rank. Algebraic and Geometric Methods In Nonlinear Control Theory, 33-43
- [49] Schützenberger, M.P.: On the definition of a family of automata. Information and Control **4**, 245-270 (1961)
- [50] Viennot, G: Algèbres de Lie Libres et Monoïdes Libres, Lecture Notes in Mathematics, (1978) 691: 94-112, January 01, 1978.
- [51] Zagier, D.: Values of zeta functions and their applications. in “First European Congress of Mathematics”, vol. 2, Birkhäuser, 497-512 (1994)