



**HAL**  
open science

## Amino-terminal sequencing of the ovine milk fat globule membrane protein butyrophilin

Lucile Montagne, Jean-Michel Girardet, Gérard Humbert

► **To cite this version:**

Lucile Montagne, Jean-Michel Girardet, Gérard Humbert. Amino-terminal sequencing of the ovine milk fat globule membrane protein butyrophilin. *Le Lait*, 1999, 79 (2), pp.275-279. hal-00929650

**HAL Id: hal-00929650**

**<https://hal.science/hal-00929650>**

Submitted on 11 May 2020

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Note

## Amino-terminal sequencing of the ovine milk fat globule membrane protein butyrophilin

Lucile Montagne<sup>a</sup>, Jean-Michel Girardet<sup>b\*</sup>, Gérard Humbert<sup>b</sup>

<sup>a</sup> Laboratoire du jeune ruminant, Inra, 65, rue de Saint-Brieuc, 35042 Rennes cedex, France

<sup>b</sup> Laboratoire des biosciences de l'aliment, unité associée à l'Inra, université Henri-Poincaré-Nancy-1, 54506 Vandœuvre-lès-Nancy cedex, France

(Received 30 April 1998; accepted 30 July 1998)

**Abstract** — Butyrophilin, a transmembrane protein of the milk fat globule membrane, was identified in cows' and ewes' milk by amino acid N-terminal sequencing. The N-terminal part of ovine butyrophilin was identical to that of the human protein. A butyrophilin-type protein with an  $M_r$  of 62 000 was also found in equimolar proportions with butyrophilin in ovine milk. This smaller  $M_r$  form was probably generated by plasmin-mediated hydrolysis during or following lactation in the epithelial cells of the mammary gland and corresponded to a smaller  $M_r$  form of butyrophilin lacking the C-terminal region. © Inra/Elsevier, Paris

**butyrophilin / milk fat globule membrane / ovine milk**

**Résumé** — **Séquence N-terminale de la butyrophiline de la membrane des globules gras du lait de brebis.** La butyrophiline, une protéine transmembranaire de la membrane des globules gras, a été identifiée dans les laits de vache et de brebis par séquençage de son extrémité N-terminale. Les 14 résidus d'acides aminés N-terminaux de la butyrophiline ovine sont identiques à ceux de la protéine humaine. De plus, une protéine de type butyrophiline et de poids moléculaire 62 000 a été mise en évidence en proportion équimolaire avec la butyrophiline dans le lait ovine. Cette forme ayant un poids moléculaire plus petit serait produit par action de la plasmine associée aux membranes dans les cellules épithéliales de la glande mammaire au cours ou après la lactation et correspondrait à de la butyrophiline dont la partie C-terminale aurait été tronquée. © Inra/Elsevier, Paris

**butyrophiline / membrane des globules gras / lait de brebis**

---

\* Correspondence and reprints. Jean-Michel.Girardet@scbiol.u-nancy.fr

## 1. INTRODUCTION

Butyrophilin is an acidic glycoprotein with apparent  $M_r$  of 67 000 associated with the membrane surrounding fat droplets in milk. It constitutes 40 % of the total milk fat globule membrane (MFGM) protein (for review, see [14]). This protein may function in the process of milk secretion because it is specifically expressed on the apical surface of mammary secretory cells during lactation. Butyrophilin and xanthine oxidase (EC 1.2.3.22) may form a supramolecular complex with low  $M_r$  guanosine triphosphate (GTP)-binding proteins in the membrane [3, 8, 15].

Butyrophilin has been detected in several species, including cow, ewe, goat, pig, human, rat, guinea pig [4] and mouse [8]. cDNA clones encoding bovine, murine and human butyrophilin were isolated and the primary structures of the three proteins deduced from the DNA sequences [8, 9, 17, 20]. The 19 N-terminal amino acid residues of guinea pig butyrophilin were sequenced [16]. It appears that butyrophilin is composed of two extracellular immunoglobulin superfamily domains (immunoglobulin V [IgV]- and immunoglobulin C1 [IgC1]-type domains), a transmembrane domain and an intracellular domain homologous to the B30.2 domain of several intracellular proteins [5, 6], including the *ret* finger protein (RFP), the 52 000  $M_r$  nuclear antigen A of Sjögren's syndrome (SS-A/Ro) and *Xenopus* nuclear factor 7 [9, 20, 21]. The N-terminal extracellular domain shares similarities to myelin/oligodendrocyte glycoprotein (MOG) and to the major histocompatibility complex chicken B blood group system (B-G) proteins [11, 21]. This particular structure of butyrophilin suggests a cell surface receptor function [20].

Butyrophilin has a potential function in lactation and a hypothetical role in autoimmune diseases such as human multiple sclerosis. In fact, the IgV N-terminal region of MOG is able to induce acute demyelinat-

ing experimental autoimmune encephalomyelitis in rats [1]. Moreover, significant correlations were found between liquid cow milk, cream or butter consumption and the prevalence of multiple sclerosis [12]. The bovine and ovine species are of economical and biotechnological interest. Ewes' milk has a higher fat content ( $70 \text{ g}\cdot\text{L}^{-1}$ ) than cows' milk ( $38 \text{ g}\cdot\text{L}^{-1}$ ), related to a high level of expression of MFGM proteins. In this study, ovine butyrophilin was separated by electrophoresis, electrotransferred onto a membrane and then characterized by amino acid N-terminal sequencing. Two forms of butyrophilin with  $M_s$  close to 66 000 and 62 000, respectively, were found in equimolar proportions in ovine milk.

## 2. MATERIALS AND METHODS

### 2.1. Extraction of MFGM

Cream was separated from 5 L of bovine or ovine milk by centrifugation at  $3\ 000 \text{ g}$  and  $30^\circ\text{C}$  for 15 min. The cream was washed three times in five volumes of  $100 \text{ mmol}\cdot\text{L}^{-1}$  Tris buffer, pH 7.2, containing  $250 \text{ mmol}\cdot\text{L}^{-1}$  sucrose,  $100 \text{ mmol}\cdot\text{L}^{-1}$   $\text{MgCl}_2$ ,  $0.24$  inhibiting U·mL<sup>-1</sup> trypsin inhibitor from egg white (Serva Feinbiochemica, D-69042 Heidelberg 1, Germany) and  $5 \text{ mmol}\cdot\text{L}^{-1}$   $\epsilon$ -aminocaproic acid. One volume of cream suspended in two volumes of Tris buffer were shaken on a laboratory shaker until butter formed. The mixture was then warmed to  $40^\circ\text{C}$  to release membrane trapped by butter granules and centrifuged at  $100\ 000 \text{ g}$  at  $4^\circ\text{C}$  for 1 h. The MFGM pellet was finally freeze-dried [7].

### 2.2. Isolation and sequencing of butyrophilin

The MFGM ( $2 \text{ mg}\cdot\text{mL}^{-1}$ ) from bovine or ovine milk was solubilized in a  $55 \text{ mmol}\cdot\text{L}^{-1}$  Tris buffer at pH 6.8 containing  $23 \text{ mg}\cdot\text{mL}^{-1}$  sodium dodecyl sulphate (SDS) and 13.3 % (v/v) 2-mercaptoethanol, and boiled for 3 min. The MFGM proteins were then separated by SDS-polyacrylamide gel electrophoresis (SDS-PAGE, [10]) and electrophoretically transferred to polyvinylidene difluoride (PVDF) membranes. The trans-


fer buffer was composed of 10 mmol·L<sup>-1</sup> CAPS (3-[cyclohexylamino]-1-propanesulfonic acid; Sigma Chemical Co., Saint-Louis, MO, USA) at pH 11, containing 10 % (v/v) methanol. N-terminal sequences of the bound proteins were carried out by Edman degradation using a 476A sequencer from Applied Biosystems (Foster City, CA, USA).

### 3. RESULTS AND DISCUSSION

On the SDS-PAGE profile of bovine MFGM proteins (*figure 1*), the principal band was located at an apparent  $M_r$  of 66 000 and corresponded to butyrophilin as demonstrated by N-terminal amino acid residue sequencing (*figure 2*). A minor component with apparent  $M_r$  62 000 was associated with bovine butyrophilin and was hardly detectable on the SDS-PAGE profile. Bovine butyrophilin represented 38 % of the total MFGM proteins (determined by densitometry).

The ovine MFGM contained also a 66 000  $M_r$  butyrophilin-type protein. Sequence alignments showed a strong identity between the two species. The N-terminal region presented only one mutation site, a proline residue instead of a glutamine residue at position 9 (*figure 2*). Recently, in the case of bovine butyrophilin, both glutamine and asparagine residues were detected at position 9 [2], due to allelic polymorphism. The N-terminal region of the ovine butyrophilin was 100 % identical to that of human butyrophilin. The whole human protein sequence is 84 % identical to the bovine sequence [20]. Both proteins share the same number of residues and contain no gaps in their alignment. Structural domains between bovine butyrophilin and proteins of the other species (human, guinea pig, mouse and ewe, most likely) are similar, which suggests a conserved function between species.

In our study, an additional 62 000  $M_r$  butyrophilin-type protein was identified in ovine MFGM (*figure 1*). The N-terminal



**Figure 1.** Electrophoresis (15 % [w/v] polyacrylamide, 0.1 % [w/v] sodium dodecyl sulphate, pH 8.8) of supernatants (Supern.) and pellets (Pellet) after ultracentrifugation of the bovine and ovine milk fat globule membranes.

**Figure 1.** Électrophorèse (polyacrylamide à 15 % (p/v), sulfate de dodécyle et de sodium à 0,1 % (p/v), pH 8,8) des surnageants (Supern.) et culots (Pellet) après ultracentrifugation des membranes des globules gras de laits bovin et ovin.

region of this low  $M_r$  butyrophilin-type protein was 100 % identical to that of ovine butyrophilin. The 66 000 and 62 000  $M_r$  proteins represented 13 and 13.6 % of the total membrane proteins, respectively. In another study [4], the SDS-PAGE profiles of bovine and ovine MFGM were similar and the low  $M_r$  butyrophilin-type protein of the ovine MFGM was a minor component. Other minor fragments of bovine butyrophilin with smaller  $M_r$  are also found in the MFGM [2]. Butyrophilin from human MFGM migrates by SDS-PAGE as a doublet of about  $M_r$  65 000 and 60 000 [20].

Smaller  $M_r$  forms of butyrophilin are probably proteolytic fragments either produced in situ by MFGM-associated plasmin or during isolation of MFGM [2, 13]. However, we did include inhibitors of plasmin

## Present work

Low $M_r$ butyrophilin-type protein of ovine MFGM	APFD VIGPPE-
Butyrophilin of ovine MFGM	APFD VIGPPEPILA-
Butyrophilin of bovine MFGM	APFD VIGPQE-

## Previous studies

	1	11	21	31	41
Bovine butyrophilin [9]	MAVFPNSCLA GCLLIFILLQ LPKLDS			.APFD VIGPQEPILA	VVGED.AELPC-
[2]	signal peptide			Q	N
Human butyrophilin [20]				.APFD VIGPPEPILA	VVGED.AELPC-
Mouse butyrophilin [8]				.....VLA	LVGSDDAELTC-
Mouse butyrophilin [17]	MAVFPNSCLL VCLLTLTVLQ LPTLDS			AAAPFD VTAPQEPVLA	LVGSD.AELTC-
	signal peptide				
Guinea pig butyrophilin [16]				..RFD VIGPTEPVLA	AVGGD.A-

**Figure 2.** N-terminal primary structures of the bovine and ovine butyrophilins and of the ovine low  $M_r$  butyrophilin-type protein compared to those of butyrophilins of various species reported in the literature. Amino acid residues different to those of bovine butyrophilin are in bold-face type.

**Figure 2.** Structures primaires N-terminales des butyrophilines bovine, ovine et de la protéine ovine de faible poids moléculaire de type butyrophiline comparées aux régions N-terminales des butyrophilines de diverses espèces rapportées dans la littérature. Les résidus d'acides aminés différents de ceux de la butyrophiline bovine sont en caractères gras.

activity during the MFGM extraction (inhibitor from egg white and  $\epsilon$ -aminocaproic acid). Proteolysis could occur earlier, i.e., during or following lactation, as in the case of proteose peptones generated by plasmin activity on  $\beta$ -casein in the epithelial cells of the mammary gland (for review, see [18]), or C-terminal fragment of component PP3 [19]. Plasmin activity might be greater in ewes' milk compared with cows' milk due to an elevated activation of the plasminogen associated with ovine MFGM.

We cannot exclude the possibility that other butyrophilin-like proteins are associ-

ated with the MFGM. In humans, two genes (BT2, BT3 or B7c) which share similarities with the canonical structure of butyrophilin have been found [15]. These genes are localized in the major histocompatibility complex region close to the butyrophilin gene. Nevertheless, putative products of these genes share only 50 % identity with the amino acid sequence of butyrophilin.

Further investigations will be carried out in order to understand why more of the low  $M_r$  form of butyrophilin is generated in ovine milk compared with bovine milk,

either by proteolysis of the 66 000  $M_r$  form of butyrophilin, or by expression of a BT gene.

## ACKNOWLEDGEMENTS

We thank J. Henry, laboratoire de biochimie, faculté de médecine, Limoges, and F. Saulnier, service commun de séquences des protéines, université Henri-Poincaré, for good advice.

## REFERENCES

- [1] Adelmann M., Wood J., Benzel P., Fiori H., Lassmann H., Matthieu J.M., Gardinier M.V., Dornmair K., Linington C., The N-terminal domain of the myelin oligodendrocyte glycoprotein (MOG) induces acute demyelinating experimental autoimmune encephalomyelitis in the Lewis rat, *J. Neuroimmunol.* 63 (1995) 17–27.
- [2] Banghart L.R., Chamberlain C.W., Velarde J., Korobko I.V., Ogg S.L., Jack L.J.W., Vakharia V.N., Mather I.H., Butyrophilin is expressed in mammary epithelial cells from a single-sized messenger RNA as a type I membrane glycoprotein, *J. Biol. Chem.* 273 (1998) 4171–4179.
- [3] Ghosal D., Ankrapp D., Keenan T.W., Low molecular mass GTP binding proteins are secreted from mammary epithelial cells in association with lipid globules, *Biochim. Biophys. Acta* 1168 (1993) 299–306.
- [4] Heid H.W., Winter S., Bruder G., Keenan T.W., Jarasch E.D., Butyrophilin, an apical plasma membrane-associated glycoprotein characteristic of lactating mammary glands of diverse species, *Biochim. Biophys. Acta* 728 (1983) 228–238.
- [5] Henry J., Ribouchon M.T., Depetris D., Mattei, M.G., Offer C., Tazi- Ahnini R., Pontarotti P., Cloning, structural analysis and mapping of the B30 and B7 multigenic families to the major histocompatibility complex (MHC) and other chromosomal regions, *Immunogenetics* 46 (1997) 383–395.
- [6] Henry J., Ribouchon M.T., Offer C., Pontarotti P., B30.2 like domain proteins: a growing family, *Biochem. Biophys. Res. Commun.* 235 (1997) 162–165.
- [7] Imam A., Laurence J.R., Neville A.M., Isolation and characterisation of a major glycoprotein from milk-fat-globule membrane of human breast milk, *Biochem. J.* 193 (1981) 47–54.
- [8] Ishii T., Aoki N., Noda A., Adachi T., Nakamura R., Matsuda T., Carboxy-terminal cytoplasmic domain of mouse butyrophilin specifically associated with a 150-kDa protein of mammary epithelial cells and milk fat globule membrane, *Biochim. Biophys. Acta* 1245 (1995) 285–292.
- [9] Jack L.J.W., Mather I.H., Cloning and analysis of cDNA encoding bovine butyrophilin, an apical glycoprotein expressed in mammary tissue and secreted in association with the milk-fat globule membrane during lactation, *J. Biol. Chem.* 265 (1990) 14481–14486.
- [10] Laemmli U.K., Favre M., Maturation of the head of bacteriophage T4. I. DNA packaging events, *J. Mol. Biol.* 80 (1973) 575–599.
- [11] Linsley P.S., Peach R., Gladstone P., Bajortah J., Extending the B7 (CD80) gene family, *Protein Sci.* 3 (1994) 1341–1343.
- [12] Malosse D., Perron H., Sascio A., Seigneurin J.M., Correlation between milk and dairy product consumption and multiple sclerosis prevalence: a worldwide study, *Neuroepidemiology* 11 (1992) 304–312.
- [13] Mather I.H., Proteins of the milk-fat-globule membrane as markers of mammary epithelial cells and apical plasma membrane, in: Neville M.C., Daniel C.W. (Eds.), *The Mammary Gland: Development, Regulation and Function*, Plenum Publication Co., New York, 1987, pp. 217–267.
- [14] Mather I.H., Jack L.J.W., A review of the molecular and cellular biology of butyrophilin, the major protein of bovine milk fat globule membrane, *J. Dairy Sci.* 76 (1993) 3832–3850.
- [15] Mondy B.L., Keenan T.W., Butyrophilin and xanthine oxidase occur in constant molar proportions in milk lipid globule membrane but vary in amount with breed and stage of lactation, *Protoplasma* 177 (1993) 32–36.
- [16] Neira L.M., Mather I.H., Biochemical and immunological comparison of bovine butyrophilin with a butyrophilin-like glycoprotein in guinea pig milk-fat-globule membrane, *Protoplasma* 159 (1990) 168–178.
- [17] Ogg S.L., Komaragiri M.V.S., Mather I.H., Structural organization and mammary-specific expression of the butyrophilin gene, *Mamm. Genome* 7 (1996) 900–905.
- [18] Pâquet D., Review: the proteose-peptone fraction of milk, *Lait* 69 (1989) 1–21.
- [19] Sørensen E.S., Petersen T.E., Phosphorylation, glycosylation and amino acid sequence of component PP3 from the proteose peptone fraction of bovine milk, *J. Dairy Sci.* 60 (1993) 535–542.
- [20] Taylor M.R., Peterson J.A., Ceriani R.L., Couto J.R., Cloning and sequence analysis of human butyrophilin reveals a potential receptor function, *Biochim. Biophys. Acta* 1306 (1996) 1–4.
- [21] Vernet C., Boretto J., Mattei M.G., Takahashi M., Jack L.J.W., Mather I.H., Rouquier S., Pontarotti P., Evolutionary study of multigenic families mapping close to the human MHC class I region, *J. Mol. Evol.* 37 (1993) 600–612.