



## Potato soil-borne diseases. A review

Fiers, Edel-Hermann, Chatot, Hingrat, Alabouvette, Steinberg

### ► To cite this version:

Fiers, Edel-Hermann, Chatot, Hingrat, Alabouvette, et al.. Potato soil-borne diseases. A review. *Agronomy for Sustainable Development*, 2012, 32 (1), pp.93-132. 10.1007/s13593-011-0035-z . hal-00930506

**HAL Id:** hal-00930506

<https://hal.science/hal-00930506>

Submitted on 11 May 2020

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

Copyright

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# Potato soil-borne diseases. A review

Marie Fiers · Véronique Edel-Hermann ·

Catherine Chatot · Yves Le Hingrat ·

Claude Alabouvette · Christian Steinberg

Accepted: 15 October 2010 / Published online: 24 June 2011  
© INRA and Springer Science+Business Media B.V. 2011

**Abstract** Potato crop is the fourth main food crop in the world and it will certainly feed a big part of the global population in the next years. The economical outlets for this crop are great; however, numerous diseases either soil- or airborne can cause huge losses in the production. Worldwide, about 40 soil-borne diseases affect potato and cause severe damages especially on tubers, the economically most important part of the plant. The occurrence and development of soil-borne diseases depend on very diverse factors affecting either the pathogen or the plant. Favorable conditions for potato diseases development are frequently the same as the conditions needed for potato growth: temperature between 10°C and 25°C, high humidity, medium pH, etc. Adapted cultural practices such as a rotation longer than 4 years, appropriate fertilization and water management, an adapted delay between

haulm killing and harvest, and dry and cool conditions for tuber storage are good ways to control potato diseases. In most cases, potato pathogens develop specific survival forms, dissemination ways and host penetration methods. The genetic variability of the pathogens implies the use of adapted diagnostic and control methods. Decision support systems developed to predict yield losses allow choosing good control methods such as the use of healthy seeds, adapted pesticides, cultural practices, and biological control agents for each potato disease. The complexity of the interactions between a pathogen and its host, influenced by biotic and abiotic factors of the environment, make the control of the diseases often very difficult. However, deep knowledge of pathosystems allows setting up integrated pest management systems allowing the production of healthy and good quality potatoes.

M. Fiers · V. Edel-Hermann · C. Alabouvette · C. Steinberg (✉)  
INRA, Université de Bourgogne UMR 1229 Microbiologie du Sol et de l'Environnement, CMSE,  
17 rue Sully, BP 86510, 21065 Dijon cedex, France  
e-mail: christian.steinberg@dijon.inra.fr

V. Edel-Hermann  
e-mail: veronique.edel@dijon.inra.fr

C. Alabouvette  
e-mail: c.ala@agrene.fr

M. Fiers · C. Chatot  
Germicopa R&D, Kerguivarch,  
29520 Châteauneuf du Faou, France

C. Chatot  
e-mail: catherine.chatot@germicopa.fr

M. Fiers  
Bretagne Plants, Roudouhir,  
29460 Hanvec, France  
e-mail: mariefiers@hotmail.com

Y. Le Hingrat  
Bretagne Plants, FNPPPT, Roudouhir,  
29460 Hanvec, France  
e-mail: yves.lehingrat@fnpppt.fr

**Keywords** Cultural practices · Decision support system · Pathogen ecology · Pedologic and climatic factors · Plant microorganism interaction · Soil · Soil suppressiveness

## Contents

Introduction . . . . .	4
I. Effects of abiotic factors on the occurrence and development of soil-borne potato diseases . . . . .	10
I. 1. Soil temperature . . . . .	10
I. 2. Soil moisture . . . . .	13
I. 3. Soil texture . . . . .	14
I. 4. Soil pH . . . . .	17
I. 5. Soil organic matter . . . . .	17
II. Effects of biotic factors on the occurrence and development of soil-borne potato diseases . . . . .	18
II. 1. Autecology of pathogens . . . . .	18
1. 1. Inoculum sources, survival and dissemination pathways . . . . .	18
1. 2. Relationship between inoculum density and disease severity . . . . .	22

I. 3. Mechanisms of infection .....	22
I. 4. Genetic variability .....	28
I. 5. Diagnosis and detection methods .....	28
II. 2. Interactions between microorganisms, organisms and pathogens .....	30
II. 3. Interactions between plants and pathogens ..	34
III. Effects of cultural practices on the occurrence and development of soil-borne potato diseases .....	35
III. 1. Rotations .....	36
III. 2. Fertilization and amendments .....	36
III. 3. Tillage management .....	41
III. 4. Planting, haulm destruction, lifting and harvesting .....	41
III. 5. Pesticides .....	42
III. 6. Organic farming versus conventional agriculture .....	42
III. 7. Handling and storage .....	43
IV. Disease management .....	43
IV. 1. Risk assessment and decision support systems ..	44
IV. 2. Control methods .....	47
Conclusions .....	50
References .....	52

## 1 Introduction

Potato crop, the world's number one non-grain food commodity, is the fourth main food crop in the world after maize, rice and wheat, with 325 million tons produced in 2007. Potatoes are grown in more than 100 countries, mainly in Asia (135 million tons) and Europe (130 million tons; FAO 2008). They have good gustative and nutritional qualities and can be grown under various climates. This is the reason why Food and Agriculture Organization (FAO) has declared the year 2008 the international year of the potato. Indeed, potato can help fulfill the first United Nations Organization's (UNO) millennium development goal that aims at eradicating extreme poverty and hunger in the world. However, potato (*Solanum tuberosum*) crop can suffer more than 40 pests and diseases caused by insects, nematodes, viruses, bacteria, and fungi. Those pathogens are air- or soil-borne and cause damages on all parts of the plant. In this review, we will focus on soil-borne fungi, bacteria, and nematodes (Table 1, Fig. 1).

Indeed, diseases caused by viruses or viroids provoke generally foliar symptoms: leaf distortion, mosaic, crinkling, leaf and vein necroses, dwarfing, and leaf rolling. Only some viruses—tobacco rattle virus (TRV), potato mop-top virus, potato virus Y, and tobacco necrosis virus—can cause damages on tubers such as blemishes or rots in tuber flesh (Table 1). They will be briefly mentioned in Table 1 as well

as the vectors (aphids, fungi, or nematodes) involved in their transmission but they will not be detailed in this review.

Soil-borne diseases affecting potato crop can be divided into two groups depending on symptoms: symptoms damaging tubers and those damaging other parts of the plant (Gudmestad et al. 2007).

Diseases affecting stems or roots affect the crop development and may lead to a reduction of the yield (Table 1). Stem lesions can be watery and may develop into the stem pith with (stem rot) or without (blackleg, white mold) the formation of sclerotia. Other lesions can appear like more discrete light brown lesions but nevertheless affecting the yield of the crop (skin spot, stem canker). Some soil-borne pathogens sometimes cause aerial symptoms like necroses or chloroses (*Phoma* leaf spot, *Verticillium* wilt) occasionally associated with wilting and rolling (bacterial ring rot). Finally, root lesions, mainly caused by nematodes feeding on the roots, lead to necroses or rots. Nematodes feeding sites are good entry points for other soil microorganisms.

Among diseases affecting tubers, symptoms can be divided into three categories: galls, blemishes, and rots (Table 1). Galls consist in outgrowth and tuber deformation. The most frequent galls are provoked by powdery scab, wart, Common Scab, root-knot nematode, and false root-knot nematode. Blemishes affect only the tuber skin but they are now economically important since consumers' habits have changed and tubers are washed before selling. Blemishes can appear on the tuber surface as spots called black dot, black scurf, skin spot or powdery scab, or as areas of atypical appearance presenting a more or less pronounced scabby (common or netted scab) or silver (silver scurf) aspect. Rots, which affect the tuber flesh more deeply, include different types such as dry rots, soft rots (charcoal rot, leak, bacterial soft rot, black leg, and stem rot), flesh discoloration (pink rot) or vascular ring discoloration (ring rot, brown rot, *Verticillium* wilt, and *Fusarium* dry rots). Dry rot diseases also damage stored potatoes.

Potato is becoming a more and more important foodstuff in the world, it is therefore essential to control diseases which cause direct yield losses and decrease of farmer's incomes due to downgrading the quality of affected tubers. Therefore, knowledge about the pathogens as well as factors influencing disease severity is needed to setup efficient control strategies. Before reviewing the different causes of occurrence and development of the main soil-borne potato diseases, it is important to recall the concepts of soil inoculum potential and soil suppressiveness which describe the complex interactions between the soil, the pathogens, and the plant. While the former evaluates what the actual indigenous pathogenic inoculums could do in the rhizosphere towards the host plants if all conditions were favorable to its pathogenic activity, the second

**Table 1** Potato soil-borne pathogens

Pathogen	Disease	Host range	Main symptoms						Pathogenicity test	Distribution	References			
			Tubers		Other parts		Stem lesions	Leaf lesions						
			Gall	Blemish	Rot									
<b>Fungi and oomycetes</b>														
<i>Colletotrichum coccodes</i>	Black dot	Moderate: 35 hosts from 13 families including <i>Cucurbitaceae</i> , <i>Fabaceae</i> and <i>Solanaceae</i>	X	X	X	X	X	X	X	Worldwide	Tsrir (2004); Aqeel et al. (2008)			
<i>Fusarium</i> spp.	Fusarium dry rots	<i>Fusarium sambucinum</i> : Wide : potato, hop, leguminous plants, cereals <i>F. coeruleum</i> : Wide : potato, cereals and many other hosts	X	X	X	X	X	X	X	Worldwide America, Europe, Asia	Peters et al. (2008)			
<i>Helminthosporium solani</i>	Silver scurf	Potato	X	X	(X)						Cunha and Rizzo (2004)			
<i>Macromomina phaeolina</i>	Charcoal rot	Wide: 284 recorded hosts both cultivated and wild												
<i>Phoma andigena</i> var. <i>andina</i>	Phoma leaf spot	Narrow : potato, <i>S. goniocalyx</i> , <i>S. medians</i> , <i>S. phureja</i> , tomato, solanaceous weeds								South America (Bolivia and Peru)				
<i>Phoma</i> spp.	Gangrene	<i>Phoma exigua</i> var. <i>exigua</i> , wide <i>Phoma exigua</i> var. <i>foveata</i> , narrow, potato and some weeds	X	X						North America, Europe, Asia, Oceania	Bain et al. (1987)			
<i>Phytophthora erythroseptica</i>	Pink rot	Narrow : potato, tomato, spinach, and tulip	X	X					X	Worldwide	Peters et al. (2004); Stamps (1978)			
<i>Polyscyathum pusulans</i>	Skin spot	Narrow : Solanaceous species	X	X					X	Europe, North America, Oceania, Asia	Vico et al. (1997)			
<i>Pythium ultimum</i> var. <i>ultimum</i>	Leak	Wide including many crops	X	X					X	Worldwide	Perez et al. (1994)			
<i>Rhizoctonia solani</i>	Black scurf/Stem canker	Narrow : Solanaceous species	X	X					X	Worldwide	Woodhall et al. (2008)			
<i>Rosellinia</i> sp.	Rosellinia black rot	Wide : plants in over 63 genera in 30 families	X	X										
<i>Sclerotinia sclerotinum</i>	White mold	Wide : approximately 400 species of dicots	X	X										
<i>Sclerotium rolfsii</i>	Stem rot	Wide : cultivated and wild plants including ferns	X	X										
<i>Spongospora subterranea</i>	Powdery scab (PMTV vector)	Wide : Solanaceous species, cabbage and related species	X	X			X		X	Worldwide	Nakayama et al. (2003); Hims and Preece (1975); Merz and Falloon (2009)			
<i>Synchytrium endobioticum</i>	Wart	Potato	X	X					X	Worldwide				
<i>Thecaphora solani</i>	Thecaphora smut	Narrow: Solanaceous species, <i>Datura stramonium</i>	X	X					X	South America and Mexico	Mordue (1988); Andrade et al. (2004)			
<i>Verticillium dahliae</i> and <i>Verticillium albo-atrum</i>	Verticillium wilt	<i>Verticillium dahliae</i> , moderate : artichoke, bell pepper, cabbage, cauliflower, chili pepper, cotton, eggplant, lettuce, mint, potato, strawberry, tomato, watermelon, etc. <i>Verticillium albo-atrum</i> , narrow : alfalfa, hops, potato	X	X	X	X	X*	X*	Worldwide	Stevenson et al. (2001), Ochiai et al. (2008)				

Table 1 (continued)

Pathogen	Disease	Host range	Main symptoms				Pathogenicity test	Distribution	Références			
			Tubers		Other parts							
			Gall	Blemish	Rot	Leaf lesions						
<b>Bacteria</b>												
<i>Clavibacter michiganensis</i> ssp. <i>sepedonicus</i>	Ring rot	Narrow : potato, sugar beet, tomato, eggplant		X	X		X	Worldwide	Nissinen (2000)			
<i>Clostridium</i> spp.	Bacterial soft rot	Wide: animals and plants		X				Worldwide				
<i>Pectobacterium atrosepticum</i> , <i>Pectobacterium carotovorum</i> subsp. <i>carotovorum</i> , <i>Dickeyeza</i> spp.	Black leg, soft rot	<i>Pectobacterium</i> spp. and <i>Pectobacterium carotovorum</i> subsp. <i>carotovorum</i> ; wide : potato, rapeseed, sugar beet, chicory, white radish, weeds <i>Pectobacterium atrosepticum</i> ; narrow : potato tomato, cabbage, weeds <i>Dickeyeza</i> spp.; potato, ornamentals, maize, chicory, white radish, turnip, carrots, parsnips, etc.		X	X	X	X	Worldwide	Franco et al. (2007); Bradbury (1977); Helias (2008)			
<i>Ralstonia solanacearum</i>	Brown rot	Wide: plants in over 200 species in 28 families		X	X		X	Asia, Africa, South America (probably worldwide)	Park et al. (2007)			
<i>Spirogyra scabiei</i> , <i>S. acidiscabiei</i> , <i>S. europeiscabiei</i>	Common and netted scab	Moderate : potato, beets, radish, rutabaga, turnip, carrot, parsnips, etc.		X	X		X	Worldwide	Bouchek-Mechiche et al. (2006); Lambert et al. (2006); Zhao et al. (2008)			
<b>Nematodes</b>												
<i>Belenolaimus longicardanus</i>	Sting nematode	Wide : vegetables (carrot, corn, crucifers, beans, potato, etc.), fruits (citrus, strawberry, etc.), agronomic crops (cotton, peanut, sorghum, soybean, etc.), turf grasses and forest crops		X			X	North America				
<i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i>	Potato rot nematode Stem and bulb nematode	Wide : almost all plants, feed also on soil fungi potato, onions, pea, beans, rye.		X	X			Europe, Africa, America	Vreugdenhil (2007)			
<i>Globodera pallida</i> , <i>Globodera rostochiensis</i>	Potato cyst nematode	Narrow : potato, tomato, eggplant, wild solanaceous weeds		X			X	Worldwide	Vreugdenhil (2007); Pylypenko et al. (2008)			
<i>Meloidogyne</i> spp.	Root-knot nematode	Wide : about 2000 species (Solanaceae, Cucurbitaceae, leguminous plants, carrots, scorsoneras, lettuces, chicory, white radish, artichokes, Swiss chards, celery, etc.)		X			X	Worldwide	Vreugdenhil (2007); Vovlas et al. (2005)			
<i>Nacobbus aberrans</i>	False root-knot nematode	Wide : potato, <i>Braisia oleacea</i> , <i>Capsicum</i> , carrots, cucumbers, lettuces, <i>Opuntia</i> spp. and other Cactaceae, sugarbeet, tomato, etc.		X				America	Iniesta et al. (2005); Stevenson et al. (2001); Vreugdenhil (2007)			

Table 1 (continued)

Pathogen	Disease	Host range	Main symptoms				Pathogenicity test	Distribution	Références
			Tubers	Gall	Blemish	Rot	Other parts		
				Stem lesions	Leaf lesions	Root lesions			
<i>Paratrichodorus</i> and <i>Trichodorus</i> spp.	Stubby root nematode (TRV vector)	<i>Pantrichodorus</i> sp; wide : alfalfa, azalea, boysenberry, vegetables, corn, tomato, potato, onion, wheat, sugarcane, rice, grasses, etc. <i>Trichodorus</i> spp.; wide : trees, shrubs, crops, turf grasses	X	X			X	X	Europe, North America
<i>Pratylenchus</i> spp.	Root-lesion nematode	Wide: a lot of fruit trees, some citrus fruits and cereals, ornamental plants, crops (potato and vine)		X			X	X	Worldwide
Virus	<b>Means of transmission</b>								
Tobacco necrosis virus (TNV)	Mechanical, Oidium brassicae	Narrow : potato, tobacco, bean, tulip	X	X					Stevenson et al. (2001)
Tobacco rattle virus (TRV)	Stubby root nematodes	Wide : potato, gladiolus, lettuce, sugar beet, tobacco, tulip, etc.	X	X					Europe, Japan, New Zealand, North America, Russia
Potato mop-top virus (PMTV)	<i>Spongopora subterranea</i>	Narrow : mainly Solanaceous species	X	X	X	X			Andean region, Canada, China, North Europe, Japan

evaluates in which ways the environmental conditions may limit *in situ* the expression of this pathogenic activity, including the saprotrophe development, if required by the inoculum (Alabouvette et al. 2006).

Plant diseases result from the compatible interactions between a susceptible host plant and a pathogen. These direct interactions are important but should not overshadow the key role of environmental factors, which influence these interactions and thereby disease incidence or severity. In contrast to aerial diseases, the soil-borne diseases are induced by pathogens which are embedded in the soil matrix. Thus, the soil interferes in many ways in the relationships between and among microorganisms, pathogens, and host plant. It can even modify the interactions among microorganisms themselves. In some soils, disease incidence or severity commonly remains low in spite of the presence of the pathogen, a susceptible host plant and favorable climatic conditions. They are called disease-suppressive soils (Messiha et al. 2007; Steinberg et al. 2007). Soil suppressiveness to diseases depends on the pathogen itself—its inoculum density and its intrinsic aggressiveness—and also on different soil factors, including both biotic and abiotic components.

In the first part of this paper, the influence of abiotic factors on disease severity will be reviewed. Then the characteristics of the inoculum and its relationships with the rest of the microbiota will be considered. Finally, risk assessment models, decision support systems, and control strategies based on collected data will be discussed.


## 2 Effects of abiotic factors on the occurrence and development of soil-borne potato diseases

Soil abiotic components such as texture, organic matter content, pH, as well as temperature and moisture greatly affect the behavior of the pathogens and determine disease incidence or severity.

### 2.1 Soil temperature

Temperature and moisture of the soil are obviously greatly dependent on the climatic conditions, but also on some cultural practices such as irrigation. Temperature is of major importance in disease development since it determines pathogen growth rate (Baljeet et al. 2005), kind of symptoms (Bouchek-Mechiche et al. 2000), and geographical distribution of the diseases. Most of the potato pathogens can grow at soil temperatures between 10°C and 25°C, the optimal potato growth temperatures (Table 2). However, gangrene, black scurf, and powdery scab are favored by mean temper-

**Fig. 1** Symptoms caused by some potato soil-borne diseases, **a** tobacco rattle virus (TRV, transmitted by nematodes), **b** root-knot nematode (*Meloidogyne incognita*), **c** common or netted scab (*Streptomyces scabies*), **d** black scurf (*Rhizoctonia solani*)



atures below 15°C (Baker 1970; Gindrat 1984; Harrison 1997); on the contrary, black dot, black leg, stem rot, and charcoal rot are favored by temperatures above 27°C. Similarly, sting and root-knot nematodes reproduce better between 25°C and 30–35°C depending on the origin of the populations.

## 2.2 Soil moisture

Soil moisture which depends on the climate and cultural practice is also determined by the soil texture (see below). In the literature dealing with interactions between soil moisture and potato diseases, many different terms are used to characterize the soil water content.

Soil moisture content, moisture-weight percentage, and water-holding capacity are used to evaluate the volume of water contained in soil. It is generally expressed as a percentage of the soil dry weight. Other publications refer to water activity which is a dimensionless quantity (between 0 and 1) describing the amount of free water in soil for biochemical reactions. Water activity, which depends on soil texture, is related to moisture content in a non-linear relationship known as a moisture sorption isotherm curve.

High soil moisture due to abundant rainfall, poor drainage, heavy soils, or irrigation, influences disease development and the opening of the lenticels which are further entry points for soil-borne pathogens into the tuber (Helias 2008). Several diseases, especially bacterial diseases, are enhanced by high moisture content (Table 2),

but few diseases are favored by low levels of moisture. This is the case for black dot, some dry rots induced by *Fusarium* spp., stem rot, wart, common scab, and sting and root-knot nematodes. High soil moisture generally has indirect effects which might favor disease severity. This is the case of flooding that provokes oxygen depletion and CO<sub>2</sub> enrichment resulting in an increase of *Spongospora subterranea* (powdery scab) development (Harrison 1997). In some cases, the influence of soil moisture on disease severity is not clearly demonstrated. Depending on the studies, black scurf, stem canker, silver scurf (*Helminthosporium solani*), and *Thecaphora* smut (*Thecaphora solani*) are either positively or negatively correlated with soil moisture (Adams et al. 1987; Hide and Firmager 1989; Sepulveda et al. 2000; El Bakali and Martin 2006; Wale et al. 2008). Conversely, high relative humidity during storage of tubers has always a negative impact (Table 2).

## 2.3 Soil texture

The soil texture describes the relative percentage of sand, loam, and clay contents. Most of fungal diseases are enhanced in light sandy soils (Table 3). Conversely, it is generally accepted that clay soils favor bacterial activity (Marshall 1975; Alabouvette et al. 1996) explaining that clay or heavy soils are conducive to bacterial soil-borne diseases (ring rot, soft rot, brown rot, and netted scab). Concerning nematodes, no general rule can be drawn up as some species are more prevalent in heavy soils (root-knot

**Table 2** Favorable climatic conditions for potato soil-borne diseases development

Pathogen	Disease	Optimal temperatures (°C)	Optimal level of humidity		Optimal light duration	References
			low	high		
<b>Fungi and oomycetes</b>						
<i>Colletotrichum coccodes</i>	Black dot	25–30; optimum: 27	X (whc <50%)	X (storage)	Colonization, sclerotia	Davet (1970); Lees (2003); Tsror (2004)
<i>Fusarium</i> spp.	Fusarium dry rot	15–20	X (9.2% whc)	X (27.9% whc)	Mycelial growth	Tivoli (1983); Kong et al. (2006)
<i>Helminthosporium solani</i>	Silver scurf	15–32	X	X (sporulation)		Adams et al. (1987); Errampalli (2001)
<i>Macrophomina phaseolina</i>	Charcoal rot	>30	X (RH>52%)	Mycelial growth	Pycnidia production, mycelial growth	Gindrat (1984); Vishwana and Sarbhoy (1989); Muthukrishnan et al. (1995); Sonani and Chaudhan (1996); Andadioba and Adisa (1999); Mehta et al. (2006); Chowdary and Govindalak (2007)
<i>Phoma andigena</i> var. <i>andina</i>	Phoma leaf spot	5–18; optimum: 10	X	Pycnidial and conidial productions	Pycnidial production	Fox et al. (1978); Gindrat (1984); Bang (1989); Coelho et al. (1997); Lo et al. (2000)
<i>Phoma</i> spp.	Gangrene	5–18; optimum: 10	X			Salas et al. (2000)
<i>Phytophthora erythroseptica</i>	Pink rot	15–30	X (waterlogged soil)			
<i>Polycephalum pustulans</i>	Skin spot	5–20	X (storage)			Hide and Cayley (1987); Vico et al. (1997)
<i>Pythium ultimum</i> var. <i>ultimum</i>	Leak	20–30	X (RH 95% in storage)	X	Sclerotia formation	Lui (2003)
<i>Rhizoctonia solani</i>	Black scurf/Stem canker	10–18	X (45% whc)			Baker (1970); Hide and Firmager (1989); Xu et al. (1997); El Bakali and Martin (2006); Panka et al. (2007)
<i>Rosellinia</i> spp.	Rosellinia black rot					
<i>Sclerotinia sclerotiorum</i>	White mold	15–27	No effect of RH			Young et al. (2004); Harikrishnan and del Rio (2006)
<i>Sclerotium rolfsii</i>	Stem rot	25–35; optimum: 30	X (30% whc)	Sclerotia production	Mycelial growth, sclerotia production	Chowdhury et al. (1993); Prithviraj et al. (2000); Blum et al. (2002); Gupta et al. (2007)
<i>Spongospora subterranea</i>	Powdery scab	Tuber galls: 12–15 Root gall: 17	X	Constant dampness		Harrison (1997) Graaf et al. (2005); Metz and Falloon (2009)
<i>Synchytrium endobioticum</i>	Wart	12–18	X	annual rainfall greater than 700 mm		Hampson and Coombes (1997); Stachewicz and Enzian (1998)
<i>Thecaphora solani</i>	Thecaphora smut	5–20	X			EPPO (1990); Sepulveda et al. (2000); Vale et al. (2008)
<i>Verticillium dahliae</i> and <i>Verticillium albo-atrum</i>	Verticillium wilt	22–26; optimum: 25	X ( $a_w=0.995$ )	Sporulation	Jong-Tae et al. (2001); Santamarina and Rosello (2006)	

**Table 2** (continued)

Pathogen	Disease	Optimal temperatures (°C)	Optimal level of humidity		Optimal light duration	References
			low	high		
<b>Bacteria</b>						
<i>Clavibacter michiganensis</i> ssp. <i>sepedonicus</i>	Ring rot	10–20	X	X		Wolf and Beckhoven (2004)
<i>Clostridium</i> spp.	Bacterial soft rot		X			Suyama et al. (1990)
<i>Pectobacterium atrosepticum</i> , <i>Pectobacterium carotovorum</i> subsp. <i>carotovorum</i>	Black leg, soft rot	15–25	X			Jeggi et al. (1991); Serfontein et al. (1991); Vries and Vuurde (1993); Latour et al. (2008); Helias (2008)
<i>Dickeya</i> spp., <i>carotovorans</i> , <i>Dickeya</i> sp.						
<i>Ralstonia solanacearum</i>	Brown rot	23 (temperate strains) 30–35 (tropical strains)	X (whc 60%)			Shekhwat and Perombelon (1991); Sunaina et al. (2000); Tomlinson et al. (2005)
<i>Streptomyces scabiei</i> , <i>S. acidiscabies</i> , <i>S. europeiscabies</i>	Common and netted scab Netted scab: 13–17	Common scab: 19–24	X	X		Adams et al. (1987); Bouchech-Mechiche et al. (2000); Pasco et al. (2005); Panka et al. (2007)
<b>Nematodes</b>						
<i>Belonolaimus longicaudatus</i>	Sting nematode	25–35	X (RH 7%)			Robbins and Barker (1974)
<i>Diaphenichthus destructor</i>	Potato rot nematode	20–37; optimum: 21	X (RH 41–66%)			Mugnier and Phillips (2007); Shojaei et al. (2006)
<i>Globodera pallida</i> , <i>Globodera rostochiensis</i>	Potato cyst nematode	10–28	No effect of soil humidity			Inserro et al. (1996); Muhammad (1996)
<i>Meloidogyne</i> spp.	Root-knot nematode					Stevenson et al. (2001); Chandell et al. (2002); Pandey et al. (2002); Wu et al. (2006)
<i>Nacobbus aberrans</i> (Paradrichodorus spp.)	False root-knot nematode	25–32	X (30% whc)			Anthoine et al. (2006)
<i>Pratylenchus</i> spp.	Stubbyroot nematode	10–25; optimum: 20				
						Jauhari and Lal (2001); Pudasaini et al. (2007)

RH relative humidity, whc water holding capacity

**Table 3** Favorable edaphic conditions for the development of potato diseases

Pathogen	Disease	Optimal soil texture	Optimal soil pH	Optimal soil nutrient content	Optimal organic matter content	References
<b>Fungi and Oomycetes</b>						
<i>Colletotrichum coccodes</i>	Black dot	X	6–7	Low nitrogen level	Kang et al. (2003); Nitzan and Tsrir (2003); Tsrir (2004)	
<i>Fusarium</i> spp.	Fusarium dry rot	X	<i>F. solani</i> >5.3 <i>F. roseum</i> , no effect	High Fe level Low Ca, borax and P levels	Variable	
<i>Helminthosporium solani</i>	Silver scurf	X	X	6.5	Lemard (1980); Lutomirska and Szutkowska (2004)	
<i>Macrophomina phascolina</i>	Charcoal rot	X	3.8–5.6	2.9–7.6‰	Singh and Kaiser (1994)	
<i>Phoma andigena</i> var. <i>andigena</i>	Phoma leaf spot	X			Tivoli et al. (1987)	
<i>Phoma</i> spp.	Gangrene	X				
<i>Phytophthora erythroseptica</i>	Pink rot					
<i>Polyscytalum pusulans</i>	Skin spot					
<i>Pythium ultimum</i> var. <i>ultimum</i>	Leak		No effect	No effect	Vivoda et al. (1991)	
<i>Rosellinia</i> spp.	Rosellinia black rot					
<i>Rhizoctonia solani</i>	Black scurf/Stem canker	X	High?	High	El Fahl and Calvert (1976); Rudkiewicz et al. (1983); Lutomirska and Szutkowska (2005)	
<i>Sclerotinia sclerotiorum</i>	White mold					
<i>Sclerotium rofsii</i>	Stem rot	X	~ 6.5	High nitrogen, organic carbon and low phosphorus and potassium levels	Sheoraj et al. (2007); Banyal et al. (2008)	
<i>Spongospora subterranea</i>	Powdery seab	X organic or over irrigated soils	X poorly drained soils	4.7–7.6	High aluminum level	
<i>Synchytrium endobioticum</i>	Wart	X		Variable	Zambolim et al. (1995); Graaf et al. (2005); Gilchrist et al. (2009); Merz and Falloon (2009)	
<i>Thecaphora solani</i>	Thecaphora smut				Hampson (1985); Hampson and Coombes (1997)	
<i>Verticillium dahliae</i> and <i>Verticillium albo-atrum</i>	Verticillium wilt	X	6–9	High salt level High Ca, low K, Mg and total soil C level	EPPO (1990) Baard and Pauer (1981); Höper and Alabouvette (1996); Davis et al. (2001)	
<b>Bacteria</b>						
<i>Clavibacter michiganensis</i> ssp. <i>sepedonicus</i>	Ring rot	X			Moffett and Wood (1984)	
<i>Clostridium</i> spp.	Bacterial soft rot					

**Table 3** (continued)

Pathogen	Disease	Optimal soil texture	Optimal soil pH	Optimal soil nutrient content	Optimal organic matter content	References
<i>Pectobacterium atrosepticum</i> , <i>Pectobacterium carotovorum</i> subsp. <i>carotovorum</i> , <i>Dickeya</i> spp. <i>Ralstonia solanacearum</i>	Black leg, soft rot	X Black leg Mainly sandy or light soils	X Soft rot Mainly clay or heavy soils		Low Ca concentration	Zielke et al. (1974); Lucke (1975); Lambert and Manzer (1991)
	Brown rot	X	X	variable	No ammonium intake	Hsu (1991); Shekhawat and Perombelon (1991); Messha et al. (2007); Michel and Mew (1998); Yi and Sul (1998); Keshwani et al. (2000); Muller et al. (2004)
<i>Streptomyces scabiei</i> , <i>S. acidiscabiei</i> , <i>S. europeiscabiei</i>	Common and netted scab	X	5.2–7	Low Mn level		Rudkiewicz et al. (1983); A labouvette et al. (1996); Loria et al. (1997); Milosevic et al. (2005); Lazarovits et al. 2007
<b>Nematodes</b>						Mashela et al. (1991)
<i>Belenolaimus longicaudatus</i>	Sting nematode	X				Pelsmacker and Coomans (1987); Ruijter and Haverkort (1999); Trifunova (2001)
<i>Ditylenchus destructor</i>	Potato rot nematode	X	6.1	Low nitrogen level	High	Kumar and Vadivelu (1996); Kandji et al. (2001); Pandey et al. (2002); Melakeberhan et al. (2004)
<i>Globodera pallida</i> , <i>Globodera rostochiensis</i>	Potato cyst nematode	X				
<i>Meloidogyne</i> spp.	Rootknot nematode	X	X	7.5	High	
<i>Nacobbus aberrans</i>	False root-knot nematode					
	Stubby-root nematode (TRV vector)	For both <i>P. pachydermus</i> and <i>T. similis</i>	Only in the case of <i>T. primitivus</i>	low	High level of Fe	Barbez (1983); Spaul and Cadet (2001)
	Root-lesion nematode	X		variable	Low level of Fe	Pelsmacker and Coomans (1987); Spaul and Cadet (2001)

nematodes) and other species in light soils (sting nematodes). Soil texture also influences soil structure, through the distribution of different pore sizes, determining the actual living space for bacteria, fungi, and predators. It also influences the water activity; water retained in pores of narrow diameter being less available for organisms that water present in large pores.

## 2.4 Soil pH

Disease development is also influenced by soil pH linked to soil nutrient availability (Table 3). Soils with extreme pH values are often highly suppressive to several plant diseases (Höper and Alabouvette 1996). However, pH fluctuations resulting from amendments influence pathogens and disease development. Decreasing pH increases the availability of phosphorus, nitrogen, and aluminum ions and decreases potato cyst nematode, brown rot, and common scab damages, respectively (Mulder et al. 1997; Michel and Mew 1998; Ruijter and Haverkort 1999; Mizuno et al. 2003). On the contrary, addition of urea in soil induces a very large increase in pH and a good control of *Synchitrium endobioticum*, the fungal pathogen causing wart (Hampson 1985).

## 2.5 Soil organic matter

Soil organic matter is both the substrate for and the result of microbial activity. In addition, together with clay, organic matter affects soil structure and thus moisture content and aeration. The quantity of organic matter in a soil has an effect on the appearance and the development of diseases but its quality is also an important point which has been too poorly addressed (Alabouvette et al. 1996).

Most physico-chemical factors are not independent from one to the others, which makes experiments and data interpretation very difficult. Soil texture can affect humidity, soil amendments impact on pH, and all those factors influence availability of chemical elements. Thus, the pathogenic inoculum present either in the soil or on the tuber surface has to find the optimal climatic and edaphic conditions to develop.

# 3 Effects of biotic factors on the occurrence and development of soil-borne potato diseases

## 3.1 Autecology of pathogens

### 3.1.1 Inoculum sources, survival and dissemination pathways

The survival of soil-borne pathogens during periods without potato crop depends on their ability to resist to unfavorable

conditions. Most of them survive in soil under the form of resistant structures able to directly infect the new host crop. Some pathogens can also survive as saprophytes on host crop residues or on alternative hosts during winter. Finally, inoculum can also be introduced into the field by the seeds; it is called seed- or tuber-borne inoculum. Inoculum sources are diverse and for many diseases several inoculum sources can play a role (Table 4). Soil-borne fungi produce different conservation structures. *Fusarium* spp. forms chlamydospores resistant to adverse conditions, *Rhizoctonia solani*, *Verticillium* spp., *Sclerotinia sclerotinum* overwinter as sclerotia. Bacteria can survive over winter with favorable moisture, temperature, and soil type (Ficke et al. 1973; Bradbury 1977; Loria et al. 2008). Nematodes can survive and persist in soil as protective cysts surrounding the eggs (*Globodera* spp.) or as juveniles in host roots (*Meloidogyne* spp.; Qian et al. 1996; Wharton and Worland 2001).

In the absence of resistant structures and of efficient saprophytic abilities, some pathogens need alternative hosts to survive in absence of potatoes. These alternative hosts frequently belong to the *Solanaceous* family and act as a long-term reservoir of the pathogen (Chang et al. 1992; Tomlinson et al. 2005).

Fungal dissemination occurs frequently as spores (conidiospores, chlamydospores, pycnidiospores, sporangiospores, oospores, and zoospores) or mycelium transported by water (rain, irrigation, and flow in soil), by soil adhering to farm equipment or introduced by contaminated seed tubers (Zambolim et al. 1995; Stevenson et al. 2001; Bae et al. 2007). Moreover, some pathogens liberate mobile dissemination forms such as zoosporangia. Zoospores of *Phytophthora erythroseptica*, *S. subterranean*, and *S. endobioticum* are responsible for short-distance dissemination of these pathogens (Wharton et al. 2007; Merz and Falloon 2009). Adult nematodes such as *Pratylenchus penetrans* are able to migrate on quite long distances better than do larvae (Pudasaini et al. 2007).

### 3.1.2 Relationship between inoculum density and disease severity

Although there is not always a clear and linear relationship, the severity of the disease generally increases with an increasing level of inoculum (Table 4). Sometimes, a minimum inoculum threshold is needed to initiate the disease development. This is the case, for instance, for potato cyst nematodes (Samaliev et al. 1998). Conversely, the disease severity of black dot does not increase any more beyond a maximum threshold of inoculum density (Nitzan et al. 2008). In fact as stated above, the relationship between inoculum density and disease severity greatly depends on the environmental factors which determine the level of soil suppressiveness.

**Table 4** Inoculum sources and correlation between inoculum density and soil-borne potato diseases severity

Pathogen	Disease	Inoculum source	Correlation between inoculum density and disease severity (minimum value used for the calculation)	References
<b>Fungi and oomycetes</b>				
<i>Colletotrichum coccodes</i>	Black dot	Soil >Seed tuber	Disease severity remains constant above a threshold of soil-borne inoculum (0.5–1.7 g inoculum per liter of soil)	Lees (2003); Nitzan et al. (2008)
<i>Fusarium</i> spp.	Fusarium dry rots	Soil, seed tuber	Positive correlation ( $10^4$ conidia·ml <sup>-1</sup> soil for <i>F. sulphureum</i> ; $10^5$ conidia·l <sup>-1</sup> soil for <i>F. coeruleum</i> )	Tivoli et al. (1987); Stevenson et al. (2001); (2005)
	Silver scurf	Seed tuber, soil	Negative correlation	Lennard (1980); Bains et al. (1996); Geary and Johnson (2006)
<i>Macrophomina phaseolina</i>	Charcoal rot			
<i>Phoma antigena</i> var. <i>andina</i>	Phoma leaf spot	Seed tubers>plant residues	Adams (1980); Tivoli et al. (1987); Carnegie (1991)	
<i>Phoma</i> spp.	Gangrene	Seed tuber	Salas et al. (2000)	
<i>Phytophthora erythroseptica</i>	Pink rot	Seed tubers; crop debris, dust in store and soil	Wale et al. (2008)	
<i>Polyscytalum pastulans</i>	Skin spot	Soil	Triki et al. (2001)	
<i>Pythium ultimum</i> var. <i>ultimum</i>	Leak	Sclerotia on seed tubers, in soil and in plant residues	Rahman et al. (1996); Tsror and Peretz-Alon (2005)	
<i>Rhizoctonia solani</i>	Black scurf/Stem canker			
<i>Rosellinia</i> sp.	Rosellinia black rot	Soil, seed tuber	Positive correlation	US Canola Association
	White mold	Soil, seed tuber	No significant/positive correlation (100 sporosori g <sup>-1</sup> soil)	Rahman et al. (1996)
	Stem rot	Soil, seed tuber, manure	Positive correlation (1/25 sporangium·g <sup>-1</sup> soil)	Zambolim et al. (1995); Graaf et al. (2005); Nakayama (2007); Merz and Falloon (2009)
	Powdery scab	Soil, seed tubers	Positive correlation	Hampson et al. (1994); Bayen et al. (2005)
<i>Sclerotinia sclerotiorum</i>	Wart	Seed tuber, soil, infested plant parts	Positive correlation	Mordue (1988); Wale et al. (2008)
<i>Sclerotium rolfsii</i>	Thecaphora smut	Soil microsclerotia, infected plant residues	Positive correlation	Nicot and Rouse (1987); Mol and Scholté (1995); Vallad et al. (2004)
<i>Spongopaspora subterranea</i>	Verticillium wilt			
<i>Synchytrium endobioticum</i>				
<i>Thecaphora solani</i>				
<i>Verticillium dahliae</i> and <i>V. albo-atrum</i>				
<b>Bacteria</b>				
<i>Clavibacter michiganensis</i> spp. <i>sepedonicus</i>	Ring rot	Seed tuber, soil, equipment	No significant correlation	Nelson (1982); Westra et al. (1994)
<i>Clostridium</i> spp.	Bacterial soft rot	Mainly seed tubers but also soil, water, insects	Positive correlation (10 <sup>3</sup> cell per tuber)	Naumann et al. (1974); Perombelon (2000); Helias (2008)
	Black leg, soft rot			
<i>Pectobacterium atrosepticum</i> , <i>Pectobacterium carotovorum</i> subsp. <i>carotovorum</i> , <i>Dickeya</i> spp.	Brown rot	Seed tuber, soil, water	Positive correlation	Hsu (1991)
<i>Ralstonia solanacearum</i>	Common and netted scab	Seed tuber and soil-borne	Positive correlation	Wilson et al. (1999); Wang and Lazarovits (2005)
<b>Nematodes</b>				
<i>Belenolaimus longicaudatus</i>	Sting nematode	Seed tuber or soil (cysts in) soil or soil-carrying seeds/seedlings/equipment (Eggs or larvae in) soil or soil-carrying seeds/seedling/equipment	Positive correlation (2 eggs·g <sup>-1</sup> soil)	Samaliev et al. (1998); Anaya et al. (2005)
<i>Ditylenchus destructor</i>	Potato rot nematode		Positive correlation (0.5 eggs cm <sup>-3</sup> soil)	Mohsin et al. (1989); Nagesh (1996); Vovlas et al. (2005)
<i>Globodera pallida</i> , <i>Globodera rostochiensis</i>	Potato cyst nematode			
<i>Meloidogyne</i> spp.	Root-knot nematode			
<i>Nacobius aberrans</i>	False root-knot nematode	Seed tuber	Positive correlation	Franco et al. (1992)
<i>Paratrichodorus</i> and <i>Trichodorus</i> spp.	Stubby root nematode (TRV vector)	Soil or soil-carrying vector	Positive correlation	Perez et al. (2000)
<i>Pratylenchus</i> spp.	Root-lesion nematode	Soil	Positive correlation (0.4 eggs g <sup>-1</sup> soil)	Holgado et al. (2009)

### 3.1.3 Mechanisms of infection

Potato plants are essentially composed of cellulose, a very solid polymer and tubers are enveloped in a protective covering called periderm made of a suberin biopolymer providing the primary barrier against diseases, insects, dehydratation, and physical intrusions (Lulai 2001). Soil-borne pathogens of potato have various ways to penetrate the host plant and break physical barriers. They enter the roots, young sprouts, underground stems, stolons, or tubers. Some pathogens cannot infect intact tuber periderm or lenticels and penetrate through wounds (Stevenson et al. 2001; Taylor et al. 2004) whereas other pathogens can penetrate either directly by mechanical and/or enzymatic degradation of the host's cells or through natural openings (stomata, lenticels, eyes) (Table 5).

Once they have penetrated the host, pathogens colonize plant tissues. Fungi grow through the parenchyma of the cortex and often reach the vascular vessels. *T. solani*, *S. endobioticum*, and *Streptomyces* spp. penetration provokes hypertrophy of the colonized tissues resulting in galls. They grow in the plant, induce cell death, and feed on them saprophytically. They secrete phytotoxins—for example thaxtomin produced by *Streptomyces* spp.—inducing the formation of several layers of suberized corky cells, creating a large lesion firmly integrated within the tuber skin (Stevenson et al. 2001; Mulder et al. 2008; Perez and Torres 2008). Compared to common scab development, powdery scab pustules formation is a relatively short process, at the end of which a single wound cork layer remains that covers the entire lesion. After hardening off, this layer can be easily removed from the lesion without any damage of the underlying tissues (Delleman et al. 2005). *Colletotrichum coccodes*, *H. solani*, *Polyscytalum pustulans*, *R. solani*, *S. subterranea*, and *Streptomyces* spp. are responsible for several superficial alterations called blemishes. Colonization by those pathogens is usually limited to superficial layers of tuber periderm (Harrison 1997; Stevenson et al. 2001; Cunha and Rizzo 2004; Lehtonen et al. 2008; Loria et al. 2008) but they can colonize other parts of the plant until they reach vascular system. *Streptomyces* spp. responsible for netted scab blemishes has pathogenic mechanisms that are assumed to not implicate thaxtomin but rather a necrotic protein (Bouchech-Mechiche et al. 2006).

Fungi and bacteria-causing rots produce a wide range of hydrolytic enzymes such as cellulases, pectinases, xylanases, and proteases (Olivieri et al. 2004). They are responsible for tissue maceration and cell death, after which the microorganisms have access to the nutritional resources of the dead plant tissues (Amadioha 1997; Aveskamp et al. 2008). *Pectobacterium* spp. develop an original pathogenic strategy based on quorum sensing, which utilizes freely

diffusible chemical signal molecules allowing pathogenic bacteria to synchronize the production of virulence factors and make the pathogenic attack more efficient (Liu et al. 2008). Finally, nematodes attacking potatoes can be classified into two categories: ectoparasites and endoparasites. Ectoparasites nematodes (*Belonolaimus longicaudatus*, *Paratrichodorus* spp., and *Trichodorus* spp.) are mobile and feed on potato roots in the area of cell division and elongation without penetrating the root (Stevenson et al. 2001; Mugniéry 2007). The endoparasitic nematodes of potato, *D. destructor* and *P. penetrans* are migrating endoparasites; they feed from cell to cell within the host, whereas *Globodera* spp., *Meloidogyne* spp., and *N. aberrans* are sedentary endoparasites, they induce specialized feeding sites in plant roots. *D. destructor* and *P. penetrans* penetrate underground parts of the plant, feed on the cortical cells, and migrate into the roots, destroying cell after cell. *Globodera pallida*, *Globodera rostochiensis*, *Meloidogyne* spp., and *N. aberrans* develop feeding cavities in host root, causing galls (Mugniéry 2007).

### 3.1.4 Genetic variability

A soil-borne disease can be caused by several species of pathogens belonging to a single genus, by one species, or even by a subgroup of a species. Each species or subspecies is adapted to particular conditions or variety. Knowledge of the genetic diversity of pathogens is useful for precise diagnosis and control of potato diseases.

Since *Erwinia* has been renamed and divided into two different genera, *Pectobacterium* and *Dickeya* (Helias 2008), bacterial soft rot previously attributed to *Erwinia carotovora*, *Erwinia atroseptica*, and *Erwinia chrysanthemi* is in fact one disease caused by several species belonging to different genera (Table 5). *Pectobacterium* spp. and *Dickeya* spp. are frequently associated with bacteria of the genus *Clostridium* which includes very numerous Gram-positive anaerobic bacteria. *Clostridium puniceum* is one of the few well-characterized pectolytic clostridia isolated from rotting potato tubers (Stevenson et al. 2001; Prescott et al. 2003).

Within a same species, the pathogen may belong to different groups with various genetic, pathogenic, and physiological traits leading to the characterization of races, biovars and, recently, genomovars—strains which are phylogenetically differentiable, but are phenotypically indistinguishable—phylotypes and sequevars—one or several strains with a given sequence (Nouri et al. 2009). Fungi without sexual reproductive stage related to the potato disease cycle, such as *Colletotrichum* spp., *Fusarium* spp., or *Verticillium* spp., are classified in vegetative compatibility groups (VCGs). Within a VCG, hyphae belonging to different isolates can anastomose and form stable heterokaryons, whereas hyphae from isolates belonging to

**Table 5** Genetic variability, strategies of conservation and attack of the pathogens and detection methods

Pathogen	Disease	Genetic variability	Conservation and overwintering ways	Main penetration ways	Detection methods	References
<b>Fungi and oomycetes</b>						
<i>Colletotrichum coccodes</i>	Black dot	6 or 7 VCG pathogenic for potato	At least 8 years at 10 cm depth in the soil as sclerotia	Mechanical	Q and RT-PCR, Fourier transform infrared (FT-IR)	Dillard and Cobb (1998); Cullen (2002); Heilmann et al. (2006); Ernkhimovich (2007); Sheolnick et al. (2007); Nitzan et al. (2008)
<i>Fusarium</i> spp.	Fusarium dry rots	13 species, especially <i>F. solani</i> var. <i>sambucinum</i> and <i>F. solani</i> var. <i>coeruleum</i> (15 VCGs)	Microconidia, chlamydospores and mycelium on plant debris	Wounds, enzymatic	Isolation and morphology, RT-PCR, PCR enzyme-linked immunosorbent assay, volatile profile	Tivoli et al. (1987); Stevenson et al. (1990); Stevenson et al. (2001); Oliveri et al. (2004); Cullen et al. (2005); Barlakoti et al. (2007); El-Hassan et al. (2007); Peters et al. (2008a); Sharifi et al. (2008); Recep et al. (2009)
<i>Helminthosporium solani</i>	Silver scurf		At least 4 years in the soil	Enzymatic	Classical detection methods, PCR	Bains et al. (1996); Errampalli (2001); Martinez et al. (2004); Geary et al. (2007)
<i>Macrophomina phaseolina</i>	Charcoal rot		Until 3 years under unfavorable climatic conditions as microsclerotia	Enzymatic		Dhingra and Sinclair (1977); Amadioha (1997)
<i>Phoma andigena</i> var. <i>andina</i>	Phoma leaf spot			Enzymatic	Conventional and RT-PCR	McDonald et al. (2000); Stevenson et al. (2001); Giebel and Dopierala (2004); Cullen et al. (2007)
<i>Phoma</i> spp.	Gangrene	2 sub-species: <i>P. exigua</i> var. <i>jovea</i> and <i>P. exigua</i> var. <i>exigua</i>		Enzymatic		
<i>Phytophthora erythroseptica</i>	Pink rot	One species with few genetic variations	Oospores	Enzymatic	RT-PCR	Lucas and Pitt (1974); Peters et al. (2004); Peters et al. (2005); Cullen et al. (2007); Taylor et al. (2008)
<i>Polycephalum pustulans</i>	Skin spot		7 years or more in soil as sclerotia	Mechanical	Conventional and RT-PCR, Conventional and RT-PCR	Lees et al. (2009)
<i>Pythium ultimum</i> var. <i>ultimum</i>	Leak		Many years in the soil and in the infected plant debris as oospores	Wounds		Cullen et al. (2007); Taylor et al. (2008)
<i>Rhizoctonia solani</i>	Black scurf/Stem canker	One species with 13 anastomosis groups pathogenic for potatoes (AG3 being predominant)	Sclerotia	Enzymatic	Classical bioassays, PCR, immunochromatographic lateral flow	Tsrir et al. (1993); Gilligan et al. (1996); Carling et al. (2002); Lees et al. (2002); Gvozdeva et al. (2006); Hughes (2008)
<i>Rosellinia</i> spp.	Rosellinia black rot	3 species: <i>R. hunadoes</i> , <i>R. necatrix</i> and <i>R. pepo</i>		Enzymatic	Conventional and Scorpion-PCR	Stevenson et al. (2001); Schena et al. (2002); Ten Hoopen and Krauss (2006)
<i>Sclerotinia sclerotiorum</i>	White mold		Sclerotia	Mechanical		Wharton, Michigan potato diseases Madalageri et al. (1991); Ohzurike and Arinze (1992)
<i>Sclerotium rofsii</i>	Stem rot			Enzymatic		Zambolim et al. (1995); Stevenson et al. (2001); Graaf et al. (2003); Ward (2004); Merz (2005); Qu et al. (2006); Nakayama (2007)
<i>Spongospora subterranea</i>	Powdery scab	For >10 years in cold areas as cistosori	Mechanical	Classical methods, conventional and RT-PCR, ELISA,		

Table 5 (continued)

Pathogen	Disease	Genetic variability	Conservation and overwintering ways	Main penetration ways	Detection methods	References
<i>Synchytrium endobioticum</i>	Wart	One species with 43 pathotypes	>30 years as winter sporangia 7 years or more in the soil	Mechanical	Conventional and RT-PCR	Boogert et al. (2005); Baeyen et al. (2006)
<i>Thecaphora solani</i>	Thecaphora smut			PCR		Andrade et al. (2004); Perez and Torres (2008)
<i>Verticillium dahliae</i> and <i>Verticillium albo-atrum</i>	Verticillium wilt	2 species: <i>V. dahliae</i> (4 VCGs) and <i>V. albo-atrum</i> (VCGs2 attacking potato)	≈63 months		Classical methods, PCR, Q-PCR	Nelson (1984); Correll et al. (1988); Joaquin and Rowe (1991); Platt and Mahuku (2000); Tsror et al. (2000); Strausbaugh et al. (1992); Zhang et al. (2005); Atallah et al. (2007)
<b>Bacteria</b>						
<i>Clavibacter michiganensis</i> ssp. <i>sepedonicus</i>	Ring rot	One species with few genetic variation	≈18 months in plain soil	Enzymatic	Immuno-fluorescence antibody, staining (IFAS), (ELISA), RT-PCR, LMW RNA profiles	Nelson (1984); Logan et al. (1987); Eichenlaub et al. (1991); Palomo et al. (2000); Smith et al. (2001); Stevenson et al. (2001); Vasinauskienė and Baranauskaitė (2003); Hukkanen et al. (2005); Gudmestad et al. (2009)
<i>Clostridium</i> spp.	Bacterial soft rot	Several species among which <i>C. puniceum</i>		Enzymatic	Conventional and RT-PCR, isolation (CVP), volatile profile, biochemical tests, ITS-RFLP profiles, 16 S rRNA analysis, ELISA	Perombelon et al. (1979); Stevenson et al. (2001); Prescott et al. (2003)
<i>Pectobacterium</i> spp., <i>Dickeyea</i> spp.	Black leg, soft rot	2 genera: <i>Pectobacterium</i> spp. among which <i>P. atrosepticum</i> and <i>P. carotovorum</i> subsp. <i>carotovorum</i> and <i>P. Dickeyea</i> spp.	Overwintering possible (on crop debris or weeds) but varying between bacteria, seasons and areas	Enzymatic	Conventional, PCR, volatile profile, biochemical tests, ITS-RFLP profiles, 16 S rRNA analysis, ELISA	Bradbury (1977); Ouellette et al. (1990); Tsror et al. (1993); Helias et al. (2000); Lazy and Lukczyc (2003); Atallah and Stevenson (2006); Latour et al. (2008); Pitman et al. (2008); Helias (2008)
<i>Ralstonia solanacearum</i>	Brown rot	One species with several biovars (1, 2, and 21) and races (1 and 3) attacking potato	Water, weeds, (soil?)	Enzymatic	Isolation, PCR, immunofluorescence and fluorescent in-situ hybridisation (FISH)	Hsu (1991); Ronda et al. (1999); Raigawani and Mahadevan (2004); Messina et al. (2007); Loria et al. (2008); Nouri et al. (2009); Smith and de Boer (2009)
<i>Streptomyces</i> spp.	Common and netted scab	Common scab: <i>S. scabies</i> , <i>S. europaeiscabiei</i> , <i>S. stelliscabiei</i> , <i>S. acidiscabiei</i> , <i>S. turquiscabiei</i> and maybe some others Netted scab: <i>S. reticuliscabiei</i> and some isolates of <i>S. europaeiscabiei</i>	Conidia	Enzymatic	Conventional and RT-PCR, RFLP, rRNA sequence analysis, eathorn source utilization, repetitive BOX profiles	Rudkiewicz and Sikorski (1984); Bouchek-Mlechica et al. (2000); Flores-Gonzalez et al. (2008); Loria et al. (2008); Mulder et al. (2008); Zhao et al. (2008)
<b>Nematodes</b>						
<i>Belonolaimus longidoratus</i>	Sting nematode			Mechanical	Centrifugal-flotation method, morphological detection	Crow et al. (2000)
<i>Ditylenchus destructor</i>	Potato rot nematode	About 4 months in favorable conditions	Mechanical	Extraction in water, morphological identification, PCR-RFLP	Shojaei et al. (2006); EPPO (2008); Il'yashenka and Ivaniuk (2008)	
<i>Globodera pallida</i> , <i>Globodera rostochiensis</i>	Potato cyst nematode	2 species: <i>G. pallida</i> and <i>G. rostochiensis</i> as cysts	Until 8 years in the soil	Mechanical and enzymatic	Soil extraction and, morphological identification, allele-specific PCR	Wharton and Worland (2001); Moxnes and Hausken (2007); Achenbach et al. (2009); Reid (2009); Rehman et al. (2009)

**Table 5** (continued)

Pathogen	Disease	Genetic variability	Conservation and overwintering ways	Main penetration ways	Detection methods	References
<i>Meloidogyne</i> spp.	Root-knot nematode	At least 7 species: <i>M. hapla</i> , <i>M. chitwoodi</i> , <i>M. fallax</i> (Mediterranean and temperate areas), <i>M. arenaria</i> , <i>M. incognita</i> , <i>M. javanica</i> and <i>M. mayagrensis</i> (Mediterranean and tropical areas)	Mechanical and enzymatic	Morphometrics, host range, biochemical and molecular (RFLP) analysis		Hlaona and Raouani (2007); Melakeberhan et al. (2007); Mugniéry (2007); Dieterich and Sommer (2009); Ozarslanidan et al. (2009)
<i>Nacobbus aberrans</i>	False root-knot nematode		Mechanical	PCR		Franco et al. (1992); Atkins et al. (2005)
<i>Paratrichodorus</i> and <i>Trichodorus</i> spp.	Stubby-root nematode (TRV vector)	7 species of <i>Paratrichodorus</i> spp. and 5 species of <i>Trichodorus</i> spp.		Morphometric and molecular analysis		Riga and Neilson (2005); Riga et al. (2007)
<i>Pratylenchus</i> spp.	Root-lesion nematode	11 species of <i>Pratylenchus</i> spp.	Enzymatic	Morphometric and molecular (PCR-RFLP) analysis		Brown et al. (1980); Saeed et al. (1998); Stevenson et al. (2001); Mugniéry and Phillips (2007)

VCG vegetative compatibility group, AG anastomosis group, RT-PCR reverse transcriptase polymerase chain reaction, Q-PCR quantitative PCR, FT/IR Fourier-transformed infrared spectroscopy, ELISA enzyme-linked immunosorbent assay

different VCGs cannot. This mechanism is the only known mechanism of genetic exchange between individuals of asexual fungi (Hiemstra and Rataj-Guranowska 2003). Hyphal anastomosis is also used to categorize the isolates of *R. solani* into anastomosis groups (AG). Presently, 13 AGs have been described, several of which being divided into subgroups. Individual AGs are not strictly associated with a specific host but rather with a family of hosts which can be in turn narrow or very broad, for example AG 1 with rice mainly and AG 8 with various cereals. AG 3 isolates, and more specifically isolates from the AG 3 PT subgroup, are often associated with potato diseases (Fiers et al. *in press*; Kuninaga et al. 2000; Carling et al. 2002). However it was shown, in Great Britain and France, that AG 2–1 and AG 5 can cause disease in potato crops but with a much lower incidence than AG 3 PT (Campion et al. 2003; Woodhall et al. 2007).

As a result of the genetic evolution of pathogens, new pathotypes are regularly discovered. Conversely, some populations such as *P. erythrocephala* and *Clavibacter michiganensis* subsp. *sepedonicus* vary slightly in pathogenicity and in genetic diversity suggesting a relatively recent introduction of a small founding population of the pathogen (Smith et al. 2001; Peters et al. 2005). Genetic evolution can be achieved by vertical or horizontal gene transfer. *Meloidogyne* populations originally did not possess the cell wall-degrading enzymes required to invade host roots. Although the mechanism of horizontal gene transfer remains largely elusive, it has been speculated that a gene coding for a cell wall-degrading enzyme was horizontally transferred from a rhizobial bacterium to the nematode and was kept in the genome of the nematode by strong selection pressures representing important initial steps facilitating the invasion of plants by nematodes (Dieterich and Sommer 2009). By genetic evolution, pathogens can adapt to the different environmental conditions they are submitted to. This enables them to skirt control measures and continuously forced farmers to use new control methods.

### 3.1.5 Diagnosis and detection methods

Rapid detection of plant parasitic pathogens enables to set up adapted control measures and avoid disease expansion and yield losses, even if the infestation level is low. Classical detection methods begin with visual observation and characterization of symptoms followed by identification using morphologic traits for nematodes (Crow et al. 2000; Riga and Neilson 2005; Melakeberhan et al. 2007; Mugniéry 2007) or isolation on selective media for fungi and bacteria. Carbon source utilization, sugar degradation, and production of specific enzymes allow the biochemical identification of bacteria (Flores-Gonzalez et al. 2008; Pitman et al. 2008). However, these classical methods are often not accurate enough to distinguish between different

strains or pathovars of the same species. Molecular biology based-diagnosis and detection methods are expected to complement classical diagnosis. The most developed detection methods are based on polymerase chain reaction (PCR), which amplifies DNA regions specific of the pathogen of interest (Table 5). The quantitative reverse transcriptase PCR is currently among the most powerful methods for the diagnosis of pathogens in complex environments. Indeed, it enables to quantify the ARN of the pathogen present in a sample. Fingerprinting methods—restriction fragment length polymorphism or amplified fragment length polymorphism—are used for intraspecific identification of pathovars or races of bacteria, fungi, or nematodes (Abeln et al. 2002; Cullen et al. 2007; Flores-Gonzalez et al. 2008; Pitman et al. 2008). Fluorescent in situ hybridisation or stable low molecular weight DNA profiles were developed to detect *R. solanacearum* and *C. michiganensis* var. *sepedonicum*, respectively (Ronda et al. 1999; Palomo et al. 2000). Immunological techniques such as immunochromatographical lateral flow, enzyme-linked immunosorbent assay and immunofluorescence are based on the recognition of specific markers at the surface of pathogenic cells to detect and identify the pathogens (Ronda et al. 1999; Merz 2005; Hughes 2008). Fungal pathogens display typical infrared spectra that differ from the spectra of substrate material such as potato; they can be early and rapidly detected by Fourier transform infrared microscopically based technique (Erukhimovitch 2007). Finally, monitoring of normal and disease-induced volatile profiles in stored potatoes or of the light reflected from plant in fields are valuable techniques to detect stress and thus potential pathogenic infections (Ouellette et al. 1990; Heath et al. 2000).

### 3.2 Interactions between microorganisms, organisms, and pathogens

Potato pathogens are not the only microorganisms living in the potato surroundings. A huge microbial biomass is associated and interacts with potatoes. About  $10^7$  bacteria colony forming units per gram of soil live in the potato rhizosphere and potato geocaulosphere which is the volume of soil surrounding the tubers (Lazarovits et al. 2007). The structure of microbial and nematode communities in the geocaulosphere varies according to the plant age and other factors related to cultivar, nutritional status, biotic and abiotic stresses, etc. (Al-Hazmi et al. 1993; Krechel et al. 2002; Ferreira et al. 2008; Desgarennes et al. 2009; Manici and Caputo 2009).

Earthworms and nematodes favor pathogen mobility by transporting them through the soil (Jensen 1978; Table 6). Nematodes enhance potato diseases because they act as vectors of the pathogens. They also enhance the diseases

either by facilitating the development of other pathogens—acting as mechanical wound agents and providers of necrotic tissues for pathogen penetration or nutrition—or by benefiting of their attacks as opportunistic micro-organisms (Jensen 1978).

The microbial or faunal interactions in the geocaulosphere are involved in disease suppressiveness of the soil. Two classical types of suppressiveness of soil are known. General suppression is related to the global activity of the whole microbial biomass in the soil. In contrast, specific suppression is due to the specific activity of certain individuals or groups of microorganisms (Alabouvette et al. 1996; Weller et al. 2002). For instance, *Serratia plymuthica*, *Pseudomonas* spp., *Bacillus* spp., *Streptomyces* spp., and *Trichoderma* spp. (Kumar and Khare 1990; Kamensky et al. 2002; Krechel et al. 2002) are able to decrease the severity of several potato diseases (Table 6). They can be considered as biological control agents. Some biological control agents can act directly against fungal pathogens by enzymatic degradation of their cell walls (Kamensky et al. 2002; Li et al. 2002), by parasitism—as it seems to be the case against numerous nematodes—(Nunez-Camargo et al. 2003; Papert et al. 2004), by antibiotics production (Grosch et al. 2005), by siderophore secretion that reduces the availability of iron needed by plant pathogens (Bharadwaj et al. 2008) or by interfering with communication between pathogens, i.e., by degrading molecules involved in the "quorum sensing" mechanisms of *Pectobacterium* spp. (Dong et al. 2004). Indirectly, biological control agents can lead to the plant strengthening and a better resistance to pathogen attacks by producing plant growth hormone or by inducing the production of plant defense molecules such as phytoalexins and PR proteins (Stevenson et al. 2001; Larkin 2008). Mycorrhizal fungi also have a beneficial effect; inoculation with arbuscular mycorrhizal fungus suppressed tuber dry rot and reduced stem canker and black scurf (Bharadwaj et al. 2008).

### 3.3 Interactions between plants and pathogens

The major method to control potato diseases is to find resistant cultivars to a majority of pathogens especially since the use of chemicals is limited (INRA and Cemagref 2005; Pailloton 2008). Different levels of resistance towards most of the soil-borne potato diseases have been observed among potato cultivars. Wild species of *Solanum* provide excellent sources of disease resistance genes that may be introgress into *S. tuberosum* genome by interspecific crossing (Jansky and Rouse 2003; Table 7) and international structures such as the International Potato Center in Peru are aiming at preserving the genetic diversity of native potatoes. Varieties of potato which contain color pigments are more and more utilized in current breeding programs

**Table 6** Detrimental beneficial and associations of microorganisms with potato soil-borne pathogens

Pathogen	Disease	Organisms enhancing diseases	Organisms reducing diseases	References
<b>Fungi and oomycetes</b>				
<i>Colletotrichum coccodes</i>	Black dot	<i>V. dahliae</i> , <i>S. subterranea</i>		Tsror (2004); Merz and Falloon (2009)
<i>Fusarium</i> spp.	Fusarium dry rots	<i>P. atrosepticum</i> , <i>Meloidogyne</i> spp. <i>D. destructor</i> ; <i>S. subterranea</i>	<i>S. plymuthica</i> , <i>D. destructor</i>	Munzert et al. (1977); Jensen (1978); Gould et al. (2008); Merz and Falloon (2009)
<i>Hemimycesporium solani</i>	Silver scurf			Elson et al. (1997); Rivera-Vargas et al. (2007)
<i>Macromomina phaeolina</i>	Charcoal rot			Kumar and Khare (1990); Gupta et al. (1999)
<i>Phoma andigena</i> var. <i>andina</i>	Phoma leaf spot			
<i>Phoma</i> spp.	Gangrene			
<i>Phytophthora erythroseptica</i>	Pink rot			
<i>Polycephalum pustulans</i>	Skin spot			
<i>Pythium ultimum</i> var. <i>ultimum</i>	Leak			
<i>Rhizoctonia solani</i>	Black scurf/stem canker	<i>G. rostochiensis</i> , <i>Meloidogyne</i> spp. + <i>V. dahliae</i> , <i>Pratylenchus neglectus</i> + <i>V. dahliae</i>		Scholtens and S'Jacob (1989); Krehel et al. (2002); Grosch et al. (2005); Back et al. (2006); Grosch et al. (2006); Santamarina and Rosello (2006); Mahmoud et al. (2008); Wilson et al. (2008)
<i>Rosellinia</i> sp.	Rosellinia black rot			Al-Chaabbi and Matrod (2002)
<i>Sclerotinia sclerotiorum</i>	White mold			Phillips (1989); Kamensky et al. (2002); Yang et al. (2008)
<i>Sclerotium rolfsii</i>	Stem rot			Kumar and Khare (1990); Dey et al. (2004)
<i>Spongopora subterranea</i>	Powdery scab			Merz and Falloon (2009)
<i>Synchytrium endobioticum</i>	Wart			Hampson and Coombes (1989)
<i>Thecaphora solani</i>	Thecaphora smut			Bazan de Segura and Carpio (1974)
<i>Verticillium dahliae</i> and <i>V. albo-atrum</i>	Verticillium wilt			Jensen (1978); Franco and Bendezeu (1985); Scholten and S'Jacob (1989); Krehel et al. (2002); Rotenberg et al. (2004); Tsror (2004); Santamarina and Rosello (2006); Bharadwaj et al. (2008)
<b>Bacteria</b>				
<i>Clavibacter michiganensis</i> spp. <i>sepedonicus</i>	Ring rot			Perombelon et al. (1979)
<i>Clostridium</i> spp.	Bacterial soft rot	<i>Pectobacterium</i> spp.		Munzert et al. (1977); Perombelon et al. (1979); Dong et al. (2004); Bharadwaj et al. (2008)
<i>Pectobacterium atrosepticum</i> , <i>Pectobacterium carotovorum</i> subsp. <i>carotovorum</i>	Black leg, soft rot	<i>Clostridium</i> spp., <i>F. solani</i> var. <i>coeruleum</i>	<i>Bacillus</i> spp., <i>Pseudomonas</i> spp	Jensen (1978); Mahmoud (2007)
<i>Ralstonia solanacearum</i>	Brown rot	<i>G. pallida</i>	<i>P. fluorescens</i> , <i>P. putida</i> , <i>B. subtilis</i>	Non pathogenic <i>Sreptomyces</i>
<i>Sreptomyces scabiei</i> , <i>S. acidiscabiei</i> , <i>S. europeescabiei</i>	Common and netted scab			Wanner (2007)

**Table 6** (continued)

Pathogen	Disease	Organisms enhancing diseases	Organisms reducing diseases	References
<b>Nematodes</b>				
<i>Belenolaimus longicaudatus</i>	Sting nematode			
<i>Ditylenchus destructor</i>	Potato rot nematode			
<i>Globodera pallida, G. rostochiensis</i>	Potato cyst nematode			
<i>Meloidogyne</i> spp.	Root-knot nematode			
<i>Nacobbus aberrans</i>	False root-knot nematode			
<i>Paratrichodorus</i> and <i>Trichodorus</i> spp.	Stubby-root nematode (TRV vector)			
<i>Pratylenchus</i> spp.	Root-lesion nematode			
		<i>V. dahliae, R. solani, V. dahliae</i>	<i>V. dahliae, F. oxysporum, P. exigua</i>	Jensen (1978); Wronkowska and Janowicz (1989); Ryan and Jones (2003); Back et al. (2006); Mugnier and Phillips (2007); IPC (1978); Scholte and S'Jacob (1989); Hafez and Sundararaj (2000); Sankaranayanan and Sundarababu (2001); Krczel et al. (2002)
		<i>P. neglectus, R. solani, V. dahliae</i>	<i>Pseudomonas</i> sp., <i>Streptomyces</i> sp., <i>Rhizobium</i> sp.; <i>Bacillus megaterium</i> var. <i>phosphaticum</i> <i>B. penerans</i> , <i>Glomus</i> <i>mossae</i>	Scholte and S'Jacob (1989); Saeed et al. (1998)
		<i>V. dahliae, R. solani</i>		

because cultivars producing anthocyanins can provide better resistance to soft rot or other diseases compared to white/yellow flesh cultivars (Wegener and Jansen 2007). Cultivars resistant to several diseases were obtained, but simultaneous resistance to all pathogens is very difficult to achieve. Moreover, for some diseases, new genotypes of pathogen appear regularly and overcome plant defense turning the former resistant cultivars into susceptible ones. Hence the levels and durability of field resistance are often highly depending on numerous abiotic and biotic factors still neither well-known or controlled.

Resistant potato cultivars counteract pathogenic attacks by plant defense reactions that generally lead to the production of suberin and antimicrobial agents, activation of defense genes and trigger hypersensitive cell death (Levine et al. 1994) delaying the pathogen development in plant tissues until a wound periderm could form. Susceptible cultivars produce non-uniform deposits of suberin making them less performing against pathogens (Finetti Sialer 1990; Ray and Hammerschmidt 1998). The anti-microbial agents produced by potatoes can be glycoalkaloids ( $\alpha$ -chaconin and  $\alpha$ -solanine), phenolic compounds and phytoalexins, antimicrobial compounds produced by the plant after pathogen attacks (Okopnyi et al. 1983; Lyon 1989; Ray and Hammerschmidt 1998; Zagorskina et al. 2006; Baker et al. 2008; Lerat et al. 2009). Plants also produce inhibitors of virulence factors (Kim et al. 2006). Another plant defense reaction called systemic acquired resistance (SAR) spreads a signal through the surrounding cells. It allows plants to become highly resistant to subsequent infection by the original pathogen but also by a wide variety of other pathogens. For example, foliar SAR-inducing applications of (benzo (1,2,3) thiadiazole-7-carbothioic acid S-methyl ester-BTH and harpin) reduce the numbers of root lesion nematodes (*Pratylenchus* spp.) and root knot nematodes (*Meloidogyne chitwoodi*; Collins et al. 2006).

#### 4 Effects of cultural practices on the occurrence and development of soil-borne potato diseases

Each technical choice made by farmers concerning the way of growing potatoes plays a predominant role on the quantitative and qualitative yield. All cultural practices may impact disease development.

##### 4.1 Rotations

The most traditional way to control diseases is to use crop rotations including a nonhost plant that can "sanitize" the soil (Alabouvette et al. 1996). Several studies show good results when potatoes are grown only once every 3 or 4 years and, as the other practices, it should be thought in a systemic approach (Table 8). The

**Table 7** Some wild potato cultivars harboring resistance towards pathogens

Cultivar	Resistance	References
<i>Solanum vernei</i>	<i>Spongospora subterranea</i>	Merz and Falloon (2009)
<i>Solanum acaule</i>	<i>Clavibacter michiganensis</i> var. <i>sepedonicus</i>	Laurila et al. (2003)
<i>Solanum commersonii</i>	<i>Ralstonia solanacearum</i>	Kim-Lee et al. (2005)
<i>Solanum bulbocastanum</i>	<i>Meloidogyne chitwoodi</i>	Nitzan et al. (2009)
Snowder ( <i>Solanum tuberosum</i> x <i>Solanum berthaultii</i> )	<i>Pythium ultimum</i> and <i>Phytophthora erythroseptica</i>	Salas et al. (2003); Thompson et al. (2007)
<i>Solanum brevideus</i>	<i>Pectobacterium</i> spp.	Ahn et al. (2001)

beneficial effect of crop rotation depends on the host range of the pathogen and its ability to survive in soil in the absence of its host plant thanks to dormant structures such as sclerotia or chlamydospores. Crop rotation must avoid including alternative hosts for the pathogen (Peters et al. 2004). Susceptible weeds—such as hairy nightshade (*Solanum sarrachoides*)—have to be eliminated as they enable the pathogen to survive during the absence of the main host (Boydston et al. 2008). Crop rotation can also fail to control highly specialized pathogens, such as *Globodera* spp., *S. endobioticum*, or *S. subterranea*. These organisms are able to survive for long periods, either saprophytically or as dormant structures, in soil, and a very low inoculum density is sufficient to induce disease (Samaliev et al. 1998; Merz and Falloon 2009). Rotations with potatoes can include very diverse crops (Table 8). If some of those crops have beneficial effects towards potato crop, other might favor pathogen development and should not enter the rotation, or at least not as the crop preceding the potatoes.

#### 4.2 Fertilization and amendments

Supplying plants with micronutrients and macronutrients can be achieved with organic or inorganic fertilizers, either through soil application, foliar spray, or seed treatment (Davis et al. 1994; Panique et al. 1997; Malakouti 2008). Adapted fertilization and amendment allow strong and healthy crops, which are less susceptible to pathogens (Khomyakov and Kostin 1981). Fertilization may also indirectly favor diseases by enhancing foliar development that maintains high level of humidity needed for example for the growth of *Pectobacterium* spp. (Rousselle et al. 1996). Amendments contribute to control diseases by modifying soil properties, especially pH (see Section 2.2) and microbial activities. That could result in specific suppression caused by the stimulated specific antagonistic populations or in general suppression caused by increased microbial activities or both (Lazarovits et al. 2001; Steinberg et al. 2007; Termorshuizen et al. 2006).

For some diseases, such as stem rot, organic fertilizers are more efficient than mineral ones in terms of disease

suppression (Amitava and Maiti 2006; Table 8). Among organic fertilizers, composts are known to have the capacity to suppress diseases, depending on their degree of maturity (organic matter content and microbial activities). The causal agents of disease suppression brought into the soil by compost amendment are complexes of bacterial and fungal populations, which invade the pile during the curing stage, although some residual activity is probably related to fungistatic compounds occurring in the composts (Raviv 2008).

#### 4.3 Tillage management

Potato cultivation traditionally involves intensive soil tillage throughout the cropping period. Mechanical tillage, ridging, and harvesting entail intensive soil disturbance and modify the environmental conditions especially the microbial characteristics of soil, both on quantitative and qualitative aspects (FAO 2008; Vian 2009). As an example, plowing contributes to redistribute vertically the inoculum, which increases the probability of infection (Taylor 2005). Over the last decades, there is a trend to replace plowing by techniques without soil inversion, i.e., no tillage or superficial tillage. It seems that this strategy could lead to some efficient disease suppression by stimulating microbial activity but conversely may limit the nutrient uptake by the plant (Klikocka 2001; Peters et al. 2004; Vian 2009). Therefore, a combination of both biotic and abiotic factors should be clearly balanced (Table 8). Indeed, rotation and conservation tillage practices can improve disease suppression by enhancing the antibiosis abilities of endophytic and root zone bacteria (Peters et al. 2003). On the other side, the plant growth and the macronutrient (N, P, K, Ca, and Mg) contents in potato plant respond positively to a deeper soil caused by plowing (Boligowa and Glen 2003; Nunes et al. 2006).

#### 4.4 Planting, haulm destruction, lifting, and harvesting

Planting, dehaulming, lifting, and harvesting are decisive for disease expression (Table 8). For example, low planting density increases the yield per plant because the foliage has more space to grow. Also, sparse plants are less exposed to

**Table 8** Cultural practices favorable to reduce disease development

Pathogen	Disease	Rotation	Fertilization and amendments	Tillage	Planting, lifting, and harvesting methods	Pesticides	Cultural systems	Storage	References
<b>Fungi and oomycetes</b>									
<i>Colletotrichum coccodes</i>	Black dot	Long rotations (>5 years) With wheat, red clover, alfalfa, rye, maize, orchard grass, fallow, barley	Mouldboard plowing at 30 cm	Avoid water stress, Short interval between haulm destruction and harvesting	Increased by oxamyl Decreased by imazalil, tolethofos-methyl, mancozeb, thiabendazole, fenpropidion and propiconazole		Dry curing and/or low temperatures below 5°C	Hide and Read (1991); Andripon et al. (1997); Denner et al. (2000); Esfahani and Bak (2004); Glaes-Varlet et al. (2004); Cwalina-Ambroziak and Czajka (2006); Nitzan et al. (2006)	
<i>Fusarium</i> spp.	Fusarium dry rots	No monoculture, minimum 3 years of rotation with red clover	Composted manure	Minimum tillage	Early harvesting Short interval between haulm destruction and harvesting, wound healing	Chlorine dioxide, fenpropidion and a mixture of thiabendazole and imazalil, mancozeb	Organic	Dry curing and/or low temperatures below 4°C	Khomyakov and Kostin (1981); Povolny (1995); Carnegie et al. (2001); Lui and Kushalappa (2002); Carter et al. (2003); Olsen et al. (2003); Peters et al. (2004); Cwalina-Ambroziak and Czajka (2006); Raviv (2008)
<i>Helminthosporium solani</i>	Silver scurf	Minimum 3 years of rotation with red clover		Minimum tillage	low planting density. Late planting and early harvesting,	Mancozeb, imazalil, prochloraz, chlorine dioxide, thiabendazole, fenpropidion, benomyl		Dry conditions, and/or temperatures below 4°C	Lennard (1980); Hide and Read (1991); Firman and Allen (1995); Carnegie et al. (1998); Carter et al. (2003); Olsen et al. (2003); Peters et al. (2004); Geary and Johnson (2006); Amadioha (1998)
<i>Macrophomina phaseolina</i> <i>Phoma andigena</i> <i>Phoma</i> spp.	Charcoal rot Phoma leaf spot var. <i>andina</i>	Gangrene			No evident effect of planting time. Early haulm destruction. Lifting at >8°C	2-aminothiobutane, thiabendazole	Organic	Wet conditions and/or temperatures above 15°C	Meredith et al. (1975); Fox and Dashwood (1979); Croke and Logan (1982); Copeland et al. (1980); Ostergaard and Henriksen (1983); Bang (1989); Povolny (1995); Carnegie et al. (1998); Peters et al. (2005); Al-Mughrab et al. (2007); Taylor et al. (2008)
<i>Phytophthora erythrosptica</i>	Pink rot	3 years with barley and red clover		Planting in well-drained fields, harvesting in cool weather, minimizing damages	Mefenoxam, metaxyl-m		Drying after harvesting	Curing in dry conditions at high temperatures	Lennard (1980); Hide and Cayley (1987); Hide and Read (1991); Carnegie et al. (1998)
<i>Polybotryum pustulans</i>	Skin spot			Early harvesting	Imazalil, prochloraz (seed), 2-aminothiobutane, benomyl, thiabendazole				
<i>Pythium ultimum</i> var. <i>ultimum</i>	Leek		Composted manure	Planting in well-drained fields, harvesting in cool weather, minimizing damages,	Mefenoxam		Drying after harvesting	Raviv (2008); Taylor et al. (2008)	

Table 8 (continued)

Table 8 (continued)

Pathogen	Disease	Rotation	Fertilization and amendments	Tillage	Pesticides	Cultural systems	Storage	References
<i>Pectobacterium atrosepticum</i> , <i>Pectobacterium carotovorum</i> subsp., <i>carotovorum</i> , <i>Dickeya</i> spp.	Black leg, soft rot	No monoculture, rotation with wheat, red clover, barley or orchard grass	No over nitrogen	Planting in well-drained fields, roto-tilling, and elimination of infected plants/tubers Limiting wounds	Chlorine dioxide, aluminum and bisulfite salts, naphthoquinone naphthalazin, kasugamycin, stable bleaching powder, streptomycin, benzoyl acid, sodium benzoate, copper oxychloride + metalaxyl, metiram, copper oxychloride + cymoxanil, klorocin	Conventional	Early efficient and quick drying after harvesting	Bushkova et al. (1981); Khomyakov and Kostin (1981); Lewoczko (1992); Saleh and Huang (1997); Karwasta and Parashar (1998); Bartz (1999); Olsen et al. (2003); Medina et al. (2004); Yeganza et al. (2004); Repšiene and Mineikienė 2006
<i>Ralstonia solanacearum</i>	Brown rot	Without solanaceous plants. With barley and flax	Without superphosphate	4 deep plowings after harvest	Tri-potassium phosphate, bleaching powder	Depends on the soil type		Kishore et al. (1996); Mahmoud (2007); Messha et al. (2007)
<i>Streptomyces scabies</i> , <i>S. acidiscabiei</i> , <i>S. europeusabiei</i>	Common and netted scab	With lupin, winter soybean, winter rye or serradilla Without: sugar beet, carrots, pasture	Ammonium lignosulfate, potassium, phosphate, compost, swine manure	Subsoiling	Increased by oxamyl, 3% boric acid, streptomycin, streptomycin sulfate, daminozide, DL-ethionine	No effect		Meredith et al. (1975); Volovik et al. (1980); Hide and Read (1991); Conn and Lazarovits (1999); Park et al. (2002); Soltani et al. (2002); Chaudhari et al. (2003); Mizano et al. (2003); Peters et al. (2004); Scholte (2005); Repšiene and Mineikienė (2006); Henriksen et al. (2007); Al-Mughabbi et al. (2008)
<b>Nematodes</b>								
<i>Belenolaimus longicaudatus</i>	String nematode	Without sorghum-sandanggrass With cotton			1,3-dichloropropene			Crow et al. (2000); Crow et al. (2001); Perez et al. (2000)
<i>Ditylenchus destructor</i>	Potato rot nematode <i>Globodera pallida</i> , <i>Globodera rostochiensis</i>	Potato cyst nematode	Long rotations With peas, flax, rye, oat or rye grass	phosphorus	Oxamyl	Conventional		Rojancovschi (1994)
<i>Meloidogyne</i> spp.	Root-knot nematode			Avoiding dissemination from infected fields with equipment	Dimethyl disulphide, 1,3-dichloropropene, aldicarb, phoxim, A.C. C. 92100, carbofuran, A.C.			Cornejo (1977); Hague et al. (1983); Mulder et al. (1997); Trifonova (1997); Ruijter and Haverkort (1999); Molendijk (1999); Minnis et al. (2004); Coosemans (2005)
<i>Nacobtus aberrans</i>	False root-knot nematode	With cotton, or black fallow Without most crops (carrot, beet, salify, red clover, cereals, vegetables,..)			64475 Methyl bromide, metham sodium, dicloropropen-cloropicrin, metham sodium + 1,3-dichloropropene, fosfiazate + metam sodium, dimethyl disulphide	No effect		Molendijk (1999); Crow et al. (2000); Carter et al. (2003); Coosemans (2005); Hafez and Sundararaj (2006); Charchar et al. (2007); Ingham et al. (2007)
					Abamectin and furatoecarb, A.C. 92100, aldicarb, carbofuran, A.C.			Cornejo (1977); Irarate et al. (1999); Main et al. (2001)
					64475			

Table 8 (continued)

Pathogen	Disease	Rotation	Fertilization and amendments	Tillage	Planting, lifting, and harvesting methods	Pesticides	Cultural systems	Storage	References
<i>Paratrichodorus</i> and <i>Trichodorus</i> spp.	Stubby-root nematode (TRV vector)	With beet, oats, grasses. Without sorghum-sudangrass or velvetbean, maize, wheat, cabbage, rape, barley				Aldicarb (+ oxamyl), 1,3-dichloropropene			Barbez (1983); Perez et al. (2000); Crow et al. (2001); Hafez and Sundararaj (2006)
<i>Pratylenchus</i> spp.	Root-lesion nematode	With wheat, ryegrass, without red clover				1,3-dichloropropene, oxamyl, fosfazate, cadusafos, carbofuran	Organic		Philis (1997); Johnston et al. (1994); Molendijk (1999); Kimpinski et al. (2001); Carter et al. (2003)

the attacks of pathogens than plants at high densities (Milic et al. 2006). Diseases can be reduced by adjusting planting, dehaulming, and harvesting dates and cultivation of early tuberizing cultivars combined with pre-harvesting desiccation of haulms and treatment of seed tubers with chemicals (Sikka and Singh 1976). Black scurf development on tubers has a positive correlation with the curing period (time between haulm destruction and harvest) because infection on tubers continues in the soil even after haulm destruction (Lakra 2000).

#### 4.5 Pesticides

Pesticides are commonly used to control various pathogens altering potato tubers. They can be applied as soil fumigant (fumigants such as carbamates are not allowed in some European countries), sprayed or powdered directly on seed tubers after harvest or applied as granular (Hide et al. 1995; Tsror et al. 2000; Errampalli et al. 2006). The chemicals have to be carefully chosen, since pathogens can adapt and become resistant (Table 8). Thiabendazole-resistance was detected in *Fusarium avenaceum*, *F. culmorum*, *F. equiseti*, and *F. sporotrichioides* (*Fusarium* dry rot; Ocamb et al. 2007), in *P. pustulans* (skin spot; Carnegie et al. 2008) and in *H. solani* (silver scurf). Mefenoxam-resistance is known for *P. erythroseptica* (pink rot) populations (Taylor et al. 2006) and numerous treatments of carbendazim select resistant mutants of *Sclerotium rolfsii* (stem rot; Solunke et al. 2001). Moreover, the use of numerous chemicals is nowadays regulated and many of them are no longer permitted in Europe.

#### 4.6 Organic farming versus conventional agriculture

Organic farming relies on agricultural techniques that exclude the use of chemical pesticides and recommend organic fertilization. As a result, the soil and tuber environment is quite different from the one caused by conventional practices and may induce disease suppression (Table 8). To reduce disease incidence or severity, the best adapted cultural system depends on the pathogen to control and varies strongly according to the soil type (Messiha et al. 2007). It has been reported that farmers who switch from conventional to organic system faced critical pest or disease problems during a transition period of about 5 years but managed to control soil-borne diseases on the long-term (Bruggen and Termorshuizen 2003). However, organic farmers generally faced more sanitary problems than conventional farmers.

#### 4.7 Handling and storage

Inappropriate manipulation of tubers at harvest or during storage can provoke wounds that increase diseases such as

black dot, *Fusarium* dry rots, silver scurf, gangrene, leak, pink rot, black leg, and soft rot (Meredith et al. 1975; Hide 1994; Vanvuurde and Devries 1994; Salas et al. 2000; Marcinkowska et al. 2005; Peters et al. 2008a, b; Table 8). Significant measures of managing potato diseases include: avoiding mechanical damage to potatoes during harvesting, shipping and sorting, curing the harmed parts thereby preventing infection and disease onset, avoiding manipulation of cold potato since potato tubers are more sensitive to injuries when cold, avoiding the exposure of table potato to light, and continuously providing stored potatoes with fresh air (Milosevic and Alovic 2006; Scheid 2006). Most of the storage diseases decrease when the tubers are cured in dry conditions and stored at temperature close to 4°C or 5°C, except gangrene (Table 8). Once again, for storage as for production, a balance between biotic and abiotic conditions should be carefully setup to preserve yield and quality. Indeed, despite they have less infection when stored in a dry atmosphere, tubers show greater weight losses than when they are stored in a humid atmosphere (Lennard 1980).

## 5 Disease management

### 5.1 Risk assessment and decision support systems

Disease occurrence and development influenced by abiotic and biotic factors are difficult to predict. However, their prediction would be very useful to assess disease risk and consequently the potential yield loss and to choose the best disease control strategy. Current methods to evaluate yield losses are based on predictive models which commonly assign a value or score to each risk factor, such as cultivar resistance, inoculum density, cultural practices, and environmental factors. The maximum score that can be assigned to each factor depends on the relative importance of the

factor in determining the disease. For example, cultivar resistance is considered to be a major determinant of powdery scab severity, so this factor has a higher score than the zinc content of soil, which is thought to be less important (Burgess and Wale 1994). Assessment of the risk for each factor and for each disease is performed by bioassays in fields or in growth chambers under controlled conditions. They are generally laborious, time consuming, and costly.


Tolerant cultivars are a particular risk factor in potato production as they can maintain and increase the inoculum level in fields (Merz and Falloon 2009). A tolerance threshold of the crop has to be determined. It takes into account the relationship between inoculum density and disease incidence or severity according to cultivar resistance (Table 4).

A score can also be attributed to each cultural practice in the equation of the model since they have various impacts on yield losses. For example, incidence and severity of *Verticillium* wilt decrease with long rotations (Johnston et al. 1994), but mint as a previous crop increases *Verticillium* wilt (Omer et al. 2008). Consequently, in the equation of the model, rotation length will be negatively correlated to yield losses whereas mint as previous crop will be positively correlated to yield losses due to *Verticillium* wilt.

On the same pattern, some predictable environmental factors such as nutrient contents and soil pH can be scored. However, abiotic environmental factors are difficult to predict. For example, at planting time, rainfall and temperature conditions occurring at the critical growth phase of the disease are almost impossible to foresee. As climatic conditions cannot be predicted at middle term, models of risk assessment are less reliable. However, no factor alone has a dramatic effect on the disease; and the beneficial reduction of a disease is usually achieved by the sum of optimized factors (Harrison 1997).

Mathematical modeling including all the data related to the environmental factors and to the results concerning

**Fig. 2** Input and output parameters of yield loss calculation models



plant resistance appeared to be helpful to evaluate risk, to overcome the scaling gap between bioassays in growth chamber and field application and to simulate scenario based on crop management (Janvier et al. 2007).

Calculation of yield losses enables to identify a damage threshold and to determine the time at which disease control must be initiated. Indeed, yield loss threshold and economic threshold are different. Economic threshold is frequently higher than yield loss threshold; because up to a certain point, losing yield is less penalizing for farmers than spending money to avoid it. Calculation of economic thresholds beyond which control of diseases is profitable takes into account a damage function drift to potato yield, pathogen population density, and crop selling prices. For example, application of control measures is found to be beneficial at an initial density of *G. rostochiensis* higher than eight eggs and larvae per gram of soil, while the damage threshold is at two eggs per gram of soil (Samaliev and Andreev 1998). Economic thresholds allow taking short-term strategic decisions such as choice of the cultivar, cultural practices, timing of crop establishment, seed treatment, planting density, etc. and long-term strategic decisions such as define research priorities, design the breeding programs, or develop integrated pest management strategies (Savary et al. 2006) (Fig. 2). Predicting models are used by farmers as decision support systems (DSS) and generally provide a theoretical yield to be obtained at the end of the cropping period, a monitoring of pest populations and comments and advices in order to increase the theoretical yield as much as possible (Been et al. 2005; Jorg et al. 2006). Some DSS are able to send real time alerts to farmers when several risk factors are combined and when control measures have to be taken immediately (Dubois and Duvauchelle 2004). DSS are environmental and farmer friendly as they enable to increase economical yields by applying the right chemical doses at the right time and when disease pressure requires it, in order to reduce unnecessary environmental pollutions and treatment cost.

## 5.2 Control methods

Ways to control diseases are evolving since the use of chemicals is supposed to be reduced. In many cases, the most efficient long-term strategy is to use resistant cultivars when available. Otherwise, management strategies consist either in exclusion, avoiding contact between plant and pathogens, or by pest eradication, and leading to complete elimination or partial reduction of pathogen populations.

For the potato crop which is multiplied vegetatively, exclusion methods begin with the use of healthy tubers. Many soil-borne pathogens can be carried on by seed tubers and the use of certified seed potatoes is a major way to control or restrict the movement of pathogens of potato

crops (Andrade et al. 2008). Seed certification programs aim at warranting seed tuber quality to potato producers and favor the diffusion of genetic progress. The certified seed production process may be 8–10 years long. Strict rules established by the national regulation institutions (i.e., National Potato Council in USA or Groupement National Interprofessionnel des Semences-Service Officiel de Contrôle in France) have to be respected and the seeds are regularly inspected for bacterial, viral, and fungal diseases, as well as varietal purity and identity. Each country is free to apply more or less severe rules. Certification systems have been developed in most of the seed producing countries to cover the production of certified seed potatoes free from pathogens and pests (McDonald 1995; Grousset and Smith 1998; Sahajdak and Uznanska 2003). An international project of commercial and phytosanitary minimal guidelines (CEE-ONU S-1) is in progress. It is intended to serve as a minimal base consensus between the various standards established at "regional" levels (EU, NATTO, etc.; UNECE 2010).

Eradication strategies aim at eliminating an established pathogen from plant propagation material or production sites. Eradication methods involve the use of pesticides, adapted cultural practices or biological control. Application of fungicides and nematicides are protecting strategies (see Section 4.5 and Table 8) whose application time and doses can be advised by DSS. However, pesticides are sometimes inefficient against pathogens, such as *Pectobacterium carotovorum* (Latour et al. 2008), or their use is limited by environmental regulations. Consequently, alternative methods based on adapted cultural practices have to be recommended (see Section 4 and Tables 3 and 8). Some crops either susceptible or resistant may serve as baiting crop, for example, resistant potato cultivars cropped just before the main potato crop decreased black scurf (Scholte 2000). Likewise, alfalfa can be used to avoid TRV transmitted by stubby root nematode, as this crop is a host for stubby root nematode but immune to TRV (Stevenson et al. 2001). Cultivar precocity can be used to avoid some diseases. Since black dot and charcoal rot damages occur late in the growing season, early cultivars are generally recommended to control these diseases (Stevenson et al. 2001). When a disease is established in a production site, its spread must be avoided as much as possible. All diseased plants have to be eliminated or burned and tools should be properly disinfected before use in another field (Salas et al. 2000; Latour et al. 2008).

Natural interactions of plants and microorganisms with the pathogens are used as biological control to protect potato crops. There is a continuum from a conducive soil to a suppressive one (Alabouvette et al. 1996) what means that in each soil, almost each pathogen can be potentially controlled by other microorganisms either by a specific

antagonism or by competition with total microbial biomass (see Section 3.2 and Table 6). Appropriate agricultural practices, thanks to the DSS, should stimulate this potential to enhance or to maintain the soil suppressiveness to potato diseases.

Another approach consists in applying biocontrol agents. However, the choice of a biological control agent must take into account the potential risks to human health. Even if *Serratia grimesii* and *Burkholderia cepacia* decrease dry rot and black scurf and stem canker, respectively, they can cause human infections and are not recommended for biological control (Table 6; Grosch et al. 2005; Gould et al. 2008). Moreover, indirect control such as strengthening of potato plants by mycorrhization increases tuber yield and allow an integrated management of potato cyst nematode and root-knot nematode (Sankaranarayanan and Sundarababu 2001; Ryan et al. 2003). Biological control may also include the use of natural toxic compounds for pathogenic agents. Fumigation of essential oils is studied to control dry rot, gangrene, black scurf, and stem canker (Bang 2007). Fish emulsion and crushed crab shell are used against *Verticillium dahliae*, *Verticillium albo-atrum*, and *S. endobioticum*, respectively (Hampson and Coombes 1995; Abbasi et al. 2006). Soil can be disinfected from pathogens by biofumigation or solar heating or both. For example, *Brassica* crops used in crop rotations and as green manure have been associated with reductions in soil-borne pests and pathogens. These reductions have been attributed to the production of volatile sulfur compounds through the process of biofumigation and to changes in soil microbial community structure (Janvier et al. 2007). Composting is also a sanitizing method which combines temperature, time, and toxic compounds to control potato diseases. The composts the most frequently used on potato crop are organic wastes (sludge, manure, tea, etc.) that have undergone long, thermophilic and, aerobic decomposition. The most effective compost composition and combinations of temperature and time have to be determined for each pathogen. As it decreases the pathogenic population and/or favors microbial enrichment of the soil, compost has generally a positive or neutral effect on disease suppression and only rarely a disease stimulating effect (Termorshuizen et al. 2006). Sanitization is also performed on tubers before planting by hot water (Janvier et al. 2007) or during storage with chemical treatments at high temperatures (Secor et al. 1988). However, heating may damage tubers resulting in fewer sprouts. Biocontrol can also be performed by disrupting pathogens molecular pathways. *P. carotovorum* quorum-sensing mechanism is controlled by a quorum-quenching strategy aiming at interrupting the quorum-sensing by using compounds or organisms able to cause interferences in the bacterial signal (Latour et al. 2008). Finally, it is also possible to

enhance plant defense reactions against soil-borne pathogens by foliar spraying with different inducers such as salicylic acid, di-potassium hydrogen phosphate, and tri-potassium phosphate (Mahmoud 2007).

The different methods that were presented above are not items that have to be taken at random. Their combination generally gives better results than each of the method applied alone.

Decision support systems developed to predict yield losses allow choosing good control methods such as the use of healthy seeds, adapted pesticides, cultural practices, and biological control agents for each potato diseases.

## 6 Conclusions

If a disease results from the interaction between the plant and a pathogen, its severity is influenced by soil abiotic and biotic factors affecting the plant, the pathogen, or both (Alabouvette et al. 1996). Biotic and abiotic factors are not independent, the abiotic factors modulating the biotic ones. They act both on the disease epidemiology, that means the environmental conditions which make the plant growing and the pathogen, present or latent on the crop, causing or not the disease. Moreover, some unfavorable factors for a given disease can be favorable to another. The multifaceted interactions between plants, pathogens and their environment make disease management complex since controlling every factor occurring in the disease development is quite impossible. Potato producers have to aim at limiting contact between plant and pathogens by using for example healthy seeds. Moreover, pathosystems are continuously changing since the pathogens genetically adapt to their hosts or to the environmental conditions implemented by human activities or not. In a system whose parameters vary continuously, the control strategies have to be adapted to each situation at every time.

This review aimed at being as exhaustive as possible about the factors impacting the occurrence and development of the soil-borne potato diseases. Such a work putting in relation numerous potato diseases and comparing their development conditions, the ecology of the causal pathogens and their abiotic and biotic interactions responds to a clear demand from both scientists, extension services, breeders, and farmers. Studies dealing with potato diseases frequently consider only one or few diseases at the same time. Thus, this review constitutes by itself a decision support system since the optimal factors limiting disease development are listed. Nevertheless, the data collected here deal more with diseases known in developed counties and those which cause severe economical losses. Knowledge about minor diseases such as *Phoma* leaf spot, *Rosellinia* black rot, and *Thecaphorpha* smut are extremely rare, probably because these diseases occur in very isolated areas. *Phoma* leaf spot was

recorded only in Bolivia and Peru, *Rosellinia* black rot was described in South America and Africa, and *Thecaphora* smut in South America and Mexico.

Moreover, soil-borne diseases are difficult to study because soil is a complex environment in which numerous interactions occur and where detection of pathogens is not easily performed. However, researches on those diseases could be beneficial at long-term in case they would spread throughout the world. It would have been rather complex to consider air-borne diseases in addition to soil-borne diseases of potato. However, air-borne diseases such as late blight caused by *Phytophthora infestans* and early blight caused by *Alternaria solani* are responsible for huge economical losses and have to be considered with as much attention as soil-borne diseases. Finally, since few years, importance of potato tuber quality raised in developed countries where tubers are washed before selling. Indeed, washing tuber makes visible some superficial blemishes that were previously hidden by adhering soil. Consumer's habits changing, blemished tubers cannot be sold anymore and the losses take seriously damaging proportions for potato market.

The previous considerations acknowledge the fact that the plant disease problem can be reduced in short term thanks to solid knowledge in epidemiology and pathogens ecology; but in longer term, control strategies must be adapted with the constant evolution of pathosystems.

**Acknowledgements** Marie Fiers was financially supported by a PhD funding from the National Association of Technical Research (ANRT) (CIFRE n°1085/2006). This work was part of a Program of Collaborative Research (PRC) between Bretagne Plants and Germicopa, subsidized by the Regional Council of Brittany.

## References

- Abbasi PA, Conn KL, Lazarovits G (2006) Effect of fish emulsion used as a preplanting soil amendment on *Verticillium* wilt, scab, and tuber yield of potato. Canadian Journal of Plant Pathology-Revue Canadienne De Phytopathologie 28:509–518
- Abeln ECA, Stax AM, de Gruyter J, van der Aa HA (2002) Genetic differentiation of *Phoma exigua* varieties by means of AFLP fingerprints. Mycol Res 106:419–427. doi:[10.1017/s0953756202005804](https://doi.org/10.1017/s0953756202005804)
- Achenbach U, Paulo J, Ilarionova E, Lubeck J, Strahwald J, Tacke E, Hofferbert HR, Gebhardt C (2009) Using SNP markers to dissect linkage disequilibrium at a major quantitative trait locus for resistance to the potato cyst nematode *Globodera pallida* on potato chromosome V. Theor Appl Genet 118:619–629. doi:[10.1007/s00122-008-0925-x](https://doi.org/10.1007/s00122-008-0925-x)
- Adams MJ (1980) The role of seed tuber and stem inoculum in the development of gangrene in potatoes. Ann Appl Biol 96:17–28
- Adams MJ, Read PJ, Lapwood DH, Cayley GR, Hide GA (1987) The effect of irrigation on powdery scab and other tuber diseases of potatoes. Ann Appl Biol 110:287–294
- Ahn YK, Kang JC, Kim HY, Lee SD, Park HG (2001) Resistance to blackleg and tuber soft rot in interspecific somatic hybrids between *S. brevidens* and *S. tuberosum* ('Superior', 'Dejima', and dihaploid of 'Superior'). J Korean Soc Horticultural Sci 42:430–434
- Al-Chaabi S, Matrod L (2002) Laboratory study to evaluate efficacy of different *Trichoderma* spp. isolates on some soil-borne pathogenic fungi. Arab J Plant Prot 20:77–83
- Al-Hazmi AS, Ibrahim AAM, Abdul-Raziq AT (1993) Distribution, frequency and population density of nematodes associated with potato in Saudi Arabia. Afro Asian J Nematology 3:107–111
- Al-Mughrabi KI, Bertheleme C, Livingston T, Burgoine A, Poirier R, Vikram A (2008) Aerobic compost tea, compost and a combination of both reduce the severity of common scab (*Streptomyces scabiei*) on potato tubers. J Plant Sci 3:168–175
- Al-Mughrabi KI, Peters RD, Platt HW, Moreau G, Vikram A, Poirier R, MacDonald I (2007) In-furrow applications of metalaxyl and phosphite for control of pink rot (*Phytophthora erythroseptica*) of potato in new brunswick. Can Plant Dis 91:1305–1309. doi:[10.1094/pdis-91-10-1305](https://doi.org/10.1094/pdis-91-10-1305)
- Alabouvette C, Hooper H, Lemanceau P, Steinberg C (1996) Soil suppressiveness to diseases induced by soilborne plant pathogens. Soil Biochem 9:371–413
- Alabouvette C, Raaijmakers J, De Boer W, Notz R, Défago G, Steinberg C, Lemanceau P (2006) Concepts and methods to assess the phytosanitary quality of soils. In: Bloem J, Hopkins DW, Benedetti A (eds) Plant-microbe interactions and soil quality handbook. CABI Publishing, Wallingford, pp 257–270
- Amadioha AC (1997) Interaction of hydrolytic enzymes produced by *Rhizoctonia bataticola* during rot development. Acta Phytopathologica et Entomologica Hungarica 32:79–87
- Amadioha AC (1998) Control of post harvest tuber rot of potato incited by *Rhizoctonia bataticola*. Arch Phytopathol Plant Prot 31:225–231
- Amadioha AC, Adisa VA (1999) Microbial deterioration of potato tubers (*Solanum tuberosum* L.) in Nigeria. Afr J Root Tuber Crops 3:40–43
- Amitava B, Maiti MK (2006) Role of host nutrition and varieties on the development of stem rot of potato. Ann Plant Prot Sci 14:479–480
- Anaya GB, Perez NJ, Rodriguez D, Crozzoli R, Greco N (2005) Response of genetically improved potato clones to the cyst nematode, *Globodera rostochiensis*, and response of a resistant clone to the nematode in microplots. Nematropica 35:145–154
- Andrade O, Munoz G, Galdames R, Duran P, Honorato R (2004) Characterization, in vitro culture, and molecular analysis of *Thecaphora solani*, the causal agent of potato smut. Phytopathology 94:875–882
- Andrade SN, Contreras MA, Castro UI (2008) Effect of using certified and uncertified potato seeds on the yield and sanitary conditions of a potato crop. Agro Sur 36:63–66
- Andrivon D, Ramage K, Guerin C, Lucas JM, Jouan B (1997) Distribution and fungicide sensitivity of *Colletotrichum coccodes* in French potato-producing areas. Plant Pathol 46:722–728
- Anthoine G, Buisson A, Gauthier JP, Mugnier D (2006) Aspects of the biology of *Nacobbus aberrans* (Thorn 1935) Thorne and Allen 1944 (Nematoda: Pratylenchidae): 2—capacities of development on hosts under in vivo and in vitro conditions. Bull OEPP 36:365–372
- Aqeel AM, Pasche JS, Gudmestad NC (2008) Variability in morphology and aggressiveness among North American vegetative compatibility groups of *Colletotrichum coccodes*. Phytopathology 98:901–909. doi:[10.1094/phyto-98-8-0901](https://doi.org/10.1094/phyto-98-8-0901)
- Atallah ZK, Bae J, Jansky SH, Rouse DI, Stevenson WR (2007) Multiplex real-time quantitative PCR to detect and quantify *Verticillium dahliae* colonization in potato lines that differ in

- response to *Verticillium* wilt. *Phytopathology* 97:865–872. doi:[10.1094/phyto-97-7-0865](https://doi.org/10.1094/phyto-97-7-0865)
- Atallah ZK, Stevenson WR (2006) A methodology to detect and quantify five pathogens causing potato tuber decay using real-time quantitative polymerase chain reaction. *Phytopathology* 96:1037–1045. doi:[10.1094/phyto-96-1037](https://doi.org/10.1094/phyto-96-1037)
- Atkins SD, Manzanilla-Lopez RH, Franco J, Peteira B, Kerry BR (2005) A molecular diagnostic method for detecting *Nacobbus* in soil and in potato tubers. *Nematology* 7:193–202
- Aveskamp MM, De Gruyter J, Crous PW (2008) Biology and recent developments in the systematics of *Phoma*, a complex genus of major quarantine significance. *Fungal Divers* 31:1–18
- Baard SW, Pauer GDC (1981) Effect of alternate drying and wetting of the soil fertilizer amendment and pH on the survival of micro sclerotia of *Verticillium dahliae*. *Phytophylactica* 13:165–168
- Baayen RP, Bonthuis H, Withagen JCM, Wander JGN, Lamers JL, Meffert JP, Coelius G, van Leeuwen GCM, Hendriks H, Heerink BGJ, van den Boogert PHF, van de Griend P, Bosch RA (2005) Resistance of potato cultivars to *Synchytrium endobioticum* in field and laboratory tests, risk of secondary infection, and implications for phytosanitary regulations. *Bull OEPP* 35:9–23
- Baayen RP, Coelius G, Hendriks H, Meffert JP, Bakker J, Bekker M, van den Boogert P, Stachewicz H, van Leeuwen GCM (2006) History of potato wart disease in Europe—a proposal for harmonisation in defining pathotypes. *Eur J Plant Pathol* 116:21–31. doi:[10.1007/s10658-006-9039-y](https://doi.org/10.1007/s10658-006-9039-y)
- Back M, Haydock P, Jenkinson P (2006) Interactions between the potato cyst nematode *Globodera rostochiensis* and diseases caused by *Rhizoctonia solani* AG3 in potatoes under field conditions. *Eur J Plant Pathol* 114:215–223. doi:[10.1007/s10658-005-5281-y](https://doi.org/10.1007/s10658-005-5281-y)
- Bae J, Atallah ZK, Jansky SH, Rouse DI, Stevenson WR (2007) Colonization dynamics and spatial progression of *Verticillium dahliae* in individual stems of two potato cultivars with differing responses to potato early dying. *Plant Dis* 91:1137–1141. doi:[10.1094/pdis-91-9-1137](https://doi.org/10.1094/pdis-91-9-1137)
- Bain RA, Lennard JH, Wastie RL (1987) The influence of cultivar and isolate on the development of gangrene (*Phoma exigua* var. *foveata*) in potato tubers. *Ann Appl Biol* 111:535–540
- Bains PS, Bisht VS, Benard DA (1996) Soil survival and thiabendazole sensitivity of *Helminthosporium solani* isolates from Alberta. *Can Potato Res* 39:23–29
- Baker KF (1970) Types of *Rhizoctonia solani* diseases and their occurrence. In: Parmeter JR (ed) *Rhizoctonia solani*, biology and pathology symposium. University of California Press, Los Angeles, pp 125–148
- Baker CJ, Whitaker BD, Mock NM, Rice CP, Roberts DP, Deahl KL, Ueng PP, Aver'yanov, AA (2008) Differential induction of redox sensitive extracellular phenolic amides in potato. *Physiological and Molecular Plant Pathology* 73:109–115. doi: [10.1016/j.pmp.2009.03.003](https://doi.org/10.1016/j.pmp.2009.03.003)
- Baljeet S, Lakra BS, Ram N, Mahender S (2005) Influence of depth of planting on development of black scurf of potato (*Rhizoctonia solani*). *Ann Biol* 21:241–244
- Bang U (1989) Cultivating measures for potatoes (*Solanum tuberosum* L.) and climatic factors affecting the gangrene pathogen *Phoma foveata* Foister, Vaxtskyddsrapporter, Avhandlingar, p 33
- Bang U (2007) Screening of natural plant volatiles to control the potato (*Solanum tuberosum*) pathogens *Helminthosporium solani*, *Fusarium solani*, *Phoma foveata* and *Rhizoctonia solani*. *Potato Res* 50:185–203. doi:[10.1007/s11540-008-9044-y](https://doi.org/10.1007/s11540-008-9044-y)
- Banyal DK, Mankotia V, Sugha SK (2008) Soil characteristics and their relation to the development of tomato collar rot caused by *Sclerotium rolfsii*. *Indian Phytopathol* 61:103–107
- Barbez D (1983) The distribution of virus-vector nematodes in seed-potato fields in Flanders (Belgium) and its relation to some biotic and abiotic factors. *Mededelingen van de Faculteit Landbouwwetenschappen Rijksuniversiteit Gent* 48:401–415
- Bardin SD, Huang HC, Moyer JR (2004) Control of *Pythium* damping-off of sugar beet by seed treatment with crop straw powders and a biocontrol agent. *Biol Control* 29:453–460. doi:[10.1016/j.biocntrol.2003.09.001](https://doi.org/10.1016/j.biocntrol.2003.09.001)
- Bartz JA (1999) Suppression of bacterial soft rot in potato fibers by application of kasugamycin. *Am J Potato Res* 76:127–136
- Bazan de Segura C, Carpio Rd (1974) Potato gangrene on the central Peruvian coast. *Informe Ministerio de Agricultura*, p 27
- Been TH, Schomaker CH, Molendijk LPG (2005) NemaDecide: a decision support system for the management of potato cyst nematodes. In: Haverkort AJ and Struik PC (eds) Potato in progress: science meets practice, Potato 2005, Emmeloord, NL, 5–7 Sept 2005, pp 143–155
- Bdliya BS, Dahiru B (2006) Efficacy of some plant extracts on the control of potato tuber soft rot caused by *Erwinia carotovora* ssp. *carotovora*. *J Plant Prot Res* 46:285–294
- Bharadwaj DP, Lundquist PO, Alstrom S (2008) Arbuscular mycorrhizal fungal spore-associated bacteria affect mycorrhizal colonization, plant growth and potato pathogens. *Soil Biol Biochem* 40:2494–2501. doi:[10.1016/j.soilbio.2008.06.012](https://doi.org/10.1016/j.soilbio.2008.06.012)
- Bisht NS (1982) Control of *Sclerotium* rot of potato. *Indian Phytopathol* 35:148–149
- Blum B, Merz U (1993) Occurrence of *Spongospora subterranea*, the causal agent of powdery scab disease of potatoes, in selected potato-producing areas. *Landwirtschaft Schweiz* 6:333–339
- Blum LEB, Prada A, Medeiros EAA, Amarante CVT (2002) Temperature, light and culture medium affecting the production of sclerotia of *Sclerotium rolfsii* and *Sclerotinia sclerotiorum*. *Revista de Ciencias Agroveterinarias* 1:27–32
- Boligowa E, Glen K (2003) Yielding and quality of potato tubers depending on the kind of organic fertilisation and tillage method. *Electronic Journal of Polish Agricultural Universities, Agronomy* 6:1–7
- Boogert PHJFVd, Gent-Pelzer MPEv, Bonants PJM, Boer SHd, Wander JGN, Levesque CA, Leeuwen GCMv, Baayen RP (2005) Development of PCR-based detection methods for the quarantine phytopathogen *Synchytrium endobioticum*, causal agent of potato wart disease. *Eur J Plant Pathol* 113:47–57. doi:[10.1007/s10658-005-0297-x](https://doi.org/10.1007/s10658-005-0297-x)
- Bouchek-Mechiche K, Gardan L, Andrivon D, Normand P (2006) *Streptomyces turgidiscabies* and *Streptomyces reticulascabiei*: one genomic species, two pathogenic groups. *Int J Syst Evol Microbiol* 56:2771–2776. doi:[10.1099/ijtsd.0.63161-0](https://doi.org/10.1099/ijtsd.0.63161-0)
- Bouchek-Mechiche K, Pasco C, Andrivon D, Jouan B (2000) Differences in host range, pathogenicity to potato cultivars and response to soil temperature among *Streptomyces* species causing common and netted scab in France. *Plant Pathol* 49:3–10
- Boydston RA, Mojtabaei H, Crosslin JM, Brown CR, Anderson T (2008) Effect of hairy nightshade (*Solanum sarrachoides*) presence on potato nematodes, diseases, and insect pests. *Weed Sci* 56:151–154. doi:[10.1614/ws-07-035.1](https://doi.org/10.1614/ws-07-035.1)
- Bradbury JF (1977) *Erwinia carotovora* var. *atroseptica*. [Descriptions of Fungi and Bacteria], IMI Descriptions of Fungi and Bacteria, Sheet 551
- Brown MJ, Riedel RM, Rowe RC (1980) Species of *Pratylenchus* associated with *Solanum tuberosum* cv Superior in Ohio. *J Nematology* 12:189–192
- Bruggen AHCv, Termorshuizen AJ (2003) Integrated approaches to root disease management in organic farming systems. *Australas Plant Pathol* 32:141–156. doi:[10.1071/ap03029](https://doi.org/10.1071/ap03029)
- Burgess PJ, Wale SJ (1994) Development of an integrated control strategy for powdery scab of potatoes. Brighton Crop Protection Conference. *Pest Dis* 1–3:301–306

- Burlakoti RR, Estrada R, Rivera VV, Boddeda A, Secor GA, Adhikari TB (2007) Real-time PCR quantification and mycotoxin production of *Fusarium graminearum* in wheat inoculated with isolates collected from potato, sugar beet, and wheat. *Phytopathology* 97:835–841. doi:10.1094/phyto-97-7-0835
- Bushkova LN, Shuvalova GV, Derzhipil'skii LM (1981) Some aspects of protection of potato and root cruciferous crops against bacterioses. *Trudy Vsesoyuznogo Nauchno-Issledovatel'skogo Instituta Zashchity Rastenii*, 71–74
- Campion C, Chatot C, Perraton B, Andrivon D (2003) Anastomosis groups, pathogenicity and sensitivity to fungicides of *Rhizoctonia solani* isolates collected on potato crops in France. *Eur J Plant Pathol* 109:983–992
- Carling DE, Baird RE, Gitaitis RD, Brainard KA, Kuninaga S (2002) Characterization of AG-13, a newly reported anastomosis group of *Rhizoctonia solani*. *Phytopathology* 92:893–899. doi:UNSP 2002-0604-01R
- Carnegie SF (1991) The role of soil, seed-potato tuber and haulm in the transmission of *Phoma foveata* from infected seed to daughter tubers. *Plant Pathol* 40:352–358
- Carnegie SF, Cameron AM, Haddon P (2001) The effect of date of haulm destruction and harvest on the development of dry rot caused by *Fusarium solani* var. *coeruleum* on potato tubers. *Ann Appl Biol* 139:209–216
- Carnegie SF, Cameron AM, Haddon P (2008) Effects of fungicide and rate of application on the development of isolates of *Polyscytalum pustulans* resistant to thiabendazole and on the control of skin spot. *Potato Res* 51:113–129. doi:10.1007/s11540-008-9094-1
- Carnegie SF, Cameron AM, Lindsay DA, Sharp E, Nevison IM (1998) The effect of treating seed potato tubers with benzimidazole, imidazole and phenylpyrrole fungicides on the control of rot and skin blemish diseases. *Ann Appl Biol* 133:343–363
- Carter MR, Kunelius HT, Sanderson JB, Kimpinski J, Platt HW, Bolinder MA (2003) Productivity parameters and soil health dynamics under long-term 2-year potato rotations in Atlantic Canada. *Soil Tillage Res* 72:153–168. doi:10.1016/s0167-1987(03)00085-0
- Chandell ST, Gaur HS, Alam MM (2002) Effect of tillage and water management on the population behaviour of root-knot nematodes *Meloidogyne triticioryae* in rice crop. *Arch Phytopathol Plant Prot* 35:195–200
- Chang RJ, Ries SM, Pataky JK (1992) Local sources of *Clavibacter michiganensis* ssp. *michiganensis* in the development of bacterial canker on tomatoes. *Phytopathology* 82:553–560
- Charchar JM, Vieira JV, Oliveira VR, Moita AW (2007) Effects of fumigant and nonfumigant nematicides to control *Meloidogyne* spp. on potato and carrot. *Nematologia Bras* 31:59–66
- Chaudhari SM, Patel RN, Khurana SMP, Patel RL, Patel NH (2003) Management of common scab of potato. *J Indian Potato Assoc* 30:135–136
- Chowdary NB, Govindaiah (2007) Influence of different abiotic conditions on the growth and sclerotial production of *Macrophomina phaseolina*. *Indian J Sericulture* 46:186–188
- Chowdhury N, Dey TK, Khan AL (1993) Effect of culture media, light, temperature and pH on the mycelial growth and sclerotia formation of *Sclerotium rolfsii* Sacc. *Bangladesh J Bot* 22:149–153
- Christ BJ (1989) Effect of planting date and inoculum level on incidence and severity of powdery scab on potato. *Potato Res* 32:419–424
- Coelho RMS, De Castro HA, Menezes M (1997) Sporulation of *Phomopsis* and *Phoma* on different culture media, temperature and luminosity conditions. *Summa Phytopathologica* 23:176–180
- Collins HP, Navare DA, Riga E, Pierce FJ (2006) Effect of foliar applied plant elicitors on microbial and nematode populations in the root zone of potato. *Commun Soil Sci Plant Anal* 37:1747–1759. doi:10.1080/00103620600710538
- Combrink NJJ, Prinsloo KP, Jandrell AC (1975) The effect of calcium, phosphate and boron on the keeping quality and quality determining tuber characteristics of potatoes. *Agroplante* 7:81–84
- Conn KL, Lazarovits G (1999) Impact of animal manures on *Verticillium* wilt, potato scab, and soil microbial populations. *Canadian Journal of Plant Pathology-Revue Canadienne De Phytopathologie* 21:81–92
- Coosemans J (2005) Dimethyl disulphide (DMDS): a potential novel nematicide and soil disinfectant. *Proceedings of the VIth International Symposium on Chemical and Non-Chemical Soil and Substrate Disinfestation*, 57–63
- Copeland RB, Logan C, Little G (1980) Fungicidal control of potato black scurf. Tests of Agrochemicals and Cultivars (A supplement to Annals of Applied Biology, Vol. 94) 1:36–37
- Cornejo QW (1977) Chemical control of *Nacobbus aberrans* and *Globodera* spp. *Nemotropica* 7:6
- Correll JC, Gordon TR, McCain AH (1988) Vegetative compatibility and pathogenicity of *Verticillium albo-atrum*. *Phytopathology* 78:1017–1021
- Croke F, Logan C (1982) The effect of humidity on potato gangrene development in naturally contaminated tubers. *Plant Pathol* 31:61–64
- Crow WT, Weingartner DP, Dickson DW (2000) Effects of potato–cotton cropping systems and nematicides on plant-parasitic nematodes and crop yields. *J Nematology* 32:297–302
- Crow WT, Weingartner DP, Dickson DW, McSorley R (2001) Effect of sorghum-sudangrass and velvetbean cover crops on plant-parasitic nematodes associated with potato production in Florida. *J Nematology* 33:285–288
- Cullen DW (2002) Detection of *Colletotrichum coccodes* from soil and potato tubers by conventional and quantitative real-time PCR. *Plant Pathol* 51:281–292
- Cullen DW, Toth IK, Boonham N, Walsh K, Barker I, Lees AK (2007) Development and validation of conventional and quantitative polymerase chain reaction assays for the detection of storage rot potato pathogens, *Phytophthora erythroseptica*, *Pythium ultimum* and *Phoma foveata*. *J Phytopathol* 155:309–315
- Cullen DW, Toth IK, Pitkin Y, Boonham N, Walsh K, Barker I, Lees AK (2005) Use of quantitative molecular diagnostic assays to investigate *Fusarium* dry rot in potato stocks and soil. *Phytopathology* 95:1462–1471. doi:10.1094/phyto-95-1462
- Cunha MG, Rizzo DM (2004) Occurrence and epidemiological aspects of potato silver scurf in California. *Horticultura Bras* 22:690–695. doi:10.1590/s0102-05362004000400005
- Cwalina-Ambroziak B, Czajka W (2006) Chemical control as a factor decreasing the infestation of potato tubers by pathogenic fungi. *Prog Plant Prot* 46:660–663
- Davet P (1970) Recherches sur le *Colletotrichum coccodes* (Wallr.) Hugues. La phase non parasitaire. *Cah ORSTOM Ser Biol* 12:83–96
- Davis JR, Huisman OC, Everson DO, Schneider AT (2001) *Verticillium* wilt of potato: a model of key factors related to disease severity and tuber yield in southeastern Idaho. *Am J Potato Res* 78:291–300
- Davis JR, Huisman OC, Westermann DT, Hafez SL, Everson DO, Sorensen LH, Schneider AT (1996) Effects of green manures on *Verticillium* wilt of potato. *Phytopathology* 86:444–453
- Davis JR, Stark JC, Sorensen LH, Schneider AT (1994) Interactive effects of nitrogen and phosphorus on *Verticillium* wilt of Russet Burbank potato. *Am Potato J* 71:467–481
- Delleman J, Mulder A, Peeten JMG, Shipper E, Turkensteen LJ (2005) Potato diseases. NIVAP Holland; Aardappelwereld magazine
- Denner FDN, Millard CP, Wehner FC (2000) Effect of soil solarisation and mouldboard ploughing on black dot of potato, caused by *Colletotrichum coccodes*. *Potato Res* 43:195–201

- Derevenko AS, Saltykova LP, Yakovleva VI, Pasechnik PS (1981) An experiment on the elimination of potato wart. Zashchita Rastenii, 44
- Desgarnnes D, Sanchez-Nava P, Pena-Santiago R, Carrion G (2009) Nematode fauna associated with the rhizosphere of potato crop (*Solanum tuberosum*) grown in the region of Cofre de Perote, Veracruz, Mexico. Revista Mexicana De Biodiversidad 80:611–614
- Dey TK, Saha AK, Rahman M, Ali MS (2004) Biocontrol potential of *Trichoderma* spp. against *Sclerotium rolfsii* causing stem and tuber rot of potato. Bangladesh J Plant Pathol 20:31–34
- Dhingra OD, Sinclair JB (eds) (1977) An annotated bibliography of *Macrophomina phaseolina* 1905–1975, Viçosa
- Dieterich C, Sommer RJ (2009) How to become a parasite—lessons from the genomes of nematodes. Trends Genet 25:203–209. doi:10.1016/j.tig.2009.03.006
- Dillard HR, Cobb AC (1998) Survival of *Colletotrichum coccodes* in infected tomato tissue and in soil. Plant Dis 82:235–238
- Dong YH, Zhang XF, Xu JL, Zhang LH (2004) Insecticidal Bacillus thuringiensis silences Erwinia carotovora virulence by a new form of microbial antagonism, signal interference. Appl Environ Microbiol 70:954–960
- Dubois L, Duvauchelle S (2004) Integrated control of potato late blight: MILPV, a new French decision support system. Phytonma 575:11–13
- Eichenlaub R, Bermpohl A, Meletzus D (1991) Genetic and physiological aspects of the pathogenic interaction of *Clavibacter michiganense* subsp. *michiganense* with the host plant. Adv Mol Genet Plant Microbe Interact 1(10):99–102
- El-Hassan KI, El-Saman MG, Mosa AA, Mostafa MH (2007) Variation among *Fusarium* spp. the causal of potato tuber dry rot in their pathogenicity and mycotoxins production. Egypt J Phytopathol 35:53–68
- El Bakali AM, Martin MP (2006) Black scurf of potato. Mycologist 20:130–132
- El Fahl AM, Calvert EL (1976) The effect of soil treatment with sulphur and lime on the incidence of potato diseases with special reference to blight. Rec Agric Res 24:7–12
- Elson MK, Schisler DA, Bothast RJ (1997) Selection of microorganisms for biological control of silver scurf (*Helminthosporium solani*) of potato tubers. Plant Dis 81:647–652
- EPPO (1990) *Thecaphora solani*. EPPO Bulletin, pp 1–3
- EPPO (2008) *Ditylenchus destructor* and *Ditylenchus dipsaci*. EPPO Bulletin, pp 363–373
- Errampalli D (2001) Emergence of silver scurf (*Helminthosporium solani*) as an economically important disease of potato. Plant Pathol 50:141–153
- Errampalli D, Peters RD, MacIsaac K, Darrach D, Boswall P (2006) Effect of a combination of chlorine dioxide and thiophanate-methyl pre-planting seed tuber treatment on the control of black scurf of potatoes. Crop Prot 25:1231–1237. doi:10.1016/j.cropro.2006.03.002
- Erukhimovich V (2007) Early and rapid detection of potato's fungal infection by Fourier transform infrared microscopy. Appl Spectrosc 61:1052–1056
- Esfahani AN, Bak AM (2004) Biological and cultural control of black dot disease of potato. J Sci Technol Agric Nat Resour 8:193–207
- Falloon R (1997) Powdery scab control. Commercial Grower 52:16–18
- FAO (2008) Available from: [www.potato2008.org](http://www.potato2008.org). Accessed on 19 Nov 2009
- Ferreira EPdB, Dusi AN, Xavier GR, Rumjanek NG (2008) Rhizosphere bacterial communities of potato cultivars evaluated through PCR-DGGE profiles. Pesqui Agropecuaria Bras 43:605–612. doi:10.1590/s0100-204x2008000500008
- Ficke W, Naumann K, Skadow K, Muller HJ, Zielke R (1973) Longevity of *Pectobacterium carotovorum* var. *atrosepticum* (van Hall) Dowson on seed material and in the soil. Arch Phytopathologie Pflanzenschutz 9:281–293
- Fiers M, Heraud C, Gautheron N, Chatot C, Le Hinrat Y, Bouchech-Mechiche K, Steinberg C. Genetic diversity of *Rhizoctonia solani* associated with potato. Mycologia (in press)
- Finetti SM (1990) Histopathological changes induced by *Nacobbus aberrans* in resistant and susceptible potato roots. Rev Nematoologie 13:155–160
- Firman DM, Allen EJ (1995) Transmission of *Helminthosporium solani* from potato seed tubers and effects of soil conditions, seed inoculum and seed physiology on silver scurf disease. J Agric Sci 124:219–234
- Flores-Gonzalez R, Velasco I, Montes F (2008) Detection and characterization of *Streptomyces* causing potato common scab in Western Europe. Plant Pathol 57:162–169. doi:10.1111/j.1365-3059.2007.01734.x
- FNPPPT, GNIS (2000) Fiches descriptives des maladies et ravageurs de la pomme de terre. Ad Hoc, Paris
- Fox RA, Dashwood EP (1979) Some effects of cultural practices for the control of potato gangrene. Bulletin, Scottish Horticultural Research Institute Association, 4–9
- Fox RA, Dashwood EP, Wilson HM (1978) Biology of potato gangrene. Scottish Horticultural Research Institute, 24th Annual Report for the year 1977, pp 68–70
- France RA, Brodie BB (1995) Differentiation of 2 New York isolates of *Pratylenchus penetrans* based on their reaction on potato. J Nematology 27:339–345
- Franco CY, Stefanova NM, Coronado Izquierdo MF (2007) Pathogenicity and maceration ability of *Pectobacterium carotovorum* and *Dickeya chrysanthemi* isolates in potato (*Solanum tuberosum* L.). Fitosanidad 11:15–18
- Franco J, Bendezu E (1985) Study of the complex *Verticillium dahliae* Kleb. and *Globodera pallida* Stone and its effect on the behaviour of some Peruvian potato cultivars. Fitopatología 20:21–27
- Franco J, Montecinos R, Ortuno N (1992) Management strategies of *Nacobbus aberrans*, Nematology from molecule to ecosystem: Proceedings Second International Nematology Congress, 11–17 August 1990, Veldhoven, the Netherlands, pp 240–248
- Garibaldi A, Gilardi G, Gullino ML (2006) First report of southern blight incited by *Sclerotium rolfsii* on potato (*Solanum tuberosum*) in northern Italy. Plant Dis 90:1114–1114. doi:10.1094/pd-90-1114
- Geary B, Johnson DA (2006) Relationship between silver scurf levels on seed and progeny tubers from successive generations of potato seed. Am J Potato Res 83:447–453
- Geary B, Johnson DA, Hamm PB, James S, Rykbost KA (2007) Potato silver scurf affected by tuber seed treatments and locations, and occurrence of fungicide resistant isolates of *Helminthosporium solani*. Plant Dis 91:315–320. doi:10.1094/pdis-91-3-0315
- Giebel J, DopieraLa U (2004) Pathogenesis of potato gangrene caused by *Phoma exigua* var. *foveata*: II. Activities of some hydrolases and dehydrogenases. J Phytopathol 152:399–403
- Gilchrist E, Jaramillo VS, Reynaldi S (2009) Effect on the powdery scab of four isolates of the fungus *Trichoderma asperellum* in three types of soils. Revista-Facultad Nacional de Agronomía Medellin 62:4783–4792
- Gilligan CA, Simons SA, Hide GA (1996) Inoculum density and spatial pattern of *Rhizoctonia solani* in field plots of *Solanum tuberosum*: effects of cropping frequency. Plant Pathol 45:232–244
- Gindrat D (1984) Storage rot of potato. II. Lesions development in relation to the parasitic species and temperature. Control measures. Revue Suisse d'Agriculture 16:313–318
- Glais-Varlet I, Bouchech-Mechiche K, Andrivon D (2004) Growth in vitro and infectivity of *Colletotrichum coccodes* on potato tubers at different temperatures. Plant Pathol 53:398–404
- Gould M, Nelson L, Waterer D, Hynes R (2008) Biocontrol of *Fusarium sambucinum*, dry rot of potato, by *Serratia plymuthica*

- 5–6. *Biocontrol Sci Technol* 18:1005–1016. doi:[10.1080/09583150802478189](https://doi.org/10.1080/09583150802478189)
- Graaf Pvd, Lees AK, Cullen DW, Duncan JM (2003) Detection and quantification of *Spongopora subterranea* in soil, water and plant tissue samples using real-time PCR. *Eur J Plant Pathol* 109:589–597. doi:[10.1023/a:1024764432164](https://doi.org/10.1023/a:1024764432164)
- Graaf Pvd, Lees AK, Wale SJ, Duncan JM (2005) Effect of soil inoculum level and environmental factors on potato powdery scab caused by *Spongopora subterranea*. *Plant Pathol* 54:22–28. doi:[10.1111/j.1365-3059.2005.01111.x](https://doi.org/10.1111/j.1365-3059.2005.01111.x)
- Grosch R, Faltin F, Lottmann J, Kofoet A, Berg G (2005) Effectiveness of 3 antagonistic bacterial isolates to control *Rhizoctonia solani* Kuhn on lettuce and potato. *Can J Microbiol* 51:345–353. doi:[10.1139/w05-002](https://doi.org/10.1139/w05-002)
- Grosch R, Scherwinski K, Lottmann J, Berg G (2006) Fungal antagonists of the plant pathogen *Rhizoctonia solani*: selection, control efficacy and influence on the indigenous microbial community. *Mycol Res* 110:1464–1474. doi:[10.1016/j.mycres.2006.09.014](https://doi.org/10.1016/j.mycres.2006.09.014)
- Grousset F, Smith IM (1998) EPPO certification scheme for seed potatoes. *Bull OEPP* 28:561–567
- Gudmestad NC, Mallik I, Pasche JS, Anderson NR, Kinzer K (2009) A real-time PCR assay for the detection of *Clavibacter michiganensis* subsp *sepedonicus* based on the cellulase A gene sequence. *Plant Dis* 93:649–659. doi:[10.1094/pdis-93-6-0649](https://doi.org/10.1094/pdis-93-6-0649)
- Gudmestad NC, Taylor RJ, Pasche JS (2007) Management of soilborne diseases of potato. *Australas Plant Pathol* 36:109–115
- Gupta CP, Sharma A, Dubey RC, Maheshwari DK (1999) *Pseudomonas aeruginosa* (GRC(1)) as a strong antagonist of *Macrophomina phaseolina* and *Fusarium oxysporum*. *Cytobios* 99:183–189
- Gupta PP, Shekhar K, Yadav BD (2007) Effect of soil temperature and moisture levels on root rot of clusterbean. *J Arid Legumes* 4:95–99
- Gvozdeva EL, Volotskaya AV, Sof'in AV, Kudryavtseva NN, Revina TA, Valueva TA (2006) Interaction of proteinases secreted by the fungal plant pathogen *Rhizoctonia solani* with natural proteinase inhibitors produced by plants. *Appl Biochem Microbiol* 42:502–507. doi:[10.1134/s0003683806050103](https://doi.org/10.1134/s0003683806050103)
- Hafez SL, Sundararaj P (2006) Efficacy of fosthiazate for the control of *Paratrichodorus* spp. and *Meloidogyne chitwoodi* on potato. *Int J Nematology* 16:157–160
- Hague NGM, Damadzadeh M, Garabedian SK, Radwan KH (1983) Methods of assessing the efficacy of non-volatile nematicides. *Pesticide Sci* 14:587–595
- Hampson MC (1985) Pathogenesis of *Synchytrium endobioticum*. 5. Wart disease suppression in potato in soils amended with urea and or ammonium-nitrate in relation to soil pH. *Plant Soil* 87:241–250
- Hampson MC, Coombes JW (1989) Pathogenesis of *Synchytrium endobioticum* 7. Earthworms as vector of wart disease of potato. *Plant Soil* 116:147–150
- Hampson MC, Coombes JW (1995) Reduction of potato wart disease with crushed crabshell—suppression or eradication. *Can J Plant Pathol Rev Can De Phytopathologie* 17:69–74
- Hampson MC, Coombes JW (1997) Pathogenesis of *Synchytrium endobioticum*. 9. Effect of irrigation regimes and soil mixes on disease incidence with pathotype 2. *Can J Plant Pathol Rev Can De Phytopathologie* 19:47–51
- Hampson MC, Coombes JW, McRae KB (1994) Pathogenesis of *Synchytrium endobioticum*. 8. Effect of temperature and resting spore density (pathotype 2) on incidence of potato wart disease. *Can J Plant Pathol Rev Can De Phytopathologie* 16:195–198
- Harikrishnan R, del Rio LE (2006) Influence of temperature, relative humidity, ascospore concentration, and length of drying of colonized dry bean flowers on white mold development. *Plant Dis* 90:946–950. doi:[10.1094/pd-06-0946](https://doi.org/10.1094/pd-06-0946)
- Harrison JG (1997) Powdery scab disease of potato—a review. *Plant Pathol* 46:1–25
- Heath WL, Haydock PPJ, Wilcox A, Evans K (2000) Monitoring the presence and distribution of potato cyst nematodes: the potential use of light reflected from the crop. *Asp Appl Biol* 56:127–130
- Heilmann LJ, Nitzan N, Johnson DA, Pasche JS, Doekkott C, Gudmestad NC (2006) Genetic variability in the potato pathogen *Colletotrichum coccodes* as determined by amplified fragment length polymorphism and vegetative compatibility group analyses. *Phytopathology* 96:1097–1107. doi:[10.1094/phyto-96-1097](https://doi.org/10.1094/phyto-96-1097)
- Helias V (2008) *Pectobacterium* spp. and *Dickeya* spp. on potato: a new nomenclature for *Erwinia* spp., symptoms, epidemiology and disease prevention. *Cahiers Agricultures* 17:349–354. doi:[10.1684/agr.2008.0216](https://doi.org/10.1684/agr.2008.0216)
- Helias V, Andrivon D, Jouan B (2000) Internal colonization pathways of potato plants by *Erwinia carotovora* ssp *atroseptica*. *Plant Pathol* 49:33–42
- Henriksen CB, Molgaard JP, Rasmussen J (2007) The effect of autumn ridging and inter-row subsoiling on potato tuber yield and quality on a sandy soil in Denmark. *Soil Tillage Res* 93:309–315. doi:[10.1016/j.still.2006.05.003](https://doi.org/10.1016/j.still.2006.05.003)
- Hide GA (1994) Effects of wounding fungicide-treated potato seed tubers on silver scurf disease on daughter tubers at harvest. *Potato Res* 37:287–290
- Hide GA, Cayley GR (1987) Effects of delaying fungicide treatment and of curing and chloropropham on the incidence of skin spot on stored potato tubers. *Ann Appl Biol* 110:617–627
- Hide GA, Firmager JP (1989) Effects of soil temperature and moisture on stem canker (*Rhizoctonia solani*) disease of potatoes. *Potato Res* 32:75–80
- Hide GA, Read PJ (1991) Effects of rotation length, fungicide treatment of seed tubers and nematicide on diseases and the quality of potato tubers. *Ann Appl Biol* 119:77–87
- Hide GA, Welham SJ, Read PJ, Ainsley AE (1995) Influence of planting seed tubers with gangrene (*Phoma foveata*) and of neighboring healthy, diseased and missing plants on the yield and size of potatoes. *J Agric Sci* 125:51–60
- Hiemstra JA, Rataj-Guranowska M (2003) Vegetative compatibility groups in *Verticillium dahliae* isolates from the Netherlands as compared to VCG diversity in Europe and in the USA. *Eur J Plant Pathol* 109:827–839
- Hims MJ, Preece TF (1975) *Spongopora subterranea* f.sp. *subterranea*. IMI Descriptions of Fungi and Bacteria, Sheet 477
- Hlaoua W, Raouani NH (2007) Effect of *Meloidogyne incognita* on the potato crop. *Nematologia Mediterr* 35:213–220
- Holgado R, Skau KAO, Magnusson C (2009) Field damage in potato by lesion nematode *Pratylenchus penetrans*, its association with tuber symptoms and its survival in storage. *Nematologia Mediterr* 37:25–29
- Höper H, Alabouvette C (1996) Importance of physical and chemical soil properties in the suppressiveness of soils to plant diseases. *Eur J Soil Biol* 32:41–58
- Hsu ST (1991) Ecology and control of *Pseudomonas solanacearum* in Taiwan. *Plant Prot Bull Taichung* 33:72–79
- Hughes J (2008) Molecular diagnostics and their importance for *Rhizoctonia solani* control in potatoes. *Outlooks Pest Manage* 19:123–126. doi:[10.1564/19jun08](https://doi.org/10.1564/19jun08)
- Hukkanen A, Karjalainen R, Nielsen SL, van der Wolf JM (2005) Epidemiology of *Clavibacter michiganensis* subsp *sepedonicus* in potato under European conditions: population development and yield reduction. *Z Fur Pflanzenkrankheiten Und Pflanzenschutz J Plant Dis Prot* 112:88–97
- Ilyashenka D, Ivaniuk V (2008) Potato stem nematode in Belarus. *Zemdirbyste Agric* 95:74–81
- Ingham RE, Hamm PB, Baune M, David NL, Wade NM (2007) Control of *Meloidogyne chitwoodi* in potato with shank-

- injected metam sodium and other nematicides. *J Nematology* 39:161–168
- INRA, Cemagref (2005) Réduire l'utilisation des pesticides et en limiter les impacts environnementaux. Ministère de l'agriculture et de la pêche, Ministère de l'énergie et du développement durable, pp 68
- Inserra RN, Chitambar JJ, Chitwood DJ, Handoo Z (2005) The Potato Pathotype of the False-Root Knot Nematode, *Nacobbus aberrans*. Working Group of the SON Exotic Nematode Plant Pest List, p 10
- Inserra RN, McSorley R, Greco N, Weingartner DP (1996) Potato cyst nematodes, a potential menace to the potato industry of Florida. Soil and Crop Science Society of Florida Proceedings 55:1–5
- IPC (1978) Annual report for 1977 International Potato Center, Lima, Peru, pp 156
- Iriarte L, Franco J, Ortuno N (1999) Influence of time of sowing on potato yield and population density of *Nacobbus aberrans*. *Fitopatología* 34:77–82
- Jaggi W, Oberholzer H, Winiger FA (1991) Effect of harvesting and storage conditions on infection of potatoes by *Erwinia carotovora*. *Kartoffelbau* 42:474–480
- Jansky SH, Rouse DI (2003) Multiple disease resistance in interspecific hybrids of potato. *Plant Dis* 87:266–272
- Janvier C, Villeneuve F, Alabouvette C, Edel-Hermann V, Mateille T, Steinberg C (2007) Soil health through soil disease suppression: which strategy from descriptors to indicators? *Soil Biol Biochem* 39:1–23
- Jauhari RK, Lal M (2001) Effect of soil moisture on the population of migratory nematode *Pratylenchus penetrans* (Cobb 1917) (Nematoda: Hoplolamidae) on tea plantations in Doon Valley. *J Parasitol Appl Anim Biol* 10:1–8
- Jensen HJ (1978) Interrelations of nematodes and other organisms in disease complexes, International Potato Center. Report of the 2nd planning conference on the developments in the control of nematode pests of potatoes, Lima, Peru, 13–17 November 1978, pp 154–160
- Joaquim TR, Rowe RC (1991) Vegetative compatibility and virulence of strains of *Verticillium dahliae* from soil and potato plants. *Phytopathology* 81:552–558
- Johnson DA, Atallah ZK (2006) Timing fungicide applications for managing *Sclerotinia* stem rot of potato. *Plant Dis* 90:755–758. doi:10.1094/pd-90-0755
- Johnston HW, Celetti MJ, Kimpinski J, Platt HW (1994) Fungal pathogens and *Pratylenchus penetrans* associated with preceding crops of clovers, winter-wheat, and annual ryegrass and their influence on succeeding potato crops on Prince Edward Island. *Am Potato J* 71:797–808
- Jong-Tae K, In-Hee P, Hyang-Burm L, Young-II H, Seung-Hun Y (2001) Identification of *Verticillium dahliae* and *V. albo-atrum* causing wilt of tomato in Korea. *Plant Pathol J* 17:222–226
- Kamensky M, Ovadis M, Chet I, Chernin L (2002) Biocontrol of *Botrytis cinerea* and *Sclerotinia sclerotiorum* in the greenhouse by a *Serratia plymuthica* strain with multiple mechanisms of antifungal activity. *Bull OILB SROP* 25:229–232
- Kandji ST, Ogor C, Albrecht A (2001) Diversity of plant-parasitic nematodes and their relationships with some soil physico-chemical characteristics in improved fallows in western Kenya. *Appl Soil Ecol* 18:143–157
- Kang HC, Park YH, Go SJ (2003) Growth inhibition of a phytopathogenic fungus, *Colletotrichum* species by acetic acid. *Microbiol Res* 158:321–326
- Karwasra SS, Parashar RD (1998) Chemical control of incipient infection, weight loss and sprouting of potato tubers during storage. *Plant Dis Res* 13:49–51
- Keshwali RL, Khare UK, Singh RP (2000) Effect of physical properties of soil on wilt incidence and population of *Ralstonia solanacearum*. *Ann Plant Prot Sci* 8:40–43
- Khomyakov MT, Kostin NP (1981) The effects of potato saturation in rotations and of dosage rates of fertilizers on the development of diseases of tubers in storage. *Trudy Vsesoyuznogo Nauchno-Issledovatel'skogo Instituta Zashchity Rastenii*, 66–70
- Kim-Lee HY, Moon JS, Hong YJ, Kim MS, Cho HM (2005) Bacterial wilt resistance in the progenies of the fusion hybrids between haploid of potato and *Solanum commersonii*. *Am J Potato Res* 82:129–137
- Kim MH, Park SC, Kim JY, Lee SY, Lim HT, Cheong H, Hahn KS, Park Y (2006) Purification and characterization of a heat-stable serine protease inhibitor from the tubers of new potato variety "Golden Valley". *Biochem Biophys Res Commun* 346:681–686. doi:10.1016/j.bbrc.2006.05.186
- Kimpinski J, Arsenault WJ, Sturz AV (2001) Differential effect of nematicide treatments on tuber yields in early- and late-maturing potato cultivars. *Plant Pathol* 50:509–514
- Kishore V, Shekhawat GS, Sunaina V (1996) Cultural practices to reduce *Pseudomonas solanacearum* in the infested soil. *J Indian Potato Assoc* 23:130–133
- Klikocka H (2001) The influence of soil tillage systems and crop cultivation methods on potato tubers infection with *Rhizoctonia solani* Kuhn. *Buletyn Instytutu Hodowli i Aklimatyzacji Roslin*, 243–247
- Kong Q, Yuan S, Wang Y, Zhu Y (2006) Study on biological characteristics of *Fusarium solani* on *Vanilla planifolia* root. *J Mt Agric Biol* 25:506–509
- Krechel A, Faupel A, Hallmann J, Ulrich A, Berg G (2002) Potato-associated bacteria and their antagonistic potential towards plant-pathogenic fungi and the plant-parasitic nematode *Meloidogyne incognita* (Kofoid & White) Chitwood. *Can J Microbiol* 48:772–786. doi:10.1139/w02-071
- Kumar S, Vadivelu S (1996) Influence of soil type on the interaction between root-knot nematode, reniform nematode and root-rot in eggplant. *Pest Manage Horticultural Ecosyst* 2:29–35
- Kumar SM, Khare MN (1990) Studies on the antagonistic relationship of soybean sporesphere microflora with *Rhizoctonia bataticola* and *Sclerotium rolfsii*. *J Biol Control* 4:72–74
- Kuninaga S, Carling DE, Takeuchi T, Yokosawa R (2000) Comparison of rDNA-ITS sequences between potato and tobacco strains in *Rhizoctonia solani* AG-3. *J Gen Plant Pathol* 66:2–11
- Lakra BS (2000) Effect of planting time and curing period on black scurf development and yield of potato. *Haryana J Horticultural Sci* 29:121–122
- Lambert DH, Manzer FE (1991) Relationship of calcium to potato scab. *Phytopathology* 81:632–636
- Lambert DH, Reeves AF, Goth RW, Grounds GS, Gigge EA (2006) Relative susceptibility of potato varieties to *Streptomyces scabiei* and *S. acidiscabies*. *Am J Potato Res* 83:67–70
- Larkin RP (2008) Relative effects of biological amendments and crop rotations on soil microbial communities and soilborne diseases of potato. *Soil Biol Biochem* 40:1341–1351. doi:10.1016/j.soilbio.2007.03.005
- Latour X, Faure D, Diallo S, Cirou A, Smadja B, Dessaux Y, Orange N (2008) Control of bacterial diseases of potato caused by *Pectobacterium* spp. (*Erwinia carotovora*). *Cahiers Agricultures* 17:355–360. doi:10.1684/agr.2008.0210
- Laurila J, Metzler MC, Ishimaru CA, Rokka VM (2003) Infection of plant material derived from *Solanum acaule* with *Clavibacter michiganensis* ssp *sepedonicus*: temperature as a determining factor in immunity of *S. acaule* to bacterial ring rot. *Plant Pathol* 52:496–504
- Lazarovits G, Hill J, Patterson G, Conn KL, Crump NS (2007) Edaphic soil levels of mineral nutrients, pH, organic matter, and cationic exchange capacity in the geocaulosphere associated with potato common scab. *Phytopathology* 97:1071–1082. doi:10.1094/phyto-97-9-1071

- Lazarovits G, Tenuta M, Conn KL (2001) Organic amendments as a disease control strategy for soilborne diseases of high-value agricultural crops. *Australas Plant Pathol* 30:111–117
- Lazy GH, Lukezic FL (2003) Pathogenic prokaryotes. In: Trigiano RN, Windham MT, Windham AS (eds) *Plant pathology*. CRC Press, New York
- Lees AK (2003) Black dot (*Colletotrichum coccodes*): an increasingly important disease of potato. *Plant Pathol* 52:3–12
- Lees AK, Cullen DW, Sullivan L, Nicolson MJ (2002) Development of conventional and quantitative real-time PCR assays for the detection and identification of *Rhizoctonia solani* AG-3 in potato and soil. *Plant Pathol* 51:293–302
- Lees AK, Sullivan L, Cullen DW (2009) A quantitative polymerase chain reaction assay for the detection of *Polyscytalum pustulans*, the cause of skin spot disease of potato. *J Phytopathol* 157:154–158. doi:[10.1111/j.1439-0434.2008.01459.x](https://doi.org/10.1111/j.1439-0434.2008.01459.x)
- Lehtonen MJ, Somervuo P, Valkonen JPT (2008) Infection with *Rhizoctonia solani* induces defense genes and systemic resistance in potato sprouts grown without light. *Phytopathology* 98:1190–1198. doi:[10.1094/phyto-98-11-1190](https://doi.org/10.1094/phyto-98-11-1190)
- Lennard JH (1980) Factors affecting the development of silver scurf (*Helminthosporium solani*) on potato-tubers. *Plant Pathol* 29:87–92
- Lerat S, Babana AH, El OM, El HA, Daayf F, Beaudoin N, Bouarab K, Beaulieu C (2009) *Streptomyces scabiei* and its toxin thaxtomin A induce scopoletin biosynthesis in tobacco and *Arabidopsis thaliana*. *Plant Cell Report* 28:1895–1903. doi:[10.1007/s00299-009-0792-1](https://doi.org/10.1007/s00299-009-0792-1)
- Levine A, Tenhaken R, Dixon R, Lamb C (1994) H<sub>2</sub>O<sub>2</sub> from the oxidative burst orchestrates the plant hypersensitive disease resistance response. *Cell* 79:583–593
- Lewocz W (1992) Activity of some fungicides as limiting factors of black leg and soft rot of potato tubers. *Biuletyn Instytutu Ziemniaka*, pp 89–96
- Li W, Roberts DP, Dery PD, Meyer SLF, Lohrke S, Lumsden RD, Hebbar KP (2002) Broad spectrum anti-biotic activity and disease suppression by the potential biocontrol agent *Burkholderia ambifaria* BC-F. *Crop Prot* 21:129–135
- Liu H, Coulthurst SJ, Pritchard L, Hedley PE, Ravensdale M, Humphris S, Burr T, Takle G, Brurberg MB, Birch PRJ, Salmon GPC, Toth IK (2008) Quorum sensing coordinates brute force and stealth modes of infection in the plant pathogen *Pectobacterium atrosepticum*. *Plos Pathogens*. doi:[10.1371/journal.ppat.1000093](https://doi.org/10.1371/journal.ppat.1000093)
- Lo C, Wang K (2000) Factors affecting pycnidial production and pycnidiospore germination of *Phoma wasabiae*, the causal agent of wasabi. *Plant Pathol Bull* 9:99–106
- Logan C, O'Neill R, McGrane P, Little G (1987) Methods for detection of the blackleg ring rot and gangrene pathogens in potato nuclear stock mother tubers and plantlets. *Bull OEPP* 17:17–24
- Loria R, Bignell DRD, Moll S, Huguet-Tapia JC, Joshi MV, Johnson EG, Seipke RF, Gibson DM (2008) Thaxtomin biosynthesis: the path to plant pathogenicity in the genus *Streptomyces*. *Antonie Van Leeuwenhoek International J Gen Mol Micro* 94:3–10. doi:[10.1007/s10482-008-9240-4](https://doi.org/10.1007/s10482-008-9240-4)
- Loria R, Bkhalid RA, Fry BA, King RR (1997) Plant pathogenicity in the genus *Streptomyces*. *Plant Disease* 81:836–846
- Lucas JA, Pitt D (1974) Immunochemical identification of a new molecular form of acid-phosphatase of host origin arising during infection of potato tubers by *Phytophthora erythrophloëtica* Pethyb. *J Gen Microbiol* 84:311–320
- Lucke W (1975) The effect of additional sprinkler irrigation and its combination with nitrogen supply on the occurrence of potato blackleg (*Pectobacterium carotovorum* (Jones) Waldee var. *atrosepticum* (van Hall) Dowson). *Arch Phytopathologie Pflanzenschutz* 11:213–223
- Lui LH (2003) Models to predict potato tuber infection by *Pythium ultimum* from duration of wetness and temperature, and leak-lesion expansion from storage duration and temperature. *Post-harvest Biol Technol* 27:313–322
- Lui LH, Kushalappa AC (2002) Response surface models to predict potato tuber infection by *Fusarium sambucinum* from duration of wetness and temperature, and dry rot lesion expansion from storage time and temperature. *Int J Food Microbiol* 76:19–25
- Lulai EC (2001) In: Stevenson WR, Loria R, Franc GD, Weingartner DP (eds) *Compendium of potato diseases*. APS Press, St. Paul, p 3
- Lutomirska B, Szutkowska M (2005) Influence of the soil type and date of irrigation on tuber infection with *Rhizoctonia solani* (Kuhn). *Prog Plant Prot* 45:865–868
- Lyon GD (1989) The biochemical basis of resistance of potatoes to soft rot *Erwinia* spp.: a review. *Plant Pathology* 38:313–339
- Madalageri BB, Dharmatti PR, Hosmani RM, Srikant K (1991) Varietal resistance of potato to wilt caused by *Sclerotium rolfsii* Sacc. *Curr Res Univ Agric Sci Bangalore* 20:26–27
- Mahmoud DAR, Mahmoud AA, Gomaa AM (2008) Antagonistic activities of potato associated bacteria via their production of hydrolytic enzymes with special reference to pectinases. *Res J Agric Biol Sci* 4:575–584
- Mahmoud SM (2007) Management of brown rot disease of potato. *Arab Universities J Agric Sci* 15:457–463
- Main G, Franco J, Ortuno N (2001) Disinfection of potato seed infected with *Nacobbus aberrans* using mild nematicides. *Manejo Integrado de Plagas*, 52–57
- Malakouti MJ (2008) The effect of micronutrients in ensuring efficient use of macronutrients. *Turk J Agric For* 32:215–220
- Manici LM, Caputo F (2009) Fungal community diversity and soil health in intensive potato cropping systems of the east Po valley, northern Italy. *Ann Appl Biol* 155:245–258. doi:[10.1111/j.1744-7348.2009.00335.x](https://doi.org/10.1111/j.1744-7348.2009.00335.x)
- Marcinkowska J, Roze-Kauzny I, Kauzny W (2005) Pathogenicity of some *Phoma exigua* var. *exigua* isolates. *Phytopathol Pol* 38; 35–44
- Marshall KC (1975) Clay mineralogy in relation to survival of soil bacteria. *Annu Rev Phytopathol* 13:357–373
- Martinez C, Rioux D, Tweddell RJ (2004) Ultrastructure of the infection process of potato tuber by *Helminthosporium solani*, causal agent of potato silver scurf. *Mycol Res* 108:828–836. doi:[10.1017/s0953756204000589](https://doi.org/10.1017/s0953756204000589)
- Mashela P, McSorley R, Duncan LW, Dunn RA (1991) Correlation of *Belonolaimus longicaudatus*, *Hoplolaimus galeatus*, and soil texture with yield of alyceclover (*Alysicarpus* spp.). *Nematropica* 21:177–184
- McDonald JE, White GP, Cote MJ (2000) Differentiation of *Phoma foveata* from *P. exigua* using a RAPD generated PCR-RFLP marker. *Eur J Plant Pathol* 106:67–75
- McDonald JG (1995) Disease control through crop certification: herbaceous crops. *Can J Plant Pathol Rev Can De Phytopathologie* 17:267–273
- Medina LFC, Stefani V, Brandelli A (2004) Use of 1,4-naphthoquinones for control of *Erwinia carotovora*. *Can J Microbiol* 50:951–956. doi:[10.1139/w04-088](https://doi.org/10.1139/w04-088)
- Mehta SK, Tripathi NN, Rakesh K, Chhabra ML (2006) Influence of light and temperature on induction of pycnidia and germination of pycnidiospores. *Ann Biol* 22:31–34
- Melakeberhan H, Dey J, Baligar VC, Carter TE (2004) Effect of soil pH on the pathogenesis of *Heterodera glycines* and *Meloidogyne incognita* on Glycine max genotypes. *Nematology* 6:585–592
- Melakeberhan H, Mennan S, Chen S, Darby B, Dudek T (2007) Integrated approaches to understanding and managing *Meloidogyne hapla* populations' parasitic variability. *Crop Prot* 26:894–902. doi:[10.1016/j.cropro.2006.08.008](https://doi.org/10.1016/j.cropro.2006.08.008)

- Meredith DS, Logan C, Copeland RB, Meijers CP, Henriksen JB, McIntosh AH (1975) Session 8C. Control of fungal diseases of potatoes. Proceedings of the Eighth British Insecticide and Fungicide Conference, Brighton, England, 17–20 November, 1975. Vols. 1 and 2, Research Reports, pp 581–623
- Merz U (2005) Improved immunological detection of *Spongopora subterranea*. Eur J Plant Pathol 111:371–379
- Merz U, Falloon RE (2009) Review: powdery scab of potato—increased knowledge of pathogen biology and disease epidemiology for effective disease management. Potato Res 52:17–37. doi:[10.1007/s11540-008-9105-2](https://doi.org/10.1007/s11540-008-9105-2)
- Messiha NAS, Bruggen AHCv, Diepeningen ADv, Vos OJd, Termorshuizen AJ, Tjou-Tam-Sin NNA, Janse JD (2007) Potato brown rot incidence and severity under different management and amendment regimes in different soil types. Eur J Plant Pathol 119:367–381. doi:[10.1007/s10658-007-9167-z](https://doi.org/10.1007/s10658-007-9167-z)
- Michel VV, Mew TW (1998) Effect of a soil amendment on the survival of *Ralstonia solanacearum* in different soils. Phytopathology 88:300–305
- Milic V, Bogdanovic M, Uric M, Kovacevic D, Crnogorac M (2006) Influence of mineral nutrition and additional space upon yield of potato. Agroznanje Agroknowledge 7:67–74
- Milosevic D, Alovic I (2006) Some most important potato diseases and their control. Radovi Poljoprivrednog Fakulteta Univerziteta u Sarajevu (Works of the Faculty of Agriculture University of Sarajevo) 51:103–120
- Milosevic D, Alovic I, Civic H (2005) Effect of some soil reclamation measures on the disease severity in potato tubers caused by common scab (*Streptomyces scabies* Thaxt.). Radovi Poljoprivrednog Fakulteta Univerziteta u Sarajevu (Works of the Faculty of Agriculture University of Sarajevo) 50, pp 33–42
- Minnis ST, Haydock PPJ, Evans K (2004) Control of potato cyst nematodes and economic benefits of application of 1,3-dichloropropene and granular nematicides. Ann Appl Biol 145:145–156
- Mizuno N, Nizamidin K, Nanzyo M, Yoshida H, Amano Y (2003) Judging conducive soils from clay mineralogical properties and soil chemical method to suppress potato common scab. Soil Microorganisms 57:97–103
- Moffett ML, Wood BA (1984) Survival of *Corynebacterium michiganense* subsp. *michiganense* within host debris in soil. Australas Plant Pathol 13:1–3
- Mohsin M, Mian IH, Zahid MI, Ali M (1989) Inoculum level of *Meloidogyne incognita* on severity of root knot and growth of jute and potato. Bangladesh J Plant Pathol 5:13–17
- Mol L, Scholte K (1995) Formation of microsclerotia of *Verticillium dahliae* Kleb on various plant parts of 2 potato cultivars. Potato Res 38:143–150
- Molendijk LPG (1999) Control strategy for nematodes works ([example of] Vredespeel). PAV-Bulletin Akkerbouw, 4–8
- Mordue JEM (1988) *Thecaphora solani*. IMI Descriptions of Fungi and Bacteria, Sheet 966
- Moxnes JF, Hausken K (2007) The population dynamics of potato cyst nematodes. Ecol Model 207:339–348. doi:[10.1016/j.ecolmodel.2007.06.020](https://doi.org/10.1016/j.ecolmodel.2007.06.020)
- Mugniéry D (2007) The canon of potato science: 15. root-knot nematodes. Potato Res 50:263–265. doi:[10.1007/s11540-008-9048-7](https://doi.org/10.1007/s11540-008-9048-7)
- Mugniéry D, Phillips MS (2007) The nematode parasites of potato. In: Vreugdenhil D, Bradshaw JE (eds) Potato biology and biotechnology. Elsevier, New York
- Muhammad Z (1996) The effects of storage conditions on subsequent hatching of *Globodera pallida* (Nematoda: Tylenchida). Pedobiologia 40:342–351
- Mulder A, Turkensteen LJ, Delleman J (eds) (2008) Aardappel ziektenboek: ziekten, plagen en beschadigingen. Aardappelwereld, Den Haag (Netherlands)
- Mulder A, Van der Wal AF, Velema RAJ, Roosjen JS (1997) Effects of soil characteristics on the relationship between potato cyst nematodes and yield. II. Acidity (soil pH). Potato Res 40:375–381
- Muller P, Abdel-Kader D, Kakau J, Pastrik KH, Seigner L (2004) Survival of *Ralstonia solanacearum* biovar 2 (race 3), the causal agent of potato brown rot, in soil and on several kinds of materials (microcosm). Gesunde Pflanzen 56:129–141
- Munzert M, Duben J, Langerfeld E (1977) On the influence of fungal and bacterial tuber rot pathogens on emergence diseases in the potato crop. Nachrichtenblatt Dtsch Pflanzenschutzdienstes 29:69–74
- Muthukrishnan K, Raguchander T, Arjunan G (1995) Factors affecting survival of *Macrophomina phaseolina* in soil. Indian J Pulses Res 8:156–161
- Nagesh M (1996) Relationship between the population densities of root-knot nematode, *Meloidogyne incognita* (Kofoid and White 1919) Chitwood 1949 and growth of potato, *Solanum tuberosum* L. Pest Manage Horticultural Ecosyst 2:9–14
- Nakayama T (2007) *Spongopora subterranea* soil contamination and its relationship to severity of powdery scab on potatoes. J Gen Plant Pathol 73:229–234
- Nakayama T, Merz U, Nakagawa A, Takehara T, Shimanuki T (2003) Differences in zoosporangial root infection of some potato varieties inoculated with Japanese and foreign field isolates of *Spongopora subterranea* f. sp. *subterranea*, Proceedings of the Fifth Symposium of the International Working Group on Plant Viruses with Fungal Vectors, Institute of Plant Sciences, Swiss Federal Institute of Technology, Zurich, Switzerland, 22–25 July, 2002, pp 115–118
- Naumann K, Ficke W, Muller HJ, Skadow K, Zielke R (1974) Transmission of potato black leg and tuber soft rot (*Pectobacterium carotovorum* var. *atrosepticum* (van Hall) Dowson) by soil, seed material and soil cultivation. Arch Phytopathologie Pflanzenschutz 10:301–316
- Nelson GA (1982) *Corynebacterium sepedonicum* in potato: effect of inoculum concentration on ring rot symptoms and latent infection. Can J Plant Pathol 4:129–133
- Nelson GA (1984) Survival of *Corynebacterium sepedonicum* in potato stems and on surfaces held at freezing and above-freezing temperatures. Am Potato J 62:23–28
- Nicot PC, Rouse DI (1987) Relationship between soil inoculum density of *Verticillium dahliae* and systemic colonization of potato stems in commercial fields over time. Phytopathology 77:1346–1355
- Nissinen R (2000) Gram positive phytopathogenic bacterium *Clavibacter michiganensis* subsp. *sepedonicus*: Virulence factors and interactions with plants. Ann Univ Turkuensis A II Biol Geogr Geol 140:1–53
- Nitzan N, Cummings TF, Johnson DA (2008) Disease potential of soil- and tuberborne inocula of *Colletotrichum coccodes* and black dot severity on potato. Plant Dis 92:1497–1502. doi:[10.1094/pdis-92-11-1497](https://doi.org/10.1094/pdis-92-11-1497)
- Nitzan N, Evans MA, Cummings TF, Johnson DA, Batchelor DL, Olsen C, Haynes KG, Brown CR (2009) Field resistance to potato stem colonization by the black dot pathogen *Colletotrichum coccodes*. Plant Dis 93:1116–1122. doi:[10.1094/pdis-93-11-1116](https://doi.org/10.1094/pdis-93-11-1116)
- Nitzan N, Tsror L (2003) Effect of temperature and pH on in vitro growth rate and sclerotial density of *Colletotrichum coccodes* isolates from different VCGs. Am J Potato Res 80:335–339
- Nitzan N, Tsror L, Johnson DA (2006) Vegetative compatibility groups and aggressiveness of North American isolates of *Colletotrichum coccodes*, the causal agent of potato black dot. Plant Dis 90:1287–1292. doi:[10.1094/pd-90-1287](https://doi.org/10.1094/pd-90-1287)
- Nouri S, Bahar M, Fegan M (2009) Diversity of *Ralstonia solanacearum* causing potato bacterial wilt in Iran and the first

- record of phylotype II/biovar 2 T strains outside South America. *Plant Pathol* 58:243–249. doi:10.1111/j.1365-3059.2008.01944.x
- Nunes JCS, Fontes PCR, Araujo EF, Sediyma C (2006) Potato plant growth and macronutrient uptake as affected by soil tillage and irrigation systems. *Pesqui Agropecuaria Bras* 41:1787–1792
- Nunez-Camargo MdC, Carrion G, Nunez-Sanchez AE (2003) Fungi associated with *Globodera rostochiensis* cysts in Mexico. *Int J Nematology* 13:151–161
- Ocamb CM, Hamm PB, Johnson DA (2007) Benzimidazole resistance of *Fusarium* species recovered from potatoes with dry rot from storages located in the Columbia basin of Oregon and Washington. *Am J Potato Res* 84:169–177
- Ochiai N, Powelson ML, Crowe FJ, Dick RP (2008) Green manure effects on soil quality in relation to suppression of *Verticillium* wilt of potatoes. *Biol Fertil Soils* 44:1013–1023. doi:10.1007/s00374-008-0289-z
- Ohazuri NC, Arinze AE (1992) The action of polygalacturonase enzyme of *Sclerotium rolfsii* on some tuber crop tissues. *Fitopatologia Bras* 17:393–398
- Olivieri FP, Maldonado S, Tonon CV, Casalongue CA (2004) Hydrolytic activities of *Fusarium solani* and *Fusarium solani* f. sp *eumartii* associated with the infection process of potato tubers. *J Phytopathol* 152:337–344
- Olsen NL, Kleinkopf GE, Woodell LK (2003) Efficacy of chlorine dioxide for disease control on stored potatoes. *Am J Potato Res* 80:387–395
- Okopnyi NS, Mavlyanov OM, Narbaev Z.N (1983) Mechanisms of resistance to rootknot nematodes in cotton. *Uzbekskii Biologicheskii Zhurnal* 6:45–47
- Omer MA, Johnson DA, Douhan LI, Hamm PB, Rowe RC (2008) Detection, quantification, and vegetative compatibility of *Verticillium dahliae* in potato and mint production soils in the Columbia Basin of Oregon and Washington. *Plant Dis* 92:1127–1131. doi:10.1094/pdis-92-7-1127
- Ostergaard SP, Henriksen JB (1983) Potato tuber rot after lifting at different soil temperatures. *Tidsskr Planteavl* 87:111–117
- Ouellette E, Raghavan GSV, Reeder RD, Greenhalgh R (1990) Volatile monitoring technique for disease detection in stored potatoes. *J Food Process Preserv* 14:279–300
- Ozarslandan A, Devran Z, Mutlu N, Elekciooglu IH (2009) First report of columbia root-knot nematode (*Meloidogyne chitwoodi*) in potato in Turkey. *Plant Disease*. doi:10.1094/pdis-93-3-0316c
- Pailletin G (2008) Ecophyto 2018. Ministère de l'agriculture et de la pêche, Paris, p 142
- Palomo JL, Velazquez E, Mateos PF, Garcia-Benavides P, Martinez-Molina E (2000) Rapid identification of *Clavibacter michiganensis* subspecies *sepedonicus* based on the stable low molecular weight RNA (LMW RNA) profiles. *Eur J Plant Pathol* 106:789–793
- Pandey RC, Dwivedi BK, Singh SP (2002) Effect of soil moisture and hydrogen ion concentration on population dynamics of *Meloidogyne incognita* at Allahabad. *Curr Nematology* 13:57–60
- Panique E, Kelling KA, Schulte EE, Hero DE, Stevenson WR, James RV (1997) Potassium rate and source effects on potato yield, quality, and disease interaction. *Am Potato J* 74:379–398
- Panka D, Sadowski C, Rolbiecki S (2007) Influence of micro irrigation on health status of chosen potato cultivars grown in very light soil. *Proceedings of the IIIrd Balkan Symposium on Vegetable and Potatoes*, 357–360
- Papert A, Kok CJ, van Elsas JD (2004) Physiological and DNA fingerprinting of the bacterial community of *Meloidogyne fallax* egg masses. *Soil Biol Biochem* 36:1843–1849. doi:10.1016/j.soilbio.2004.04.038
- Park EJ, Lee SD, Chung EJ, Lee MH, Um HY, Murugaiyan S, Moon BJ, Lee SW (2007) MicroTom—a model plant system to study bacterial wilt by *Ralstonia solanacearum*. *Plant Pathol* 23:239–244
- Park Y, Kang H, Been C, Choi Y, Choi Y (2002) Chemical control of potato common scab (*Streptomyces scabies*). *Korean J Horticultural Sci Technol* 20:319–324
- Pasco C, Jouan B, Andrivon D (2005) Resistance of potato genotypes to common and netted scab-causing species of *Streptomyces*. *Plant Pathol* 54:383–392. doi:10.1111/j.1365-3059.2005.01178.x
- Pelsmaeker Md, Coomans A (1987) Nematodes in potato fields and the relation to some biotic and abiotic factors. *Mededelingen van de Faculteit Landbouwwetenschappen, Rijksuniversiteit Gent* 52:561–569
- Perez EE, Weingartner DP, McSorley R (2000) Correlation between *Paratrichodorus minor* population levels and corky ringspot symptoms on potato. *Nematropica* 30:247–251
- Perez W, Gutarra L, French E (1994) *Pythium ultimum* Trow. causing watery rot in potato tubers in Peru. *Fitopatologia* 29:191–195
- Perez W, Torres H (2008) Potato smut, *Thecaphora solani*. In: Elsevier (ed) Diseases, pests and disorders of potatoes Wale, Stuart., Platt, Bud., Platt, Harold William., Cattlin, Nigel D., pp. 66–68. ISBN 0123736706, 9780123736703
- Perombelon MCM (2000) Blackleg risk potential of seed potatoes determined by quantification of tuber contamination by the causal agent and *Erwinia carotovora* subsp. *atroseptica*: a critical review. *Bulletin OEPP* 30
- Perombelon MCM, Gullingshandley J, Kelman A (1979) Population dynamics of *Erwinia carotovora* and pectolytic *Clostridium* spp. in relation to decay of potatoes. *Phytopathology* 69:167–173
- Peters JC, Lees AK, Cullen DW, Sullivan L, Stroud GP, Cunningham AC (2008a) Characterization of *Fusarium* spp. responsible for causing dry rot of potato in Great Britain. *Plant Pathol* 57:262–271
- Peters RD, Clark RJ, Coffin AD, Sturz AV, Lambert DH, Miller JS (2005) Limited genetic diversity in North American isolates of *Phytophthora erythroseptica* pathogenic to potato based on RAPD analysis. *Plant Dis* 89:380–384. doi:10.1094/pd-89-0380
- Peters RD, MacLeod C, Seifert KA, Martin RA, Hale LR, Grau CR, MacInnis S (2008b) Pathogenicity to potato tubers of *Fusarium* spp. isolated from potato, cereal and forage crops. *Am J Potato Res* 85:367–374. doi:10.1007/s12230-008-9037-z
- Peters RD, Sturz AV, Carter MR, Sanderson JB (2003) Developing disease-suppressive soils through crop rotation and tillage management practices. *Soil Tillage Res* 72:181–192
- Peters RD, Sturz AV, Carter MR, Sanderson JB (2004) Influence of crop rotation and conservation tillage practices on the severity of soil-borne potato diseases in temperate humid agriculture. *Can J Soil Sci* 84:397–402
- Philis J (1997) Effect of cadusafos and carbofuran against *Pratylenchus penetrans* and some ectoparasitic nematodes infesting potato in Cyprus. *Nematologia Mediterr* 25:169–172
- Phillips AJL (1989) Fungi associated with sclerotia of *Sclerotinia sclerotiorum* in South Africa and their effects on the pathogen. *Phytophylactica* 21:135–140
- Pitman AR, Wright PJ, Galbraith MD, Harrow SA (2008) Biochemical and genetic diversity of pectolytic enterobacteria causing soft rot disease of potatoes in New Zealand. *Australas Plant Pathol* 37:559–568. doi:10.1071/ap08056
- Platt HW, Mahuku G (2000) Detection methods for *Verticillium* species in naturally infested and inoculated soils. *Am J Potato Res* 77:271–274
- Povolny P (1995) Influence of exposure to light and cultivation system on resistance to *Phoma foiveata* and *Fusarium solani* var. *coeruleum* in potato tubers—a pilot study. *Swed J Agric Res* 25:47–50
- Prescott L, Harley JP, Klein DA, Bacq-Calberg C-M, Dusart J (2003) La classe des Clostridia. In: Prescott L, Harley JP, Klein DA, Bacq-Calberg C-M, Dusart J (eds) *Microbiologie*, De Boeck, pp 523–525
- Prithiviraj B, Sarma BK, Srivastava JS, Singh UP (2000) Effect of  $\text{Ca}^{2+}$ ,  $\text{Mg}^{2+}$ , light, temperature and pH on aethelial stage

- formation in *Sclerotium rolfsii* Sacc. Z Fur Pflanzenkrankheiten Und Pflanzenschutz J Plant Dis Prot 107:274–278
- Pudasaini MP, Viaene N, Moens M (2007) The influence of host and temperature on the vertical migration of *Pratylenchus penetrans*. Nematology 9:437–447
- Pylypenko LA, Phillips MS, Blok VC (2008) Characterisation of two Ukrainian populations of *Globodera pallida* in terms of their virulence and mtDNA, and the biological assessment of a new resistant cultivar Vales Everest. Nematology 10:585–590
- Qian Z, Ma C, Gu Z, Qian G, Zhang F, He C, Gu A (1996) Biology, identification and control of root knot nematode *Meloidogyne hapla* on *Aconitum carmichaeli*. Acta Agriculturae Shanghai 12:81–86
- Qu XS, Kavanagh JA, Egan D, Christ BJ (2006) Detection and quantification of *Spongopora subterranea* f. sp *subterranea* by PCR in host tissue and naturally infested soils. Am J Potato Res 83:21–30
- Rahman ML, Hossain MM, Ashrafuzzaman M, Islam T (1996) Effect of inoculum levels of *Rhizoctonia solani* on the incidence of black scurf disease of potato. Bangladesh J Plant Pathol 12:21–22
- Rangaswami G, Mahadevan A (2004) Disease of crop plants in India. Fourth Edition, ISBN: 978-81-203-1247-0, p 548
- Raviv M (2008) The use of compost in growing media as suppressive agent against soil-borne diseases. Proceedings of the International Symposium on Growing Media, 39–49
- Ray H, Hammerschmidt R (1998) Responses of potato tuber to infection by *Fusarium sambucinum*. Physiol Mol Plant Pathol 53:81–92
- Recep K, Fikrettin S, Erkol D, Cafer E (2009) Biological control of the potato dry rot caused by *Fusarium* species using PGPR strains. Biol Control 50:194–198. doi:10.1016/j.bioccontrol.2009.04.004
- Rehman S, Butterbach P, Popejus H, Overmars H, Davis EL, Jones JT, Goverse A, Bakker J, Smart G (2009) Identification and characterization of the most abundant cellulases in stylet secretions from *Globodera rostochiensis*. Phytopathology 99:194–202. doi:10.1094/phyto-98-2-0194
- Reid A (2009) PCR detection of potato cyst nematode. Meth Mol Biol. doi:10.1007/978-1-59745-062-1\_22
- Repsiene R, Mineikiene EV (2006) The influence of meteorological conditions and different agricultural systems on the spreading of potato cv. 'Mirta' tuber diseases and their yield. Zemes ukio Mokslai, 16–25
- Riga E, Karanastasi E, Oliveira CMG, Neilson R (2007) Molecular identification of Two stubby root nematode species. Am J Potato Res 84:161–167
- Riga E, Neilson R (2005) First report of the stubby-root nematode, *Paratrichodorus teres*, from potato in the Columbia basin of Washington State. Plant Dis 89:1361–1361
- Rivera-Varas VV, Freeman TA, Gudmestad NC, Secor GA (2007) Mycoparasitism of *Helminthosporium solani* by *Acremonium strictum*. Phytopathology 97:1331–1337. doi:10.1094/phyto-97-10-1331
- Robbins RT, Barker KR (1974) Effects of soil type, particle size, temperature, and moisture on reproduction of *Belonolaimus longicaudatus*. J Nematology 6:1–6
- Rojancovschi E (1994) Researches concerning the effectiveness of some pesticides for the protection of potato crop against the potato rot nematode, *Ditylenchus destructor* Thorne. BPP, Buletinul de Protectia Plantelor, 29–31
- Ronda BHNAM, van Beuningen AR, Gorkink RFJ, Zwaardemaker NG, Janse JD (1999) Evaluation of a PCR for detection of *Ralstonia (Pseudomonas) solanacearum* (race 3, biovar 2) and *Clavibacter michiganensis* subsp. *sepedonicus* and comparison with immuno-fluorescence microscopy, plating on semi-selective SMSA medium, pathogenicity test and fluorescent in-situ hybridisation. Mededelingen Faculteit Landbouwkundige en Toegepaste Biologische Wetenschappen Universiteit Gent 64:583–591
- Rotenberg D, MacGuidwin AE, Saeed IAM, Rouse DI (2004) Interaction of spatially separated *Pratylenchus penetrans* and *Verticillium dahliae* on potato measured by impaired photosynthesis. Plant Pathol 53:294–302. doi:10.1111/j.1365-3059.2004.01005.x
- Rousselle P, Robert Y, Crosnier J-C (eds) (1996) La pomme de terre: Production, amélioration, ennemis et maladies, utilisations, Paris
- Rudkiewicz F, Sikorski J (1984) Effect of the incidence of common scab (*Streptomyces* spp.) on seed potatoes on plant growth and tuber yield. Biuletyn Instytutu Ziemiaka, 129–138
- Rudkiewicz F, Sikorski J, Slazak J (1983) Effect of the soil type, fertilization and control of *Phytophthora infestans* on the development of some diseases of potato plants and tubers. Biuletyn Instytutu Ziemiaka, 157–170
- Ruijter FJD, Haverkort AJ (1999) Effects of potato-cyst nematodes (*Globodera pallida*) and soil pH on root growth, nutrient uptake and crop growth of potato. Eur J Plant Pathol 105:61–76. doi:10.1023/a:1008641511688
- Ryan NA, Deliopoulos T, Jones P, Haydock PPJ (2003) Effects of a mixed-isolate mycorrhizal inoculum on the potato–potato cyst nematode interaction. Ann Appl Biol 143:111–119
- Ryan NA, Jones PW (2003) Effect of tuber-borne micro-organisms on hatching activity of potato root leachate towards potato cyst nematodes. Nematology 5:55–63
- Saeed IAM, Macguidwin AE, Rouse DI (1998) Effect of initial nematode population density on the interaction of *Pratylenchus penetrans* and *Verticillium dahliae* on 'Russet Burbank' potato. J Nematology 30:100–107
- Sahajdak A, Uznanska B (2003) Potato health requirements arising from EU regulations and the Treaty of Accession. Ochrona Roslin 47:20–23
- Salas B, Secor GA, Taylor RJ, Gladmestad NC (2003) Assessment of resistance of tubers of potato cultivars to *Phytophthora erythroseptica* and *Pythium ultimum*. Plant Dis 87:91–97
- Salas B, Stack RW, Secor GA, Gudmestad NC (2000) The effect of wounding, temperature, and inoculum on the development of pink rot of potatoes caused by *Phytophthora erythroseptica*. Plant Dis 84:1327–1333
- Saleh OI, Huang JS (1997) Bacterial soft rot disease of tomato fruits in Florida, USA: identification, response of some American and Egyptian cultivars of solanaceous plants and chemical control. Assiut J Agric Sci 28:11–26
- Samaliev H, Andreev R (1998) Relationship between initial population density of potato cyst nematode *Globodera pallida* Thorne and the yield of partially resistant potato varieties. Bulgarian J Agric Sci 4:421–427
- Samaliev H, Grigorov P, Samalieva A (1998) Influence of population density of *Globodera rostochiensis* (Nematoda: Heteroderidae) on potato yield. Rasteniev"n Nauki 35:235–238
- Sankaranarayanan C, Sundarababu R (2001) Influence of moisture and pH on the efficiency of VA-mycorrhiza, *Glomus mosseae* (Nicol and Gerd.) Gerd. and Trappe against *Meloidogyne incognita* (Kofoid and White) Chitw. on blackgram (*Vigna mungo* L.) Hepper. J Biol Control 15:69–72
- Santamarina MP, Rosello J (2006) Influence of temperature and water activity on the antagonism of *Trichoderma harzianum* to *Verticillium* and *Rhizoctonia*. Crop Prot 25:1130–1134. doi:10.1016/j.cropro.2006.02.006
- Savary S, Teng PS, Willocquet L, Nutter FW (2006) Quantification and modeling of crop losses: a review of purposes. Annu Rev Phytopathol 44:89–112. doi:10.1146/annurev.phyto.44.070505.143342
- Scheid L (2006) What to do against storage diseases? Kartoffelbau, 362–365
- Schena L, Nigro F, Ippolito A (2002) Identification and detection of *Rosellinia necatrix* by conventional and real-time Scorpion-PCR. Eur J Plant Pathol 108:355–366

- Schisler DA, Slininger PJ, Miller JS, Woodell LK, Clayton S, Olsen N (2009) Bacterial antagonists, zoospore inoculum retention time and potato cultivar influence pink rot disease development. Am J Potato Res 86:102–111. doi:10.1007/s12230-008-9066-7
- Scholte K (2000) Effect of potato used as a trap crop on potato cyst nematodes and other soil pathogens and on the growth of a subsequent main potato crop. Ann Appl Biol 136:229–238
- Scholte K (2005) Netted scab In: Delleman J, Mulder A, Peeten JMG, Schippers B, Turkensteen LJ (eds) Potato diseases. Diseases, pests and defects, NIVAP Holland, Aardappelwereld magazine, pp 83–84
- Scholte K, S'Jacob JJ (1989) Synergistic interactions between *Rhizoctonia solani* Kuhn, *Verticillium dahliae* Kleb., *Meloidogyne* spp. and *Pratylenchus neglectus* (Rensch) Chitwood & Oteifa, in potato. Potato Res 32:387–395
- Secor GA, Debuhr L, Gudmestad NC (1988) Susceptibility of *Corynebacterium sepedonicum* to disinfectants *in vitro*. Plant Dis 72:585–588
- Sepulveda RP, Lopez TH, Nunez LD (2000) Effect of different soil humidity on the development of potato smut (*Angiosorus solani*) in two potato varieties (*Solanum tuberosum*) under greenhouse conditions. Agricultura Tec 60:313–319
- Serfontein S, Logan C, Swanepoel AE, Boeema BH, Theron DJ (1991) A potato wilt disease in South Africa caused by *Erwinia carotovora* subspecies *carotovora* and *Erwinia chrysanthemi*. Plant Pathol 40:382–386
- Sharifi K, Zare R, Rees-George J (2008) Vegetative compatibility groups among *Fusarium solani* isolates causing potato dry rot. J Biol Sci 8:374–379
- Shcolnick S, Dinoor A, Tsror L (2007) Additional vegetative compatibility groups in *Colletotrichum coccodes* subpopulations from Europe and Israel. Plant Dis 91:805–808. doi:10.1094/pdis-91-7-0805
- Shekhawat GS, Perombelon MCM (1991) Factors affecting survival in soil and virulence of *Pseudomonas solanacearum*. Z Fur Pflanzenkrankheiten Und Pflanzenschutz J Plant Dis Prot 98:258–267
- Sheoraj S, Prajapati RK, Srivastava SSL, Pandey RK (2007) Influence of soil texture, temperature and relative humidity on development of collar rot of lentil. Indian Phytopathol 60:273–274
- Shojaei M, Karegar A, Deljoo A (2006) Aspects of biology of, and responses of several potato cultivars to, potato rot nematode. Iran J Plant Pathol 42:Pe577–Pe595, en165–en168
- Sikka LC, Singh AK (1976) Factors influencing quality of ware potatoes. J Indian Potato Assoc 3:24–28
- Singh RDN, Kaiser SAKM (1994) Effect of different culture media and pH levels on growth and cultural characteristics of charcoal rot pathogen (*Macrophomina phaseolina*) infecting maize. Crop Res Hisar 7:282–287
- Slack SA, Westra AAG (1998) Evaluation of flusulfamide for the control of bacterial ring rot of potato. Am J Potato Res 75:225–230
- Smith DS, De Boer SH (2009) Implementation of an artificial reaction control in a TaqMan method for PCR detection of *Ralstonia solanacearum* race 3 biovar 2. Eur J Plant Pathol 124:405–412. doi:10.1007/s10658-008-9427-6
- Smith NC, Hennessy J, Stead DE (2001) Repetitive sequence-derived PCR profiling using the BOX-A1R primer for rapid identification of the plant pathogen *Clavibacter michiganensis* subspecies *sepedonicus*. Eur J Plant Pathol 107:739–748
- Soltani N, Conn KL, Abbasi PA, Lazarovits G (2002) Reduction of potato scab and *Verticillium* wilt with ammonium lignosulfonate soil amendment in four Ontario potato fields. Can J Plant Pathol Rev Can De Phytopathologie 24:332–339
- Solunke BS, Karepaa BM, Gangawane LV (2001) Survival ability of carbendazim resistant *Sclerotium rolfsii* in mixed population. Indian Phytopathol 54:486–487
- Soman AK, Chauhan RKS (1996) Potato tuber rots in Gwalior, Madhya Pradesh. J Indian Potato Assoc 23:144–148
- Spaull VW, Cadet P (2001) Nematodes and nutrients: association between plant-parasitic nematodes and soil chemicals. Proceedings of the Annual Congress—South African Sugar Technologists' Association, 116–117
- Stachewicz VH, Enzian S (1998) Do temperature and rainfall limit the occurrence of potato wart in Germany? Nachrichtenblatt Dtsch Pflanzenschutzdienstes 50:105–111
- Stamps DJ (1978) *Phytophthora erythroseptica*. IMI Descriptions of Fungi and Bacteria, Sheet 593
- Steinberg C, Edel-Hermann V, Alabouvette C, Lemanceau P (2007) Soil suppressiveness to plant diseases. In: van Elsas JD, Jansson J, Trevors JT (eds) Modern soil microbiology. CRC, New York, pp 455–478
- Stevenson WR, Loria R, Franc GD, Weingartner DP (2001) Compendium of potato diseases. The American Phytopathological Society, St. Paul, MN, USA
- Strausbaugh CA, Schroth MN, Weinhold AR, Hancock JG (1992) Assessment of vegetative compatibility of *Verticillium dahliae* tester strains and isolates from California potatoes. Phytopathology 82:61–68
- Sunaina V, Kishore V, Shekhawat GS, Kumar M (2000) Persistence of *Ralstonia solanacearum* in naturally infested soil under changing environment. Potato, Global Research & Development. Proceedings of the Global Conference on Potato, New Delhi, India, 6–11 December, 1999: vol 1, pp 444–447
- Suyama K, Tjahjono B, Fujii H (1990) Occurrence of slimy rot disease on the stored potato tubers. Ann Phytopathological Soc Jpn 56:577–583
- Taylor RJ (2005) Influence of tillage and method of metam sodium application on distribution and survival of *Verticillium dahliae* in the soil and the development of *Verticillium* wilt of potato. Am J Potato Res 82:451–461
- Taylor RJ, Pasche JS, Gudmestad NC (2006) Biological significance of mefenoxam resistance in *Phytophthora erythroseptica* and its implications for the management of pink rot of potato. Plant Dis 90:927–934. doi:10.1094/pd-90-0927
- Taylor RJ, Pasche JS, Gudmestad NC (2008) Susceptibility of eight potato cultivars to tuber infection by *Phytophthora erythroseptica* and *Pythium ultimum* and its relationship to mefenoxam-mediated control of pink rot and leak. Ann Appl Biol 152:189–199. doi:10.1111/j.1744-7348.2007.00203.x
- Taylor RJ, Salas B, Gudmestad NC (2004) Differences in etiology affect mefenoxam efficacy and the control of pink rot and leak tuber diseases of potato. Plant Dis 88:301–307
- Ten Hoopen GM, Krauss U (2006) Biology and control of *Rosellinia bunodes*, *Rosellinia necatrix* and *Rosellinia pepo*: a review. Crop Prot 25:89–107
- Termorshuizen AJ, van Rijn E, van der Gaag DJ, Alabouvette C, Chen Y, Lagerlof J, Malandrakis AA, Paplomatas EJ, Ramert B, Ryckeboer J, Steinberg C, Zmora-Nahum S (2006) Suppressiveness of 18 composts against 7 pathosystems: variability in pathogen response. Soil Biol Biochem 38:2461–2477. doi:10.1016/j.soilbio.2006.03.002
- Thompson AL, Taylor RJ, Pasche JS, Novy RG, Gudmestad NC (2007) Resistance to *Phytophthora erythroseptica* and *Pythium ultimum* in a potato clone derived from *S-berthaultii* and *S-tuberosum*. Am J Potato Res 84:149–160
- Tivoli B (1983) Action of temperature and humidity on the behavior in soil of 3 *Fusarium* species or varieties causing dry-rot in potato-tubers. Agronomie 3:1001–1009
- Tivoli B, Corbiere R, Lemarchand E (1990) Relations between the Ph of soils and their level of conduciveness to *Fusarium solani* var *Coeruleum* and *Fusarium roseum* var *sambucinum* responsible for the dry rot of potato-tubers. Agronomie 10:63–68
- Tivoli B, Tika N, Lemarchand E (1987) Observations on the conduciveness of soils to fungi causing dry rot of potato-tubers—*Fusarium*

- solanum* var. *coeruleum*, *F. roseum* var. *sambucinum* and *Phoma exigua* var. *foveata*. Agronomie 7:531–538
- Tomlinson DT, Elphinstone JG, El-Fatah HA, Agag SHA, Kamal M, Soliman MY, El-Aliem MMA, El-Ghany HA, El-Haddad SA, Fawzi FG, Janse JD (2005) Survival of the potato brown rot bacterium (*Ralstonia solanacearum* biovar 2) in Egyptian soils. Potato in Progress: Science Meets Practice, 233–238
- Trifonova Z (1997) Effect of population density on cyst production of *Globodera rostochiensis* Woll. 1923. Bulgarian Journal of Agricultural Science 3:755–758
- Trifonova Z (1923) (2001) Effect of inorganic fertilizers and soil type on the growth of potato and reproduction of *Globodera rostochiensis* Woll. Macedonian Agric Rev 48:57–60
- Triki MA, Priou S, El MM, Baudry A (2001) Leek syndrome of potato in Tunisia caused by *Pythium aphanidermatum* and *Pythium ultimum*. Potato Res 44:221–231
- Tsror L (2004) Effect of light duration on severity of black dot caused by *Colletotrichum coccodes* on potato, Plant Pathology 53, 288–293
- Tsror L, Erlich O, Cahlon Y, Hadar A, Cohen Y, Klein L, Peretz-Alon I (2000) Control of *Verticillium dahliae* prior to potato production by soil fumigation with chloropicrin. Proceedings of the International Symposium on Chemical and Non-Chemical Soil and Substrate Disinfestation, 201–204
- Tsror L, Nachmias A, Erlich O, Aharon M, Perombelon MCM (1993) A 9-year monitoring study of diseases on potato seed tubers imported to Israel. Phytoparasitica 21:321–328
- Tsror L, Peretz-Alon I (2005) The influence of the inoculum source of *Rhizoctonia solani* on development of black scurf on potato. J Phytopathol 153:240–244
- Tsror L, Shlevin E, Peretz-Alon I (2005) Efficacy of metam sodium for controlling *Verticillium dahliae* prior to potato production in sandy soils. Am J Potato Res 82:419–423
- UNECE (2010) Working party on agricultural quality standards. Available from: <http://www.unece.org/trade/agr/>. Accessed on 15 April 2010.
- Vallad GE, Qin QM, Subbarao KV (2004) *Verticillium* wilt of cool season vegetable crops: their distribution, impact and management. Recent research developments in plant pathology 3:189–210
- Vanvurde JW, Devries PM (1994) Population dynamics of *Erwinia carotovora* subsp. *atroseptica* on the surface of intact and wounded seed potatoes during storage. J Appl Bacteriol 76:568–575
- Vasinauskiene M, Baranauskaite L (2003) Occurrence of *Clavibacter michiganensis* subsp. *sepedonicus* in Lithuania. Zemes ukio Mokslai, 30–33
- Vian J (2009) Comparaison de différentes techniques de travail du sol en agriculture biologique: effet de la structure et de la localisation des résidus sur les microorganismes du sol et leurs activités de minéralisation du carbone et de l'azote. Institut des Sciences et Industries du Vivant et de l'Environnement (Agro Paris Tech). Agro Paris Tech, Paris, p 130
- Vico I, Krstic B, Stojanovic G (1997) The occurrence of *Polyscytalum pustulans* on potatoes in Yugoslavia. Acta Horticulturae 462:339–343
- Vishwa D, Sarbhoy AK (1989) Studies on the germination and longevity of pycnidiospores of *Macrophomina phaseolina*. Indian Phytopathol 42:123–127
- Vivoda E, Davis RM, Nunez JJ, Guerard JP (1991) Factors affecting the development of cavity spot of carrot. Plant Dis 75:519–522
- Volovik AS, Borisenok AV, Shuiskaya NG (1980) Agrotechniques in the control of diseases. Zashchita Rastenii, 26
- Vovlas N, Mifsud D, Landa BB, Castillo P (2005) Pathogenicity of the root-knot nematode *Meloidogyne javanica* on potato. Plant Pathol 54:657–664. doi:[10.1111/j.1365-3059.2005.01244.x](https://doi.org/10.1111/j.1365-3059.2005.01244.x)
- Vreugdenhil D (2007) Potato biology and biotechnology—advances and perspectives. Elsevier, Amsterdam
- Vries PMd, Vuurde JWV (1993) Survival of *Erwinia carotovora* subsp. *atroseptica* on seed potato. Gewasbeschermering 24:103–108
- Wale S, Platt HWB, Cattlin ND (2008) Diseases, pests and disorders of potatoes. Academic, London
- Wang AX, Lazarovits G (2005) Role of seed tubers in the spread of plant pathogenic *Streptomyces* and initiating potato common scab disease. Am J Potato Res 82:221–230
- Wanner LA (2007) High proportions of nonpathogenic *Streptomyces* are associated with common scab-resistant potato lines and less severe disease. Can J Microbiol 53:1062–1075. doi:[10.1139/w07-061](https://doi.org/10.1139/w07-061)
- Ward LI (2004) A real-time PCR assay based method for routine diagnosis of *Spongospora subterranea* on potato tubers. J Phytopathol 152:633–638
- Wegener CB, Jansen G (2007) Soft-rot resistance of coloured potato cultivars (*Solanum tuberosum* L.): the role of anthocyanins. Potato Res 50:31–44. doi:[10.1007/s11540-007-9027-4](https://doi.org/10.1007/s11540-007-9027-4)
- Weller DM, Raaijmakers JM, Gardener BBM, Thomashow LS (2002) Microbial populations responsible for specific soil suppressiveness to plant pathogens. Ann Rev Phytopathol 40:309–+. doi:[10.1146/annurev.phyto.40.030402.110010](https://doi.org/10.1146/annurev.phyto.40.030402.110010)
- Westra AAG, Arneson CP, Slack SA (1994) Effect of interaction of inoculum dose, cultivar, and geographic location on the development of foliar symptoms of bacterial ring rot of potato. Phytopathology 84:410–415
- Wharton DA, Worland MR (2001) Water relations during desiccation of cysts of the potato-cyst nematode *Globodera rostochiensis*. J Comp Physiol B Biochem Syst Environ Physiol 171:121–126
- Wharton PS, Kirk WW, Berry D, Tumbalam P (2007) Seed treatment application-timing options for control of *Fusarium* decay and sprout rot of cut seedpieces. Am J Potato Res 84:237–244
- Wilson CR, Ransom LM, Pemberton BM (1999) The relative importance of seed-borne inoculum to common scab disease of potato and the efficacy of seed tuber and soil treatments for disease control. J Phytopathol Phytopathologische Z 147:13–18
- Wilson PS, Ahvenniemi PM, Lehtonen MJ, Kukkonen M, Rita H, Valkonen JPT (2008) Biological and chemical control and their combined use to control different stages of the *Rhizoctonia* disease complex on potato through the growing season. Ann Appl Biol 153:307–320. doi:[10.1111/j.1744-7348.2008.00292.x](https://doi.org/10.1111/j.1744-7348.2008.00292.x)
- Wolf JMvd, Beckhoven JRCMv (2004) Factors affecting survival of *Clavibacter michiganensis* subsp. *sepedonicus* in water. J Phytopathol 152:161–168. doi:[10.1111/j.1439-0434.2004.00820.x](https://doi.org/10.1111/j.1439-0434.2004.00820.x)
- Wolf JMvd, Beckhoven JRCMv, Hukkanen A, Karjalainen R, Muller P (2005) Fate of *Clavibacter michiganensis* subsp. *sepedonicus*, the causal organism of bacterial ring rot of potato, in weeds and field crops. J Phytopathol 153:358–365. doi:[10.1111/j.1439-0434.2005.00985.x](https://doi.org/10.1111/j.1439-0434.2005.00985.x)
- Woodhall JW, Lees AK, Edwards SG, Jenkinson P (2007) Characterization of *Rhizoctonia solani* from potato in Great Britain. Plant Pathol 56:286–295
- Woodhall JW, Lees AK, Edwards SG, Jenkinson P (2008) Infection of potato by *Rhizoctonia solani*: effect of anastomosis group. Plant Pathol 57:897–905. doi:[10.1111/j.1365-3059.2008.01889.x](https://doi.org/10.1111/j.1365-3059.2008.01889.x)
- Wronkowska H, Janowicz K (1989) Examination of *Globodera rostochiensis* cysts infected with strains of *Fusarium oxysporum* in vitro. Zesz Problemowe Postepow Nauk Rolniczych 358:57–61
- Wu Q, Wang X, Liao J, Qin G, Li S (2006) Effect of light and temperature on culture of root-knot nematode on *Solanum tuberosum*. Plant Prot 32:27–29
- Xu J, Zhou B, Liu Y, Li M, Li J (1997) Biological characteristics of *Rhizoctonia solani* Kuhnzheng. Plant Prot 23:29–30
- Yaganza ES, Rioux D, Simard M, Arul J, Tweddell RJ (2004) Ultrastructural alterations of *Erwinia carotovora* subsp *atroseptica* caused by treatment with aluminum chloride and sodium metabisulfite. Appl Environ Microbiol 70:6800–6808. doi:[10.1128/aem.70.11.6800-6808.2004](https://doi.org/10.1128/aem.70.11.6800-6808.2004)

- Yang LP, Xie JT, Jiang DH, Fu YP, Li GQ, Lin FC (2008) Antifungal substances produced by *Penicillium oxalicum* strain PY-1-potential antibiotics against plant pathogenic fungi. World J Microbiol Biotechnol 24:909–915. doi:[10.1007/s11274-007-9626-x](https://doi.org/10.1007/s11274-007-9626-x)
- Yi Y, Sul K (1998) Control strategy of acidified nutrient solution on bacterial wilt of tomato plants. Korean J Plant Pathol 14:744–746
- Young CS, Clarkson JP, Smith JA, Watling M, Phelps K, Whippes JM (2004) Environmental conditions influencing *Sclerotinia sclerotiorum* infection and disease development in lettuce. Plant Pathol 53:387–397. doi:[10.1111/j.1365-3059.2004.01018](https://doi.org/10.1111/j.1365-3059.2004.01018)
- Zagoskina NV, Goncharuk EA, Dubravina GA, Kalashnikova EA (2006) Effect of exometabolites of the fungus *Rhizoctonia solani* on the cell cultures of potato various genotypes. Biotehnologiya 5:19–22
- Zambolim L, Parizzi P, Matsuoka K, Xavier F, Vale RD, Chaves GM (1995) Powdery scab of potato. Fitopatologia Bras 20:5–12
- Zhang Z, Chen R, Wang Y, Wang K, Zheng X (2005) Molecular detection of *Verticillium albo-atrum* by PCR based on its sequences. Agric Sci China 4:760–766
- Zhao WQ, Liu DQ, Yu XM (2008) First report of potato scab caused by *Streptomyces turgidiscabies* in China. Plant Dis 92:1587. doi:[10.1094/pdis-92-11-1587c](https://doi.org/10.1094/pdis-92-11-1587c)
- Zielke R, Muller HJ, Ficke W, Naumann K, Skadow K (1974) The effect of soil and climate on the occurrence of blackleg and tuber soft rot of potato, Archiv fur Phytopathologie und Pflanzenschutz 10:245–253
- Zimny L, Wacawowicz R, Oliwa T (2006) Tuber infestation by *Rhizoctonia solani* in relation to cultivation systems of potato. Prog Plant Prot 46:388–394