



## Erratum to: Management of soil nitrate heterogeneity resulting from crop rows in a lettuce-tomato rotation under a greenhouse

François Lecompte

### ► To cite this version:

François Lecompte. Erratum to: Management of soil nitrate heterogeneity resulting from crop rows in a lettuce-tomato rotation under a greenhouse. *Agronomy for Sustainable Development*, 2012, 32 (2), pp.609-610. 10.1007/s13593-011-0064-7 . hal-00930541

HAL Id: hal-00930541

<https://hal.science/hal-00930541>

Submitted on 11 May 2020

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

## Erratum to: Management of soil nitrate heterogeneity resulting from crop rows in a lettuce–tomato rotation under a greenhouse

François Lecompte

Published online: 21 December 2011  
© INRA and Springer-Verlag, France 2011

**Erratum to: Agron. Sustain. Dev.**  
**DOI 10.1007/s13593-011-0047-8**

In the original article Table 3 is not correct. The calculated values of leaching for the “L” system is 54 kg/ha in the inter-row (IR), and 25 kg/ha in the row (R), and NOT 54 and 75 kg/ha as published in the online version of the article. We reprint the complete corrected table here below.

Everything else in the paper remains correct.

---

The online version of the original article can be found at <http://dx.doi.org/10.1007/s13593-011-0047-8>.

F. Lecompte (✉)  
UR 1115 Plantes et Systèmes de Cultures Horticoles, INRA,  
Domaine Saint Paul, Site Agroparc,  
84914, Avignon Cedex, France  
e-mail: francois.lecompte@avignon.inra.fr

**Table 3** N balance components for lettuce at two positions (Row R and Inter-row IR) and for two management systems “C” (current farm practices) and “L” (limited water and nutrient inputs)

	kgN/ha	C		L	
		IR	R	IR	R
2006–2007 (L1)	Initial stock ( $S_i$ ) <sup>a</sup>	160	106	59	34
	Final stock ( $S_f$ )	30	16	13	10
	$S_f - S_i$	-130	-90	-46	-24
	Crop uptake ( $U$ )	99	76	58	38
	Mineralization ( $M$ ) <sup>b</sup>	39	39	39	39
	Leaching ( $L$ ) <sup>c</sup>	70	53	27	25
	Initial stock ( $S_i$ ) <sup>a</sup>	156	96	78	34
2007–2008 (L2)	Final stock ( $S_f$ )	25	21	19	18
	$S_f - S_i$	-131	-75	-59	-16
	Crop uptake ( $U$ )	87	63	47	33
	Mineralization ( $M$ ) <sup>b</sup>	42	42	42	42
	Leaching ( $L$ ) <sup>c</sup>	86	54	54	25

<sup>a</sup>Including the N fertilization before lettuce transplanting in system C

<sup>b</sup>Estimated from stock variations between September and November 2008

<sup>c</sup> $L = M - U - (S_f - S_i)$ . Part of this term may include unaccounted denitrification or volatilization