



**HAL**  
open science

# The question of interior blow-up for an elliptic Neumann problem: the critical case

Olivier Rey

► **To cite this version:**

Olivier Rey. The question of interior blow-up for an elliptic Neumann problem: the critical case. Journal de Mathématiques Pures et Appliquées, 2002, 81, pp.655-696. hal-00935381

**HAL Id: hal-00935381**

**<https://hal.science/hal-00935381v1>**

Submitted on 23 Jan 2014

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# The question of interior blow-up points for an elliptic Neumann problem : the critical case

Olivier REY

Centre de Mathématiques, École Polytechnique  
U.M.R. 7640 du C.N.R.S., 91128 Palaiseau Cedex

## Abstract

In contrast with the subcritical case, we prove that for any bounded domain  $\Omega$  in  $\mathbb{R}^3$ , the Neumann elliptic problem with critical nonlinearity

$$-\Delta u + \mu u = u^5, \quad u > 0 \text{ in } \Omega; \quad \partial u / \partial \nu = 0 \text{ on } \partial \Omega$$

has no solution blowing up at only interior points as  $\mu$  goes to infinity.

## 1 Introduction and Results

Wondering about the mechanisms of pattern formation in biology, Turing [42] made the very important discovery that contrary to the intuition, which associates diffusion phenomena to a smoothing of initial data, spatial concentration structures may result from the interaction of two substances with different diffusion rates. Since that time many biological patterns have been explained in such a way. Models describing the evolution of the two involved substances concentrations, as those proposed by Keller and Segel, or Gierer and Meinhardt, consist in a system of two coupled nonlinear parabolic equations. Under some further assumptions, finding stationary solutions to the system reduces to solving a single nonlinear elliptic equation with Neumann boundary conditions [32]

$$(P_\mu) \begin{cases} -\Delta u + \mu u & = u^p, \quad u > 0 & \text{in } \Omega \\ \frac{\partial u}{\partial \nu} & = 0 & \text{on } \partial \Omega \end{cases}$$

where  $p > 1$ ,  $\mu > 0$  are fixed parameters, and  $\Omega$  is a smooth bounded domain in  $\mathbb{R}^n$ . Note that setting  $v = \mu^{-1/(p-1)}u$ ,  $d^2 = 1/\mu$ , problem  $(P_\mu)$  is equivalent to

$$(P'_d) \begin{cases} -d^2 \Delta v + v & = v^p, \quad v > 0 & \text{in } \Omega \\ \frac{\partial v}{\partial \nu} & = 0 & \text{on } \partial \Omega. \end{cases}$$

Since the works of Lin, Ni and Takagi [31, 32, 36, 37], many papers have been devoted to the study of  $(P'_d)$ , under the assumption that  $p$  is subcritical, i.e.  $n = 2$ , or  $n \geq 3$  and  $p < (n+2)/(n-2)$ . A natural question is to know whether the results which hold for subcritical exponents are true, or not, for critical

or supercritical exponents. The study of the critical case made similarities and differences appear with respect to the subcritical case. For example, it was proved that when  $p$  is subcritical, the only solution to  $(P_\mu)$  for small  $\mu$  is constant [32], and the same holds when  $n = 3$  and  $p$  is critical [52, 49]. However, if  $n = 4, 5, 6$ ,  $\Omega$  is a ball and  $p$  is critical,  $(P_\mu)$  has at least one nonconstant radial solution for small  $\mu$  [6].

For large  $\mu$ , it is known that in both subcritical and critical cases,  $(P_\mu)$  has solutions which concentrate at some points of the domain as  $\mu$  goes to infinity (alternatively,  $d$  goes to zero in  $(P'_d)$ ). The next question is to characterize such concentration points. In both cases, the least energy solutions have, for large  $\mu$ , exactly one maximum point which lies on the boundary of the domain, and which goes, as  $\mu$  goes to infinity, to a maximum point of the mean curvature of the boundary [36, 37] [4, 5, 35, 44, 40].

For subcritical exponents, higher energy solutions exist which blow up at one or several points of the boundary [21, 29, 49, 17] as  $\mu$  goes to infinity. Solutions also exist which blow up at one or several points in the interior of the domain [48, 19, 16, 23, 27, 20, 51, 10]. In particular,  $(P_\mu)$  has single interior spike solutions which blow up at a local maximum point of the distance function  $d(x, \partial\Omega)$ ,  $x \in \Omega$ . (Solutions have also been built which blow up at interior and boundary points at the same time [26].) For critical exponent, all the existence results concern solutions which blow up at one or several points of the boundary as  $\mu$  goes to infinity [1-6, 12, 38, 40, 43-47, 33, 22, 24-25]. Hence the question : do solutions blowing up at interior points exist ?

The only known result is partial and negative [14] : for  $n \geq 5$  and critical  $p$ ,  $(P_\mu)$  has no solution  $u_\mu$  such that

$$\|u_\mu - \alpha_n \sum_{i=1}^k U_{\lambda_\mu^i, y_\mu^i}\|_{H^1(\Omega)} \rightarrow 0 \quad \text{as } \mu \rightarrow \infty$$

with  $\alpha_n = (n(n-2))^{(n-2)/4}$ ,  $k \in \mathbb{N}^*$

$$U_{\lambda, y}(x) = \frac{\lambda^{(n-2)/2}}{(1 + \lambda^2|x-y|^2)^{(n-2)/2}} \quad \lambda \in \mathbb{R}_+^*, \quad x, y \in \mathbb{R}^n \quad (1.1)$$

and  $\lambda_\mu^i \rightarrow \infty$ ,  $y_\mu^i \rightarrow y^i$  in  $\Omega$  as  $\mu \rightarrow \infty$ ,  $y^i \neq y^j$  if  $i \neq j$ .

Such a result could have also been derived from the arguments in [38] and, in the case  $n = 3$  and  $k = 1$ , from [40] (see the final remark at the end of Section 3). A recent paper shows that the case  $k = 1$  cannot happen in any dimension [18].

Our aim in this paper is to consider the question of interior blow-up points, for  $n = 3$  and critical  $p$ , without any assumption neither on the number of those points, nor on the distance between them, which may be zero.

$(u_\mu)_{\mu \geq \mu_0}$  being a sequence of nonconstant solutions to  $(P_\mu)$ , there are several and equivalent ways to define blow-up points of  $(u_\mu)$ . For example,  $y \in \bar{\Omega}$  will be said to be a blow-up point of  $(u_\mu)$  if and only if

$$\liminf_{r \rightarrow 0} \limsup_{\mu \rightarrow \infty} \int_{B(y,r) \cap \Omega} |\nabla u_\mu|^2 \left( \text{or } \int_{B(y,r) \cap \Omega} u_\mu^6 \right) > 0.$$

Our main result is :

**Theorem 1** *Let  $(u_\mu)_{\mu \geq \mu_0}$  a sequence, bounded in  $H^1(\Omega)$ , of solutions to  $(P_\mu)$ . There exists at least one blow-up point which lies on the boundary of  $\Omega$ .*

We notice that, in contrast with the subcritical case, the existence of solutions with a finite number of blow-up points all lying in the interior of the domain, is excluded. We emphasize that the main difficulty in this work is to eliminate the possibility of *multiple* interior peaks without a *a priori* assumption on the location of those peaks, which may be very close from each other, or even centered at the same point.

The next section is devoted to an *a priori* analysis of the solutions to  $(P_\mu)$  as  $\mu$  goes to infinity. This analysis, mixing together energy-dependent and energy-independent estimates, provides us with informations about the shape of solutions, which allow us to prove, through variational methods, the theorem in Section 3.

## 2 Blow-up analysis

### 2.1 Energy-independent estimates

We begin with an energy-independent description of the nonconstant solutions to  $(P_\mu)$  as  $\mu$  goes to infinity. We have the following proposition :

**Proposition 2.1** *Let  $(u_\mu)_{\mu \geq \mu_0}$  be a sequence of nonconstant solutions to  $(P_\mu)$ . Let  $\varepsilon > 0$ ,  $R > 1$ . For  $\mu$  large enough,  $u_\mu$  has  $N_\mu \in \mathbb{N}^*$  local maximum points  $x_\mu^i \in \bar{\Omega}$ ,  $1 \leq i \leq N_\mu$ , such that :*

- (i)  $\frac{\mu^{1/4}}{u_\mu(x_\mu^i)} < \varepsilon$
- (ii)  $\left\| \frac{1}{u_\mu(x_\mu^i)} u_\mu \left( \frac{x}{u_\mu^2(x_\mu^i)} + x_\mu^i \right) - \frac{1}{(1 + \frac{|x|^2}{3})^{1/2}} \right\|_{C^2(B(0,2R) \cap \Omega_\mu^i)} < \varepsilon$   
with  $\Omega_\mu^i = u_\mu^2(x_\mu^i)(\Omega - x_\mu^i)$
- (iii)  $x_\mu^i \in \partial\Omega$  or  $u_\mu^2(x_\mu^i)d(x_\mu^i, \partial\Omega) > \frac{1}{\varepsilon}$
- (iv)  $\bar{B}(x_\mu^i, r_\mu^i) \cap \bar{B}(x_\mu^j, r_\mu^j) = \emptyset$  for  $i \neq j$ ,  $r_\mu^i = \frac{R}{u_\mu^2(x_\mu^i)}$
- (v)  $\left( d(x, \{x_\mu^i, 1 \leq i \leq N_\mu\}) \right)^{1/2} u_\mu(x) \leq C(\varepsilon, R)$ .

(i) says that the maxima increase faster than  $\mu^{1/4}$  as  $\mu$  goes to infinity; (ii) describes the shape of  $u_\mu$  in a neighbourhood of a maximum point  $x_\mu^i$ ; (iii) shows that either the maximum points  $x_\mu^i$  are on the boundary, or are not too close from the boundary with respect to the height of the maximum; (iv) shows that these maximum points are not too close from each other with respect to their heights, and (v) provides us with a global bound for  $u_\mu$  in  $\Omega$ .

Such a proposition relies on arguments initially developed by Schoen [41], in the context of the Yamabe problem. The proof, which follows the same scheme as in [30, Proposition 5.1], with the convenient additional arguments, is given in Appendix A.

Let now  $x_\mu$  be any point in  $\bar{\Omega}$ . We set

$$v_\mu(y) = \frac{1}{\mu^{1/4}} u_\mu\left(\frac{y}{\mu^{1/2}} + x_\mu\right) \quad y \in \Omega_\mu = \mu^{1/2}(\Omega - x_\mu) \quad (2.1)$$

which satisfies

$$-\Delta v_\mu + v_\mu = v_\mu^5, \quad v_\mu > 0 \text{ in } \Omega_\mu; \quad \frac{\partial v_\mu}{\partial \nu} = 0 \text{ on } \partial\Omega_\mu. \quad (2.2)$$

Proposition 2.1 is equivalent to :

**Proposition 2.2** *Let  $\varepsilon > 0$ ,  $R > 1$ . For  $\mu$  large enough,  $v_\mu$  has  $N_\mu \in \mathbb{N}^*$  local maximum points  $y_\mu^i = \mu^{1/2}(x_\mu^i - x_\mu) \in \bar{\Omega}$ ,  $1 \leq i \leq N_\mu$ , such that :*

- (i)  $v_\mu(y_\mu^i) > \frac{1}{\varepsilon}$
- (ii)  $\left\| \frac{1}{v_\mu(y_\mu^i)} v_\mu\left(\frac{y}{v_\mu^2(y_\mu^i)} + y_\mu^i\right) - \frac{1}{(1 + \frac{|y|^2}{3})^{1/2}} \right\|_{C^2(B(0,2R) \cap \Omega_\mu^i)} < \varepsilon$   
with  $\Omega_\mu^i = v_\mu^2(y_\mu^i)(\Omega_\mu - y_\mu^i) = u_\mu^2(x_\mu^i)(\Omega - x_\mu^i)$
- (iii)  $y_\mu^i \in \partial\Omega$  or  $v_\mu^2(y_\mu^i)d(y_\mu^i, \partial\Omega) > \frac{1}{\varepsilon}$
- (iv)  $\bar{B}(y_\mu^i, s_\mu^i) \cap \bar{B}(y_\mu^j, s_\mu^j) = \emptyset$  for  $i \neq j$ ,  $s_\mu^i = \frac{R}{v_\mu^2(x_\mu^i)}$
- (v)  $\left( d(y, \{y_\mu^i, 1 \leq i \leq N_\mu\}) \right)^{1/2} v_\mu(y) \leq C(\varepsilon, R)$ .

The interest of considering  $v_\mu$  instead of  $u_\mu$  is that  $v_\mu$  solves a nonlinear elliptic equation (2.2) whose linear part has constant, hence bounded coefficients. This fact allows us to use techniques and results of Li, concerning the scalar curvature problem [28], Li and Zhu concerning Yamabe type equations [30].

In view of Proposition 2.2, we set :

**Definition 1**  $\bar{y} \in \mathbb{R}^3$  is called an isolated blow-up point of  $(v_\mu)$  if there exist  $\bar{r} > 0$ ,  $C \in \mathbb{R}$  and a sequence  $(y_\mu)$  in  $\Omega_\mu$ , converging to  $\bar{y}$ , such that  $y_\mu$  is a local maximum of  $v_\mu$ ,  $v_\mu(y_\mu) \rightarrow \infty$  and

$$v_\mu(y) \leq \frac{C}{|y - y_\mu|^{1/2}} \quad y \in B(y_\mu, \bar{r}) \cap \Omega_\mu.$$

$\bar{y}$  is called a simple isolated blow-up point if moreover there exists  $r_0 > 0$  such that, for  $\mu$  large enough,  $r^{1/2}\bar{v}_\mu(r)$  has exactly one critical point in  $(0, r_0)$ , with

$$\bar{v}_\mu(r) = \frac{1}{|\partial B(y_\mu, r) \cap \Omega_\mu|} \int_{\partial B(y_\mu, r) \cap \Omega_\mu} v_\mu \quad 0 < r < \bar{r}.$$

Then, we can state :

**Proposition 2.3**

(i) Assume that  $\bar{y}$  is an isolated blow-up point, such that for large  $\mu$

$$d(y_\mu, \partial\Omega_\mu) \geq \rho \tag{2.3}$$

for some  $\rho > 0$ . Then,  $\bar{y}$  is a simple isolated blow-up point.

(ii) The same conclusion holds, without assuming (2.3), provided that  $\Omega$  is convex.

Proposition 2.3(i) follows directly from [28, Section 3] or [30, Section 4], and (ii) follows from [52, Section 2.2]. Moreover, we know that under assumption (2.3), simple isolated blow-up points are such that

$$v_\mu(y_\mu)v_\mu(y) \leq \frac{C}{|y - y_\mu|} \quad y \in B(y_\mu, r_0) \cap \Omega_\mu \tag{2.4}$$

with  $C$  some positive constant independent of  $\mu$ , and the same is true without assuming (2.3) if  $\Omega$  is convex (see [28, Prop. 2.3][30, Prop. 3.1][52, Prop. 2.1]).

In view of our further needs, we consider the points  $y_\mu^i$  defined in Proposition 2.2, and we prove the following proposition, which is of crucial interest in the sequel :

**Proposition 2.4** Let  $y_\mu^i$  be as in Proposition 2.2.

(i) Assume that there exists  $\rho > 0$  such that, for large  $\mu$ ,  $d(y_\mu^i, \partial\Omega_\mu) \geq \rho$ ,  $1 \leq i \leq N_\mu$ . Then, there exists  $\delta > 0$  such that, for  $\mu$  large enough

$$|y_\mu^i - y_\mu^j| \geq \delta \quad \forall i, j \quad i \neq j. \tag{2.5}$$

(ii) If  $\Omega$  is convex, (2.5) holds with  $\rho = 0$ .

This proposition still follows from [28, 30, 52]. Let us sketch the argument for (i). Assuming that the proposition is false, we may suppose, without loss of generality

$$\delta_\mu = |y_\mu^1 - y_\mu^2| = \min_{i \neq j} |y_\mu^i - y_\mu^j| \rightarrow 0 \text{ as } \mu \rightarrow \infty. \quad (2.6)$$

We set

$$w_\mu(z) = \delta_\mu^{1/2} v_\mu(\delta_\mu z + y_\mu^1) \quad z \in \tilde{\Omega}_\mu = (\Omega_\mu - y_\mu^1)/\delta_\mu.$$

$w_\mu$  satisfies

$$-\Delta w_\mu + \delta_\mu^2 w_\mu = w_\mu^5, \quad w_\mu > 0 \text{ in } \tilde{\Omega}_\mu; \quad \frac{\partial w_\mu}{\partial \nu} = 0 \text{ on } \partial \tilde{\Omega}_\mu \quad (2.7)$$

and, denoting  $z_\mu^i = (y_\mu^i - y_\mu^1)/\delta_\mu$ ,  $1 \leq i \leq N_\mu$ , we know that

$$w_\mu(0), w_\mu(z_\mu^2) > R^{1/2} \quad \text{because of Prop. 2.2}(iv) \quad (2.8)$$

$$|z_\mu^i - z_\mu^j| \geq 1 \quad i \neq j \quad \text{because of (2.6)} \quad (2.9)$$

$$\left( d(y, \{z_\mu^i, 1 \leq i \leq N_\mu\}) \right)^{1/2} w_\mu(y) \leq C \quad \text{because of Prop. 2.2}(v). \quad (2.10)$$

Up to a subsequence, we may assume that  $z_\mu^2 \rightarrow z^2$  as  $\mu \rightarrow \infty$ ,  $|z^2| = 1$ . We claim that  $w_\mu(0)$  and  $w_\mu(z_\mu^2)$  go to infinity as  $\mu$  goes to infinity.

Indeed, let us assume that  $w_\mu(z_\mu^2)$  stays bounded. Prop. 2.2(ii) then implies that  $w_\mu$  stays bounded in a fixed neighbourhood of  $z^2$ . Taking also account of (2.9-10), we see that  $w_\mu$  stays bounded in any ball  $B(z^2, r)$ ,  $0 < r < 1$ . If  $w_\mu(0)$  goes to infinity, 0 is an isolated blow-up point, hence a simple isolated blow-up point, and an inequality as (2.4) holds for  $w_\mu$ , i.e.

$$w_\mu(0)w_\mu(z) \leq \frac{C}{|z|} \quad z \in B(0, r'_0) \cap \tilde{\Omega}_\mu \quad (2.11)$$

Consequently, at a small and fixed distance of 0,  $w_\mu$  goes to zero. Therefore, Harnack inequality applied to (2.7) shows that  $w_\mu$  goes uniformly to zero in  $B(z^2, r)$ ,  $0 < r < 1$ , in contradiction with (2.8).

If  $w_\mu(0)$  stays bounded, either there is some  $z_\mu^i$ ,  $i > 2$ , such that  $|z_\mu^i|$  stays bounded and  $w_\mu(z_\mu^i)$  goes to infinity, and we can repeat the previous argument with  $z_\mu^i$  instead of 0, whence again a contradiction; or  $w_\mu$  stays bounded in any ball centered in 0. Then, elliptic theory shows that, up to a subsequence,  $w_\mu$  goes in  $C_{loc}^2(\mathbb{R}^3)$  to a limit  $w$  which satisfies  $-\Delta w = w^5$ ,  $w \geq 0$  in  $\mathbb{R}^3$ ,  $w \not\equiv 0$ ,  $\nabla w(0) = \nabla w(z^2) = 0$ . According to [13], such a  $w$  does not exist, hence again a contradiction. Therefore, 0 and  $z^2$  are two simple isolated blow-up points.

Up to a reindexation and passing to a subsequence, we may assume that for  $i \geq 2$ , either  $z_\mu^i \rightarrow z^i$ , or  $|z_\mu^i| \rightarrow \infty$  as  $\mu \rightarrow \infty$ . Because of (2.9),  $|z^i - z^j| \geq 1$  if  $i \neq j$ . If  $\bar{z}$  is a blow-up point for  $w_\mu$ ,  $\bar{z} = z^i$  for some index  $i$ , because of (2.10). Let  $S$  be the set of these blow-up points, which are isolated and simple (note

that  $d(z_\mu^i, \partial\tilde{\Omega}_\mu) \geq \rho/\delta_\mu \rightarrow \infty$ , and the equivalent of Prop. 2.3(i) holds for the solutions of (2.7), as [52] shows). We consider

$$\xi_\mu(z) = w_\mu(0)w_\mu(z)$$

which satisfies

$$-\Delta\xi_\mu + \delta_\mu^2\xi_\mu = w_\mu^4\xi_\mu, \quad \xi_\mu > 0 \text{ in } \tilde{\Omega}_\mu; \quad \frac{\partial\xi_\mu}{\partial\nu} = 0 \text{ on } \partial\tilde{\Omega}_\mu. \quad (2.12)$$

From (2.10-11) and Harnack inequality applied to (2.7), we know that  $w_\mu$  goes uniformly to zero in any compact set  $K \subset \mathbb{R}^3 \setminus S$  (note that  $K \subset \tilde{\Omega}_\mu$  for  $\mu$  large enough, since  $d(0, \partial\tilde{\Omega}_\mu) \geq \rho/\delta_\mu$ ). From (2.11) and Harnack inequality applied to (2.12), we know that  $\xi_\mu$  stays uniformly bounded in any compact set  $K \subset \mathbb{R}^3 \setminus S$ . Then, elliptic theory ensures that, along some subsequence,  $\xi_\mu$  converges in  $C_{loc}^2(\mathbb{R}^3 \setminus S)$  to a limit  $\xi$ , which is a positive regular harmonic function in  $\mathbb{R}^3 \setminus S$ . Therefore, we can write

$$\xi(z) = \frac{a}{|z|} + \frac{b}{|z - z^2|} + h(z)$$

with  $a \geq 0$ ,  $b \geq 0$  and  $h$  is regular positive harmonic function in  $\mathbb{R}^3 \setminus (S - \{0, z^2\})$ .  $0$  being a simple isolated blow-up point of  $(w_\mu)$ ,  $r \mapsto r^{1/2}\bar{\xi}_\mu(r)$  has a unique critical point in  $(0, r'_0)$ , and Prop. 2.2(ii) shows that this function has a maximum point which goes to zero as  $\mu$  goes to infinity. Therefore,  $r \mapsto r^{1/2}\bar{\xi}(r)$  is nonincreasing in  $(0, r'_0)$ . Then, either  $\xi \equiv 0$ , or  $a > 0$ . Integrating (2.12) on  $B_{r'_0}$ , we find

$$\int_{\partial B_{r'_0}} \frac{\partial\xi_\mu}{\partial\nu} + \delta_\mu^2 \int_{B_{r'_0}} \xi_\mu = w_\mu(0) \int_{B_{r'_0}} \xi_\mu^5.$$

From Prop. 2.2(ii), we have

$$\begin{aligned} w_\mu(0) \int_{B_{r'_0}} \xi_\mu^5 &\geq \frac{w_\mu(0)}{2} \int_{B(0, 2R/w_\mu^2(0))} \frac{w_\mu^5(0)}{(1 + \frac{w_\mu^4(0)|y|^2}{3})^{5/2}} dy \\ &\geq \frac{1}{2} \int_{B(0, 2R)} \frac{dy}{(1 + \frac{|y|^2}{3})^{5/2}} \\ &\geq \tau \end{aligned}$$

with  $\tau$  a strictly positive constant. On the other hand, (2.11) implies that

$$\delta_\mu^2 \int_{B_{r'_0}} \xi_\mu \leq C\delta_\mu^2 \int_{B_{r'_0}} \frac{dy}{|y|} = o(1)$$

as  $\mu$  goes to infinity. Moreover, if  $\xi \equiv 0$ , we have also

$$\int_{\partial B_{r'_0}} \frac{\partial\xi_\mu}{\partial\nu} = o(1)$$

hence a contradiction. Consequently,  $a > 0$ . In the same way  $b > 0$ . The classical Pohozaev identity for (2.12) provides us with the equality

$$-\delta_\mu^2 \int_{B_\sigma(0)} \xi_\mu^2 = \int_{\partial B_\sigma(0)} \left( \frac{1}{2} \xi_\mu \frac{\partial \xi_\mu}{\partial \nu} - \frac{\sigma}{2} |\nabla \xi_\mu|^2 + \sigma \left( \frac{\partial \xi_\mu}{\partial \nu} \right)^2 \right) - \frac{1}{2} \int_{\partial B_\sigma(0)} \left( \frac{\delta_\mu^2}{2} \xi_\mu^2 - \frac{1}{6w_\mu^4(0)} \xi_\mu^6 \right)$$

for any small  $\sigma > 0$ . As  $\delta_\mu$  goes to zero and  $w_\mu(0)$  goes to infinity, (2.11) implies that the left hand side, and the last integral on the right hand side, go to zero as  $\mu$  goes to infinity. In the same time, a straightforward computation shows that

$$\lim_{\sigma \rightarrow 0} \lim_{\mu \rightarrow \infty} \int_{\partial B_\sigma(0)} \left( \frac{1}{2} \xi_\mu \frac{\partial \xi_\mu}{\partial \nu} - \frac{\sigma}{2} |\nabla \xi_\mu|^2 + \sigma \left( \frac{\partial \xi_\mu}{\partial \nu} \right)^2 \right) \rightarrow -2\pi a(b + h(0)) < 0$$

whence a contradiction.

(ii), which is not necessary for our further purposes, may be proved in the same way, using the analysis of boundary blow-up points performed in [52].

**Remark.** If  $(u_\mu)$  is assumed to be bounded in  $H^1(\Omega)$ ,  $N_\mu$  is also bounded, since

$$\|u_\mu\|_{H^1(\Omega)} \geq N_\mu \tau$$

where  $\tau > 0$  is some fixed constant, as Prop. 2.1(ii) shows. Then, up to a reindexation and passing to a subsequence, we may assume that for  $\mu$  large enough

$$N_\mu = k_1 + k_2 = k \in \mathbb{N}^*$$

$$d(y_\mu^i, \partial\Omega_\mu) \geq \rho \quad \text{for some } \rho > 0 \quad 1 \leq i \leq k_1$$

$$d(y_\mu^i, \partial\Omega_\mu) \rightarrow 0 \quad \text{as } \mu \rightarrow \infty \quad k_1 + 1 \leq i \leq k$$

Setting  $\delta_\mu = \min_{\substack{1 \leq i, j \leq k_1 \\ i \neq j}} |y_\mu^i - y_\mu^j|$ , the same arguments as previously show the existence of  $\delta > 0$  such that  $|y_\mu^i - y_\mu^j| > \delta$ ,  $i \neq j$ ,  $1 \leq i, j \leq k_1$ .

## 2.2 Energy-dependent estimates

We turn now to an energy-dependent blow-up analysis of  $(u_\mu)$ , whose comparison with the previous results will provide us with the informations that we need to prove the theorem in the next section. First, as in [40], we define for  $\lambda \in \mathbb{R}_+^*$  and  $a \in \mathbb{R}^3$  the function

$$V_{\mu, \lambda, a}(x) = U_{\lambda, a}(x) - \varphi_{\mu, \lambda, a}(x) \quad x \in \mathbb{R}^3 \quad (2.13)$$

where  $U_{\lambda,a}$  is given by (1.3), i.e.  $U_{\lambda,a} = \lambda^{1/2}(1 + \lambda^2|x - a|^2)^{-1/2}$ , and

$$\varphi_{\mu,\lambda,a}(x) = \frac{1 - e^{-\mu^{1/2}|x-a|}}{\lambda^{1/2}|x-a|}. \quad (2.14)$$

$V_{\mu,\lambda,a}$ , which satisfies in  $\mathbb{R}^3$

$$-\Delta(3^{1/4}V_{\mu,\lambda,a}) + \mu(3^{1/4}V_{\mu,\lambda,a}) = (3^{1/4}V_{\mu,\lambda,a})^5 + \mu 3^{1/4}(U_{\lambda,a} - \frac{1}{\lambda^{1/2}|x-a|}) \quad (2.15)$$

is an improved approximate solution to  $(P_\mu)$  with respect to  $3^{1/4}U_{\lambda,a}$ , as  $\mu^{1/2}/\lambda$  goes to zero - see [40]. For  $\mu^{1/2}/\lambda$  small,  $\varphi_{\mu,\lambda,a}$  acts as a perturbation of  $U_{\lambda,a}$  in  $H^1(\Omega)$ , since

$$\|\varphi_{\mu,\lambda,a}\|_{H^1(\Omega)}^2 = O(\mu^{1/2}/\lambda) \quad (2.16)$$

as integral estimates show [40]. Now, we can state :

**Proposition 2.5** *Let  $(u_\mu)_{\mu \geq \mu_0}$  be a sequence of nonconstant solutions to  $(P_\mu)$ , bounded in  $H^1(\Omega)$ . There exist  $k \in \mathbb{N}^*$ ,  $(\lambda_\mu^i)$  and  $(a_\mu^i)$  sequences in  $\mathbb{R}_+^*$  and  $\bar{\Omega}$  respectively such that, for some subsequence*

$$\|u_\mu - 3^{1/4} \sum_{i=1}^k V_{\mu,\lambda_\mu^i,a_\mu^i}\|_{H^1(\Omega)} \rightarrow 0 \quad \text{as } \mu \rightarrow \infty \quad (2.17)$$

with

$$\mu^{1/2}/\lambda_\mu^i \rightarrow 0 \quad (2.18)$$

$$\lambda_\mu^i d(a_\mu^i, \partial\Omega) \rightarrow 0 \text{ or } \infty \quad (2.19)$$

$$\lambda_\mu^i/\lambda_\mu^j + \lambda_\mu^j/\lambda_\mu^i + \lambda_\mu^i \lambda_\mu^j |a_\mu^i - a_\mu^j|^2 \rightarrow \infty \quad \text{if } i \neq j \quad (2.20)$$

as  $\mu \rightarrow \infty$ .

Note that Proposition 2.5 holds for Palais-Smale sequences as well, whereas the previous one applies to exact solutions of the equation only. Such an analysis is performed for the first time in [8], and [11]. A proof of it, following Bahri's arguments, is given in Appendix B.

In view of Proposition 2.5 we may assume, extracting some subsequence, that each of the sequences  $(a_\mu^i)$  converges to a limit  $a^i \in \bar{\Omega}$ . It is easily seen that the  $a^i$ 's are exactly the blow-up points of this subsequence.

### 2.3 The shape of solutions with only interior blow-up points

Let us assume that all the  $a^i$ 's which occur in Proposition 2.5 lie in the interior of  $\Omega$ . In order to prove the theorem, we have to prove that such a case cannot occur.

Comparing Proposition 2.5 and Proposition 2.2, Prop. 2.2(ii) implies that to each  $x_\mu^i$  corresponds some  $a_\mu^{j(i)}$  such that,  $\varepsilon$  being small

$$\lambda_\mu^{j(i)}/u_\mu^2(x_\mu^i) \text{ is close to } 1; \quad \lambda_\mu^{j(i)}|a_\mu^{j(i)} - x_\mu^i| \text{ is close to } 0 \quad (2.21)$$

with  $j(i_1) \neq j(i_2)$  if  $i_1 \neq i_2$ .

Conversely, Prop. 2.2(v) implies that to each  $a_\mu^j$  corresponds some  $x_\mu^{i(j)}$  such that  $\lambda_\mu^j|a_\mu^j - x_\mu^{i(j)}|$  is bounded. We claim that for  $\mu$  large enough,  $i(j_1) \neq i(j_2)$  if  $j_1 \neq j_2$ .

Otherwise, up to subsequences and reindexations, we may assume that  $j = 1, \dots, p$  are the  $p$  indices,  $p \geq 2$ , such that  $i(j) = 1$ . We consider  $v_\mu$  defined by (2.1), with  $x_\mu = x_\mu^1$ . Through the change of variable

$$x = \frac{y}{\mu^{1/2}} + x_\mu^1$$

$x = x_\mu^1$  is sent to  $y = 0$ , and  $x = a_\mu^i$  is sent to  $y = \mu^{1/2}(a_\mu^i - x_\mu^1)$ . We know that 0 is an isolated blow-up point of  $(v_\mu)$ . Since  $x_\mu^1$  goes to  $a^1 = \dots = a^p$  which lies in the interior of  $\Omega$ ,  $d(0, \Omega_\mu) \rightarrow \infty$ , and Proposition 2.3 implies that 0 is a simple isolated blow-up point.

On the other hand, we notice that the  $\lambda_\mu^j$ 's are not of the same order as  $\mu$  goes to infinity, that is

$$\lambda_\mu^i/\lambda_\mu^j + \lambda_\mu^j/\lambda_\mu^i \rightarrow \infty \quad 0 \leq i < j \leq p.$$

Otherwise,  $\lambda_\mu^j|a_\mu^j - x_\mu^1|$  being bounded,  $\lambda_\mu^i\lambda_\mu^j|a_\mu^i - a_\mu^j|^2$  would also be bounded, and (2.20) could not be satisfied.

Finally, we notice that the boundedness of  $\lambda_\mu^j|a_\mu^j - x_\mu^1|$  implies, through (2.18), that  $\mu^{1/2}(a_\mu^j - x_\mu^1)$  goes to 0 as  $\mu$  goes to infinity. It follows that  $r \rightarrow r^{1/2}\bar{v}_\mu$ ,  $r = |y - x_\mu^1|$ , has several maximum points in any fixed interval  $(0, r_0)$ , for  $\mu$  large enough. This contradicts the fact that 0 is a simple blow-up point.

Once we know that there is a correspondance one to one between the  $x_\mu^i$ 's and the  $a_\mu^j$ 's, we infer from (2.21) and Proposition 2.4 that there exists  $\gamma > 0$  such that

$$|a_\mu^i - a_\mu^j| > \frac{\gamma}{\mu^{1/2}} \quad i \neq j$$

for  $\mu$  large enough. As a consequence, we know that a sequence  $(u_\mu)$ , bounded in  $H^1(\Omega)$ , of solutions to  $(P_\mu)$  whose all blow-up points lie in the interior of  $\Omega$  writes as

$$u_\mu = 3^{1/4} \sum_{i=1}^k V_{\mu, \lambda_\mu^i, a_\mu^i} + v_\mu \quad k \in \mathbb{N}^* \quad (2.22)$$

with

$$\frac{\mu^{1/2}}{\lambda_\mu^i} \rightarrow 0; \quad a_\mu^i \rightarrow a^i \in \Omega \quad |a_\mu^i - a_\mu^j| > \frac{\gamma}{\mu^{1/2}} \text{ if } i \neq j \quad (2.23)$$

and

$$v_\mu \rightarrow 0 \quad \text{in } H^1(\Omega) \quad (2.24)$$

as  $\mu$  goes to infinity.

We are going to prove, in the next section, that such a  $u_\mu$  cannot solve  $(P_\mu)$  for large  $\mu$  - hence the theorem.

### 3 Proof of the theorem

We adopt in this section a variational approach of the problem. We define the functional

$$J_\mu(u) = \frac{1}{2} \int_\Omega (|\nabla u|^2 + \mu u^2) - \frac{1}{6} \int_\Omega u^6 \quad u \in H^1(\Omega) \quad (3.1)$$

whose strictly positive critical points are exactly the solutions to  $(P_\mu)$ .

#### 3.1 A parametrization of the variational problem

This subsection is devoted to a parametrization of the variational problem in a neighbourhood of the eventual solutions to  $(P_\mu)$  defined by (2.22-24).  $k \in \mathbb{N}^*$  and  $\rho > 0$  being fixed, for  $\varepsilon > 0$  we set

$$\mathcal{V}_{\varepsilon, \mu} = \left\{ u \in H^1(\Omega) / \exists (\lambda_i)_{1 \leq i \leq k} \in (\mathbb{R}_+^*)^k, \mu^{1/2}/\lambda_i < \varepsilon, \exists (a_i)_{1 \leq i \leq k} \in (\Omega_\rho)^k, \right. \\ \left. |a_i - a_j| > \frac{\gamma}{4\mu^{1/2}} \text{ if } i \neq j, \text{ s.t. } |\nabla(u - 3^{1/4} \sum_{i=1}^k V_{\mu, \lambda_i, a_i})|_2 < \varepsilon \right\}$$

with  $\Omega_\rho = \{x \in \Omega, d(x, \partial\Omega) > \rho\}$ . Defining also

$$B_{\varepsilon, \mu} = \left\{ (\alpha, \lambda, a) \in \mathbb{R}^k \times (\mathbb{R}_+^*)^k \times (\Omega_{\rho-\varepsilon})^k \text{ s.t. } 3^{1/4}/2 < \alpha_i < 2 \cdot 3^{1/4}, \right. \\ \left. \mu^{1/2}/\lambda_i < \varepsilon, |a_i - a_j| > \frac{\gamma - \varepsilon}{4\mu^{1/2}} \text{ if } i \neq j \right\}$$

we have :

**Lemma 3.1** *There exist  $\mu_0 > 0$ ,  $\varepsilon_0 > 0$  such that for any  $\mu \geq \mu_0$ , any  $\varepsilon$ ,  $0 < \varepsilon \leq \varepsilon_0$ , and any  $u \in \mathcal{V}_{\varepsilon, \mu}$ , the infimum*

$$\inf_{(\alpha, \lambda, y) \in B_{4\varepsilon, \mu}} |\nabla(u - 3^{1/4} \sum_{i=1}^k V_{\mu, \lambda_i, a_i})|_2$$

*is achieved at only one point, which lies in  $B_{2\varepsilon, \mu}$ .*

Such a lemma is proved, for  $k = 1$ , in [40, App. A]. The result extends easily to the case  $k > 1$ , proceeding as in [8, Prop. 7][38, App. A]. Then, for  $\lambda \in (\mathbb{R}_+^*)^k$  and  $a \in \Omega^k$ , we set

$$E_{\lambda,a,\mu} = \left\{ v \in H^1(\Omega) / \int_{\Omega} \nabla v \cdot \nabla V_{\mu}^i = \int_{\Omega} \nabla v \cdot \nabla \frac{\partial V_{\mu}^i}{\partial \lambda_i} = \int_{\Omega} \nabla v \cdot \nabla \frac{\partial V_{\mu}^i}{\partial (a_i)_l} = 0 \right. \\ \left. 1 \leq i \leq k, 1 \leq l \leq 3 \right\} \quad (3.2)$$

with  $V_{\mu}^i = V_{\mu,\lambda_i,a_i}$ , for sake of simplicity. For  $\mu \geq \mu_0$ , Lemma 3.1 induces a map  $\Phi$  from the open subset  $\mathcal{V}_{\varepsilon_0,\mu}$  of  $H^1(\Omega)$  to the manifold

$$\mathcal{M}_{\mu} = \left\{ (\alpha, \lambda, a, v) \in \mathbb{R}^k \times (R_+^*)^k \times \Omega^k \times H^1(\Omega) \text{ s.t. } (\alpha, \lambda, a) \in B_{2\varepsilon_0,\mu}, \right. \\ \left. v \in E_{\lambda,a,\mu}, |\nabla v|_2 < \varepsilon \right\}$$

where  $(\alpha(u), \lambda(u), a(u))$  is the unique point in  $B_{2\varepsilon_0,\mu}$  at which the infimum of  $|\nabla(u - 3^{1/4} \sum_{i=1}^k V_{\mu,\lambda_i,a_i})|_2$  is achieved, and  $v(u) = u - \sum_{i=1}^k \alpha_i(u) V_{\mu,\lambda_i,a_i}(u)$ . This map is open, and induces a diffeomorphism between  $\mathcal{V}_{\varepsilon_0,\mu}$  and its image, which contains

$$\mathcal{N}_{\mu} = \left\{ (\alpha, \lambda, a, v) \in \mathbb{R}^k \times (R_+^*)^k \times \Omega^k \times H^1(\Omega) \text{ s.t. } |\alpha_i - 3^{1/4}| < \eta_0, \right. \\ \left. \mu^{1/2}/\lambda_i < \eta_0, |a_i - a_j| > \frac{\gamma}{2\mu^{1/2}} \text{ if } i \neq j, v \in E_{\lambda,a,\mu} \text{ and } |\nabla v|_2 < \varepsilon \right\} \quad (3.3)$$

for some  $\eta_0 > 0$  small enough. Setting

$$K_{\mu} : \quad \mathcal{N}_{\mu} \rightarrow \mathbb{R} \\ (\alpha, \lambda, a, v) \mapsto J_{\mu}(\sum_{i=1}^k \alpha_i V_{\mu,\lambda_i,a_i} + v) \quad (3.4)$$

we know that  $(\alpha, \lambda, a, v) \in \mathcal{N}_{\mu}$  is a critical point of  $K_{\mu}$  if and only if  $u = \sum_{i=1}^k \alpha_i V_{\mu,\lambda_i,a_i} + v$  is a critical point of  $J_{\mu}$ . Let us notice that, for  $\mu$  large enough,  $u_{\mu}$  given by (2.22-24) is in  $\mathcal{V}_{\varepsilon_0,\mu}$ . Moreover, setting

$$\Phi(u_{\mu}) = (\tilde{\alpha}_{\mu}, \tilde{\lambda}_{\mu}, \tilde{a}_{\mu}, \tilde{v}_{\mu})$$

it follows from [8,38, Lemma A.1] that

$$\tilde{\alpha}_{\mu}^i \rightarrow 3^{1/4} \quad \mu^{1/2}/\tilde{\lambda}_{\mu} \rightarrow 0 \quad \tilde{a}_{\mu}^i \rightarrow a_i \quad |\tilde{a}_{\mu}^i - \tilde{a}_{\mu}^j| > \frac{\gamma}{2\mu^{1/2}} \text{ if } i \neq j$$

and

$$\tilde{v}_{\mu} \rightarrow 0 \text{ in } H^1(\Omega).$$

In particular,  $\Phi(u_{\mu}) \in \mathcal{N}_{\mu}$  for  $\mu$  large enough. We are going to show that  $(\tilde{\alpha}_{\mu}, \tilde{\lambda}_{\mu}, \tilde{a}_{\mu}, \tilde{v}_{\mu})$  cannot be a critical point of  $K_{\mu}$  for  $\mu$  large enough, whence the theorem.

### 3.2 The $v$ -derivative of $K_\mu$

In this subsection, we estimate the  $H^1$ -norm of  $v_\mu$  as  $\mu$  goes to infinity. In view of (3.3-4), expanding  $K_\mu$  with respect to  $v$  in a neighbourhood of  $v = 0$ , we find

$$K_\mu(\alpha, \lambda, a, v) = K_\mu(\alpha, \lambda, a, 0) + f_{\alpha, \lambda, a, \mu}(v) + Q_{\alpha, \lambda, a, \mu}(v) + R_{\alpha, \lambda, a, \mu}(v) \quad (3.5)$$

with

$$f_{\alpha, \lambda, a, \mu}(v) = \mu \int_{\Omega} \left( \sum_{i=1}^k \alpha_i V_\mu^i \right) v - \int_{\Omega} \left( \sum_{i=1}^k \alpha_i V_\mu^i \right)^5 v \quad (3.6)$$

$$Q_{\alpha, \lambda, a, \mu}(v) = \frac{1}{2} \int_{\Omega} (|\nabla u|^2 + \mu u^2) - \frac{5}{2} \int_{\Omega} \left( \sum_{i=1}^k \alpha_i V_\mu^i \right)^4 v^2 \quad (3.7)$$

$$R_{\alpha, \lambda, a, \mu}(v) = O(\|v\|_{H^1(\Omega)}^3). \quad (3.8)$$

Moreover, choosing some  $\eta_0 > 0$  sufficiently small, there exist  $\kappa > 0$ ,  $\kappa' > 0$  such that for  $\mu$  large enough and any  $(\alpha, \lambda, a, v) \in \mathcal{N}_\mu$

$$\kappa \int_{\Omega} (|\nabla u|^2 + \mu u^2) \leq Q_{\alpha, \lambda, a, \mu}(v) \leq \kappa' \int_{\Omega} (|\nabla u|^2 + \mu u^2). \quad (3.9)$$

The second inequality is a direct consequence of Hölder inequality, Sobolev embedding theorem and estimate (C.3) in appendix. The coercivity property follows from [40, Lemma 3.2] in the case  $k = 1$ . The result extends to the case  $k > 1$  using the arguments of [7, Prop. 3.1], which are valid provided that  $\lambda_i \lambda_j |a_i - a_j|^2$ ,  $i \neq j$ , is large enough. But  $(\alpha, \lambda, a, v) \in \mathcal{N}_\mu$  implies that  $\lambda_i \lambda_j |a_i - a_j|^2 > \gamma^2 / 4\eta_0^2$ , whence the desired result choosing  $\eta_0$  small enough.

On the other hand, we claim that there exists  $C > 0$  such that, for  $\mu$  large enough and any  $(\alpha, \lambda, a, v) \in \mathcal{N}_\mu$ , we have

$$|f_{\alpha, \lambda, a, \mu}(v)| \leq \left( \frac{1}{\mu^{1/4} |\lambda|^{1/2}} + \frac{\mu^{1/2}}{|\lambda|} \right) \left( \frac{1}{2} \int_{\Omega} (|\nabla u|^2 + \mu u^2) \right)^{1/2} \quad (3.10)$$

with  $|\lambda| = (\sum_{i=1}^k \lambda_i^2)^{1/2}$ . Let us assume that the claim is true. Then, we deduce from (3.8-10) and the implicit functions theorem the following proposition :

**Proposition 3.1** *There exist  $\eta_1 > 0$ ,  $\eta_2 > 0$  such that, for  $\mu$  large enough, there exists a smooth map*

$$\begin{aligned} \tilde{\mathcal{N}}_\mu = \left\{ (\alpha, \lambda, a) \in \mathbb{R}^k \times (R_+^*)^k \times \Omega_p^k \text{ s.t. } |\alpha_i - 3^{1/4}| < \eta_1, \right. \\ \left. \begin{aligned} & \mu^{1/2} / \lambda_i < \eta_1, \quad |a_i - a_j| > \frac{\gamma}{2\mu^{1/2}} \text{ if } i \neq j \end{aligned} \right\} \rightarrow E_{\lambda, a, \mu} \\ (\alpha, \lambda, a) \mapsto \bar{v}_\mu(\alpha, \lambda, a) \end{aligned}$$

such that  $\bar{v}_\mu(\alpha, \lambda, a)$  is the unique point  $v \in E_{\lambda, a, \mu}$ ,  $|\nabla v|_2^2 + \mu |v|_2^2 < \eta_2$ , satisfying

$$\frac{\partial K_\mu}{\partial v}(\alpha, \lambda, a, \bar{v}_\mu(\alpha, \lambda, a)) = 0 \quad \text{in } T_{(\alpha, \lambda, a, \bar{v}_\mu)} \mathcal{N}_\mu. \quad (3.11)$$

Moreover, there exists  $C > 0$  such that

$$\int_{\Omega} |\nabla \bar{v}_{\mu}|^2 + \mu \int_{\Omega} \bar{v}_{\mu}^2 \leq C \left( \frac{1}{\mu^{1/2}|\lambda|} + \frac{\mu}{|\lambda|} \right). \quad (3.12)$$

We notice that (3.11) means that there exist  $(A, B, C) \in \mathbb{R}^k \times \mathbb{R}^k \times (\mathbb{R}^3)^k$  such that, for  $w \in H^1(\Omega)$

$$\begin{aligned} & \frac{\partial K_{\mu}}{\partial v}(\alpha, \lambda, a, \bar{v}_{\mu}) \cdot w \\ &= \sum_{i=1}^k \left( A_i \int_{\Omega} \nabla V_{\mu}^i \cdot \nabla w + B_i \int_{\Omega} \nabla \frac{\partial V_{\mu}^i}{\partial \lambda_i} \cdot \nabla w + \sum_{l=1}^3 C_{il} \int_{\Omega} \nabla \frac{\partial V_{\mu}^i}{\partial (a_i)_l} \cdot \nabla w \right). \end{aligned}$$

Taking respectively  $w = V_{\mu}^i, \frac{\partial V_{\mu}^i}{\partial \lambda_i}, \frac{\partial V_{\mu}^i}{\partial (a_i)_l}, 1 \leq i \leq k, 1 \leq l \leq 3$ , we see that the  $A_i, B_i, C_{il}$ 's solve a linear system which is nearly diagonal since, using the integral estimates in [7, 40] and Appendix C, we have

$$\begin{aligned} \int_{\Omega} \nabla V_{\mu}^i \cdot \nabla V_{\mu}^j &= \frac{3\pi^2}{4} \delta_{ij} + O\left(\frac{\mu^{1/2}}{\lambda_i^{1/2} \lambda_j^{1/2}}\right) \\ \int_{\Omega} \nabla V_{\mu}^i \cdot \nabla \frac{\partial V_{\mu}^j}{\partial \lambda_j} &= O\left(\frac{\mu^{1/2}}{\lambda_i^{1/2} \lambda_j^{3/2}}\right) \\ \int_{\Omega} \nabla V_{\mu}^i \cdot \nabla \frac{\partial V_{\mu}^j}{\partial (a_j)_l} &= O\left(\frac{\mu^{1/2} \lambda_j^{1/2}}{\lambda_i^{1/2}}\right) \\ \int_{\Omega} \nabla \frac{\partial V_{\mu}^i}{\partial \lambda_i} \cdot \nabla \frac{\partial V_{\mu}^j}{\partial \lambda_j} &= \frac{15\pi^2}{64} \frac{\delta_{ij}}{\lambda_i^2} + O\left(\frac{\mu^{1/2}}{\lambda_i^{3/2} \lambda_j^{3/2}}\right) \\ \int_{\Omega} \nabla \frac{\partial V_{\mu}^i}{\partial \lambda_i} \cdot \nabla \frac{\partial V_{\mu}^j}{\partial (a_j)_l} &= O\left(\frac{\mu^{1/2} \lambda_j^{1/2}}{\lambda_i^{3/2}}\right) \\ \int_{\Omega} \nabla \frac{\partial V_{\mu}^i}{\partial (a_i)_l} \cdot \nabla \frac{\partial V_{\mu}^j}{\partial (a_j)_m} &= \frac{15\pi^2}{64} \lambda_i^2 \delta_{ij} \delta_{lm} + O\left(\frac{\mu^{1/2} \lambda_j^{1/2}}{\lambda_i^{3/2}}\right). \end{aligned} \quad (3.13)$$

On the other hand

$$\begin{aligned} \frac{\partial K_{\mu}}{\partial v}(\alpha, \lambda, a, \bar{v}_{\mu}) \cdot V_{\mu}^i &= \frac{1}{\alpha_i} \frac{\partial K_{\mu}}{\partial \alpha_i}(\alpha, \lambda, a, \bar{v}_{\mu}) = O\left(|\alpha_i - 3^{1/4}| + \frac{\mu^{1/2}}{|\lambda|}\right) \\ \frac{\partial K_{\mu}}{\partial v}(\alpha, \lambda, a, \bar{v}_{\mu}) \cdot \frac{\partial V_{\mu}^i}{\partial \lambda_i} &= \frac{1}{\alpha_i} \frac{\partial K_{\mu}}{\partial \lambda_i}(\alpha, \lambda, a, \bar{v}_{\mu}) = O\left(\frac{\mu^{1/2}}{\lambda_i |\lambda|}\right) \\ \frac{\partial K_{\mu}}{\partial v}(\alpha, \lambda, a, \bar{v}_{\mu}) \cdot \frac{\partial V_{\mu}^i}{\partial (a_i)_l} &= \frac{1}{\alpha_i} \frac{\partial K_{\mu}}{\partial (a_i)_l}(\alpha, \lambda, a, \bar{v}_{\mu}) = O\left(\frac{\mu^{1/2} \lambda_i}{|\lambda|}\right) \end{aligned}$$

as it follows from Proposition C.1 in appendix. Then, solving the linear system

provides us with the estimates

$$\left. \begin{aligned} A_i &= O\left(|\alpha_i - 3^{1/4}| + \frac{\mu^{1/2}}{|\lambda|}\right) \\ B_i &= O\left(\frac{\mu^{1/2}\lambda_i}{|\lambda|}\right) \\ C_{il} &= O\left(\frac{\mu^{1/2}}{\lambda_i|\lambda|}\right) \end{aligned} \right\} \quad (3.14)$$

Before ending this subsection, we prove claim (3.10). From the proof of Lemma 3.1 in [40], we know that there exists  $C > 0$  such that, for  $\mu$  large enough and any  $(\alpha, \lambda, a, v) \in \mathcal{N}_\mu$

$$\begin{aligned} & \left| \mu \int_{\Omega} \alpha_i V_\mu^i v - \int_{\Omega} (\alpha_i V_\mu^i)^5 v \right| \\ & \leq C \left( \frac{1}{\mu^{1/4}|\lambda|^{1/2}} + \frac{\mu^{1/2}}{|\lambda|} \right) \left( \frac{1}{2} \int_{\Omega} (|\nabla u|^2 + \mu u^2) \right)^{1/2} \end{aligned} \quad (3.15)$$

$1 \leq i \leq k$ . On the other hand, if  $i \neq j$ , the Hölder inequality and the Sobolev embedding theorem yield

$$\int_{\Omega} (V_\mu^i)^4 V_\mu^j v \leq C \|v\|_{H^1(\Omega)} \left( \int_{\Omega} |V_\mu^i|^{24/5} |V_\mu^j|^{6/5} \right)^{5/6}. \quad (3.16)$$

(3.15-16) and estimate (C.16) in Appendix show that (3.10) is satisfied.

### 3.3 The $\alpha$ -derivative of $K_\mu$

For  $(\alpha, \lambda, a) \in \tilde{\mathcal{N}}_\mu$ , we set

$$\tilde{K}_\mu(\alpha, \lambda, a) = K_\mu(\alpha, \lambda, a, \bar{v}_\mu(\alpha, \lambda, a)).$$

From the definition (3.1) of  $E_{\lambda, a, \mu}$ , it follows that the partial derivative of  $\bar{v}_\mu$  with respect to  $\alpha_i$  is also in  $E_{\lambda, a, \mu}$ . Therefore, we deduce from (3.11) that

$$\frac{\partial \tilde{K}_\mu}{\partial \alpha_i}(\alpha, \lambda, a) = \frac{\partial K_\mu}{\partial \alpha_i}(\alpha, \lambda, a, \bar{v}_\mu(\alpha, \lambda, a))$$

that is, according to Proposition C.1 in appendix

$$\frac{\partial \tilde{K}_\mu}{\partial \alpha_i}(\alpha, \lambda, a) = \frac{\pi^2}{4} \alpha_i (3 - \alpha_i^4) + O\left(\frac{\mu^{1/2}}{|\lambda|}\right).$$

In the same way, we have

$$\frac{\partial^2 \tilde{K}_\mu}{\partial \alpha_i^2}(\alpha, \lambda, a) = \frac{\partial^2 K_\mu}{\partial \alpha_i^2}(\alpha, \lambda, a, \bar{v}_\mu(\alpha, \lambda, a))$$

whence, according to (C.22)

$$\frac{\partial^2 \tilde{K}_\mu}{\partial \alpha_i^2}(\alpha, \lambda, a) = \frac{\pi^2}{4}(3 - 5\alpha_i^4) + O\left(\frac{\mu^{1/2}}{|\lambda|}\right).$$

Consequently, we obtain :

**Proposition 3.2** *For  $\mu$  large enough, there exists a smooth map*

$$\begin{aligned} \tilde{\mathcal{N}}_\mu = \left\{ (\lambda, a) \in (R_+^*)^k \times \Omega_\rho^k \text{ s.t. } \mu^{1/2}/\lambda_i < \eta_1 \right. \\ \left. |a_i - a_j| > \frac{\gamma}{2\mu^{1/2}} \text{ if } i \neq j \right\} &\rightarrow \mathbb{R}_+^k \\ (\lambda, a) &\mapsto \bar{\alpha}_\mu(\lambda, a) \end{aligned}$$

such that  $\bar{\alpha}_\mu(\lambda, a)$  is the unique point  $\alpha \in \mathbb{R}_+^k$ ,  $|\alpha_i - 3^{1/4}| < \eta_1$ , satisfying

$$\frac{\partial \tilde{K}_\mu}{\partial \alpha}(\bar{\alpha}_\mu(\lambda, a), \lambda, a) = 0.$$

Moreover,  $\bar{\alpha}_\mu(\lambda, a)$  satisfies

$$\bar{\alpha}_\mu^i(\lambda, a) = 3^{1/4} + O\left(\frac{\mu^{1/2}}{|\lambda|}\right). \quad (3.17)$$

### 3.4 The $\lambda$ -derivative of $K_\mu$

This last subsection will provide us with the contradiction which proves the theorem. For  $(\lambda, a) \in \tilde{\mathcal{N}}_\mu$ , we set

$$\tilde{K}_\mu(\lambda, a) = \tilde{K}_\mu(\bar{\alpha}_\mu(\lambda, a), \lambda, a, \bar{v}_\mu(\bar{\alpha}_\mu(\lambda, a), \lambda, a)).$$

From (3.11) we know that

$$\begin{aligned} \frac{\partial \tilde{K}_\mu}{\partial \lambda_i}(\lambda, a) &= \frac{\partial \tilde{K}_\mu}{\partial \lambda_i}(\bar{\alpha}_\mu, \lambda, a, \bar{v}_\mu) \\ &+ \sum_{j=1}^k \left( A_j \int_{\Omega} \nabla V_\mu^j \cdot \nabla \frac{\partial \bar{v}_\mu}{\partial \lambda_i} + B_j \int_{\Omega} \nabla \frac{\partial V_\mu^j}{\partial \lambda_j} \cdot \nabla \frac{\partial \bar{v}_\mu}{\partial \lambda_i} + \sum_{l=1}^3 C_{jl} \int_{\Omega} \nabla \frac{\partial V_\mu^j}{\partial (a_i)_l} \cdot \nabla \frac{\partial \bar{v}_\mu}{\partial \lambda_i} \right). \end{aligned}$$

Since  $\bar{v}_\mu \in E_{\lambda, a, \mu}$ , we have

$$\begin{aligned} \int_{\Omega} \nabla V_\mu^j \cdot \nabla \frac{\partial \bar{v}_\mu}{\partial \lambda_i} &= - \int_{\Omega} \nabla \frac{\partial V_\mu^j}{\partial \lambda_i} \cdot \nabla \bar{v}_\mu = 0 \\ \int_{\Omega} \nabla \frac{\partial V_\mu^j}{\partial \lambda_j} \cdot \nabla \frac{\partial \bar{v}_\mu}{\partial \lambda_i} &= - \int_{\Omega} \nabla \frac{\partial^2 V_\mu^j}{\partial \lambda_j \partial \lambda_i} \cdot \nabla \bar{v}_\mu = O\left(\delta_{ij} \left| \nabla \frac{\partial^2 V_\mu^i}{\partial \lambda_i^2} \right|_2 |\nabla \bar{v}_\mu|_2\right) \\ \int_{\Omega} \nabla \frac{\partial V_\mu^j}{\partial (a_j)_l} \cdot \nabla \frac{\partial \bar{v}_\mu}{\partial \lambda_i} - \int_{\Omega} \nabla \frac{\partial^2 V_\mu^j}{\partial (a_j)_l \partial \lambda_i} \cdot \nabla \bar{v}_\mu &= O\left(\delta_{ij} \left| \nabla \frac{\partial^2 V_\mu^i}{\partial (a_i)_l \partial \lambda_i} \right|_2 |\nabla \bar{v}_\mu|_2\right). \end{aligned}$$

Then, (3.12,14) and estimates (C.8) yield

$$\frac{\partial \tilde{K}_\mu}{\partial \lambda_i}(\lambda, a) = \frac{\partial \tilde{K}_\mu}{\partial \lambda_i}(\bar{\alpha}_\mu, \lambda, a, \bar{v}_\mu) + O\left(\frac{1}{\lambda_i}\left(\frac{\mu^{1/4}}{|\lambda|^{3/2}} + \frac{\mu}{|\lambda|^2}\right)\right)$$

whence, according to (3.17) and (C.20)

$$\frac{\partial \tilde{K}_\mu}{\partial \lambda_i}(\lambda, a) = -2\pi \frac{\mu^{1/2}}{\lambda_i^2} + o\left(\frac{\mu^{1/2}}{\lambda_i^{3/2}|\lambda|^{1/2}}\right). \quad (3.18)$$

On the other hand, Propositions 3.1 and 3.2 show that, necessarily

$$\tilde{v}_\mu = \bar{v}_\mu(\tilde{\alpha}_\mu, \tilde{\lambda}_\mu, \tilde{a}_\mu) \quad \tilde{\alpha}_\mu = \bar{\alpha}_\mu(\tilde{\lambda}_\mu, \tilde{a}_\mu)$$

for  $\mu$  large enough where, according to Subsection 3.1,  $(\tilde{\alpha}_\mu, \tilde{\lambda}_\mu, \tilde{a}_\mu, \tilde{v}_\mu) = \Phi(u_\mu)$ . Moreover,  $u_\mu$  being a critical point of  $J_\mu$ ,  $(\tilde{\lambda}_\mu, \tilde{a}_\mu)$  satisfies

$$\frac{\partial \tilde{K}_\mu}{\partial \lambda_i}(\tilde{\lambda}_\mu, \tilde{a}_\mu) = 0 \quad 1 \leq i \leq k. \quad (3.19)$$

However, up to a subsequence and a reindexation, we may assume, without loss of generality, that  $\tilde{\lambda}_\mu^1 = \min_{1 \leq i \leq k} \tilde{\lambda}_\mu^i$  for  $\mu$  large enough. Then, (3.18) implies that

$$\frac{\partial \tilde{K}_\mu}{\partial \lambda_1}(\tilde{\lambda}_\mu, \tilde{a}_\mu) = -2\pi \frac{\mu^{1/2}}{(\tilde{\lambda}_\mu^1)^2} + o\left(\frac{\mu^{1/2}}{(\tilde{\lambda}_\mu^1)^2}\right)$$

in contradiction with (3.19). This completes the proof of the theorem.

**Remarks.**

(1) If  $a_i$  were on the boundary of  $\Omega$ , some additional term would occur in (3.18), involving the mean curvature of the frontier at  $a_i$ , and (3.19) would not lead to a contradiction - see [40].

(2) If, instead of  $(P_\mu)$ , we consider the subcritical problem

$$-\Delta u + \mu u = u^{5-\varepsilon_\mu}, \quad u > 0 \text{ in } \Omega; \quad \frac{\partial u}{\partial \nu} = 0 \text{ on } \partial\Omega$$

$\varepsilon_\mu > 0$ ,  $\varepsilon_\mu \ln \mu \rightarrow 0$  as  $\mu \rightarrow \infty$ , and the corresponding modified functional, an additional term would occur in (3.18) which, up to a strictly positive constant, would be equal to  $\varepsilon_\mu/\lambda_i$  - see [9, 39]. Then, the derivative of  $\tilde{K}_\mu$  with respect to  $\lambda_i$  would vanish for some  $\lambda_i \sim \mu^{1/2}/\varepsilon_\mu$ . Therefore, the obstruction to the fulfillment of (3.19) disappears in the subcritical case, in accordance with the known results.

(3) In [14] are considered problems as

$$-\Delta u + \mu u = u^{(n+2)/(n-2)} + a(x)u^q, \quad u > 0 \text{ in } \Omega; \quad \frac{\partial u}{\partial \nu} = 0 \text{ on } \partial\Omega$$

$\Omega \in \mathbb{R}^n$ ,  $n \geq 3$ ,  $1 < q < (n+2)/(n-2)$ . We notice that the additional term with respect to  $(P_\mu)$  would introduce in (3.18) a quantity as  $a(a_i)/\lambda_i^{(q+3)/2}$  if  $q < 2$ ,  $a(a_i)(\ln \lambda_i)/\lambda_i^{5/2}$  if  $q = 2$ ,  $a(a_i)/\lambda_i^{(7-q)/2}$  if  $q > 2$ . Then, we see that (3.19) could be satisfied for  $a(a_i) > 0$  and  $q > 3$ . This agrees with Theorem 1.1 in [14].

(4) In the special case  $k = 1$ , the result could be easily derived from [40]. Namely, considering the functional

$$I_\mu(u) = \frac{\int_\Omega (|\nabla u|^2 + \mu u^2)}{(\int_\Omega u^6)^{1/3}}$$

[40] provides us with the following expansion for the equivalent of  $\tilde{K}_\mu$ , that we denote in the same way :

$$\begin{aligned} \frac{\partial \tilde{K}_\mu}{\partial \lambda}(\lambda, a) &= 2\pi^{1/3} \left( -2\frac{\mu^{1/2}}{\lambda^2} + \frac{H(y)}{\lambda^2} \left( \ln \frac{\lambda}{2\mu^{1/2}} - \gamma - \frac{1}{2} \right) \right) \\ &\quad + O \left( \frac{1}{\lambda^2 \mu^{1/2}} + \frac{\mu}{\lambda^3} + \frac{\mu^{1/2}}{\lambda^3} \ln \frac{\lambda}{\mu^{1/2}} \right) \end{aligned}$$

for a concentration point  $a$  on the boundary. If  $a$  lies in the interior of the domain, similar and easier computations would have given

$$\frac{\partial \tilde{K}_\mu}{\partial \lambda}(\lambda, a) = -4\pi^{1/3} \frac{\mu^{1/2}}{\lambda^2} + O \left( \frac{1}{\lambda^2 \mu^{1/2}} \frac{\mu}{\lambda^3} \right)$$

which cannot vanish for large  $\lambda$ .

For  $n \geq 5$ , computations in [38] provide us with the expansion

$$\frac{\partial \tilde{K}_\mu}{\partial \lambda_i}(\lambda, a) = C_n \frac{H(a_i)}{\lambda_i^2} - C'_n \frac{\mu}{\lambda_i^3} + \text{lower order terms}$$

(with  $C_n$  and  $C'_n$  strictly positive constant) when the concentrations points are on the boundary. For interior points we would have

$$\frac{\partial \tilde{K}_\mu}{\partial \lambda_i}(\lambda, a) = -2C'_n \frac{\mu}{\lambda_i^3} + \text{lower order terms.}$$

Again, these quantities cannot vanish for large  $\lambda_i$ 's, whence the equivalent of the Theorem under the assumption that the concentration points stay far from each other - as in [14].

Treating the case  $n = 4$  in the same way would require, in order to obtain convenient expansions, to consider suitable approximate solutions, as we did in the case  $n = 3$ .

## APPENDIX

### A Proof of Proposition 2.1

The proof relies on the following lemma :

**Lemma A.1** *Let  $(u_\mu)_{\mu \geq \mu_0}$  be a sequence of nonconstant solutions to  $(P_\mu)$ . Let  $\varepsilon > 0$ ,  $R > 1$ . There exists a constant  $C_0 = C_0(R)$  such that, for  $\mu$  large enough and any compact set  $K$*

$$\max_{x \in \bar{\Omega} \setminus K} d(x, K)^{1/2} u_\mu(x) \geq C_0$$

(with  $d(x, K) = 1$  if  $K = \emptyset$ ) implies the existence of  $x_\mu$ , local maximum point of  $u_\mu$  in  $\bar{\Omega}$ , such that

$$\begin{aligned} (i) \quad & \frac{\mu^{1/4}}{u_\mu(x_\mu)} < \varepsilon \\ (ii) \quad & \left\| \frac{1}{u_\mu(x_\mu)} u_\mu \left( \frac{x}{u_\mu^2(x_\mu)} + x_\mu \right) - \frac{1}{(1 + \frac{|x|^2}{3})^{1/2}} \right\|_{C^2(B(0, 2R) \cap \Omega_\mu)} < \varepsilon \\ & \text{with } \Omega_\mu = u_\mu^2(x_\mu)(\Omega - x_\mu) \\ (iii) \quad & u_\mu^2(x_\mu) d(x_\mu, K) > R \\ & x_\mu \in \partial\Omega \quad \text{or} \quad u_\mu^2(x_\mu) d(x_\mu, \partial\Omega) > \frac{1}{\varepsilon}. \end{aligned}$$

**Proof of Proposition 2.1.** Lemma A.1, applied with  $K = \emptyset$ , provides us with  $x_\mu = x_\mu^1$  such that (i) (ii) (iii) of Proposition 3.1 are satisfied. Then, we set

$$K = \bar{B}(x_\mu^1, r_\mu^1) \quad \text{with } r_\mu^1 = \frac{R}{u_\mu^2(x_\mu^1)}.$$

If  $\max_{x \in \bar{\Omega} \setminus K} d(x, K)^{1/2} u_\mu(x) < C_0$ , there is nothing more to prove. If not, Lemma A.1 provides us with a new  $x_\mu = x_\mu^2$  such that (i) (ii) (iii) of Proposition 2.1 are again satisfied, and  $\bar{B}(x_\mu^1, r_\mu^1) \cap \bar{B}(x_\mu^2, r_\mu^2) = \emptyset$ , with  $r_\mu^2 = \frac{R}{u_\mu^2(x_\mu^2)}$ . The process must stop after a finite number  $N_\mu$  of steps since, because of (ii)

$$\int_{B(x_\mu^i, r_\mu^i)} |\nabla u_\mu|^2 \geq \tau > 0$$

with  $\tau$  a constant which does not depend on  $i, \mu$ . (Note that if  $(u_\mu)$  is assumed to be bounded in  $H^1(\Omega)$ ,  $N_\mu$  is also bounded as  $\mu$  goes to infinity.) We have

$$\max_{x \in \bar{\Omega} \setminus \cup_{i=1}^{N_\mu} B(x_\mu^i, r_\mu^i)} \left( d(x, \cup_{i=1}^{N_\mu} B(x_\mu^i, r_\mu^i)) \right)^{1/2} u_\mu(x) < C_0.$$

Consequently, if  $x \notin \cup_i B(x_\mu^i, 2r_\mu^i)$

$$\left( d(x, \{x_\mu^i, 1 \leq i \leq N_\mu\}) \right)^{1/2} u_\mu(x) \leq \left( 2d(x, \cup_i B(x_\mu^i, r_\mu^i)) \right)^{1/2} u_\mu(x) < \sqrt{2}C_0$$

and, if  $x \in \cup_i B(x_\mu^i, 2r_\mu^i)$ , using (ii)

$$\begin{aligned} & \left( d(x, \{x_\mu^i, 1 \leq i \leq N_\mu\}) \right)^{1/2} u_\mu(x) \\ & \leq |x - x_\mu^i|^{1/2} u_\mu(x) \\ & \leq |x - x_\mu^i|^{1/2} u_\mu(x) \left( \left( 1 + \frac{1}{3} u_\mu^4(x_\mu^i) |x - x_\mu^i|^2 \right)^{-1/2} + \varepsilon \right) \\ & \leq 2^{-1/2} \mathfrak{S}^{1/4} + 2^{1/2} \varepsilon R^{1/2}. \end{aligned}$$

Therefore, (v) is satisfied and the proof of Proposition 2.1 is complete.

**Proof of Lemma A.1.** Arguing by contradiction, we assume that there exist a sequence  $(u_\mu)$  of nonconstant solutions  $(P_\mu)$  and a sequence  $(K_\mu)$  of compact sets such that

$$\max_{x \in \bar{\Omega} \setminus K_\mu} d(x, K_\mu)^{1/2} u_\mu(x) \rightarrow \infty \quad (\text{with } d(x, K_\mu) = 1 \text{ if } K_\mu = \emptyset)$$

and there is no  $x_\mu$ , as specified in the lemma. Let  $\tilde{x}_\mu \in \Omega \setminus K_\mu$  be a global maximum point of  $d(x, K_\mu)^{1/2} u_\mu(x)$  in  $\bar{\Omega} \setminus K_\mu$ . We set

$$v_\mu(x) = \frac{1}{u_\mu(\tilde{x}_\mu)} u_\mu\left(\frac{x}{u_\mu^2(\tilde{x}_\mu)} + \tilde{x}_\mu\right) \quad x \in \tilde{\Omega}_\mu = u_\mu^2(\tilde{x}_\mu)(\Omega - \tilde{x}_\mu)$$

and

$$R_\mu = \frac{1}{4} d(\tilde{x}_\mu, K_\mu) u_\mu^2(\tilde{x}_\mu).$$

By assumption,  $R_\mu \rightarrow \infty$  as  $\mu \rightarrow \infty$ . Since

$$d\left(\frac{x}{u_\mu^2(\tilde{x}_\mu)} + \tilde{x}_\mu, K_\mu\right) \geq \frac{1}{2} d(\tilde{x}_\mu, K_\mu) \quad \forall x \in B(0, R_\mu)$$

we have, for  $x \in B(0, R_\mu) \cap \tilde{\Omega}_\mu$

$$\left( \frac{1}{2} d(\tilde{x}_\mu, K_\mu) \right)^{1/2} v_\mu(x) \leq (d(\tilde{x}_\mu, K_\mu))^{1/2}$$

whence

$$v_\mu(x) \leq \sqrt{2} \quad \forall x \in B(0, R_\mu).$$

On the other hand,  $v_\mu$  satisfies

$$-\Delta v_\mu + \frac{\mu}{u_\mu^4(\tilde{x}_\mu)} v_\mu = v_\mu^5, \quad v_\mu > 0 \text{ in } \tilde{\Omega}_\mu; \quad \frac{\partial v_\mu}{\partial \nu} = 0 \text{ on } \partial \tilde{\Omega}_\mu.$$

Let us assume that  $\mu u_\mu^{-4}(\tilde{x}_\mu)$  goes to infinity as  $\mu$  goes to infinity.  $w_\mu$  defined as

$$w_\mu = v_\mu\left(\frac{x}{\alpha_\mu}\right) \quad x \in \alpha_\mu \tilde{\Omega}_\mu \quad \text{with} \quad \alpha_\mu = \frac{\mu^{1/2}}{u_\mu^2(\tilde{x}_\mu)}$$

satisfies

$$-\Delta w_\mu + w_\mu = \frac{1}{\alpha_\mu^2} w_\mu^5, \quad w_\mu > 0 \text{ in } \alpha_\mu \tilde{\Omega}_\mu; \quad \frac{\partial w_\mu}{\partial \nu} = 0 \text{ on } \partial(\alpha_\mu \tilde{\Omega}_\mu)$$

and

$$w_\mu(x) \leq \sqrt{2} \quad \forall x \in B(0, \alpha_\mu R_\mu).$$

Elliptic theory shows that, up to a subsequence and a space rotation,  $w_\mu$  converges in  $C_{loc}^2(\mathbb{R}_T^3)$  to a limit  $w$  which satisfies

$$-\Delta w + w = 0, \quad 0 \leq w \leq \sqrt{2} \text{ in } \mathbb{R}_T^3; \quad \frac{\partial w}{\partial \nu} = 0 \text{ on } \partial \mathbb{R}_T^3$$

with

$$\mathbb{R}_T^3 = \{x = (x_1, x_2, x_3) \in \mathbb{R}^3, x_3 > -T\} \quad 0 \leq T \leq \infty.$$

This implies  $w \equiv 0$ , in contradiction with  $w(0) = 1$ . Therefore,  $\mu u_\mu^{-4}(\tilde{x}_\mu)$  is bounded.

Up to a subsequence, we can assume

$$\mu u_\mu^{-4}(\tilde{x}_\mu) \rightarrow \theta \quad \text{as } \mu \rightarrow \infty \quad 0 \leq \theta < \infty$$

and, up to a space rotation,  $v_\mu$  converges in  $C_{loc}^2(\mathbb{R}_S^3)$  to a limit  $v$  which satisfies

$$-\Delta v + \theta v = v^5, \quad 0 \leq v \leq \sqrt{2} \text{ in } \mathbb{R}_S^3; \quad \frac{\partial v}{\partial \nu} = 0 \text{ on } \partial \mathbb{R}_S^3.$$

Note that if  $S < \infty$ ,  $v$  may be continued by reflection in a bounded solution of the equation in whole  $\mathbb{R}^3$ . As a consequence, if  $\theta > 0$ ,  $v$  is constant, i.e.  $v \equiv 0$  or  $v = \theta^{1/4}$ . Since  $v(0) = 1$ ,  $\theta = 1$  and  $v \equiv 1$ . This implies that  $u_\mu$  is constant for  $\mu$  large enough, in contradiction with the initial assumption.

Before proving this fact, let us complete the proof of the lemma. If  $\theta = 0$ , we know that  $v$  writes as - see [13]

$$v(x) = \frac{\lambda^{1/2}}{(1 + \lambda^2 \frac{|x-x_0|^2}{3})^{1/2}} \quad \text{for some } x_0 \in \mathbb{R}^3, \lambda \in \mathbb{R}_+^*.$$

$v \leq \sqrt{2}$  and  $v(0) = 1$  imply that

$$\lambda \leq \sqrt{2} \quad \lambda^{1/2} (1 + \lambda^2 \frac{|x_0|^2}{3})^{-1/2} = 1$$

whence

$$1 \leq \lambda \leq \sqrt{2} \quad |x_0|^2 = 3(\lambda - 1)/\lambda^2 \leq 3/4.$$

Moreover, if  $S < \infty$ , necessarily  $x_0 \in \partial\mathbb{R}_S^3$ . From the shape of  $v$  and the convergence of  $v_\mu$  to  $v$  in  $C_{loc}^2(\mathbb{R}_S^3)$ , we deduce the existence, for large  $\mu$ , of a local maximum point  $z_\mu$  of  $v_\mu$ , which goes to  $x_0$  as  $\mu$  goes to infinity. Then

$$x_\mu = \frac{z_\mu}{u_\mu^2(\tilde{x}_\mu)} + \tilde{x}_\mu$$

is a local maximum point of  $u_\mu$ , and

$$\frac{u_\mu(x_\mu)}{u_\mu(\tilde{x}_\mu)} = v_\mu(z_\mu) \rightarrow \lambda^{1/2} \in [1, \sqrt{2}]$$

whence  $\mu u_\mu^{-4}(\tilde{x}_\mu) \rightarrow 0$  as  $\mu \rightarrow \infty$ . Moreover,  $|x_\mu - \tilde{x}_\mu| \leq \sqrt{3}u_\mu^{-2}(\tilde{x}_\mu)$  for  $\mu$  large enough. As  $d(\tilde{x}_\mu, K_\mu)u_\mu^2(\tilde{x}_\mu) \rightarrow \infty$ , we know that for large  $\mu$ ,  $x_\mu \in \bar{\Omega} \setminus K_\mu$ , and  $d(x_\mu, K_\mu)u_\mu^2(x_\mu) \rightarrow \infty$  as  $\mu$  goes to infinity.

Then, we can repeat the same argument as previously, with  $x_\mu$  instead of  $\tilde{x}_\mu$ . In this case 0, local maximum point of  $v_\mu$ , is a critical point of  $v$ . As the only critical point of  $v$  is  $x_0$ , we obtain  $x_0 = 0$ , whence  $S = 0$  or  $S = \infty$ , i.e.

$$u_\mu^2(x_\mu)d(x_\mu, \partial\Omega) \rightarrow 0 \quad \text{or} \quad u_\mu^2(x_\mu)d(x_\mu, \partial\Omega) \rightarrow \infty.$$

$v(0) = 1$  gives  $\lambda = 1$ , and the convergence of  $v_\mu$  to  $v$  in  $C_{loc}^2(\mathbb{R}_S^3)$

$$\left\| \frac{1}{u_\mu(x_\mu)} u_\mu\left(\frac{x}{u_\mu^2(x_\mu)} + x_\mu\right) - \frac{1}{(1 + \frac{|x|^2}{3})^{1/2}} \right\|_{C^2(B(0,2R) \cap \Omega_\mu \cap \mathbb{R}_S^3)} = o(1) \quad (\text{A.1})$$

for any  $R > 0$ , with  $\Omega_\mu = u_\mu^2(x_\mu)(\Omega - x_\mu)$ .

Actually, (A.1) holds in  $C^2(B(0,2R) \cap \Omega_\mu)$ . Indeed, if  $S = 0$ , up to a subsequence  $x_\mu$  goes to some  $\bar{x} \in \partial\Omega$  as  $\mu$  goes to infinity. Up to a space rotation, we may assume that the tangent space to  $\partial\Omega$  at  $\bar{x}$  is parallel to the plane  $x_3 = 0$ . There exist  $\mathcal{U}$  neighbourhood of  $\bar{x}$  in  $\mathbb{R}^3$ ,  $r > 0$  and  $\Psi : \mathcal{U} \rightarrow B(0, R) \subset \mathbb{R}^3$  a  $C^2$ -diffeomorphism such that

$$\Psi(\mathcal{U} \cap \Omega) = B^+(0, r) = \{x \in \mathbb{R}^3, |x| < r, x_3 > 0\}$$

and

$$\Psi(\bar{x}) = 0 \quad \nabla\Psi(\bar{x}) = 0.$$

Then, for any  $r'$ ,  $0 < r' < r$

$$\bar{v}_\mu(y) = \frac{1}{u_\mu(x_\mu)} u_\mu\left(\Psi^{-1}\left(\frac{y}{u_\mu^2(x_\mu)} + y_\mu\right)\right) \quad \text{with } y_\mu = \Psi(x_\mu)$$

satisfies, for  $\mu$  large enough

$$-\sum_{i,j} a_{ij}(y) \frac{\partial^2 \bar{v}_\mu}{\partial y_i \partial y_j} + \sum_i b_i(y) \frac{\partial \bar{v}_\mu}{\partial y_i} + \frac{\mu}{u_\mu^4(x_\mu)} \bar{v}_\mu = v_\mu^5$$

in  $B(0, r') \cap \{y_3 > -u_\mu^2(x_\mu)(y_\mu)_3\}$ , and

$$\frac{\partial \bar{v}_\mu}{\partial y_3} = 0$$

on  $B(0, r') \cap \{y_3 = -u_\mu^2(x_\mu)(y_\mu)_3\}$ , with

$$a_{ij}(y) = \delta_{ij} + o(1) \quad b_i(y) = o(1) \quad \text{uniformly as } \mu \rightarrow \infty.$$

$\bar{v}_\mu$ , continued by reflection in  $B(0, r'/2)$ , satisfies in  $B(0, r'/2)$  the same equation. The same arguments as previously show that the continuation of  $\bar{v}_\mu$  converges in  $C_{loc}^2(\mathbb{R}^3)$  to  $v(y) = (1 + |y|^2/3)^{-1/2}$ . Consequently, (A.1) holds in  $C^2(B(0, 2R) \cap \Omega_\mu)$ , as announced.

Finally, we show that if  $S = 0$ ,  $x_\mu \in \partial\Omega$  for  $\mu$  large enough. Indeed, assume that  $x_\mu \notin \partial\Omega$ . Since  $x_\mu$  is a local maximum of  $v_\mu$ ,  $y_\mu = \Psi(x_\mu)$  is a local maximum of  $\bar{v}_\mu$ , whence  $\nabla \bar{v}_\mu(y_\mu) = 0$ . Let  $y'_\mu$  be the orthogonal projection of  $y_\mu$  on the plane  $x_3 = 0$ . We have

$$\frac{\partial \bar{v}_\mu}{\partial y_3}(y_\mu) = \frac{\partial \bar{v}_\mu}{\partial y_3}(y'_\mu) = 0$$

Consequently, there exist  $y''_\mu = ty_\mu + (1-t)y'_\mu$ ,  $0 < t < 1$ , such that  $\frac{\partial^2 \bar{v}_\mu}{\partial y_3^2}(y''_\mu) \geq 0$ . On the other hand

$$\frac{\partial^2 \bar{v}_\mu}{\partial y_3^2}(y''_\mu) \rightarrow \frac{\partial^2 v}{\partial y_3^2}(0) < 0$$

according to the shape of  $v$ , whence a contradiction.

To complete the proof of the lemma, it only remains to prove that

$$\theta_\mu = \mu u_\mu^{-4}(\tilde{x}_\mu) \rightarrow 1 \quad \text{as } \mu \rightarrow \infty$$

implies that  $u_\mu \equiv \mu^{1/4} \theta_\mu^{-1/4}$  for large  $\mu$ , in contradiction with the assumption that  $u_\mu$  is nonconstant. Let  $a_\mu$  be a point in the closure of  $B(0, R_\mu/2) \cap \tilde{\Omega}_\mu$ , such that

$$|v_\mu(a_\mu) - \theta_\mu^{1/4}| = \|v_\mu - \theta_\mu^{1/4}\|_{L^\infty(B(0, R_\mu/2) \cap \tilde{\Omega}_\mu)}.$$

$\tilde{v}_\mu$  defined as  $\tilde{v}_\mu = v_\mu(x + a_\mu)$ , satisfies

$$-\Delta \tilde{v}_\mu + \theta_\mu \tilde{v}_\mu = \tilde{v}_\mu^5, \quad \tilde{v}_\mu > 0 \text{ in } \tilde{\Omega}_\mu - a_\mu; \quad \frac{\partial \tilde{v}_\mu}{\partial \nu} = 0 \text{ on } \partial(\tilde{\Omega}_\mu - a_\mu)$$

and

$$\tilde{v}_\mu(x) \leq \sqrt{2} \quad \forall x \in B(0, R_\mu/2) \cap (\tilde{\Omega}_\mu - a_\mu)$$

since

$$\left( B(0, R_\mu/2) \cap (\tilde{\Omega}_\mu - a_\mu) \right) \subset \left( (B(0, R_\mu) \cap (\tilde{\Omega}_\mu - a_\mu)) - a_\mu \right).$$

Up to a subsequence and a space rotation,  $\tilde{v}_\mu$  converges in  $C_{loc}^2(\mathbb{R}_{T'}^3)$ ,  $0 \leq T' \leq \infty$ , to a limit  $\tilde{v}$  which satisfies

$$-\Delta \tilde{v} + \tilde{v} = \tilde{v}^5, \quad 0 \leq \tilde{v} \leq \sqrt{2} \text{ in } \mathbb{R}_{T'}^3; \quad \frac{\partial \tilde{v}}{\partial \nu} = 0 \text{ on } \partial \mathbb{R}_{T'}^3.$$

As a consequence,  $\tilde{v} \equiv 0$  or  $\tilde{v} \equiv 1$ . Therefore,  $|v_\mu(a_\mu) - \theta_\mu^{1/4}| = \|v_\mu - \theta_\mu^{1/4}\|_{L^\infty(B(0, R_\mu/2) \cap \tilde{\Omega}_\mu)}$  goes to a limit  $l$ ,  $l = 1$  if  $\tilde{v} \equiv 0$ , and  $l = 0$  if  $\tilde{v} \equiv 1$ .

Let us assume that  $l = 1$ . In this case,  $v_\mu(a_\mu) = \tilde{v}_\mu(0)$  goes to zero as  $\mu$  goes to infinity. We also know that  $v_\mu(0) = 1$ , whence the existence, for  $\mu$  large enough, of  $b_\mu \in (B(0, R_\mu/2) \cap \tilde{\Omega}_\mu)$  such that  $v_\mu(b_\mu) = 1/2$ . Setting  $\tilde{v}'_\mu(x) = v_\mu(x + b_\mu)$ , and repeating the same argument as above, we see that up to a subsequence and a space rotation,  $\tilde{v}'_\mu$  converges in  $C_{loc}^2(\mathbb{R}_{T''}^3)$ ,  $0 \leq T'' \leq \infty$ , to a limit  $\tilde{v}'$ . As previously,  $\tilde{v}'$  has to be identically 0 or 1, in contradiction with  $\tilde{v}'(0) = 1/2$ . Therefore,  $l = 0$ .

Finally, let us assume that  $v_\mu \not\equiv \theta_\mu^{1/4}$  in  $B(0, R_\mu/2) \cap \tilde{\Omega}_\mu$ . We set

$$\tilde{\tilde{v}}_\mu(x) = \frac{\tilde{v}_\mu(x) - \theta_\mu^{1/4}}{\tilde{v}_\mu(0) - \theta_\mu^{1/4}}$$

which satisfies

$$-\Delta \tilde{\tilde{v}}_\mu + \theta_\mu \tilde{\tilde{v}}_\mu = \tilde{\tilde{v}}_\mu(\theta_\mu^4 + \theta_\mu^3 \tilde{\tilde{v}}_\mu + \theta_\mu^2 \tilde{\tilde{v}}_\mu^2 + \theta_\mu \tilde{\tilde{v}}_\mu^3 + \tilde{\tilde{v}}_\mu^4) \quad \text{in } \tilde{\Omega}_\mu - a_\mu$$

and

$$|\tilde{\tilde{v}}_\mu| \leq 1 \quad \text{in } B(0, R_\mu/2) \cap (\tilde{\Omega}_\mu - a_\mu).$$

Therefore, up to a subsequence,  $\tilde{\tilde{v}}_\mu$  converges in  $C_{loc}^2(\mathbb{R}_{T'}^3)$  to a limit  $\tilde{\tilde{v}}$  which satisfies

$$-\Delta \tilde{\tilde{v}} = 4\tilde{\tilde{v}} \text{ in } \mathbb{R}_{T'}^3; \quad \frac{\partial \tilde{\tilde{v}}}{\partial \nu} = 0 \text{ on } \partial \mathbb{R}_{T'}^3,$$

and  $\|\tilde{\tilde{v}}_\mu\|_{L^\infty(\mathbb{R}_{T''}^3)} \leq 1$ . It follows that  $\tilde{\tilde{v}} \equiv 0$ , in contradiction with  $|\tilde{\tilde{v}}(0)| = 1$ . Therefore,  $v_\mu$  is constant in  $B(0, R_\mu/2) \cap \tilde{\Omega}_\mu$  and, actually, in whole  $\tilde{\Omega}_\mu$ . This means that  $u_\mu$  is constant in  $\Omega$ , a contradiction.

## B Proof of Proposition 2.5

Let  $u_\mu$  be a sequence, bounded in  $H^1(\Omega)$ , of solutions to  $(P_\mu)$ . On one hand, we have

$$\int_{\Omega} (|\nabla u_\mu|^2 + \mu u_\mu^2) = \int_{\Omega} u_\mu^6. \quad (\text{B.1})$$

On the other hand, the continuous embedding of  $H^1(\Omega)$  into  $L^6(\Omega)$  yields

$$\left( \int_{\Omega} u_\mu^6 \right)^{2/3} \leq C \left( \int_{\Omega} (|\nabla u_\mu|^2 + \mu u_\mu^2) \right).$$

Therefore,  $\|u_\mu\|_{H^1(\Omega)} \geq C^{-3/2}$ , and  $|u_\mu|_6^6 \geq C^{-3}$ . Passing to a subsequence, we may assume that  $|u_\mu|_6^6 \rightarrow l > 0$  as  $\mu \rightarrow \infty$ . We set

$$Q_\mu(t) = \sup_{x \in \Omega} \int_{(x+tB(0,1)) \cap \Omega} u_\mu^6 \quad t > 0.$$

$Q_\mu$  is continuous and increasing. Let  $R$  be such that  $\Omega \subset x_0 + B(0, R)$ , for some  $x_0 \in \Omega$ . Consequently,  $Q_\mu(R) = |u_\mu|_6^6 = l + o(1)$ . Choosing  $\rho > 0$  such that

$$\rho < \min\left(\frac{l}{2}, A^{3/2}\right) \quad (\text{B.2})$$

with

$$A = \inf_{u \in H^1(\Omega)} \frac{\|u_\mu\|_{H^1(\Omega)}}{|u_\mu|_6^2}$$

for large  $\mu$  there exist  $\varepsilon_\mu$ ,  $0 < \varepsilon_\mu < R$ , and  $a_\mu \in \bar{\Omega}$  such that

$$Q_\mu(\varepsilon_\mu) = \int_{(a_\mu + \varepsilon_\mu B(0,1)) \cap \Omega} u_\mu^6 = \rho. \quad (\text{B.3})$$

We set

$$\tilde{u}_\mu(x) = \varepsilon_\mu^{1/2} u_\mu(\varepsilon_\mu x + a_\mu) \quad x \in \Omega_\mu = (\Omega - a_\mu)/\varepsilon_\mu$$

which satisfies

$$-\Delta \tilde{u}_\mu + \mu \varepsilon_\mu^2 \tilde{u}_\mu = \tilde{u}_\mu^5, \quad \tilde{u}_\mu > 0 \text{ in } \Omega_\mu; \quad \frac{\partial \tilde{u}_\mu}{\partial \nu} = 0 \text{ on } \partial \Omega_\mu. \quad (\text{B.4})$$

Up to some subsequence, one of the three cases occur :

- (i)  $\mu \varepsilon_\mu^2 \rightarrow \infty$
- (ii)  $\mu \varepsilon_\mu^2 \rightarrow \alpha > 0$
- (iii)  $\mu \varepsilon_\mu^2 \rightarrow 0$ .

We are going to prove that, actually, (iii) is the only possible case.

Let us assume, first, that  $\mu \varepsilon_\mu^2$  goes to infinity. (B.1) and the boundedness of  $(u_\mu)$  in  $H^1(\Omega)$  implies that  $\mu |u_\mu|_2^2$  is bounded, whence

$$\int_{\Omega_\mu} \tilde{u}_\mu^2 = \frac{1}{\mu \varepsilon_\mu^2} \int_{\Omega} u_\mu^2 \rightarrow 0 \quad \text{as } \mu \rightarrow \infty. \quad (\text{B.5})$$

Let  $\zeta$  be a smooth positive function from  $\mathbb{R}^3$  to  $\mathbb{R}$ , with  $S = \text{supp} \zeta \subset b + B(0, 1)$  for some  $b \in \mathbb{R}^3$ . We have

$$\int_{\Omega_\mu} \nabla \tilde{u}_\mu \cdot \nabla (\zeta^2 \tilde{u}_\mu) + \mu \varepsilon_\mu^2 \int_{\Omega_\mu} \tilde{u}_\mu (\zeta^2 \tilde{u}_\mu) = \int_{\Omega_\mu} \tilde{u}_\mu^5 (\zeta^2 \tilde{u}_\mu).$$

Therefore

$$\int_{\Omega_\mu} |\nabla (\zeta \tilde{u}_\mu)|^2 - \int_{\Omega_\mu} |\nabla \zeta|^2 \tilde{u}_\mu^2 + \mu \varepsilon_\mu^2 \int_{\Omega_\mu} (\zeta \tilde{u}_\mu)^2 = \int_{\Omega_\mu} \tilde{u}_\mu^4 (\zeta \tilde{u}_\mu)^2.$$

Noticing that

$$\inf_{v \in H^1(\Omega_\mu)} \frac{|\nabla v|_2^2 + \varepsilon_\mu^2 |v|_2^2}{|v|_6^2} = \inf_{u \in H^1(\Omega)} \frac{\|u\|_{H^1}}{|u|_6^2} = A \quad (\text{B.6})$$

taking account of (B.5) and using Hölder inequality, we obtain

$$A \left( \int_{\Omega_\mu} (\zeta \tilde{u}_\mu)^6 \right)^{1/3} \leq \left( \int_{S \cap \Omega_\mu} \tilde{u}_\mu^6 \right)^{2/3} \left( \int_{\Omega_\mu} (\zeta \tilde{u}_\mu)^6 \right)^{1/3} + o(1).$$

In view of (B.3-4)

$$\int_{S \cap \Omega_\mu} \tilde{u}_\mu^6 \leq \int_{(a_\mu + \varepsilon_\mu b + \varepsilon_\mu B(0,1)) \cap \Omega} u_\mu^6 \leq Q_\mu(\varepsilon_\mu) < A^{3/2}$$

so that, for any smooth  $\zeta$  such that  $S = \text{supp}\zeta \subset b + B(0,1)$ ,  $b \in \mathbb{R}^3$

$$\int_{\Omega_\mu} (\zeta \tilde{u}_\mu)^6 \rightarrow 0 \quad \text{as } \mu \rightarrow \infty \quad (\text{B.7})$$

in contradiction with (B.3).

Let us assume now that  $\mu \varepsilon_\mu^2$  is bounded. In particular,  $\varepsilon_\mu$  goes to zero and, up to a subsequence and a space rotation,  $\Omega_\mu$  goes to  $\mathbb{R}_T^3$ , with

$$\mathbb{R}_T^3 = \{x = (x_1, x_2, x_3) \in \mathbb{R}^3, x_3 > -T\} \quad T = \lim_{\mu \rightarrow \infty} d(a_\mu, \partial\Omega)/\varepsilon_\mu.$$

Noticing that  $|\tilde{u}_\mu|_{L^6(\Omega_\mu)} = |u_\mu|_{L^6(\Omega)}$  and  $|\nabla \tilde{u}_\mu|_{L^2(\Omega_\mu)} = |\nabla u_\mu|_{L^2(\Omega)}$ , we may assume that there exist  $\tilde{u} \in L^6(\mathbb{R}_T^3)$ ,  $\tilde{v} \in L^2(\mathbb{R}_T^3)$  such that for any compact set  $K \in \mathbb{R}_T^3$  ( $K \in \Omega_\mu$  for  $\mu$  large enough)

$$\tilde{u}_\mu \rightharpoonup \tilde{u} \text{ in } L^6(K) \quad \nabla \tilde{u}_\mu \rightharpoonup \tilde{v} \text{ in } L^2(K).$$

Moreover, for any  $\varphi \in C_0^\infty(\mathbb{R}_T^3)$

$$\int_{\Omega_\mu} \nabla \tilde{u}_\mu \cdot \nabla \varphi \rightarrow \int_{\mathbb{R}_T^3} \tilde{v} \cdot \nabla \varphi \quad \text{and} \quad \int_{\Omega_\mu} \nabla \tilde{u}_\mu \cdot \nabla \varphi \rightarrow - \int_{\mathbb{R}_T^3} \tilde{u} \Delta \varphi$$

whence  $\tilde{v} = \nabla \tilde{u}$ .

Assuming that case (ii) occurs,  $\tilde{u}$  satisfies

$$-\Delta \tilde{u} + \alpha \tilde{u} = \tilde{u}^5 \quad \text{in } \mathbb{R}_T^3. \quad (\text{B.8})$$

Moreover,  $\tilde{u} \geq 0$ , since, along some subsequence,  $\tilde{u}_\mu \rightarrow \tilde{u}$  almost everywhere.

If  $T = \infty$ ,  $\mathbb{R}_T^3 = \mathbb{R}^3$ , and the only solutions to (B.8) are  $\tilde{u} \equiv 0$  or  $\tilde{u} \equiv \alpha^{1/4}$ .  $\tilde{u} \equiv \alpha^{1/4}$  is excluded, since  $\tilde{u} \in L^6(\mathbb{R}^3)$ . If  $\tilde{u} \equiv 0$ ,  $\tilde{u}_\mu$  goes to zero in  $L^2(K)$ , for any compact set  $K$ . Then, using this result instead of (B.5), the same argument as previously leads to (B.7), that is a contradiction.

If  $T < \infty$ , we first notice that the normal derivative of  $\tilde{u}$  vanishes on  $\partial\mathbb{R}_T^3$ . Indeed, from (B.4) we know that for any  $\varphi \in C_0^\infty(\mathbb{R}^3)$

$$\int_{\Omega_\mu} \nabla \tilde{u}_\mu \cdot \nabla \varphi + \mu \varepsilon_\mu^2 \int_{\Omega_\mu} \tilde{u}_\mu \varphi = \int_{\Omega_\mu} \tilde{u}_\mu^5 \varphi$$

from which we easily deduce that

$$\int_{\mathbb{R}_T^3} \nabla \tilde{u} \cdot \nabla \varphi + \alpha \int_{\mathbb{R}_T^3} \tilde{u} \varphi = \int_{\mathbb{R}_T^3} \tilde{u}^5 \varphi$$

whence

$$\frac{\partial \tilde{u}}{\partial \nu} = 0 \quad \text{on } \partial\mathbb{R}_T^3.$$

Then, we can continue  $\tilde{u}$  by reflection with respect to  $x_3 = -T$ , and the continuation, still denoted  $\tilde{u}$ , satisfies (B.8) in whole  $\mathbb{R}^3$ , whence  $\tilde{u} \equiv 0$  or  $\tilde{u} \equiv \alpha^{1/4}$ .  $\tilde{u} \equiv \alpha^{1/4}$  is again impossible, since  $\tilde{u} \in L^6(\mathbb{R}^3)$ .  $\tilde{u} \equiv 0$  implies, as previously, that  $\tilde{u}_\mu$  goes to zero in  $L^2(K)$ , for any compact set  $K \in \mathbb{R}_T^3$ .

If  $\bar{B}(0, 1) \in \mathbb{R}_T^3$  (i.e.  $T > 1$ ), we can repeat the argument which leads to (B.7), hence a contradiction. If  $0 \leq T \leq 1$ ,  $\tilde{u}_\mu$  may be continued in  $B(0, 2) \setminus \Omega_\mu$  in such a way that,  $\tilde{u}'_\mu$  denoting the continuation of  $\tilde{u}_\mu$  in  $\Omega_\mu \cup B(0, 2)$

$$\|\tilde{u}'_\mu\|_{H^1(B(0,2))} \leq C \|\tilde{u}_\mu\|_{H^1(B(0,2) \cap \Omega_\mu)} \leq C'$$

$C$  and  $C'$  independent of  $\mu$ . Then, along some subsequence

$$\tilde{u}'_\mu \rightharpoonup \tilde{u}' \quad \text{in } H^1(B(0, 2)) \quad \tilde{u}'_\mu \rightarrow \tilde{u}' \quad \text{in } L^2(B(0, 2))$$

with  $\tilde{u}' = \tilde{u} = 0$  in  $B(0, 2) \cap \mathbb{R}_T^3$ . Consequently

$$\int_{\Omega_\mu \cap B(0,2)} \tilde{u}_\mu^2 \leq 2 \int_{\Omega_\mu \cap B(0,2)} (\tilde{u}_\mu - \tilde{u}'_\mu)^2 + 2 \int_{\Omega_\mu \cap B(0,2)} \tilde{u}'_\mu^2 \rightarrow 0 \quad \text{as } \mu \rightarrow \infty$$

and we are still able to prove (B.7), hence again a contradiction.

As announced, (iii) is the only possible case, and  $\tilde{u}$  satisfies

$$-\Delta \tilde{u} = \tilde{u}^5, \quad \tilde{u} \geq 0 \quad \text{in } \mathbb{R}_T^3; \quad \frac{\partial \tilde{u}}{\partial \nu} = 0 \quad \text{on } \partial\mathbb{R}_T^3 \quad (\text{B.9})$$

with  $\nabla \tilde{u} \in L^2(\mathbb{R}_T^3)$ ,  $\tilde{u} \in L^6(\mathbb{R}_T^3)$ . If  $\tilde{u} \equiv 0$ , reasoning as previously provides us with a contradiction. Therefore, there exist  $\lambda \in \mathbb{R}_+^*$  and  $a \in \partial\mathbb{R}^3$  such that, according to [13]

$$\tilde{u}(x) = 3^{1/4} \frac{\lambda^{1/2}}{(1 + \lambda^2 |x - a|^2)^{1/2}}$$

and either  $T = \infty$ , or  $T < \infty$  and  $a \in \partial\mathbb{R}_T^3$ . Then, we set

$$u_\mu^{(1)}(x) = u_\mu(x) - 3^{1/4} V_{\mu, \lambda/\varepsilon_\mu, a_\mu + \varepsilon_\mu a}(x) \quad x \in \Omega \quad (\text{B.10})$$

where  $V$  is defined by (2.13-14). We notice that

$$\frac{\mu}{(\lambda/\varepsilon_\mu)^2} \rightarrow 0 \quad (\text{B.11})$$

and

$$(\lambda/\varepsilon_\mu)d(a_\mu + \varepsilon_\mu a, \partial\Omega) \rightarrow \infty \quad (T = \infty) \quad \text{or} \quad (\lambda/\varepsilon_\mu)d(a_\mu + \varepsilon_\mu a, \partial\Omega) \rightarrow 0 \quad (T < \infty). \quad (\text{B.12})$$

Indeed,  $a \in \partial\mathbb{R}^3$  means that  $a$  is limit of points of  $\partial\Omega_\mu$ , so that we may write

$$a = \frac{x_\mu - a_\mu}{\varepsilon_\mu} + y_\mu \quad x_\mu \in \partial\Omega \quad y_\mu \rightarrow 0 \quad \text{in } \mathbb{R}^3$$

and

$$(\lambda/\varepsilon_\mu)d(a_\mu + \varepsilon_\mu a, \partial\Omega) \leq \lambda|y_\mu| \rightarrow 0 \quad \text{as } \mu \rightarrow \infty.$$

We claim that

$$\int_\Omega |\nabla u_\mu^{(1)}|^2 = \int_\Omega |\nabla u_\mu|^2 - \left( S^{3/2} \text{ if } T = \infty ; S^{3/2}/2 \text{ if } T < \infty \right) + o(1) \quad (\text{B.13})$$

where

$$S = \inf_{u \in H_0^1(\Omega)} \frac{|\nabla u|_2^2}{|u|_6^2} = \frac{3\pi^{4/3}}{2^{4/3}}$$

is the Sobolev constant, and

$$-\Delta u_\mu^{(1)} + \mu u_\mu^{(1)} = (u_\mu^{(1)+})^5 + f_\mu \quad \text{in } \Omega \quad u_\mu^{(1)+} = \max(u_\mu^{(1)}, 0) \quad (\text{B.14})$$

with

$$(f_\mu, \varphi)_{H^{-1}(\Omega), H^1(\Omega)} = o(\|\varphi\|_{H^1(\Omega)}) \quad \text{uniformly for } \varphi \in H^1(\Omega)$$

as  $\mu$  goes to infinity.

If  $u_\mu^{(1)}$  does not go to zero in  $H^1(\Omega)$ , we can apply to  $u_\mu^{(1)}$  the same arguments as those that we used concerning  $u_\mu$ . It is easily checked that the presence of  $f_\mu$  does not affect the situation; in the same way, (B.12) ensures that the normal derivative of  $u_\mu^{(1)}$  on  $\partial\Omega$  is sufficiently small for our purposes. In particular

$$\int_{\Omega_\mu} \left| \frac{\partial \tilde{u}_\mu^{(1)}}{\partial \nu} \right| \tilde{u}_\mu^{(1)} \rightarrow 0 \quad \text{and} \quad \int_{\Omega_\mu} \frac{\partial \tilde{u}_\mu^{(1)}}{\partial \nu} \varphi \rightarrow 0$$

for any  $\varphi \in C_0^\infty(\mathbb{R}^3)$ , as  $\mu \rightarrow \infty$ . Then, we obtain some  $u_\mu^{(2)}$  which either goes to zero in  $H^1(\Omega)$ , or may be treated as previously to define some  $u_\mu^{(3)}$ , and so on. The process must stop after a finite number of steps since, according to (B.13),  $|\nabla u|_2^2$  loses each time some fixed amount. In the end, we obtain that  $u_\mu$  writes as

$$u_\mu = 3^{1/4} \sum_{i=1}^k V_{\mu, \lambda_\mu^i, a_\mu^i} + v_\mu \quad v_\mu \rightarrow 0 \quad \text{in } H^1(\Omega) \quad (\text{B.15})$$

with

$$\begin{aligned}\mu^{1/2}/\lambda_\mu^i &\rightarrow 0 \\ \lambda_\mu^i d(a_\mu^i, \partial\Omega) &\rightarrow 0 \quad \text{or} \quad \lambda_\mu^i d(a_\mu^i, \partial\Omega) \rightarrow \infty\end{aligned}$$

as  $\mu \rightarrow \infty$ . Moreover, in view of (B.13), we have

$$|\nabla u_\mu|_2^2 = 3^{1/2} \sum_{i=1}^k |\nabla V_{\mu, \lambda_\mu^i, a_\mu^i}|_2^2 + o(1). \quad (\text{B.16})$$

As, on the other hand, we deduce from computations in [7] that

$$\int_{\Omega} \nabla V_{\mu, \lambda_\mu^i, a_\mu^i} \cdot \nabla V_{\mu, \lambda_\mu^j, a_\mu^j} \geq C \left( \frac{\lambda_\mu^i}{\lambda_\mu^j} + \frac{\lambda_\mu^j}{\lambda_\mu^i} + \lambda_\mu^i \lambda_\mu^j |a_\mu^i - a_\mu^j|^2 \right)^{1/2} \quad (\text{B.17})$$

where  $C$  is a strictly positive constant, (B.15-16) imply that the right hand side in (B.17) goes to zero if  $i \neq j$ , and the proof of Proposition 2.5 is complete.

**Proof of (B.13).** In view of (B.10), we have

$$\int_{\Omega} |\nabla u_\mu^{(1)}|^2 = \int_{\Omega} |\nabla u_\mu|^2 - 2 \cdot 3^{1/4} \int_{\Omega} \nabla u_\mu \cdot \nabla V_\mu + 3^{1/2} \int_{\Omega} |\nabla V_\mu|^2$$

with  $V_\mu = V_{\mu, \lambda/\varepsilon_\mu, a_\mu + \varepsilon_\mu a}$ , for sake of simplicity. On one hand, it follows from (2.16) that

$$\int_{\Omega} |\nabla V_\mu|^2 = \int_{\Omega} |\nabla U_\mu|^2 + o(1)$$

with  $U_\mu = U_{\mu, \lambda/\varepsilon_\mu, a_\mu + \varepsilon_\mu a}$ , and standard computations yield

$$\begin{aligned}\int_{\Omega} |\nabla U_\mu|^2 &= S^{3/2} + o(1) \quad \text{if} \quad \frac{1}{\varepsilon_\mu} d(a_\mu + \varepsilon_\mu a, \partial\Omega) \rightarrow \infty \\ \int_{\Omega} |\nabla U_\mu|^2 &= \frac{S^{3/2}}{2} + o(1) \quad \text{if} \quad \frac{1}{\varepsilon_\mu} d(a_\mu + \varepsilon_\mu a, \partial\Omega) \rightarrow 0.\end{aligned}$$

On the other hand, still using (2.16) and the boundedness of  $u_\mu$  in  $H^1(\Omega)$

$$\begin{aligned}\int_{\Omega} \nabla u_\mu \cdot \nabla V_\mu &= \int_{\Omega} \nabla u_\mu \cdot \nabla U_\mu + o(1) \\ &= \int_{\Omega_\mu} \nabla \tilde{u}_\mu \cdot \nabla \tilde{u} + o(1) \\ &= 3^{1/4} \int_{\mathbb{R}_T^3} |\nabla U_{\lambda, a}|^2 + o(1)\end{aligned}$$

and

$$\begin{aligned}\int_{\mathbb{R}_T^3} |\nabla U_{\lambda, a}|^2 &= S^{3/2} \text{ if } T = \infty \quad (\text{i.e. } \frac{1}{\varepsilon_\mu} d(a_\mu + \varepsilon_\mu a, \partial\Omega) \rightarrow \infty) \\ \int_{\mathbb{R}_T^3} |\nabla U_{\lambda, a}|^2 &= \frac{S^{3/2}}{2} + o(1) \text{ if } T < \infty, \quad a \in \partial\mathbb{R}_T^3 \quad (\frac{1}{\varepsilon_\mu} d(a_\mu + \varepsilon_\mu a, \partial\Omega) \rightarrow 0)\end{aligned}$$

whence (B.13).

**Proof of (B.14).** From (B.10) and (2.15) we know that

$$-\Delta u_\mu^{(1)} + \mu u_\mu^{(1)} = u_\mu^5 - (3^{1/4}V_\mu)^5 + \mu 3^{1/4} \left( U_\mu - \frac{1}{(\lambda/\varepsilon_\mu)^{1/2}|x - a_\mu - \varepsilon_\mu a|} \right)$$

in  $\Omega$ . Let  $\varphi \in H^1(\Omega)$ . We notice that

$$\begin{aligned} & \mu \int_{\Omega} \left( U_\mu - \frac{1}{(\lambda/\varepsilon_\mu)^{1/2}|x - a_\mu - \varepsilon_\mu a|} \right) \varphi \\ & \leq \mu |\varphi|_2 \left( \int_{\Omega} \left( U_\mu - \frac{1}{(\lambda/\varepsilon_\mu)^{1/2}|x - a_\mu - \varepsilon_\mu a|} \right)^2 \right)^{1/2} \\ & \leq C \mu \varepsilon_\mu^2 |\varphi|_2 \end{aligned}$$

as a direct computation shows. Secondly, setting  $\varphi_\mu = \varphi_{\mu, \lambda/\varepsilon_\mu, a_\mu + \varepsilon_\mu a}$ , we have in view of (2.16)

$$\begin{aligned} \int_{\Omega} (V_\mu^5 - U_\mu^5) \varphi & \leq \int_{\Omega} (U_\mu^4 \varphi_\mu + \varphi_\mu^5) |\varphi| \\ & \leq C' |\varphi_\mu|_6 \|\varphi\|_{H^1(\Omega)} \end{aligned}$$

using Hölder inequality, the embedding of  $H^1(\Omega)$  in  $L^6(\Omega)$ , the boundedness of  $(U_\mu)$  in  $L^6(\Omega)$  and the fact that  $\varphi_\mu$  goes to zero in  $L^6(\Omega)$  as  $\mu \varepsilon_\mu^2$  goes to zero. Consequently, we are left to show that

$$\left| \int_{\Omega} \left( (u_\mu^{(1)+})^5 - u_\mu^5 + (3^{1/4}U_\mu)^5 \right) \varphi \right| \leq C_\mu \|\varphi\|_{H^1(\Omega)} \quad (\text{B.18})$$

with  $C_\mu$  independent of  $\varphi$ ,  $C_\mu \rightarrow 0$  as  $\mu \rightarrow \infty$ . We proceed as in [15]. Let  $I_\mu$  denote the left hand side integral in (B.18). Setting  $\bar{U}_\mu = 3^{1/4}U_\mu$ , we have

$$\begin{aligned} I_\mu & = \int_{\Omega} \varphi \int_0^{\bar{U}_\mu} \frac{\partial}{\partial t} \left[ ((u_\mu - t)^+)^5 + t^5 \right] dt \\ & = -5 \int_{\Omega} \varphi \int_0^{\bar{U}_\mu} \left[ ((u_\mu - t)^+)^4 + t^4 \right] dt \\ & = \int_0^1 \left( \int_{\Omega} \left[ ((u_\mu - s\bar{U}_\mu)^+)^4 - (1-s)^4 \bar{U}_\mu^4 \right] \bar{U}_\mu \varphi \right) ds. \end{aligned}$$

Therefore

$$\begin{aligned} |I_\mu| & \leq 5 |\varphi|_6 \max_{0 \leq s \leq 1} \int_{\Omega} \left| ((u_\mu - s\bar{U}_\mu)^+)^4 - (1-s)^4 \bar{U}_\mu^4 \right|^{6/5} \bar{U}_\mu^{6/5} \\ & \leq C |\varphi|_{H^1(\Omega_\mu)} \int_{\Omega} \left| ((\tilde{u}_\mu - s\tilde{u})^+)^4 - (1-s)^4 \tilde{u}^4 \right|^{6/5} \tilde{u}^{6/5}. \end{aligned}$$

We know that  $\|\tilde{u}_\mu\|_{H^1(\Omega)}$  is bounded, and that  $\tilde{u}_\mu \rightarrow \tilde{u}$  in  $H^1(K)$ ,  $\tilde{u}_\mu \rightarrow \tilde{u}$  in  $L^q(\Omega)$ ,  $1 \leq q < 6$ , for any compact set  $K \in \mathbb{R}^3$ . The conclusion follows easily.

## C Estimates

In this last part, we collect the integral estimates which are used in Section 3. First, we recall that in [40] the following is proved, for  $V = V_{\mu, \lambda, a}$

$$\int_{\Omega} |\nabla V|^2 = \frac{3\pi^2}{4} - 6\pi \frac{\mu^{1/2}}{\lambda} + O\left(\frac{1}{\lambda\mu^{1/2}}\right) \quad (\text{C.1})$$

$$\int_{\Omega} V^2 = \frac{2\pi}{\lambda\mu^{1/2}} + O\left(\frac{1}{\lambda\mu^{3/2}}\right) \quad (\text{C.2})$$

$$\int_{\Omega} V^6 = \frac{\pi^2}{4} - 8\pi \frac{\mu^{1/2}}{\lambda} + O\left(\frac{\mu^{1/2}}{\lambda^2}\right) \quad (\text{C.3})$$

as  $a$  does not approach the boundary of  $\Omega$ , and  $\mu^{1/2}/\lambda$  goes to infinity. Actually,  $a$  is assumed in [40] to be on  $\partial\Omega$ . The above results follow from the same computations, made easier by the fact that a boundary effect has not to be considered. In the same way, we have

$$\int_{\Omega} \nabla V \cdot \nabla \frac{\partial V}{\partial \lambda} = 3\pi \frac{\mu^{1/2}}{\lambda^2} + O\left(\frac{1}{\lambda^2\mu^{1/2}}\right) \quad \int_{\Omega} \nabla V \cdot \nabla \frac{\partial V}{\partial a_j} = O(\mu^{1/2}) \quad (\text{C.4})$$

$$\int_{\Omega} V \frac{\partial V}{\partial \lambda} = -\frac{\pi}{\lambda^2\mu^{1/2}} + O\left(\frac{1}{\lambda^2\mu^{3/2}}\right) \quad \int_{\Omega} V \frac{\partial V}{\partial a_j} = O\left(\frac{1}{\mu^{1/2}}\right) \quad (\text{C.5})$$

$$\int_{\Omega} V^5 \frac{\partial V}{\partial \lambda} = \frac{4\pi\mu^{1/2}}{3\lambda^2} + O\left(\frac{\mu^{1/2}}{\lambda^3}\right) \quad \int_{\Omega} V^5 \frac{\partial V}{\partial a_j} = O(\mu^{1/2}) \quad (\text{C.6})$$

and

$$\int_{\Omega} \left| \nabla \frac{\partial V}{\partial \lambda} \right|^2 = \frac{15\pi^2}{64\lambda^2} + O\left(\frac{\mu^{1/2}}{\lambda^3}\right) \quad \int_{\Omega} \left| \nabla \frac{\partial V}{\partial a_j} \right|^2 = \frac{15\pi^2}{64}\lambda^2 + O(\lambda\mu^{1/2}) \quad (\text{C.7})$$

$$\int_{\Omega} \left| \nabla \frac{\partial^2 V}{\partial \lambda^2} \right|^2 = O\left(\frac{1}{\lambda^2}\right) \quad \int_{\Omega} \left| \nabla \frac{\partial^2 V}{\partial \lambda \partial a_j} \right|^2 = O(1). \quad (\text{C.8})$$

In view of (3.13), we have also to estimate integrals in which both  $V_i$  and  $V_j$  occur,  $i \neq j$ . Let us prove, for example

$$\int_{\Omega} \nabla V_i \cdot \nabla V_j = O\left(\frac{\mu^{1/2}}{\lambda_i^{1/2}\lambda_j^{1/2}}\right) \quad (\text{C.9})$$

as  $\mu^{1/2}/\lambda_i$ ,  $\mu^{1/2}/\lambda_j$  go to infinity,  $a_i$  and  $a_j$  do not approach the boundary of  $\Omega$  and  $|a_i - a_j| > \gamma/2\mu^{1/2}$ . From (2.15) we deduce that

$$\int_{\Omega} \nabla V_i \cdot \nabla V_j = \int_{\partial\Omega} \frac{\partial V_i}{\partial \nu} V_j + \int_{\Omega} \left( 3U_i^5 - \frac{\mu e^{-\mu^{1/2}|x-a_i|}}{\lambda^{1/2}|x-a_i|} \right) V_j. \quad (\text{C.10})$$

It follows from the definition (2.13-14) of  $V$  that,  $a_i$  and  $a_j$  staying far from  $\partial\Omega$

$$V_j = o\left(\frac{1}{\lambda_j^{1/2}}\right) \quad |\nabla V_i| = o\left(\frac{\mu^{1/2}}{\lambda_i^{1/2}}\right) \quad \text{on } \partial\Omega$$

whence

$$\int_{\partial\Omega} \frac{\partial V_i}{\partial \nu} V_j = o\left(\frac{\mu^{1/2}}{\lambda_i^{1/2} \lambda_j^{1/2}}\right). \quad (\text{C.11})$$

Turning now to the integral on  $\Omega$ , we notice that

$$V_j = O\left(\frac{\mu^{1/2}}{\lambda_j^{1/2}}\right) \quad \text{outside of } B_j = B(a_j, \gamma/4\mu^{1/2}). \quad (\text{C.12})$$

On the other hand, taking  $R' > 0$  such that  $\Omega \subset B(0, R')$

$$\int_{\Omega} U_i^5 \leq \frac{4\pi}{\lambda_i^{1/2}} \int_0^{2R'\lambda_i} \frac{r^2}{(1+r^2)^{5/2}} dr = O\left(\frac{1}{\lambda_i^{1/2}}\right) \quad (\text{C.13})$$

and

$$\int_{\Omega} \frac{\mu e^{-\mu^{1/2}|x-a_i|}}{\lambda^{1/2}|x-a_i|} dx \leq \frac{4\pi}{\lambda_i^{1/2}} \int_0^{2R'} \mu r e^{-\mu^{1/2}r} dr = O\left(\frac{1}{\lambda_i^{1/2}}\right).$$

Therefore

$$\int_{\Omega \setminus B_j} \left(3U_i^5 - \frac{\mu e^{-\mu^{1/2}|x-a_i|}}{\lambda^{1/2}|x-a_i|}\right) V_j = O\left(\frac{\mu^{1/2}}{\lambda_i^{1/2} \lambda_j^{1/2}}\right). \quad (\text{C.14})$$

Finally, we notice that

$$U_i^5 = O\left(\frac{\mu^{5/2}}{\lambda_i^{5/2}}\right) \quad \frac{\mu e^{-\mu^{1/2}|x-a_i|}}{\lambda^{1/2}|x-a_i|} = O\left(\frac{\mu^{3/2}}{\lambda_i^{1/2}}\right) \quad \text{in } B_j$$

and

$$\begin{aligned} \int_{B_j} |V_j| &\leq \frac{4\pi}{\lambda_j^{5/2}} \int_0^{\frac{\gamma\lambda_j}{4\mu^{1/2}}} \left| \frac{1}{(1+r^2)^{1/2}} - \frac{1 - e^{-(\mu^{1/2}/\lambda_j)r}}{r} \right| r^2 dr \\ &\leq \frac{4\pi}{\lambda_j^{5/2}} \left( \int_0^{\frac{\gamma\lambda_j}{4\mu^{1/2}}} \left(r - \frac{r^2}{(1+r^2)^{1/2}}\right) dr + \int_0^{\frac{\gamma\lambda_j}{4\mu^{1/2}}} r e^{-(\mu^{1/2}/\lambda_j)r} dr \right). \end{aligned}$$

Consequently

$$\int_{B_j} |V_j| = O\left(\frac{\ln(\lambda_j/\mu^{1/2})}{\lambda_j^{5/2}} + \frac{1}{\lambda_j^{1/2} \mu}\right) \quad (\text{C.15})$$

and

$$\int_{B_j} \left(3U_i^5 - \frac{\mu e^{-\mu^{1/2}|x-a_i|}}{\lambda^{1/2}|x-a_i|}\right) V_j = O\left(\frac{\mu^{1/2}}{\lambda_i^{1/2} \lambda_j^{1/2}}\right).$$

This estimate, joined to (C.10-11,16), yield (C.9).

The other quantities in (3.13), with  $i \neq j$ , may be estimated in the same way. Similar computations also yield

$$\left( \int_{\Omega} |V_i|^{24/5} |V_j|^{6/5} \right)^{5/6} = O\left(\frac{\mu^{1/2}}{\lambda_i^{1/2} \lambda_j^{1/2}}\right). \quad (\text{C.16})$$

Indeed, we have

$$\int_{\Omega} |V_i|^{24/5} \leq \frac{4\pi}{\lambda_i^{3/5}} \int_0^{2R'\lambda_i} \left| \frac{1}{(1+r^2)^{1/2}} - \frac{1 - e^{-(\mu^{1/2}/\lambda_i)r}}{r} \right|^{24/5} r^2 dr$$

and the integral on the right hand side goes to a finite limit as  $\mu^{1/2}/\lambda_i$  goes to zero, whence

$$\left( \int_{\Omega} |V_i|^{24/5} \right)^{5/6} = O\left(\frac{1}{\lambda_i^{1/2}}\right)$$

and, taking account of (C.12)

$$\left( \int_{\Omega \setminus B_j} |V_i|^{24/5} |V_j|^{6/5} \right)^{5/6} = O\left(\frac{\mu^{1/2}}{\lambda_i^{1/2} \lambda_j^{1/2}}\right). \quad (\text{C.17})$$

On the other hand

$$\begin{aligned} \int_{B_j} |V_j|^{6/5} &\leq \frac{4\pi}{\lambda_j^{5/2}} \int_0^{\frac{\gamma\lambda_j}{4\mu^{1/2}}} \left| \frac{1}{(1+r^2)^{1/2}} - \frac{1 - e^{-(\mu^{1/2}/\lambda_j)r}}{r} \right|^{6/5} r^2 dr \\ &= O\left(\frac{1}{\lambda_j^{12/5}} \int_0^{\frac{\gamma\lambda_j}{4\mu^{1/2}}} \left( \frac{1}{r^3} + \frac{e^{-(\mu^{1/2}/\lambda_j)r}}{r} \right)^{6/5} r^2 dr \right) \end{aligned}$$

and

$$\int_0^{\frac{\gamma\lambda_j}{4\mu^{1/2}}} r^{4/5} e^{-(6\mu^{1/2}/5\lambda_j)r} dr = \left(\frac{\lambda_j}{\mu^{1/2}}\right)^{9/5} \int_0^{\gamma/4} t^{4/5} e^{-6t/5} dt = O\left(\frac{\lambda_j^{9/5}}{\mu^{9/10}}\right).$$

Therefore

$$\left( \int_{\Omega} |V_j|^{6/5} \right)^{5/6} = O\left(\frac{1}{\lambda_j^{1/2} \mu^{3/4}}\right)$$

and (C.12) applied to  $V_i$  in  $B_j$  provides us with

$$\left( \int_{B_j} |V_i|^{24/5} |V_j|^{6/5} \right)^{5/6} = O\left(\frac{\mu^{1/2}}{\lambda_i^{1/2} \lambda_j^{1/2}} \cdot \frac{\mu^{3/4}}{\lambda_i^{3/2}}\right). \quad (\text{C.18})$$

(C.17-18) prove (C.16).

We state now :

**Proposition C.1** *Let  $(\alpha, \lambda, a) \in \tilde{\mathcal{N}}_\mu$  (i.e.  $|\alpha_i - 3^{1/4}| < \eta_1$ ,  $\mu^{1/2}/\lambda_i < \eta_1$ ,  $d(a_i, \partial\Omega) > \rho$ ,  $|a_i - a_j| > \frac{\gamma}{2\mu^{1/2}}$  if  $i \neq j$ ). We have*

$$\frac{\partial K_\mu}{\partial \alpha_i}(\alpha, \lambda, a, \bar{v}_\mu(\alpha, \lambda, a)) = \frac{3\pi^2}{4} \alpha_i (3 - \alpha_i^4) + O\left(\frac{\mu^{1/2}}{|\lambda|}\right) \quad (\text{C.19})$$

$$\begin{aligned} \frac{\partial K_\mu}{\partial \lambda_i}(\alpha, \lambda, a, \bar{v}_\mu(\alpha, \lambda, a)) &= \pi(2 - \frac{4}{3}\alpha^4) \frac{\mu^{1/2}}{\lambda_i^2} \\ &+ O\left(\frac{\mu^{1/2}}{\lambda_i^{3/2}|\lambda|^{1/2}} \sum_l |3 - \alpha_l^4|\right) + o\left(\frac{\mu^{1/2}}{\lambda_i^{3/2}|\lambda|^{1/2}}\right) \end{aligned} \quad (\text{C.20})$$

$$\frac{\partial K_\mu}{\partial (a_i)_j}(\alpha, \lambda, a, \bar{v}_\mu(\alpha, \lambda, a)) = O\left(\frac{\mu^{1/2}\lambda_i}{|\lambda|}\right) \quad (\text{C.21})$$

$$\frac{\partial^2 K_\mu}{\partial \alpha_i^2}(\alpha, \lambda, a, \bar{v}_\mu(\alpha, \lambda, a)) = \frac{3\pi^2}{4} (3 - 5\alpha_i^4) + O\left(\frac{\mu^{1/2}}{|\lambda|}\right) \quad (\text{C.22})$$

as  $\mu \rightarrow \infty$  and  $\mu^{1/2}/\lambda_i \rightarrow \infty$ ,  $1 \leq i \leq k$ .

Let us prove (C.19). According to (3.3-4), we have for  $(\alpha, \lambda, a, v) \in \mathcal{N}_\mu$

$$\begin{aligned} \frac{\partial K_\mu}{\partial \alpha_i}(\alpha, \lambda, a, v) &= \int_\Omega \nabla(\sum_j \alpha_j V_j) \cdot \nabla V_i + \mu \int_\Omega (\sum_j \alpha_j V_j + v) V_i \\ &\quad - \int_\Omega (\sum_j \alpha_j V_j + v)^5 V_i. \end{aligned}$$

Let us expand  $(\sum_j \alpha_j V_j + v)^5$ . (3.15-16) and (C.16) provide us with

$$\begin{aligned} &\left| \mu \int_\Omega V_i v - \int_\Omega (\sum_j \alpha_j V_j + v)^4 V_i v \right| \\ &\leq C \frac{\mu^{1/2}}{\lambda_i^{1/2}} \left( \frac{1}{|\lambda|^{1/2}} + \frac{1}{\mu^{3/4}} \right) \left( \int_\Omega (|\nabla v|^2 + \mu v^2) \right)^{1/2}. \end{aligned}$$

Moreover, Hölder inequality, Sobolev embedding theorem and (C.3) show that the contribution to the integral of the terms in which  $v$  occurs with an exponent larger or equal to 2 is dominated by  $\|v\|_{H^1(\Omega)}^2$ . As a consequence, if  $(\alpha, \lambda, a) \in \tilde{\mathcal{N}}_\mu$  and  $v = \bar{v}_\mu(\alpha, \lambda, a)$ , (3.12) yields

$$\begin{aligned} \frac{\partial K_\mu}{\partial \alpha_i}(\alpha, \lambda, a, \bar{v}_\mu) &= \frac{\partial K_\mu}{\partial \alpha_i}(\alpha, \lambda, a, 0) \\ &+ O\left(\frac{\mu^{1/2}}{\lambda_i^{1/2}|\lambda|^{3/2}} + \frac{1}{\mu^{1/2}\lambda_i^{1/2}|\lambda|^{1/2}}\right). \end{aligned} \quad (\text{C.23})$$

In view of (C.1-3), we have

$$\alpha_i \int_{\Omega} |\nabla V_i|^2 + \mu \alpha_i \int_{\Omega} V_i^2 - \alpha_i^5 \int_{\Omega} V_i^6 = \frac{\pi^2}{4} \alpha_i (3 - \alpha_i^4) + O\left(\frac{\mu^{1/2}}{\lambda_i}\right).$$

We recall that, according to (C.9), we have also

$$\int_{\Omega} \nabla V_i \cdot \nabla V_j = O\left(\frac{\mu^{1/2}}{\lambda_i^{1/2} \lambda_j^{1/2}}\right).$$

Similar computations show that

$$\int_{\Omega} V_i^5 V_j = O\left(\frac{\mu^{1/2}}{\lambda_i^{1/2} \lambda_j^{1/2}}\right)$$

and

$$\int_{\Omega} \left( \left( \sum_j \alpha_j V_j \right)^5 V_i - \sum_j \alpha_j^5 V_j^5 V_i \right) = o\left(\frac{\mu^{1/2}}{\lambda_i^{1/2} |\lambda|^{1/2}}\right).$$

Lastly, from (C.12,15) we deduce that

$$\mu \int_{\Omega} V_i V_j = O\left(\frac{\mu^{1/2}}{\lambda_i^{1/2} \lambda_j^{1/2}}\right).$$

Collecting these results, (C.19) is established.

(C.21-22) follow from the same kind of integral estimates. The only result which requires to be more careful is (C.20). From (3.3-4), we have

$$\begin{aligned} \frac{1}{\alpha_i} \frac{\partial K_{\mu}}{\partial \lambda_i}(\alpha, \lambda, a, v) &= \int_{\Omega} \nabla \left( \sum_j \alpha_j V_j \right) \cdot \nabla \frac{\partial V_i}{\partial \lambda_i} \\ &\quad + \mu \int_{\Omega} \left( \sum_j \alpha_j V_j + v \right) \frac{\partial V_i}{\partial \lambda_i} - \int_{\Omega} \left( \sum_j \alpha_j V_j + v \right)^5 \frac{\partial V_i}{\partial \lambda_i} \end{aligned}$$

for  $(\alpha, \lambda, a, v) \in \mathcal{N}_{\mu}$ . The same arguments as in the proof of Lemma 3.1 in [40] show that

$$\begin{aligned} \left| \mu \int_{\Omega} \frac{\partial V_i}{\partial \lambda_i} v - \alpha_i^4 \int_{\Omega} V_i^4 \frac{\partial V_i}{\partial \lambda_i} v \right| \\ \leq C \frac{\mu^{1/2}}{\lambda_i^{3/2}} \left( \frac{1}{|\lambda|^{1/2}} + \frac{1}{\mu^{3/4}} \right) \left( \int_{\Omega} (|\nabla v|^2 + \mu v^2) \right)^{1/2}. \end{aligned}$$

We have also, if  $i \neq j$

$$\left| \int_{\Omega} V_i^4 \frac{\partial V_i}{\partial \lambda_i} v \right| \leq C \|v\|_{H^1(\Omega)} \left( \int_{\Omega} |V_j|^{24/5} \left| \frac{\partial V_i}{\partial \lambda_i} \right|^{6/5} \right)^{5/6}$$

and proceeding as in the proof of (C.16), we find

$$\left( \int_{\Omega} |V_j|^{24/5} \left| \frac{\partial V_i}{\partial \lambda_i} \right|^{6/5} \right)^{5/6} = O\left( \frac{\mu^{1/2}}{\lambda_i^{3/2} \lambda_j^{1/2}} \right).$$

Lastly, using Hölder inequality, Sobolev embedding theorem and integral estimates as the previous one, we see that the terms in which  $v$  occurs with an exponent larger or equal to 2 have a contribution to the integral which is dominated by  $\|v\|_{H^1(\Omega)}^2/\lambda_i$ . Therefore, similarly to (C.23) we obtain, using (3.12)

$$\frac{1}{\alpha_i} \frac{\partial K_{\mu}}{\partial \alpha_i}(\alpha, \lambda, a, \bar{v}_{\mu}) = \frac{1}{\alpha_i} \frac{\partial K_{\mu}}{\partial \alpha_i}(\alpha, \lambda, a, 0) + O\left( \frac{\mu^{1/2}}{\lambda_i^{3/2} |\lambda|^{3/2}} + \frac{1}{\mu^{1/2} \lambda_i^{3/2} |\lambda|^{1/2}} \right)$$

for any  $(\alpha, \lambda, a) \in \tilde{\mathcal{N}}_{\mu}$ . We notice that, according to (C.4-6)

$$\begin{aligned} \int_{\Omega} \nabla V_i \cdot \nabla \frac{\partial V_i}{\partial \lambda_i} + \mu \int_{\Omega} V_i \frac{\partial V_i}{\partial \lambda_i} - \alpha_i^4 \int_{\Omega} V_i^5 \frac{\partial V_i}{\partial \lambda_i} \\ = \pi \left( 2 - \frac{4}{3} \alpha^4 \right) \frac{\mu^{1/2}}{\lambda_i^2} + O\left( \frac{1}{\lambda_i^2 \mu^{1/2}} \sum_l |3 - \alpha_l^4| \right). \end{aligned}$$

In addition, using (2.15), we have

$$\begin{aligned} \int_{\Omega} \nabla(\alpha_j V_j) \cdot \nabla \frac{\partial V_i}{\partial \lambda_i} + \mu \int_{\Omega} \alpha_j V_j \frac{\partial V_i}{\partial \lambda_i} - \int_{\Omega} (\alpha_j V_j)^5 \frac{\partial V_i}{\partial \lambda_i} \\ = \alpha_j \left( \int_{\partial\Omega} \frac{\partial V_j}{\partial \nu} \frac{\partial V_i}{\partial \lambda_i} + \int_{\Omega} \left( 3U_j^5 - \alpha_j^4 V_j^5 + \mu \left( U_j - \frac{1}{\lambda_j^{1/2} |x - a_j|} \right) \right) \frac{\partial V_i}{\partial \lambda_i} \right). \end{aligned}$$

As  $a_i$  and  $a_j$  do not approach the boundary of  $\Omega$ , the definition of  $V_i, V_j$  shows that

$$\frac{\partial V_j}{\partial \nu} = o\left( \frac{\mu^{1/2}}{\lambda_j^{1/2}} \right) \quad \frac{\partial V_i}{\partial \lambda_i} = o\left( \frac{1}{\lambda_i^{3/2}} \right) \quad \text{uniformly on } \partial\Omega$$

whence

$$\int_{\partial\Omega} \frac{\partial V_j}{\partial \nu} \frac{\partial V_i}{\partial \lambda_i} = O\left( \frac{\mu^{1/2}}{\lambda_i^{3/2} \lambda_j^{1/2}} \right).$$

On the other hand, similarly to (C.2), we have

$$\int_{\Omega} \left( \frac{\partial V_i}{\partial \lambda_i} \right)^2 = O\left( \frac{1}{\lambda_i^3 \mu^{1/2}} \right)$$

and

$$\int_{\Omega} \left( U_j - \frac{1}{\lambda_j^{1/2} |x - a_j|} \right)^2 \leq \frac{4\pi}{\lambda_j^2} \int_0^{2R'\lambda_j} \left( \frac{1}{(1+r^2)^{1/2}} - \frac{1}{r} \right)^2 r^2 dr = O\left( \frac{1}{\lambda_j^2} \right)$$

whence, using Schwarz inequality

$$\mu \int_{\Omega} \left( U_j - \frac{1}{\lambda_j^{1/2} |x - a_j|} \right) \frac{\partial V_i}{\partial \lambda_i} = O\left(\frac{\mu^{3/4}}{\lambda_i^{3/2} \lambda_j}\right).$$

Lastly, we write

$$\int_{\Omega} (3U_j^5 - \alpha_j^4 V_j^5) \frac{\partial V_i}{\partial \lambda_i} = 3 \int_{\Omega} (U_j^5 - V_j^5) \frac{\partial V_i}{\partial \lambda_i} + (3 - \alpha_j^4) \int_{\Omega} V_j^5 \frac{\partial V_i}{\partial \lambda_i}.$$

Proceeding as in the proof of (C.10), we find

$$\int_{\Omega} V_j^5 \frac{\partial V_i}{\partial \lambda_i} = O\left(\frac{\mu^{1/2}}{\lambda_i^{1/2} \lambda_j^{1/2}}\right).$$

Concerning the last integral that we have to estimate, setting as previously  $B_i = B(a_i, \gamma/4\mu^{1/2})$ , we have, similarly to (C.15)

$$\int_{B_i} \left| \frac{\partial V_i}{\partial \lambda_i} \right| = O\left(\frac{\ln(\lambda_i/\mu^{1/2})}{\lambda_i^{7/2}} + \frac{1}{\lambda_i^{3/2} \mu}\right)$$

and

$$\begin{aligned} & \int_{\Omega} |U_j^5 - V_j^5| \\ &= O\left(\int_{\Omega} (U_j^4 \varphi_j + \varphi_j^5)\right) \\ &= O\left(\frac{1}{\lambda_j^{1/2}} \int_0^{2R'\lambda_j} \frac{1 - e^{-(\mu^{1/2}/\lambda_j)r}}{(1+r^2)^2} r^2 dr + \frac{\mu}{\lambda_j^{5/2}} \int_0^{2R'\mu^{1/2}} \frac{(1 - e^{-r})^5}{r^3} dr\right) \\ &= O\left(\frac{\mu^{1/2}}{\lambda_j^{3/2}}\right). \end{aligned}$$

As we have also

$$\frac{\partial V_i}{\partial \lambda_i} = O\left(\frac{\mu^{1/2}}{\lambda_j^{3/2}}\right) \text{ outside of } B_i \quad U_j^5 - V_j^5 = O\left(\frac{\mu^{5/2}}{\lambda_j^{5/2}}\right) \text{ in } B_i$$

we obtain

$$\int_{\Omega} (U_j^5 - V_j^5) \frac{\partial V_i}{\partial \lambda_i} = O\left(\frac{\mu}{\lambda_i^{3/2} \lambda_j^{3/2}}\right)$$

and (C.20) is proved.

## References

- [1] Adimurthi, G. Mancini - *The Neumann problem for elliptic equations with critical nonlinearity*, "A tribute in honour of G. Prodi", Scuola Norm. Sup. Pisa (1991), 9-25.
- [2] Adimurthi, G. Mancini - *Geometry and topology of the boundary in the critical Neumann problem*, J. reine angew. Math. **456** (1994), 1-18.
- [3] Adimurthi, G. Mancini, S.L. Yadava - *The role of the mean curvature in semilinear Neumann problem involving critical exponent*, Comm. Part. Diff. Equ. **20** (1995), 591-631.
- [4] Adimurthi, F. Pacella, S.L. Yadava - *Interaction between the geometry of the boundary and positive solutions of a semilinear Neumann problem with critical nonlinearity*, J. Funct. Anal. **113** (1993), 318-350.
- [5] Adimurthi, F. Pacella, S.L. Yadava - *Characterization of concentration points and  $L^\infty$ -estimates for solutions of a semilinear Neumann problem involving the critical Sobolev exponent*, Diff. and Integ. Equ. **8** (1995), 41-68.
- [6] Adimurthi, S.L. Yadava - *Existence and nonexistence of positive radial solutions of Neumann problems with critical Sobolev exponents*, Arch. Rat. Mech. Anal. **115** (1991), 275-296.
- [7] A. Bahri - *Critical points at infinity in some variational problems*, Pitman Research Notes in Math. Series **182**, Longman (1989).
- [8] A. Bahri, J.M. Coron - *On a nonlinear elliptic equation involving the critical Sobolev exponent : the effect of the topology of the domain*, Comm. Pure Appl. Math. **41** (1988), 255-294.
- [9] A. Bahri, Y.Y. Li, O. Rey - *On a variational problem with lack of compactness : the topological effect of the critical point at infinity*, Calc. Var. PDE **3** (1995), 67-93.
- [10] P. Bates, G. Fusco - *Equilibria with many nuclei for the Cahn-Hilliard equation*, J. Diff. Equ. **160** (2000), 283-356.
- [11] H. Brezis, J.M. Coron - *Convergence of solutions of H-systems or how to blow bubbles*, Arch. Rational Mech. Anal. **89** (1985), 21-56 .
- [12] C. Budd, M.C. Knaap, L.A. Peletier - *Asymptotic behaviour of solutions of elliptic equations with critical exponent and Neumann boundary conditions*, Proc. Roy. Soc. Edinburgh **117 A** (1991), 225-250.
- [13] L. Caffarelli, B. Gidas, J. Spruck - *Asymptotic symmetry and local behavior of semilinear elliptic equations with critical Sobolev growth*, Comm. Pure Appl. Math. **42** (1989), 271-297.

- [14] D. Cao, E.S. Noussair, S. Yan - *Existence and nonexistence of interior-peaked solution for a nonlinear Neumann problem*, to appear.
- [15] G. Cerami, D. Fortunato, M. Struwe - *Bifurcation and multiplicity results for nonlinear elliptic problems involving critical Sobolev exponents*, Ann. Inst. H. Poincaré, Anal. non lin. **1** (1984), 341-350.
- [16] G. Cerami, J. Wei - *Multiplicity of multiple interior peaks solutions for some singularly perturbed Neumann problems*, Intern. Math. Res. Notes **12** (1998), 601-626.
- [17] E.N. Dancer, S. Yan - *Multipeak solutions for a singularly perturbed Neumann problem*, Pacific J. Math. **189** (1999), 241-262.
- [18] N. Ghoussoub, C. Gui, M. Zhu - *On a singularly perturbed Neumann problem with the critical exponent*, to appear.
- [19] M. Grossi, A. Pistoia - *On the effect of critical points of distance function in superlinear elliptic problems*, Adv. Diff. Equ. **5** (2000), 1397-1420.
- [20] M. Grossi, A. Pistoia, J. Wei - *Existence of multipeak solutions for a semilinear elliptic problem via nonsmooth critical point theory*, Calc. Var. PDE **11** (2000), 143-175.
- [21] C. Gui - *Multi-peak solutions for a semilinear Neumann problem*, Duke Math. J. **84** (1996), 739-769.
- [22] C. Gui, N. Ghoussoub - *Multi-peak solutions for a Neumann problem involving the critical Sobolev exponent*, Math. Z. **229** (1998), 443-474.
- [23] C. Gui, C.S. Lin - *Estimates for boundary bubbling solutions to an elliptic Neumann problem*, to appear in Comm. Part. Diff. Equ.
- [24] C. Gui, J. Wei - *Multiple interior peak solutions for some singularly perturbed Neumann problems*, J. Diff. Equ. **158** (1999), 1-27.
- [25] C. Gui, J. Wei - *On the existence of arbitrary number of bubbles for some semilinear elliptic equations with critical Sobolev exponent*, to appear.
- [26] C. Gui, J. Wei - *Multiple mixed boundary and interior peak solutions for some singularly perturbed Neumann problems*, to appear in Canadian J. Math.
- [27] M. Kowalczyk - *Multiple spike layers in the shadow Gierer-Meinhardt system : existence of equilibria and quasi-invariant manifold*, Duke Math. J. **98** (1999), 59-111.
- [28] Y.Y. Li - *Prescribing scalar curvature on  $S^n$  and related problems, Part. I*, J. Diff. Equ. **120** (1995), 319-410.

- [29] Y.Y. Li - *On a singularly perturbed equation with Neumann boundary condition*, Comm. Part. Diff. Equ. **23** (1998), 487-545.
- [30] Y.Y. Li, M. Zhu - *Yamabe type equations on three dimensional Riemannian manifolds*, Comm. Contemp. Math. **1** (1999), 1-50.
- [31] C.S. Lin, W.M. Ni - *On the Diffusion Coefficient of a Semilinear Neumann Problem*, Springer Lecture Notes **1340**, Springer, New York, Berlin (1986).
- [32] C.S. Lin, W.N. Ni, I. Takagi - *Large amplitude stationary solutions to a chemotaxis system*, J. Diff Equ. **72** (1988), 1-27.
- [33] S. Maier-Paape, K. Schmitt, Z.Q. Wang - *On Neumann problems for semi-linear elliptic equations with critical nonlinearity : existence and symmetry of multi-peaked solutions*, Comm. Part. Diff. Equ. **22** (1997), 1493-1527.
- [34] H. Meinhardt - *Models of biological pattern formation*, Academic Press, London, New York (1982).
- [35] W.N. Ni, X.B. Pan, I. Takagi - *Singular behavior of least-energy solutions of a semi-linear Neumann problem involving critical Sobolev exponents*, Duke Math. J. **67** (1992), 1-20.
- [36] W.N. Ni, I. Takagi - *On the shape of least-energy solutions to a semi-linear problem Neumann problem*, Comm. Pure Appl. Math. **44** (1991), 819-851.
- [37] W.M. Ni, I. Takagi - *Locating the peaks of least-energy solutions to a semi-linear Neumann problem*, Duke Math. J. **70** (1993), 247-281.
- [38] O. Rey - *Boundary effect for an elliptic Neumann problem with critical nonlinearity*, Comm. Part. Diff. Equ. **22** (1997), 1055-1139.
- [39] O. Rey - *The topological impact of critical points at infinity ia a variational problem with lack of compactness : the dimension 3*, Adv. Diff. Equ. **4** (1999), 581-616.
- [40] O. Rey - *An elliptic Neumann problem with critical nonlinearity in three dimensional domains*, Comm. Contemp. Math. **1** (1999), 405-449.
- [41] R. Schoen - *On the number of constant scalar curvature metrics in a conformal class*, in *Differential Geometry : A symposium in honor of Manfredo Do Carmo* (H.B. Lawson and K. Tenenblat eds.), Wiley, New York (1991), 311-320.
- [42] A.M. Turing - *The chemical basis of morphogenesis*, Phil. Trans. R. Soc. **B 237** (1952), 37-52 (see also *Morphogenesis* in Coll. Works of A.M. Turing (P.T. Saunders ed.), North-Holland Publishing Co., Amsterdam (1992).
- [43] X.J. Wang - *Neumann problem of semilinear elliptic equations involving critical Sobolev exponents*, J. Diff. Equ. **93** (1991), 283-310.

- [44] Z.Q. Wang - *Remarks on a nonlinear Neumann problem with critical exponent*, Houston J. Math. **20** (1994), 671-684.
- [45] Z.Q. Wang - *The effect of domain geometry on the number of positive solutions of Neumann problems with critical exponents*, Diff. and Integ. Equ. **8** (1995), 1533-1554.
- [46] Z.Q. Wang - *High energy and multi-peaked solutions for a nonlinear Neumann problem with critical exponent*, Proc. Roy. Soc. Edimburgh **125 A** (1995), 1003-1029.
- [47] Z.Q. Wang - *Construction of multi-peaked solution for a nonlinear Neumann problem with critical exponent*, J. Nonlinear Anal. TMA **27** (1996), 1281-1306.
- [48] J. Wei - *On the interior spike layer solutions of singularly perturbed semi-linear Neumann problems*, Tohoku Math. J. **50** (1998), 159-178.
- [49] J. Wei, M. Winter - *Stationary solutions for the Cahn-Hilliard equation*, Ann. Inst. H. Poincaré, Anal. non lin. **15** (1998), 459-482.
- [50] J. Wei, X. Xu - *Uniqueness and a priori estimates for some nonlinear elliptic Neumann equations in  $\mathbb{R}^3$* , to appear.
- [51] S. Yan - *On the number of interior multipeak solutions for singularly perturbed Neumann problems*, Top. Meth. Nonlin. Anal. **12** (1998), 61-78.
- [52] M. Zhu - *Uniqueness results through a priori estimates, I. A three dimensional Neumann problem*, J. Diff. Equ. **154** (1999), 284-317.