



**HAL**  
open science

## Draft genome sequence of *Lactobacillus hominis* strain CRBIP 24.179 $\tau$ , isolated from human intestine

Sylvie Cousin, Sophie Creno, Laurence Ma, Dominique Clermont, Valentin  
Loux, Chantal Bizet, Christiane Bouchier

### ► To cite this version:

Sylvie Cousin, Sophie Creno, Laurence Ma, Dominique Clermont, Valentin Loux, et al.. Draft genome sequence of *Lactobacillus hominis* strain CRBIP 24.179 $\tau$ , isolated from human intestine. *Genome Announcements*, 2013, 1 (4), pp.1. 10.1128/genomeA.00662-13 . hal-00939673

**HAL Id: hal-00939673**

**<https://hal.science/hal-00939673>**

Submitted on 30 Jan 2014

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# Draft Genome Sequence of *Lactobacillus hominis* Strain CRBIP 24.179<sup>T</sup>, Isolated from Human Intestine

Sylvie Cousin,<sup>a</sup> Sophie Creno,<sup>b</sup> Laurence Ma,<sup>b</sup> Dominique Clermont,<sup>a</sup> Valentin Loux,<sup>c</sup> Chantal Bizet,<sup>a</sup> Christiane Bouchier<sup>b</sup>

Institut Pasteur, Centre de Ressources Biologiques de l'Institut Pasteur, Département de Microbiologie, Paris, France<sup>a</sup>; Institut Pasteur, Plate-forme Génomique, Département Génomes et Génétique, Paris, France<sup>b</sup>; INRA, UR1077 Mathématique, Informatique et Génome, Jouy-en-Josas, France<sup>c</sup>

**We report the draft genome sequence of the strain *Lactobacillus hominis* CRBIP 24.179<sup>T</sup>, isolated from a human clinical sample. The total length of the 28 contigs is about 1.9 Mb, with a G+C content of 37% and 1,983 coding sequences.**

Received 26 July 2013 Accepted 29 July 2013 Published 22 August 2013

**Citation** Cousin S, Creno S, Ma L, Clermont D, Loux V, Bizet C, Bouchier C. 2013. Draft genome sequence of *Lactobacillus hominis* strain CRBIP 24.179<sup>T</sup>, isolated from human intestine. *Genome Announc.* 1(4):e00662-13. doi:10.1128/genomeA.00662-13.

**Copyright** © 2013 Cousin et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 3.0 Unported license](http://creativecommons.org/licenses/by/3.0/).

Address correspondence to Christiane Bouchier, [bouchier@pasteur.fr](mailto:bouchier@pasteur.fr).

*Lactobacillus hominis* strain 61D<sup>T</sup>, renamed *L. hominis* CRBIP 24.179<sup>T</sup>, was isolated in the early 1960s from a clinical isolate (1). Some species of the genus *Lactobacillus* have been isolated from clinical patients with a variety of clinical problems (2, 3). Most of those strains (but not all) belong to the species *Lactobacillus rhamnosus*, *Lactobacillus casei*, *Lactobacillus paracasei*, and *Lactobacillus plantarum*. These species are also among those most frequently found in human intestinal flora. Nevertheless, there is only limited information concerning the risk that is linked to potential virulence factors present in *Lactobacillus* strains (4).

Here, we report the genome sequence of *L. hominis* CRBIP 24.179<sup>T</sup>, obtained using a whole-genome strategy based on Illumina paired-end sequencing, with an insert length of 380 bp (Illumina genome analyzer HiSeq 2000). In order to test assembly *de novo* tools, quality-filtered reads (116,085,864 reads, 99 bases mean read length, ~5,190-fold coverage) were assembled using two softwares, VelvetOptimiser version 2.2.0 (5) and ABySS version 1.2.6 (6). The resulting contigs were compared using Mauve version 2.3.1 (7). The two scaffolds were somewhat similar (28 Velvet contigs and 37 ABySS contigs, with maximum lengths of 365,062 and 267,206 bases, respectively). Nevertheless, differences related to the number and the maximum length of contigs and Mauve assessment support the *de novo* assembled contigs obtained using Velvet.

The draft genome consists of 36 contigs, with a total length of 1,927,726 nucleotides (nt) and a G+C content of 37%. The sizes of the contigs were between 701 bases for the shortest and 365,062 bases for the longest. The contigs were annotated with the AGMIAL platform (8), an integrated bacterial genome annotation system. The prediction of coding sequences used the self-training gene detection software SHOW based on hidden Markov models (<http://genome.jouy.inra.fr/ssb/SHOW/>). tRNAs and rRNAs were detected using tRNAscan-SE (9) and RNAmmer (10) softwares, respectively. There were 1,983 coding sequences (CDSs) predicted (1,938 complete), as well as one rRNA operon with 1 copy each of the 23S, 5S, and 16S genes. Fifty-four tRNA genes were also predicted.

**Nucleotide sequence accession numbers.** The strain is publicly available in two European collections under the no. CRBIP

24.179<sup>T</sup> and DSM 23910<sup>T</sup>. The draft of this whole-genome sequencing project has been deposited in EMBL under the accession no. [CAKE01000001](https://www.ebi.ac.uk/ena/record/CAKE01000001) to [CAKE01000036](https://www.ebi.ac.uk/ena/record/CAKE01000036). The version described in this paper is the first version.

## ACKNOWLEDGMENTS

We thank N. Joly (Biology IT Center, Institut Pasteur, Paris) for the programs used for the quality filtering of the reads.

This communication is an initiative of the European Consortium of Microbial Resource Centers (EMbaRC), supported by the European Commission's Seventh Framework Programme (FP7, 2007–2013), Research Infrastructures action under the grant agreement no. FP7-228310.

## REFERENCES

1. Cousin S, Motreff L, Gulat-Okalla ML, Gouyette C, Spröer C, Schumann P, Begaud E, Bouchier C, Clermont D, Bizet C. 2013. *Lactobacillus pasteurii* sp. nov. and *Lactobacillus hominis* sp. nov. *Int. J. Syst. Evol. Microbiol.* 63:53–59.
2. Aguirre M, Collins MD. 1993. Lactic acid bacteria and human clinical infection. *J. Appl. Bacteriol.* 75:95–107.
3. Bourne KA, Beebe JL, Lue YA, Ellner PD. 1978. Bacteremia due to *Bifidobacterium*, *Eubacterium* or *Lactobacillus*, twenty-one cases and review of the literature. *Yale J. Biol. Med.* 51:505–512.
4. Gasser F. 1994. Safety of lactic acid bacteria and their occurrence in human clinical infections. *Bull. Inst. Pasteur* 92:45–67.
5. Zerbino DR, Birney E. 2008. Velvet: algorithms for *de novo* short read assembly using de Bruijn graphs. *Genome Res.* 18:821–829.
6. Simpson JT, Wong K, Jackman SD, Schein JE, Jones SJ, Birol I. 2009. ABySS: a parallel assembler for short read sequence data. *Genome Res.* 19:1117–1123.
7. Darling AC, Mau B, Blattner FR, Perna NT. 2004. Mauve: multiple alignment of conserved genomic sequence with rearrangements. *Genome Res.* 14:1394–1403.
8. Bryson K, Loux V, Bossy R, Nicolas P, Chaillou S, van de Guchte M, Penaud S, Maguin E, Hoebeke M, Bessières P, Gibrat JF. 2006. AGMIAL: implementing an annotation strategy for prokaryote genomes as a distributed system. *Nucleic Acids Res.* 34:3533–3545.
9. Lowe TM, Eddy SR. 1997. tRNAscan-SE: a program for improved detection of transfer RNA genes in genomic sequence. *Nucleic Acids Res.* 25:955–964.
10. Lagesen K, Hallin P, Rødland EA, Staerfeldt HH, Rognes T, Ussery DW. 2007. RNAmmer: consistent and rapid annotation of ribosomal RNA genes. *Nucleic Acids Res.* 35:3100–3108.