



HAL
open science

Improving the MicroResp (TM) substrate-induced respiration method by a more complete description of CO₂ behavior in closed incubation wells

Pierre P. Renault, Meriem M. Ben Sassi, Annette A. Berard

► To cite this version:

Pierre P. Renault, Meriem M. Ben Sassi, Annette A. Berard. Improving the MicroResp (TM) substrate-induced respiration method by a more complete description of CO₂ behavior in closed incubation wells. *Geoderma*, 2013, 207, pp.82-91. 10.1016/j.geoderma.2013.05.010 . hal-00939766v1

HAL Id: hal-00939766

<https://hal.science/hal-00939766v1>

Submitted on 29 May 2020 (v1), last revised 1 Aug 2016 (v2)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Improving the MicroResp™ substrate-induced respiration method by a more complete description of CO₂ behavior in closed incubation wells

P. Renault ^{a,*}, M. Ben-Sassi ^{a,b,1}, A. Bérard ^{a,1}

^a INRA, UMR 1114 EMMAH, Domaine St-Paul, Site Agroparc, 84914 Avignon Cedex 9, France

^b ADEME, 20 avenue du Grésillé, BP 90406, F-49004 Angers Cedex 01, France

A B S T R A C T

The MicroResp™ method allows soil respiration and microbial community physiological profiles to be determined colorimetrically in microplates. This method, however, neglects CO₂ storage in the agar gel carrying the colorimetric indicator, and calcite dissolution associated with CO₂-induced change in soil solution pH. Our objective was to improve the method by taking into account CO₂ in the gel in the calculation of microbial respiration, describing the effect of microbial CO₂ on the pH of the soil solution and calcite dissolution, and checking whether CO₂ distribution among calcite, soil solution, air and gel is near equilibrium after incubation. We propose a thermodynamic equilibrium model describing (a) distribution of CO₂ among calcite, soil solution, gel and air, (b) dissociations of water, carbonic acid, cresol red, and substrates in the gel and soil solution, (c) exchange of adsorbed cations with H₃O⁺ in the gel, and (d) calcite dissolution in soil. In-gel experiments were designed to calibrate the model, quantify the rate of CO₂ exchange with air, and compare conservation procedures. On-soil experiments were designed to check whether calcite dissolution is near equilibrium and whether the model predicts the effect of CO₂ on the pH of the solution. In-microplate experiments were designed to assess the effects of incubation period and soil quantity on estimated microbial respiration. The model can describe the distribution and speciation of CO₂ in the gel, the soil solution and the air space of each microplate well. Initial properties of the gel vary with storage: soda lime partly extracts CO₂ supplied as NaHCO₃, and dries out the gel, which can skew the calibration. When incubation is over, the proportion of microbial CO₂ in the gel is higher at lower microbial respiration. Incubations shorter than 4 h underestimate microbial respiration due to the slow diffusion of CO₂ in the gel. CO₂ in the soil solution cannot be overlooked; it decreases the soil pH and may promote calcite dissolution in calcareous soil. It is important to precisely estimate initial CO₂ air fraction and to control temperature, which affects both thermodynamic constants and microorganisms.

Keywords:

Soil
Substrate-induced respiration
Carbon dioxide
Geochemistry
MicroResp
Cresol red

1. Introduction

Soil microbial respiration that consumes O₂ may regulate CH₄ and N₂O greenhouse gas emissions and reduction of metals (Lahlah et al., 2009; Parry et al., 2000), organic matter turnover (Schlesinger and Andrews, 2000), and acid-base, complexation and precipitation/dissolution reactions (Dassonville et al., 2004; Lahlah et al., 2009). Profiling microbial

respiration on various substrates provides an insight into microbial functional diversity (Chapman et al., 2007; Degens and Harris, 1997).

Microbial respiration has been assessed by O₂ consumption (Garland et al., 2003), by CO₂ production (Cheng and Coleman, 1989), and by coupling the two (Sierra and Renault, 1995). Respiration is more often characterized by CO₂ measurements, which are easier and sensitive (Dilly, 2001). However, many characterizations suffer from common limitations (long gas analysis, large volumes required, complex set-up).

The MicroResp™ method is a miniaturized substrate-induced respiration method that overcomes these limitations (Campbell et al., 2003), and offers a wide range of applications (Ben Sassi et al., 2012; Tlili et al., 2011): it couples the microplate format of the Biolog™ test restricted to cultivable micro-organisms (Garland and Mills, 1991; Stefanowicz, 2006) with the measurement of CO₂ air fraction according to the work of Rowell (1995) on indicator dyes in agar gel. In each closed well of a 96-well microplate, moist soil with or without C substrates is incubated for 6 h in the presence of an agar gel carrying cresol red as indicator dye (Campbell et al.,

* Corresponding author at: INRA, UMR 1114 EMMAH (Environnement Méditerranéen et Modélisation des Agro-Hydrosystèmes), Domaine Saint-Paul, Site Agroparc, 84914 Avignon Cedex 9, France. Tel.: +33 4 32 72 22 23; fax: +33 4 32 72 22 12.

E-mail addresses: pierre.renault@avignon.inra.fr (P. Renault), meriem.bensassi@avignon.inra.fr (M. Ben-Sassi), annette.berard@avignon.inra.fr (A. Bérard).

¹ Fax: +33 4 32 72 22 12.

2003). The method only takes into account CO₂ in the well air space, which is assumed to be of microbial origin (Campbell et al., 2003). The CO₂ air fraction is estimated from gel absorbance at 570 nm (Rowell, 1995). The MicroResp™ method has been widely used on soils with pH < 7 (Burton et al., 2010; Campbell et al., 2003; Gonzalez-Quiñones et al., 2009; Lalor et al., 2007; Macdonald et al., 2009; Wakelin et al., 2008; Yao et al., 2011; Zhou et al., 2011) and occasionally used on soils with higher pH and calcite (Bérard et al., 2011, 2012; Saul-Tcherkas and Steinberger, 2009; Wakelin et al., 2008), although the technical manual recommends restricting it to soil with pH < 7 (Cameron, 2007). The method was modified by Oren and Steinberger (2008a) to include CO₂ in the soil solution and the effects of substrates on the soil solution pH and calcite dissolution, but without considering the effects of CO₂ on calcite dissolution and on the pH of the solution (Stumm and Morgan, 1996). Their improvements, which require measurements on sterilized soil and evaluation of substrate impact on soil pH, have not been widely adopted to date (García-Palacios et al., 2011; Oren and Steinberger, 2008b). To our knowledge, CO₂ in the agar gel has never been taken into account.

Thus the MicroResp™ method still suffers from limitations. First, the volume of in-well gel (0.15 mL) cannot be neglected given the volumes of soil solution (about 0.12 mL) and air space (about 1 mL). The pH of the gel (from 7 to 9 or more) magnifies the problem with more HCO₃⁻ and sometimes CO₃²⁻ than H₂CO₃* (i.e., aqueous CO₂ and actual H₂CO₃) in the gel solution (Stumm and Morgan, 1996) and in solutions of calcareous soils (Ström et al., 2001). Second, failing to allow for the fact that increasing the CO₂ air fraction decreases the pH of the soil solution (Stumm and Morgan, 1996) ultimately overestimates the amount of CO₂ in the solution. For calcareous soils, increasing the CO₂ air fraction may also induce calcite dissolution (Stevenson and Verburg, 2006; Stumm and Morgan, 1996; Tamir et al., 2011), leading to an abiotic production of total CO₂ in the soil solution as H₂CO₃*, HCO₃⁻ and CO₃²⁻, even without acidic substrates. Third, no check has been made to determine whether the transfers between calcite, soil solution, air and gel can be considered to be in equilibrium. Failure to approach equilibrium would make calibration impossible. In addition, although tackled by Oren and Steinberger (2008a), it remains important to know whether most of the calcite dissolution caused by acidic substrate occurs before or after microplate clamping.

Accordingly, the objective of this study was to improve the MicroResp™ method by (i) including CO₂ in the gel carrying the colorimetric indicator, (ii) refining microbial CO₂ in soil by describing its effects along with those of substrates and calcite on the pH of the solution, and quantifying calcite dissolution, and (iii) checking whether CO₂ distribution among calcite, soil solution, air space and the gel is near-balanced after 6 h incubation.

2. Materials and methods

The study combines modeling of CO₂ distribution during MicroResp™ incubations based on geochemical equilibrium with measurements performed on the gel, on a calcareous soil, and on both combined in the MicroResp™ experimental design. For certain incubations, the soil was sterilized and/or supplied with substrate (e.g., glucose, glucosamine-HCl and Na₂-malate as neutral, acid and alkaline substrates, respectively).

2.1. Background of the MicroResp™ method

The MicroResp™ experimental design consists of two 96-well microtiter plates placed face-to-face (Campbell et al., 2003). One of the plates, with a capacity of 300 μL · well⁻¹, holds 150 μL · well⁻¹ of an agar gel (10 g · L⁻¹) enriched in KCl (0.15 mol · L⁻¹), NaHCO₃ (2.5 mmol · L⁻¹) and cresol red dye (32.7 μmol · L⁻¹) to estimate

CO₂ air fraction based on gel absorbance at 570 nm. After preparation, this plate is generally conserved for 7 d in a closed environment with soda lime and water, without and with a protective Parafilm during the first day and the 6 following days, respectively. The other plate, with a capacity of 1.2 mL · well⁻¹, holds about 0.45 g · well⁻¹ of moist soil with or without substrate. Just before incubation, the plate containing the gel is read with an absorbance microplate reader. The two plates are then sealed together with a silicone rubber gasket with interconnecting holes. After 6 h of incubation, the plates are separated and the plate containing the gel is immediately re-read. We have adapted the method to 24-, 12- and 6-well microplates with silicone rubber gaskets manufactured in our lab to seal identical plates.

2.2. Modeling of in-well CO₂ distribution and its effect on gel absorbance

The modeling study aimed to describe the relationships between gel absorbance, CO₂ air fraction, amounts of total CO₂ in both gel and soil solution (H₂CO₃* (i.e., aqueous CO₂ and actual H₂CO₃), HCO₃⁻, CO₃²⁻), and calcite dissolution. Thermodynamic equilibrium was considered for transfers between soil, air space and gel, for acid-base reactions in solutions, for exchange of adsorbed cations with H₃O⁺ in the gel solution, and for calcite dissolution. For the gel and the soil solution, mass action laws were combined with equations relating the balance of ionic charges and the total quantity of either cresol red or substrate. Mass action laws were written with *K'* constants combining H₃O⁺ activity and concentrations of other species. In soil solution where ionic strength *I* ≈ 15 mmol · L⁻¹ (Table 3), activities are almost equal to concentrations, so *K'* constants were assumed to be equal to *K* constants combining activities only. In gel solution where *I* ≈ 150 mmol · L⁻¹, *K'* constants were estimated from *K** constants combining concentrations only or *K* constants using species activities only. Deviations between *K* and *K** are consistent with independent estimates of activity coefficients.

CO₂ distribution among air space, gel and soil solution was described by Henry's law. Henry constants, *k_H* for water and soil solution and *k_H'* for saline solution and gel, were estimated from the equations of Harned and Davis (1943) and Weiss (1974), respectively (Table 1). Deviation between *k_H* and *k_H'* is consistent with the increase in H₂CO₃* activity coefficient in the gel calculated by the Pytkowicz's (1975) equation (Table 1). The fugacity coefficient of CO₂ in air was set to 1 (DOE, 1994). H₂O activity in gel and soil solution was set to 1 (Stumm and Morgan, 1996). *K_w* and *K_w'* constants for H₂O ion product were estimated from the equations of Harned and Owen (1958) and Millero (1995), respectively (Table 1). *K_{a1}* and *K_{a2}* constants for the first and second dissociation of H₂CO₃*, respectively, were estimated from the equations of Weiss (1974) (Table 1), and *K_{a1}'* and *K_{a2}'* constants were estimated from the equations of Millero et al. (2007) for NaCl solutions (Table 1), since they differ from those for seawaters (Millero et al., 2006). In the gel, cresol red, symbolized by H₂CR, dissociates into HCR⁻ and CR²⁻. The first dissociation is complete, as its p*K_{CR1}* constant is about 1.1 (El Nahhal et al., 2012; French et al., 2002; Heger et al., 2006; Smith and Matachek, 2002), and the *K_{CR2}* constant of the second dissociation was estimated by the equation of French et al. (2002) (Table 1). *K_{CR2}'* was deduced from *K_{CR2}* and HCR⁻ and CR²⁻ activity coefficients. An exchange of adsorbed cations with protons in the gel was taken into account empirically (see Eq. (7) below). Activity coefficients of ions in the gel solution were calculated by the Davies equation (Table 1) (Pankow, 1991), except for CR²⁻ where the two negative charges have to be considered independently, as is the case with other diprotic acids used as indicator dyes (Salvatore et al., 1986). We thus extended the relationship that uses specific interaction theory for bromophenol blue (Salvatore et al., 1986) in order to assess realistic variations in the p*K_{CR2}* of the second dissociation of cresol red at 25 °C (Table 1). In soil, dissociations of acid and alkaline substrates

Comment citer ce document :

Renault, P., Ben Sassi, M., Berard, A. (2013). Improving the MicroResp (TM) substrate-induced respiration method by a more complete description of CO₂ behavior in closed incubation wells. Geoderma, 207, 82-91. DOI : 10.1016/j.geoderma.2013.05.010

Table 1
 Thermodynamic parameters involved in the calculation of geochemical equilibrium. R is the gas constant ($8.31441 \text{ J} \cdot \text{mol}^{-1} \cdot \text{K}^{-1}$), T the temperature (K), P the air pressure (Pa), and S the salinity of the agar gel solution (expressed as equivalent g NaCl $\cdot \text{kg}^{-1}$).

Constant	Equation	References
k_{H} : Henry's constant for CO_2 solubilization in pure water	$\frac{1}{k_{\text{H}}} = 10^{(-((\frac{-282.38}{T}) + (-0.0178471 \times T) + 15.5873)) \times \frac{1000 \times R \times T}{P}}$	Harned and Davis (1943)
k_{H}' : Henry's constant CO_2 solubilization in saline water	$\frac{1}{k_{\text{H}}'} = \exp \left(\frac{9345.17}{T} + 23.3585 \times \ln \left(\frac{T}{100} \right) - 60.2409 + S \times \left(0.023517 + 0.00470356 \times \left(\frac{T}{100} \right)^2 - 0.023656 \times \left(\frac{T}{100} \right) \right) \right) \times \frac{1000 \times R \times T}{P}$	Weiss (1974)
$\gamma_{\text{H}_2\text{CO}_3^*}$: activity of H_2CO_3^* in saline water	$\gamma_{\text{H}_2\text{CO}_3^*} = 10^{0.00035863 + (0.00196297 \times S)}$	Pytkowicz (1975)
K_{w} : ion product of water in pure water	$\text{p}K_{\text{w}} = -\log_{10}(K_{\text{w}}) = \frac{4470.99}{T} - 6.0875 + 0.01706 \times T$	Harned and Owen (1958)
K_{w}' : ion product of water in saline water	$\text{p}K_{\text{w}}' = -\log(K_{\text{w}}') = -\log \left(\exp \left(\frac{148.9802 + \left(\frac{-13847.26}{T} \right) + (-23.6521 \times \ln(T))}{+ \left(\left(-5.977 + \frac{118.67}{T} + 1.0495 \times \ln(T) \right) \times (S^{0.5}) \right) + (-0.01615 \times S)} \right) \right)$	Millero (1995)
$K_{\text{a}1}$: first dissociation constant of carbonic acid in pure water	$\text{p}K_{\text{a}1} = -\log(K_{\text{a}1}) = -114.3106 + \left(\frac{5773.67}{T} \right) + (17.779524 \times \ln(T))$	Weiss (1974)
$K_{\text{a}1}^*$: first dissociation constant of carbonic acid in saline water	$\text{p}K_{\text{a}1}^* = -\log(K_{\text{a}1}^*) = \text{p}K_{\text{a}1} + A_1 + \frac{B_1}{T} + (C_1 \times \ln(T))$ with : $A_1 = 35.2911 \times m^{0.5} + 0.8491 \times m - 0.32 \times m^{1.5} + 0.055 \times m^2$ $B_1 = -1583.09 \times m^{0.5}$ $C_1 = -5.4366 \times m^{0.5}$	Millero et al. (2007)
$K_{\text{a}2}$: second dissociation constant of carbonic acid in pure water	$\text{p}K_{\text{a}2} = -\log(K_{\text{a}2}) = -83.2997 + \left(\frac{4821.38}{T} \right) + (13.5962 \times \ln(T))$	Weiss (1974)
$K_{\text{a}2}^*$: second dissociation constant of carbonic acid in saline water	$\text{p}K_{\text{a}2}^* = -\log(K_{\text{a}2}^*) = \text{p}K_{\text{a}2} + A_2 + \frac{B_2}{T} + (C_2 \times \ln(T))$ with : $A_2 = 38.2746 \times m^{0.5} + 1.6057 \times m - 0.647 \times m^{1.5} + 0.113 \times m^2$ $B_2 = -1738.16 \times m^{0.5}$ $C_2 = -6.0346 \times m^{0.5}$	Millero et al. (2007)
$K_{\text{CR}2}$: second dissociation constant of cresol red in pure water	$\text{p}K_{\text{CR}2} = -\log(\text{p}K_{\text{CR}2}) = 2.049 + \left(\frac{913.4}{T} \right) + (1.266 \times \log(T))$	French et al. (2002)
$\gamma_{\text{CR}^{2-}}$: activity coefficient for cresol red anion CR^{2-} with 2 e^- charges	$\gamma_{\text{CR}^{2-}} = \left(2 \times \left(-0.5107 \times \left(\frac{\sqrt{I}}{1 + (1.5 \times \sqrt{I})} \right) \right) \right) - (-0.178 \times m)$	Empirical equation deduced from
γ_i^* : Davis equation of activity coefficients for ionic compound i , having a charge z_i , except for CR^{2-}	$\log(\gamma_i) = -A \times z_i^2 \times \left(\left(\frac{\sqrt{I}}{1 + \sqrt{I}} \right) - 0.2 \times I \right)$	Pankow (1991)

were taken into account: reactions considered for substrates containing more than one acid or alkaline functional group were chosen based on their $\text{p}K$ and the pH of the soil (Table 2). For calcareous soils, the solubility constant K_{S} for calcite was set to 8.3, mirroring data reported by Stumm and Morgan (1996) at 25 °C. The simulated increase of $[\text{Ca}^{2+}]$ during incubations makes it possible to assess the release of abiotic CO_2 . Geochemical simulations were performed on soils at various CO_2 air fractions for different substrates and amounts of substrates. pH was adjusted so as to cancel the sum of charges of ions in solution. The variation $\Delta[\text{CO}_2]_{\text{tot}}$ in total CO_2 concentration in the solution (H_2CO_3^* , HCO_3^- and CO_3^{2-}) minus the increase $\Delta[\text{Ca}^{2+}]$ in Ca^{2+} amount for calcareous soils between the considered and 0.04% CO_2 air fractions was then fitted by empirical functions:

$$\Delta[\text{CO}_2]_{\text{tot}} - \Delta[\text{Ca}^{2+}] = a \times ([\text{CO}_2]_{\text{a}})^b - c, \quad (1)$$

Table 2
 Retained acid-base reactions for substrates used in MicroResp™ measurements for soils whose $\text{pH}(\text{water})$ is about 7.5–8.5.

Substrate supplied	Acid form	Charge of the acid form	Charge of the alkaline form	$\text{p}K_{\text{a}}$
Glucosamine-HCl	$\text{C}_6\text{H}_{13}\text{NO}_5\text{-HCl}$	+1	0	11.5
Na_2 -malate	$\text{Na}_2\text{-C}_4\text{H}_4\text{O}_5$	-1	-2	5.1
Alanine	$\text{C}_3\text{H}_7\text{NO}_2$	0	-1	9.71
Glycine	$\text{C}_2\text{H}_5\text{NO}_2$	0	-1	9.58

where the three parameters a , b and c are specific to each combination of soil, substrate type and substrate amount. Considering initial and final CO_2 air fractions eliminates the reference to 0.04% CO_2 that was introduced to facilitate the adjustment of empirical functions. For each soil and each substrate, parameters a , b and c can be obtained by fitting Eq. (1) to geochemical simulations based on equations proposed in this paper. It requires having first at 0.04% CO_2 rough estimates of the sum of charges of ions other than H_3O^+ , HO^- , HCO_3^- and CO_3^{2-} in the soil solution and, for calcareous soils, the contribution of Ca^{2+} to this sum. These values can be estimated from chemical analysis and by fitting simulated pH to pH measured on soil slurries at different CO_2 partial pressures.

The measured dimensionless absorbance A_{pH} is the sum of the absorbances of empty microplate A_{e} , agar gel without cresol red A_{g} , and cresol red A_{CR} :

$$A_{\text{pH}} = A_{\text{e}} + A_{\text{g}} + A_{\text{CR}}. \quad (2)$$

A_{CR} may be expressed as a function of HCR^- and CR^{2-} concentrations:

$$A_{\text{CR}} = \varepsilon_{\text{HCR}^-} \times l_{\text{g}} \times [\text{HCR}^-] + \varepsilon_{\text{CR}^{2-}} \times l_{\text{g}} \times [\text{CR}^{2-}], \quad (3)$$

where l_{g} is the path length of light through the gel (m), and $\varepsilon_{\text{HCR}^-}$ and $\varepsilon_{\text{CR}^{2-}}$ the molar absorptivities of HCR^- and CR^{2-} ($\text{L} \cdot \text{mol}^{-1} \cdot \text{m}^{-1}$), respectively. Combining Eqs. (2) and (3) with a mass action law for the second dissociation of cresol red and a related equation for the

stability of total cresol red amount leads to the following estimate of gel solution pH:

$$\text{pH} = \text{p}K_{\text{CR}2'} + \log_{10} \left(\frac{A_{\text{pH}} - (A_e + A_g) - (A_{\text{max}} \times \left(\frac{\varepsilon_{\text{HCR}^-}}{\varepsilon_{\text{CR}2^-}} \right))}{A_{\text{max}} + (A_e + A_g) - A_{\text{pH}}} \right), \quad (4)$$

where $A_{\text{max}} (= \varepsilon_{\text{CR}2^-} \times l_g \times [\text{CR}_{\text{tot}}])$ is the absorbance of cresol red when it is in $\text{CR}2^-$ form only. The ratio $\varepsilon_{\text{HCR}^-} / \varepsilon_{\text{CR}2^-}$ was set to 0.0019, in line with Smith and Matachek (2002) at 574 nm. The concentration of H_2CO_3^* in the gel solution may then be estimated as:

$$[\text{H}_2\text{CO}_3^*] = \frac{\Delta q_g + \left(\frac{10^{-\text{pH}}}{\gamma_{\text{H}_3\text{O}^+}} \right) + 10^{\text{pH} - \text{p}K_w'}}{10^{\text{pH} - \text{p}K_{\text{a}1'}} + (2 \times 10^{2\text{pH} - (\text{p}K_{\text{a}1}' + \text{p}K_{\text{a}2}')})}, \quad (5)$$

where Δq_g is the net charge of ions other than H_3O^+ , HO^- , HCO_3^- and CO_3^{2-} in the gel solution ($\text{mol c}^+ \cdot \text{L}^{-1}$). It would be equal to the concentration $[\text{Na}^+]_{\text{NaHCO}_3}$ of Na^+ supplied as NaHCO_3 ($\text{mol} \cdot \text{L}^{-1}$) if agar did not affect the solution, which is not the case (Ferreira et al., 2012; Lahaye and Rochas, 1991; Scholten and Pierik, 1998):

$$\Delta q_g = [\text{Na}^+]_{\text{NaHCO}_3} + \Delta c^+, \quad (6)$$

where Δc^+ is the net concentration of charges ($\text{mol c}^+ \cdot \text{L}^{-1}$) of ions released by agar. Δc^+ increases with decreasing pH of the gel, as in other media (Renault et al., 2009), so we assumed the following equation:

$$\Delta c^+ = \Delta c_{\text{min}}^+ + \left(\Delta c_{\text{max}}^+ \times \left(1 - \frac{\left(\frac{\text{pH}}{\text{pH}_{1/2}} \right)^\alpha}{1 + \left(\frac{\text{pH}}{\text{pH}_{1/2}} \right)^\alpha} \right) \right), \quad (7)$$

where Δc_{min}^+ is the concentration of charges released by agar at a high pH ($\text{mol c}^+ \cdot \text{L}^{-1}$), Δc_{max}^+ the maximum concentration of charges released during acidification ($\text{mol c}^+ \cdot \text{L}^{-1}$), $\text{pH}_{1/2}$ the pH at which one half of the initially-adsorbed charges are released, and α an empirical constant related to the sharp or gentle release of charges with changes in pH around $\text{pH} \approx \text{pH}_{1/2}$. $Q_g(\text{CO}_2)$, i.e., the amount of CO_2 stored in the gel (mol) as H_2CO_3^* , HCO_3^- and CO_3^{2-} is then easily estimated:

$$Q_g(\text{CO}_2) = V_g \times [\text{H}_2\text{CO}_3^*] \times \left(1 + 10^{\text{pH} - \text{p}K_{\text{a}1}'} + (2 \times 10^{2\text{pH} - (\text{p}K_{\text{a}1}' + \text{p}K_{\text{a}2}')}) \right), \quad (8)$$

where V_g is the volume of the gel (L). While the CO_2 air fraction is initially the lab CO_2 air fraction not in equilibrium with the gel, its final value is assumed to be in equilibrium with the gel and estimated from Henry's law (Table 1):

$$[\text{CO}_2]_a = k_H' \times [\text{H}_2\text{CO}_3^*]_g, \quad (9)$$

where subscripts a and g stand for air and gel, respectively. The amount of CO_2 stored in the air space is estimated as:

$$Q_a(\text{CO}_2) = V_a \times [\text{CO}_2]_a, \quad (10)$$

where V_a is volume of the air space (L). Thus variations in CO_2 quantities in both gel and air space are estimated from the amounts of CO_2 in these compartments at the start and the end of incubations, assuming that the gel is not in equilibrium with the air space or the soil solution at the start. By contrast, empirical functions (Eq. (1)) are used to estimate the variations $\Delta Q_s(\text{CO}_2)$ of amount of CO_2 in soil solution (mol) minus CO_2 released by calcite dissolution:

$$\Delta Q_s(\text{CO}_2) = V_s \times (\Delta[\text{CO}_2]_{\text{tot}, t=6\text{h}} - [\text{CO}_2]_{\text{tot}, t=0}), \quad (11)$$

where V_s is the volume of the soil solution (L). A negative value means that abiotic CO_2 production exceeds the total amount of CO_2 stored in the soil solution.

In addition, we assume that the variability in A_{pH} between wells of a given plate results from the variability in light path length l_g . Initial and final absorbances A_{pH} of each well were therefore replaced by absorbances A_{pH}' by taking into account the initial absorbance of the target well $A_{\text{pH}}(t=0)$ and the mean absorbance $\overline{A_{\text{pH}}}(t=0)$ across all the wells in a plate:

$$A_{\text{pH}}' = A_e + \left(\left(\frac{\overline{A_{\text{pH}}}(t=0) - A_e}{A_{\text{pH}}(t=0) - A_e} \right) \times (A_{\text{pH}} - A_e) \right). \quad (12)$$

In this way, A_{pH} , A_g and A_{max} can be replaced by A_{pH}' , $\overline{A_g}$ and $\overline{A_{\text{max}}}$, respectively, in Eq. (4). When $A_e \approx 0$, Eq. (12) approaches the correction of Campbell et al. (2003):

$$A_{\text{pH}}' \approx \left(\frac{\overline{A_{\text{pH}}}(t=0)}{A_{\text{pH}}(t=0)} \right) \times A_{\text{pH}}. \quad (13)$$

At about 80 °C, the gel is distributed in greater amounts than desired, i.e., 0.20, 0.94, 1.74 and 4.09 mL instead of 0.15, 0.664, 1.314 and 3.27 mL for 96-, 24-, 12- and 6-well plates, respectively, and mean gel thickness $\overline{l_g}$ (mm) varies with plates, i.e., 5.51, 4.68, 4.38 and 4.13 mm, for 96-, 24-, 12- and 6-well plates, respectively. Therefore, $\overline{A_g}$ and $\overline{A_{\text{max}}}$ were estimated from measured values on 96-well microplates according to the following equations:

$$\overline{A_g}(n_w = i) = \left(\frac{\overline{l_g}(n_w = i)}{\overline{l_g}(n_w = 96)} \right) \times \overline{A_g}(n_w = 96), \quad (14a)$$

$$\overline{A_{\text{max}}}(n_w = i) = \left(\frac{\overline{l_g}(n_w = i)}{\overline{l_g}(n_w = 96)} \right) \times \overline{A_{\text{max}}}(n_w = 96), \quad (14b)$$

where n_w is the number of wells of the microplate considered.

Thus estimating the amount of microbial CO_2 produced during MicroResp™ incubations hinges on knowing the values of three parameters concerning absorbance (A_e , $\overline{A_g}$ and $\overline{A_{\text{max}}}$) and four parameters concerning the exchange properties of the gel (Δc_{min}^+ , Δc_{max}^+ , $\text{pH}_{1/2}$, and α). All other parameters may be estimated from the literature.

2.3. Experimental approach

2.3.1. The soil

Measurements were performed on a calcareous cambisol (FAO classification) from the INRA Saint-Paul experimental station (43°91' N, 4°88' E) near Avignon. It was cultivated with peas in 2008, but since summer 2008 the soil lay bare. Annual rainfall is about 650 mm. About 20 kg of this soil was sampled in the first 10 cm depth at the edge of a 0.0075 ha experimental field on 8 December 2011. Its moisture was about 21.3 % wt/wt. The soil was air-dried for 4 d in the lab (14 % wt/wt residual moisture), mechanically crushed, sieved at 2 mm for soil analyses and at 2–3 mm for experiments, and stored at 4 °C in hermetically-sealed bags until the beginning of the experiments. Its properties, measured at the Laboratoire d'Analyse des Sols (LAS-INRA) in Arras (France), were as follows: 347 g kg^{-1} CaCO_3 and, after decarbonation, 323 g kg^{-1} clay; 259 g kg^{-1} silt; 41 g kg^{-1} sand; 13.2 g kg^{-1} organic C; 1.54 g kg^{-1} total N; 1.4 mg kg^{-1} N-NH_4^+ ; 101 mg kg^{-1} N-NO_3^- . Soil pH(water) and pH(KCl 1 M) were 8.51 and 7.85, respectively. Chemical properties of solutions for soil-to-solution mass ratios of 1, 2.5 and 5 (Table 3) were used to assess composition of the soil solution in MicroResp™ incubations. For geochemical simulations on this soil, we assumed that the sum of charges of ions other than H_3O^+ , HO^- , HCO_3^- and CO_3^{2-} was equal to 4.6 mmol $\text{c}^+ \cdot \text{L}^{-1}$, including the initial concentration of Ca^{2+} equal to 6 mmol $\text{c}^+ \cdot \text{L}^{-1}$.

Comment citer ce document :

Table 3

Chemical properties of the solutions for mixtures of soil-to-solution mass ratios of 1, 2.5 and 5.

Dilution	pH(water)	Ca ²⁺	Mg ²⁺	K ⁺	NO ₃ ⁻	Cl ⁻
mmol · L ⁻¹						
5	7.56	0.47	0.049	0.212	0.28	0.019
2.5	7.84	0.71	0.078	0.252	0.54	0.025
1	7.88	1.38	0.173	0.360	1.12	0.057
Limit for ω = 31% ^a	7.97	2.96	0.426	0.523	3.06	0.120

^a Linear regression of Ln values.

2.3.2. Experiments on the gel

The agar product used in this work was the Merk 1614 product. Its properties are presented and compared with those of other market products in Scholten and Pierik (1998).

2.3.2.1. Calibration and assessment of the relationship between A_{pH} and CO_2 air fraction. We used an EL800 absorbance microplate reader (BioTek Instruments Inc., Winooski, VT) to measure (i) absorbance of empty microplates (A_e), (ii) mean absorbance of microplates filled with gel prepared as in the MicroResp™ method (Campbell et al., 2003), but excluding cresol red ($A_e + \overline{A_g}$), and (iii) mean absorbance of microplates filled with gel with or without added cresol red and in which NaHCO₃ 2.5 mmol · L⁻¹ was replaced by NaOH 10 mmol · L⁻¹ to ensure a pH > 11. The mean maximum absorbance of cresol red as CR²⁻ ($\overline{A_{max}}$) was estimated by the difference in absorbances at pH > 11. Just after reading the absorbances, gel pH was measured in 8 wells of each microplate using a 16-gauge micro-combination needle pH probe (MI-414B, Microelectrodes, Inc., Bedford, NH) gently touching the bottom of the wells and connected to a Bioblock Scientific 93327 pH-EH meter (HANNA Instruments, Smithfield, RI). Additional measurements were performed to assess the relationship between A_{pH} and the pH of the gel solution: for agar gel with or without added cresol red and buffered at pH ≈ pK_{CR2} by replacing NaHCO₃ 2.5 mmol · L⁻¹ with H₃BO₃ 0.50 mol · L⁻¹ and NaOH 0.65 mol · L⁻¹, and for microplates prepared according to the MicroResp™ method after 24 h in sealed bags containing either dry soda lime and water in distinct capsules or air enriched with 1.5% CO₂. To estimate Δc_{min}^+ , Δc_{max}^+ , pH_{1/2} and α , we set out to obtain the relationship between Δq_g and gel pH after equilibrating the gel with air enriched by 0.5%, 1% and 1.5% CO₂. To better circumvent the effect of the gel, we simultaneously measured the pH of a solution of the same composition, but without agar, equilibrated with CO₂-enriched air. However, the calculations are highly sensitive to gel pH, and 0.5% or more CO₂ bars insight into pHs higher than 7.6. Therefore, parameters Δc_{min}^+ , Δc_{max}^+ , pH_{1/2} and α were empirically refined to ensure that (i) the calculated CO₂ disappearing from the air space of empty wells over 6 h balanced against the CO₂ stored in the gel of these wells in 96-well microplates, and (ii) the respiration rate estimated for 6 h incubation in wells of 24-well microplates filled with either 0.6, 1.2, 1.8 or 2.4 g of moist soil did not depend on the final CO₂ air fraction in the wells. The relationship between gel absorbance and CO₂ air fraction was checked for microplates filled with gel previously equilibrated 5 d with CO₂ air fractions of 0.5, 1, 1.5 and 2% at about 22.5 °C after being stored uncovered in sealed plastic bags with soda lime and water for 6 d.

2.3.2.2. Kinetics of CO₂ exchange between the air space and the gel. Just after being filled with gel, two 96-well microplates were stored for 1 d in sealed plastic bags that contained dry soda lime and water in distinct containers, and two other microplates in sealed plastic bags with air enriched by 1.5% CO₂. Just after opening a bag, one of the two microplates was used for absorbance measurements every 2 to 5 min for 6 h at about 22.5 °C, while the other was used to record the pH of the gel every 1 to 5 min in a single well, using the 16-gauge micro-combination needle pH probe. Beyond 6 h, drying led to cracks in the gel of some wells.

2.3.2.3. Effect of microplate storage on the initial state of the gel. Since soda lime partly dries the gel and extracts CO₂ supplied as NaHCO₃, we checked whether absorbance varied between microplates prepared simultaneously and kept for 7 d in a poorly-closed desiccator with dry soda lime and water in separate capsules, in a sealed plastic bag with dry soda lime and water in separate capsules, and in a sealed plastic bag with soda lime and water in the same capsule. In addition, we monitored the kinetics of gel desiccation in microplates stored for 7 d in sealed plastic bags with dry soda lime and water in separate capsules, one of them being covered with a plastic film after 1 d of storage, the other remaining exposed.

2.3.3. Experiment on the soil

2.3.3.1. Ability of the model to simulate the pH of soil solutions for various CO₂ air fractions. Since retained reactions greatly simplify the complex geochemistry of soils, we checked whether they permit the pH of a soil solution to be simulated and whether the calcite dissolution that results from an increase in CO₂ air fraction or the supply of an acid substrate could be described by a thermodynamic equilibrium. The relationship between CO₂ air fraction and pH of the soil solution was assessed on soil slurries with a water-to-soil mass ratio of 1, 2.5 and 5 after equilibrating with either lab air (between 0.04 and 0.12% CO₂) or air enriched by 0.5, 1, 1.5 and 2% CO₂, respectively, by sparging with air and stirring the slurry.

2.3.3.2. Abiotic CO₂ emissions after supplying acid substrate to the soil. Soil was autoclaved at 121 °C and 0.1 MPa for 1 h, incubated for 2 d, then autoclaved a second time for 1 h at the same temperature and pressure to eliminate any microorganisms that were not destroyed in the initial autoclaving (Skipper et al., 1996). Then 24 wells of a 96-well microplate were filled with sterilized soil, while 24 other wells were filled with non-sterilized soil. For each modality, 6 wells each were supplemented with either 0.05 mL of water or with a mixture of 0.025 mL water plus 0.025 mL of a 120 mg · mL⁻¹ solution of glucose, glucosamine-HCl or Na₂-malate, respectively. Abiotic emissions due to glucosamine-HCl supply were estimated from 6 h incubation and compared with biotic emissions on unsterilized soil.

2.3.4. MicroResp™ incubations of various times and for various amounts of soil

Three MicroResp™ incubation sets were performed to assess and illustrate the new MicroResp™ data analysis. In a first set of incubations, the wells of four 24-well microplates were filled with 0.6, 1.2, 1.8 or 2.4 g of the same soil at 19 wt.% soil moisture supplemented with 0.033, 0.066, 0.099 or 0.133 mL of water, respectively, and 0.033, 0.066, 0.099 or 0.133 mL of a 120 mg · mL⁻¹ solution of glucose. The microplates were then incubated at 22.5 °C for 6 h. Results served to estimate the parameters α and pH_{1/2} and to assess the relative contributions of gel, air space and soil solution to CO₂ storage. In a second set of incubations, the wells of five 96-well microplates were either left empty or filled with about 0.38 g of soil initially at 19 wt.% soil moisture supplemented with 0.05 mL of water or 0.025 mL of water and 0.025 mL of a 120 mg · mL⁻¹ solution of glucose, Na₂-malate or glucosamine-HCl. The microplates were then incubated at 22.5 °C for 1, 2, 4, 6 and 8 h. Results served to check whether estimated microbial respiration was incubation time-dependent. A third set of experiments was performed solely to illustrate variations in microbial CO₂ production with changes in substrates. Only one 96-well microplate was used; the wells were filled with about 0.38 g of dry soil initially at 19 wt.% soil moisture and supplemented with 0.05 mL of water or with 0.025 mL of water and 0.025 mL of a 120 mg · mL⁻¹ solution of glucose, sucrose, trehalose, mannose, cellobiose, dextrin, glucosamine-HCl, alanine, glycine or Na₂-malate.

Comment citer ce document :

Renault, P., Ben Sassi, M., Berard, A. (2013). Improving the MicroResp (TM) substrate-induced respiration method by a more complete description of CO₂ behavior in closed incubation wells. *Geoderma*, 207, 82-91. DOI : 10.1016/j.geoderma.2013.05.010

3. Results and discussion

3.1. Calibration of the relationship between A_{pH} and the CO_2 air fraction

The means and standard deviations of the absorbances of empty microplates (A_e) were 0.030 and 0.001, respectively. The means and standard deviations of the absorbances of the wells filled with gel without cresol red ($A_e + A_g$) were 0.171 and 0.009, respectively. Thus an estimate of \bar{A}_g is 0.141, the variability in A_e can be neglected given its low contribution to A_{pH} , and the standard deviation of A_g is nearly 0.009, leading to a variation coefficient of A_g of about 6.2%. Similar estimates for A_g mean and standard deviation were obtained for agar gel without cresol red and for $pH \approx 11.7$ and $pH = pK_{CR2}'$, although slightly lower when borate was used to buffer the gel solution. The means and standard deviations of the absorbances of microplates with gel carrying cresol red and having a $pH > 11$ were 0.987 and 0.030, respectively, while the means and standard deviations of absorbances of microplates with gel without cresol red and having a $pH > 11$ were 0.162 and 0.008, respectively. Thus an estimate of \bar{A}_{max} is 0.825. Although an exact estimation of the A_{max} variation coefficient is impossible, it should be lower than that of A_g in our experiment. Using these values and the calculated pK_{CR2}' to simulate variations in the absorbance A_{pH} as a function of the pH of the gel solution correctly reflects experimental data (Fig. 1). The pH generally decreases from 9 to 10 at the beginning of incubation (depending on the efficiency of the soda lime to remove CO_2 from the gel) to more than 7.1, a value rarely reached, which corresponds approximately to 2% of CO_2 in the air space.

The initial vs. final absorbances of empty wells (i.e., $A_{pH}(t = 0)$ and $A_{pH}(t = 6 \text{ h})$, respectively) strongly suggest that Δc_{min}^+ is close to or slightly lower than zero, and that there is no exchange of adsorbed cations with H_3O^+ in the gel solution at $pH \geq 8$: the opposite (i.e., $\Delta c_{min}^+ \geq 0$ and/or $\Delta c^+ \gg 0$) would require an unrealistically high initial CO_2 air fraction in the air space of the wells to explain the change in absorbance measured for these wells (results not shown). The pHs of the solution of the same composition as the gel in the MicroResp™ method but without agar and in equilibrium with 0.5%, 1% and 1.5% CO_2 were 7.49, 7.16 and 6.98, respectively. These values correspond to a net Δq_g of about 3.05, 2.85 and 2.82 $mmol \text{ c}^+ \cdot L^{-1}$, respectively, which is near the 2.5 $mmol \cdot L^{-1}$ of Na^+ supplied as $NaHCO_3$. By contrast, the pHs of the agar gel in equilibrium with 0.5%, 1% and 1.5% CO_2 were 7.66, 7.41 and 7.22, respectively. These values correspond to a net Δq_g of about 4.50,




Fig. 1. Relationship between pH of the agar gel and mean of the measured absorbance A_{pH} ($A_{pH} = A_e + A_g + A_{CR}$) over the 96 wells of a microplate. The X coordinates of the experimental points are directly-measured pH. The Y coordinates of the experimental points are directly-measured absorbances when the gel matched the composition used in the MicroResp™ method, whereas for non-matched gels they were recalculated to replace the absorbance of the modified gel by the actual absorbance of the gel matched to the composition used in the MicroResp™ method.




Fig. 2. Relationship between means of measured absorbance A_{pH} ($A_{pH} = A_e + A_g + A_{CR}$) over the 96 wells of a microplate and CO_2 air fraction in equilibrium with the microplates.

5.03 and 4.89 $mmol \text{ c}^+ \cdot L^{-1}$, respectively, suggesting that Δq_g increases with a decrease in pH in this pH range with a Δc_{max}^+ probably higher than 2. We set Δc_{max}^+ to 3.0 $mmol \text{ c}^+ \cdot L^{-1}$ in order to obtain a good fit of the relationship between the air CO_2 fraction and the absorbance at 570 nm on experimental data for 0.5%, 1%, 1.5% and 2% CO_2 (Fig. 2). Parameters $pH_{1/2}$ and α were set to 7.7 and 80, respectively, by minimizing differences in the respiration rates of various amounts of the same soil (see below), making it possible to simulate a majority of cation exchange for pH values between 7.4 and 8.

3.2. Effects of soil alkalinity, substrate and calcite on soil pH and microbial CO_2 in solution

While solute concentrations were highly dependent on water-to-soil mass ratio (Table 3), pH measured on soil slurries in equilibrium with a range of CO_2 air fractions was only slightly dependent on water-to-soil mass ratio (1, 2.5 and 5 in this work), but highly dependent on CO_2 air fraction (Fig. 3). Simulations performed for soil solutions in equilibrium with calcite and having an initial $[Ca^{2+}]$ near the value extrapolated for soil solutions at the start of MicroResp™ incubations were used to simulate slurry behavior. Although this was not the objective of these measurements, the simulations confirmed that pH(water) (8.51 for our soil) differs from the pH of the soil solution in equilibrium with lab air (pH slightly lower than 8 here), since alkaline soil solution can trap a large amount of gaseous CO_2 , but only very slowly in the absence of facilitated transfer (vigorous stirring and bubbling).

Empirical functions were then easily obtained from geochemical modeling to estimate the amount of microbial CO_2 stored in the soil




Fig. 3. Measured pH of soil slurry as a function of CO_2 air fraction in equilibrium with it by simultaneous air bubbling and slurry stirring. Simulations were performed for soil at weight moistures as in MicroResp™ incubations.

Comment citer ce document :




Fig. 4. Simulated amount of total CO_2 ($\text{mol} \cdot \text{L}^{-1}$) minus CO_2 derived from calcite dissolution as a function of air space CO_2 fraction for a soil solution in equilibrium with calcite and initially having a sum of charges other than H_3O^+ , HO^- , HCO_3^- and CO_3^{2-} equal to $-4.60 \cdot 10^{-3} \text{ mol c}^+ \cdot \text{L}^{-1}$ and $+6.00 \cdot 10^{-3} \text{ mol c}^+ \cdot \text{L}^{-1} \text{Ca}^{2+}$, and for different amounts of Na_2 -malate substrate (0, 10, 20, 30 and 42 $\text{g} \cdot \text{L}^{-1}$). Zero values indicate that total CO_2 in the soil solution is equal to CO_2 release by calcite dissolution. Negative values mean that calcite dissolution releases more CO_2 than was initially contained in the soil solution, indicating that CO_2 is partly transferred to the air and agar gel in the well.

solution (H_2CO_3^* , HCO_3^- and CO_3^{2-}) minus the amount of total abiotic CO_2 emitted by calcite dissolution. The variation of $\Delta[\text{CO}_2]_{\text{tot}}$ in total CO_2 concentration in the solution (H_2CO_3^* , HCO_3^- and CO_3^{2-}) minus the increase $\Delta[\text{Ca}^{2+}]$ in Ca^{2+} concentration for calcareous soils between the considered CO_2 air fraction and the 0.04% CO_2 air fraction was then fitted by empirical functions. An example of these functions for several amounts of Na_2 -malate is proposed in Fig. 4, and empirical coefficients defined in Eq. (1) for these functions are presented in Table 4 for the relevant soil.

3.3. Kinetics of transfer and calcite dissolution and the effects on MicroResp™ estimation

Microbial respiration rate was assessed in a soil sterilized and subsequently supplied with a slightly acidic substrate (glucosamine-HCl) to indirectly check whether calcite dissolution can be described by a thermodynamic equilibrium system. The results showed that abiotic CO_2 microbial respiration can be neglected, as absolute values were small ($3.50 \cdot 10^{-1} \mu\text{g C-CO}_2 \cdot \text{g}^{-1} \text{soil} \cdot \text{h}^{-1}$) compared with unsterilized soil enriched with the same substrate ($3.23 \mu\text{g C-CO}_2 \cdot \text{g}^{-1} \text{soil} \cdot \text{h}^{-1}$), and positive or negative sign was dependent on the initial in-lab CO_2 air fraction.

Fig. 5a–b illustrates the changes in the pH of the gel solution and in the absorbance A_{pH} of microplates initially stored in sealed plastic

Table 4

Retained coefficients a , b and c used in Eq. (1) to describe variations in amount of microbial CO_2 in the soil solution (i.e., the total amount of $\text{H}_2\text{CO}_3^* + \text{HCO}_3^- + \text{CO}_3^{2-}$ reduced by the contribution of calcite dissolution to this pool) with regard to microbial CO_2 at 0.04% CO_2 air fraction:

Substrate	Concentration	a	b	c
–		$1.83 \cdot 10^{-3}$	$6.45 \cdot 10^{-1}$	$2.28 \cdot 10^{-4}$
Glycine		$5.77 \cdot 10^{-2}$	$2.57 \cdot 10^{-2}$	$5.25 \cdot 10^{-2}$
Alanine		$5.20 \cdot 10^{-3}$	$2.50 \cdot 10^{-1}$	$2.28 \cdot 10^{-3}$
Glucosamine-HCl		$1.85 \cdot 10^{-3}$	$6.40 \cdot 10^{-1}$	$2.33 \cdot 10^{-4}$
Na_2 -malate	10	$1.93 \cdot 10^{-3}$	$6.38 \cdot 10^{-1}$	$2.46 \cdot 10^{-4}$
	20	$2.03 \cdot 10^{-3}$	$6.31 \cdot 10^{-1}$	$2.64 \cdot 10^{-4}$
	30	$2.13 \cdot 10^{-3}$	$6.24 \cdot 10^{-1}$	$2.83 \cdot 10^{-4}$
	42	$2.25 \cdot 10^{-3}$	$6.17 \cdot 10^{-1}$	$3.28 \cdot 10^{-4}$




Fig. 5. Changes (a) in absorbance A_{pH} and (b) in the pH of the gel solution of microplates initially stored in sealed plastic bags enriched with either 1.5% CO_2 or with soda lime and water in separate capsules, and exposed to lab air.

bags either enriched with 1.5% CO_2 or containing soda lime and water in distinct capsules and exposed to lab air. Superimposed yellow and pink color layers (corresponding to CO_2 -enriched air and CO_2 -free air, respectively) in microplate well gel were clearly visible for long periods of these experiments. These observations indicate that CO_2 transfer between the air space and the gel is low and that equilibrium is reached only after about 4 h. This is mainly because low CO_2 transfer in the gel prevents a quick redistribution of CO_2 when it accumulates or disappears in the gel near the interface with air. For MicroResp™ microplate wells where microbial CO_2 production was continuous over the 6 h of incubation, it is reasonable to consider that equilibrium is never reached and that the estimated microbial respiration a priori underestimates actual respiration. However, the relative bias resulting from the equilibrium hypothesis will decrease with increasing incubation times. Staggered incubations (1, 2, 4, 6 and 8 h) of soil enriched with glucose suggest that 4 h is the minimum time requirement for incubations and that the 6 h incubation proposed in the MicroResp™ method is reasonable (Fig. 6).




Fig. 6. Microbial respiration rates ($\mu\text{g C-CO}_2 \cdot \text{g}^{-1} \text{dry soil} \cdot \text{h}^{-1}$) estimated from soil incubation in 96-well microplates for 1, 2, 4, 6 and 8 h, without substrate (Y coordinates on the right) and with glucose (Y coordinates on the left).

Comment citer ce document :

Initial state of the gel may vary with initial storage time and conditions. For example, the initial absorbance of 96-well microplates varies between simultaneously-prepared microplates stored for 7 d in a poorly-closed desiccator with dry soda lime and water in separate capsules ($\overline{A}_{pH}(t=0) = 0.90$), a sealed plastic bag with dry soda lime and water in separate capsules ($\overline{A}_{pH}(t=0) = 1.04$), and a sealed plastic bag with soda lime and water in the same capsule ($\overline{A}_{pH}(t=0) = 0.97$). For plates stored in sealed plastic bags, differences in $\overline{A}_{pH}(t=0)$ between dry soda lime and water in separate capsules and in the same capsule may have resulted from differences in gel desiccation, which would explain why $\overline{A}_{pH}(t=0)$ exceeds the maximum simulated absorbance ($A_e + \overline{A}_g + A_{max} = 0.996$). The kinetics of gel desiccation in microplates stored for 7 d in sealed plastic bags with dry soda lime and water in separate capsules varies greatly with the presence or absence of a plastic film covering the wells (Fig. 7). The procedure proposed in the MicroResp™ method therefore appears to have been adapted to protect the gel from desiccation. However, the initial absorbance may still vary with the amount of CO₂ extracted by soda lime during plate storage. This amount has to be taken into account in microbial respiration estimates. However, the worst-case scenario would be the gel drying out, which would ultimately modify the value of the sum Δq_g , and skew the calibration. For this reason, it is essential to take all feasible precautions to protect the gel from partial drying. Repeated use of the same microplate gels “regenerated” between consecutive incubations by exposure to soda lime should be ruled out. Finally, we note that microplates with an initial absorbance $\overline{A}_{pH}(t=0)$ of about 0.90 would correspond to a gel of about pH 9.1 and about 0.0073% CO₂ air fraction, indicating that the NaOH partially replaces NaHCO₃ within the gel during storage due to CO₂ extraction by the soda lime.

3.4. The effects of soil mass and substrate on respiration estimated from 6 h incubation

The 24-well microplates that were filled with 0.6, 1.2, 1.8 and 2.4 g of the same soil at 19 wt.% soil moisture and supplemented with 0.033, 0.066, 0.099 and 0.133 mL of water and 0.033, 0.066, 0.099 and 0.133 mL of a 120 mg · mL⁻¹ solution of glucose, respectively, were then incubated at 22.5 °C for 6 h. Although the results partly contributed to calibration (pH_{1/2} and α were estimated so as to minimize differences in estimated microbial respirations between these four soil weights), they show that estimated soil microbial respiration is not dependent on soil weight and therefore not dependent on CO₂ air fraction at the end of the incubations (Fig. 8). This is all the more interesting as there is a variation in the relative proportions of total CO₂ in the gas phase in the soil solution and in the gel (Fig. 9). The accumulation of CO₂ during the 6 h of incubation is always stronger in the gel than in the air space and the soil solution, especially




Fig. 7. Change in the weight of microplates filled with agar gel during its initial 7-day storage in sealed bags with or without Parafilm anti-desiccant.




Fig. 8. Microbial respiration rates ($\mu\text{g C-CO}_2 \cdot \text{g}^{-1} \text{ dry soil} \cdot \text{h}^{-1}$) estimated for different amounts of soil (0.6, 1.2, 1.8 and 2.4 g of fresh soil at 19 wt.% soil moisture) supplied with water and glucose.

when final CO₂ content is low. Low final levels of CO₂ air fraction are common for soils incubated without substrates or with inefficient, poorly-used substrates. These results clearly demonstrate that we cannot neglect variations in the amount of CO₂ in the gel. The fraction of CO₂ stored in the soil solution cannot be neglected, but it can be estimated with less precision in view of its low contribution to the total accumulation of microbial CO₂.

We then tested this new mathematical procedure for estimating microbial respiration from MicroResp™ incubation on various substrates (Fig. 10). As already observed in Fig. 9, the new method leads to estimates of microbial respiration rates 2 to 4 times higher than estimates considering only CO₂ accumulation in the air space of the wells. Microbial respiration rates decreased in the following order of added substrate: sucrose \geq glucose \geq malate \geq glucosamine-HCl \geq cellobiose \geq alanine \geq mannose \geq dextrin \geq trehalose, and microbial respiration rates were between 5 and 15 times stronger in the soil with substrate than in the soil with water only. The coefficients of variation of individual measurements (i.e., in a well) ranged from 6.3% (soil with water) to 53% (soil + trehalose).

4. Conclusion

Here we propose a novel and more complete description of the fate of CO₂ during soil incubation in the wells of microplates used in the MicroResp™ method; for the first time, the storage of CO₂ (as H₂CO₃^{*}, HCO₃⁻ and CO₃²⁻) in the gel carrying the colorimetric indicator and the effects of CO₂ on the pH of the soil solution and, in calcareous soils, the dissolution of calcite are now taken into account. The comparison between the experimental data and simulations shows that this improved model makes it possible to simulate the main geochemical processes involved in the fate of CO₂. It yields appropriate estimates of microbial CO₂ production, as shown by 6 h-plus incubations of 0.6 to 2.4 g of soil (in 24-well microplates). An incubation time of 6 h is a good compromise between the need for short incubations to prevent microbial growth/selection and the need for long incubations to reduce the gradient in CO₂ distribution within the gel and between the gel and the air space. The final proportion of CO₂ stored in the indicator gel is far from negligible, especially when the final CO₂ air fraction is low, as is the case for incubation of soil without substrate or with substrates that are difficult to consume. Incubation for less than 4 h may underestimate microbial respiration, since CO₂ dissolution and transfer within the gel are both slow processes. The amount of CO₂ in the gel at the start of incubation depends on the level of CO₂ (supplied as NaHCO₃) extraction by soda lime while the gel is in storage. Extracted CO₂ varies with microplate distance to soda lime, duration of the exposure to soda lime, etc. Storage may partially desiccate the gel and modify the




Fig. 9. Distribution of microbial production of CO₂ in microplate well gel, air space and soil solution as a function of (a) 0.6 g, (b) 1.2 g, (c) 1.8 g and (d) 2.4 g of fresh soil (19wt.% soil moisture) supplied with water and glucose, corresponding to estimated final CO₂ air fractions equal to 0.32, 0.57, 0.99 and 1.74% CO₂. For the calculations, abiotic CO₂ originating from calcite CO₂ was assumed to be in the soil solution only. This slightly minimizes the contribution of soil solution to CO₂ storage based on calculations assuming even CO₂ distribution between soil solution, air space and gel.

relationship between CO₂ partial pressure and gel absorbance, and so repeated use of the same microplates “regenerated” between consecutive incubations by exposure to soda lime should be avoided. CO₂ stored in the soil cannot be neglected, but an increase in CO₂ partial pressure decreases the pH of the soil solution and, in the case of calcareous soils, generally promotes calcite dissolution. Neglecting this decrease in pH or the dissolution of calcite would lead to an overestimation of biotic CO₂ production. Since microbial CO₂ production in soils without substrate often leads to a final CO₂ fraction of less than 0.4%, it is important to have a good estimate of the initial CO₂ partial pressure (varying between 0.04 and more than 0.1% in lab air), possibly by having empty wells with gel. It is also vital to keep firm control over




Fig. 10. Example illustrating microbial respiration rate for several added substrates. Microbial respiration rates have been estimated by taking into account either CO₂ accumulation in air only or CO₂ accumulation in the gel, air and soil solution minus the CO₂ issued from calcite dissolution.

temperature, which can affect all the thermodynamic constants as well as microbial activity.

Authors can provide on request an Excel file to perform all calculations.

Acknowledgments

This work received support from the CleanWast Project that features as part of the French ANR [National Research Agency] PRECODD Programme. We thank the INRA's Département Environnement et Agronomie and the ADEME for providing support through M. Ben-Sassi's PhD grant. We also thank Ghislain Sévenier and Cécile Péru for valuable technical assistance, and Glen McCulley for English editing and proofreading.

References

Ben Sassi, M., Dolinger, J., Renault, P., Tlili, A., Bérard, A., 2012. The FungiResp method: an application of the MicroResp™ method to assess fungi in microbial communities as soil biological indicators. *Ecological Indicators* 23, 482–490.

Bérard, A., Ben Sassi, M., Renault, P., Gros, R., 2012. Severe drought-induced community tolerance to heat wave. An experimental study on soil microbial processes. *Journal of Soils and Sediments* 12, 513–518.

Bérard, A., Bouchet, T., Sévenier, G., Pablo, A.L., Gros, R., 2011. Resilience of soil microbial communities impacted by severe drought and high temperature in the context of Mediterranean heat-waves. *European Journal of Soil Biology* 47, 333–342.

Burton, J., Chen, C., Xu, Z., Ghadiri, H., 2010. Soil microbial biomass, activity and community composition in adjacent native and plantation forests of subtropical Australia. *Journal of Soils and Sediments* 10, 1267–1277.

Cameron, C., 2007. *MicroResp™ Technical Manual – A Versatile Soil Respiration System*. Macaulay Institute, Craigiebuckler, Aberdeen, Scotland, UK.

Campbell, C.D., Chapman, S.J., Cameron, C.M., Davidson, M.S., Potts, J.M., 2003. A rapid microtiter plate method to measure carbon dioxide evolved from carbon substrate amendments so as to determine the physiological profiles of soil microbial communities by using whole soil. *Applied and Environmental Microbiology* 69, 3593–3599.

Chapman, S.J., Campbell, C.D., Artz, R.R.E., 2007. Assessing CLPPs using MicroResp™ – a comparison with Biolog and multi-SIR. *Journal of Soils and Sediments* 7, 406–410.

Cheng, W., Coleman, D.C., 1989. A simple method for measuring CO₂ in a continuous air-flow system: modifications to the substrate-induced respiration technique. *Soil Biology & Biochemistry* 21, 385–388.

Dassonville, F., Renault, P., Vallès, V., 2004. A model describing the interactions between anaerobic microbiology and geochemistry in a soil amended with glucose and nitrate. *European Journal of Soil Science* 55, 29–45.

Degens, B.P., Harris, J.A., 1997. Development of a physiological approach to measuring the catabolic diversity of soil microbial communities. *Soil Biology & Biochemistry* 29, 1309–1320.

Dilly, O., 2001. Microbial respiratory quotient during basal metabolism and after glucose amendment in soils and litter. *Soil Biology & Biochemistry* 33, 117–127.

DOE, 1994. *Handbook of methods for the analysis of the various parameters of the carbon dioxide system in sea water; version 2*. In: Dickson, A.G., Goyet, C. (Eds.), ORNL/CDIAC-74.

Comment citer ce document :

Renault, P., Ben Sassi, M., Berard, A. (2013). Improving the MicroResp (TM) substrate-induced respiration method by a more complete description of CO₂ behavior in closed incubation wells. *Geoderma*, 207, 82-91. DOI : 10.1016/j.geoderma.2013.05.010

- El Nahhal, I.M., Zourab, S.M., Kodeh, F.S., Babonneau, F., Hegazy, W., 2012. Sol-gel encapsulation of cresol red in presence of surfactants. *Journal of Sol-Gel Science and Technology*. <http://dx.doi.org/10.1007/s10971-012-2693-y>.
- Ferreira, L.G., Noseda, M.D., Gonçalves, A.G., Ducatti, D.R.B., Fujii, M.T., Duarte, M.E.R., 2012. Chemical structure of the complex pyruvylated and sulfated agar from the red seaweed *Palisada flagellifera* (Ceramiales, Rhodophyta). *Carbohydrate Research* 347, 83–94.
- French, C.R., Carr, J.J., Dougherty, E.M., Eidson, L.A.K., Reynolds, J.C., De Grandpre, M.D., 2002. Spectrophotometric pH measurements of freshwater. *Analytica Chimica Acta* 453, 13–20.
- García-Palacios, P., Bowker, M.A., Chapman, S.J., Maestre, F.T., Soliveres, S., Gallardo, A., Valladares, F., Guerrero, C., Escudero, A., 2011. Early-successional vegetation changes after roadside prairie restoration modify processes related with soil functioning by changing microbial functional diversity. *Soil Biology & Biochemistry* 43, 1245–1253.
- Garland, J.L., Mills, A.L., 1991. Classification and characterization of heterotrophic microbial communities on the basis of patterns of community-level sole-carbon-source utilization. *Applied and Environmental Microbiology* 57, 2351–2359.
- Garland, J.L., Roberts, M.S., Levine, L.H., Mills, A.L., 2003. Community-level physiological profiling performed with an oxygen-sensitive fluorophore in a microtiter plate. *Applied and Environmental Microbiology* 69, 2994–2998.
- Gonzalez-Quinones, V., Banning, N.C., Ballesta, R.J., Murphy, D.V., 2009. Influence of cold storage on soil microbial community level physiological profiles and implications for soil quality monitoring. *Soil Biology & Biochemistry* 41, 1574–1576.
- Harned, H.S., Davis Jr., R.D., 1943. The ionization constant of carbonic acid in water and the solubility of carbon dioxide in water and aqueous salt solutions from 0 to 50 °C. *Journal of the American Chemical Society* 65, 2030–2037.
- Harned, H.S., Owen, B.B., 1958. *The Physical Chemistry of Electrolytic Solutions*, 3rd ed. Van Nostrand Reinhold, New York (803 pp.).
- Heger, D., Klánová, J., Klán, P., 2006. Enhanced protonation of cresol red in acidic aqueous solutions caused by freezing. *The Journal of Physical Chemistry. B* 110, 1277–1287.
- Lahaye, M., Rochas, C., 1991. Chemical structure and physico-chemical properties of agar. *Hydrobiologia* 221, 137–148.
- Lahlah, J., Renault, P., Cazevielle, P., Buzet, A., Hazemann, J.L., Womes, M., Cambier, P., 2009. Geochemical resilience of a ferralsol subjected to anoxia and organic matter amendment. *Soil Science Society of America Journal* 73, 1958–1971.
- Lalor, B.M., Cookson, W.R., Murphy, D.V., 2007. Comparison of two methods that assess soil community level physiological profiles in a forest ecosystem. *Soil Biology & Biochemistry* 39, 454–462.
- Macdonald, C.A., Thomas, N., Robinson, L., Tate, K.R., Ross, D.J., Dando, J., Singh, B.K., 2009. Physiological, biochemical and molecular responses of the soil microbial community after afforestation of pastures with *Pinus radiata*. *Soil Biology & Biochemistry* 41, 1642–1651.
- Millero, F.J., 1995. Thermodynamics of the carbon dioxide system in the oceans. *Geochimica et Cosmochimica Acta* 59, 661–677.
- Millero, F.J., Graham, T.B., Huang, F., Bustos-Serrano, H., Pierrot, D., 2006. Dissociation constants of carbonic acid in seawater as a function of salinity and temperature. *Marine Chemistry* 100, 80–94.
- Millero, F., Huang, F., Graham, T., Pierrot, D., 2007. The dissociation of carbonic acid in NaCl solutions as a function of concentration and temperature. *Geochimica et Cosmochimica Acta* 71, 46–55.
- Oren, A., Steinberger, Y., 2008. Coping with artifacts induced by CaCO₃-CO₂-H₂O equilibria in substrate utilization profiling of calcareous soils. *Soil Biology & Biochemistry* 40, 2569–2577.
- Oren, A., Steinberger, Y., 2008. Catabolic profiles of soil fungal communities along a geographic climatic gradient in Israel. *Soil Biology & Biochemistry* 40, 2578–2587.
- Pankow, J.F., 1991. *Thermodynamic principles*. Aquatic Chemistry Concepts. Lewis Publishers, Inc., Chelsea, Michigan, pp. 17–50.
- Parry, S., Renault, P., Chadoeuf, J., Chenu, C., Lensi, R., 2000. Particulate organic matter as a source of variation in denitrification in clods of soil. *European Journal of Soil Science* 51, 271–281.
- Pytkowicz, R.M., 1975. Activity coefficients of bicarbonates and carbonates in seawater. *Lynnology and Oceanography* 20, 971–975.
- Renault, P., Cazevielle, P., Verdier, J., Lahlah, J., Clara, C., Favre, F., 2009. Variations in the cation exchange capacity of a ferralsol supplied with vinasse, under changing aeration conditions. Comparison between CEC measuring methods. *Geoderma* 154, 101–110.
- Rowell, M.J., 1995. Colorimetric method for CO₂ measurement in soils. *Soil Biology & Biochemistry* 27, 373–375.
- Salvatore, F., Ferri, D., Palombi, R., 1986. Salt effect on the dissociation constant of acid-base indicators. *Journal of Solution Chemistry* 15, 423–431.
- Saul-Tcherkas, V., Steinberger, Y., 2009. Substrate utilization patterns of desert soil microbial communities in response to xeric and mesic conditions. *Soil Biology & Biochemistry* 41, 1882–1893.
- Schlesinger, W.H., Andrews, J.A., 2000. Soil respiration and the global carbon cycle. *Biogeochemistry* 48, 7–20.
- Scholten, H.J., Pierik, R.L.M., 1998. Agar as a gelling agent: chemical and physical analysis. *Plant Cell Reports* 17, 230–235.
- Sierra, J., Renault, P., 1995. Oxygen consumption by soil microorganisms as affected by oxygen and carbon dioxide levels. *Applied Soil Ecology* 2, 175–184.
- Skipper, H.D., Arthur, G., Wollum, I.L., Turco, R.F., Wolf, D.C., 1996. Microbiological aspects on environmental fate studies of pesticides. *Weed Technology* 10, 174–190.
- Smith, E.T., Matachek, J.R., 2002. A colorful investigation of a diprotic acid: a general chemistry laboratory exercise. *The Chemical Educator* 7, 359–363.
- Stefanowicz, A., 2006. The Biolog plates technique as a tool in ecological studies of microbial communities. *Polish Journal of Environmental Studies* 15, 669–676.
- Stevenson, B.A., Verburg, P.S.J., 2006. Effluxed CO₂-¹³C from sterilized and unsterilized treatments of a calcareous soil. *Soil Biology & Biochemistry* 38, 1727–1733.
- Ström, L., Godbold, D.L., Jones, D.L., 2001. Procedure for determining the biodegradation of radiolabeled substrates in a calcareous soil. *Soil Science Society of America Journal* 65, 347–351.
- Stumm, W., Morgan, J.J., 1996. Dissolved carbon dioxide. In: Shnoor, J.L., Zehnder, A.J.B. (Eds.), *Aquatic chemistry. Chemical Equilibria and Rates in Natural Waters*, 3rd edition. John Wiley & Sons, New York, pp. 148–205.
- Tamir, G., Shenker, M., Heller, H., Bloom, P.R., Fine, P., Bar-Tal, A., 2011. Can soil carbonate dissolution lead to overestimation of soil respiration? *Soil Science Society of America Journal* 75, 1414–1422.
- Tlili, A., Marechal, M., Montuelle, B., Volat, B., Dorigo, U., Bérard, A., 2011. Use of the MicroResp™ method to assess pollution-induced community tolerance for lotic biofilms. *Environmental Pollution* 159, 18–24.
- Wakelin, S.A., Macdonald, L.M., Rogers, S.L., Gregg, A.L., Bolger, T.P., Baldock, J.A., 2008. Habitat selective factors influencing the structural composition and functional capacity of microbial communities in agricultural soils. *Soil Biology & Biochemistry* 40, 803–813.
- Weiss, R.F., 1974. Carbon dioxide in water and seawater: the solubility of a non-ideal gas. *Marine Chemistry* 2, 203–215.
- Yao, H., Campbell, C.D., Qiao, X., 2011. Soil pH controls nitrification and carbon substrate utilization more than urea or charcoal in some highly acidic soils. *Biology and Fertility of Soils* 47, 515–522. <http://dx.doi.org/10.1007/s00374-011-0554-4>.
- Zhou, X., Liu, X., Rui, Y., Chen, C., Wu, H., Xu, Z., 2011. Symbiotic nitrogen fixation and soil N availability under legume crops in an arid environment. *Journal of Soils and Sediments* 11, 762–770.

Comment citer ce document :

Renault, P., Ben Sassi, M., Berard, A. (2013). Improving the MicroResp (TM) substrate-induced respiration method by a more complete description of CO₂ behavior in closed incubation wells.

Geoderma, 207, 82-91. DOI : 10.1016/j.geoderma.2013.05.010