

### Quantification of the lithogenic carbon pump following a simulated dust-deposition event in large mesocosms

Matthieu Bressac, Cécile Guieu, David Doxaran, François Bourrin, Karine Desboeufs, Nathalie Leblond, Céline Ridame

### ▶ To cite this version:

Matthieu Bressac, Cécile Guieu, David Doxaran, François Bourrin, Karine Desboeufs, et al.. Quantification of the lithogenic carbon pump following a simulated dust-deposition event in large mesocosms. Biogeosciences, 2014, 11, pp.1007-1020. 10.5194/bg-11-1007-2014 . hal-00952676

### HAL Id: hal-00952676 https://hal.science/hal-00952676

Submitted on 13 Jan 2016  $\,$ 

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

Biogeosciences, 11, 1007–1020, 2014 www.biogeosciences.net/11/1007/2014/ doi:10.5194/bg-11-1007-2014 © Author(s) 2014. CC Attribution 3.0 License.





# Quantification of the lithogenic carbon pump following a simulated dust-deposition event in large mesocosms

M. Bressac<sup>1,2</sup>, C. Guieu<sup>2</sup>, D. Doxaran<sup>2</sup>, F. Bourrin<sup>3</sup>, K. Desboeufs<sup>4</sup>, N. Leblond<sup>5</sup>, and C. Ridame<sup>6</sup>

<sup>1</sup>ACRI-ST, BP 234, 06904 Sophia-Antipolis, France

<sup>2</sup>CNRS-INSU, Université Pierre et Marie Curie-Paris 6, UMR7093, Laboratoire d'Océanographie de Villefranche/Mer, Observatoire Océanologique, 06230 Villefranche-sur-Mer, France

<sup>3</sup>Université de Perpignan Via Domitia, CEFREM, CNRS-UPVD, UMR5110, 52 avenue Paul Alduy, 66860 Perpignan, France <sup>4</sup>Université Paris Diderot, Sorbonne Paris Cité, Université Paris Est Créteil, UMR CNRS 7583, Laboratoire Interuniversitaire des Systèmes Atmosphériques (LISA), 61 av. Gal de Gaulle, 94010 C France

<sup>5</sup>Observatoire Océanologique de Villefranche/Mer, CNRS, Université Pierre et Marie Curie-Paris 6, 06230 Villefranche-sur-Mer, France

<sup>6</sup>CNRS-INSU-IRD, Université Pierre et Marie Curie, UMR7159, LOCEAN: Laboratoire d'Océanographie et du Climat: Expérimentation et Approches Numériques, 4 Place Jussieu – 75252 Paris Cedex 05, France

Correspondence to: M. Bressac (bressac@obs-vlfr.fr)

Received: 8 July 2013 – Published in Biogeosciences Discuss.: 19 August 2013 Revised: 26 November 2013 – Accepted: 29 December 2013 – Published: 21 February 2014

Abstract. Lithogenic particles, such as desert dust, have been postulated to influence particulate organic carbon (POC) export to the deep ocean by acting as mineral ballasts. However, an accurate understanding and quantification of the POC-dust association that occurs within the upper ocean is required in order to refine the "ballast hypothesis". In the framework of the DUNE (a DUst experiment in a low-Nutrient, low-chlorophyll Ecosystem) project, two artificial seedings were performed seven days apart within large mesocosms. A suite of optical and biogeochemical measurements were used to quantify surface POC export following simulated dust events within a low-nutrient, low-chlorophyll ecosystem. The two successive seedings led to a 2.3-6.7fold higher POC flux than the POC flux observed in controlled mesocosms. A simple linear regression analysis revealed that the lithogenic fluxes explained more than 85 % of the variance in POC fluxes. On the scale of a dust-deposition event, we estimated that 42-50 % of POC fluxes were strictly associated with lithogenic particles (through aggregation and most probably sorption processes). Lithogenic ballasting also likely impacted the remaining POC fraction which resulted from the fertilization effect. The observations support the "ballast hypothesis" and provide a quantitative estimation of the surface POC export abiotically triggered by dust deposition. In this work, we demonstrate that the strength of such a "lithogenic carbon pump" depends on the biogeochemical conditions of the water column at the time of deposition. Based on these observations, we suggest that this lithogenic carbon pump could represent a major component of the biological pump in oceanic areas subjected to intense atmospheric forcing.

#### 1 Introduction

The magnitude of downward particulate organic carbon (POC) export and its subsequent sequestration depends on the flux generated within the euphotic zone, the velocity at which it sinks, and the rate at which it decomposes (De La Rocha and Passow, 2007). Early studies showed strong correlations in the deep ocean between mineral (biogenic calcite, biogenic opal, and lithogenic particles) and POC fluxes (e.g., Deuser et al., 1981, 1983). To explain this predictive power, a "ballast hypothesis" was proposed, and it has been suggested that the downward POC flux is partly driven by associations with denser mineral particles that increase the overall sinking velocity for aggregates and that provide physical protection against the remineralization of labile organic

matter (OM) (Ittekkot, 1993; Hedges et al., 2000; Armstrong et al., 2002; Francois et al., 2002; Klaas and Archer, 2002). Although little evidence for a direct protective mechanism has emerged from in situ observations (Ingalls et al., 2006; Ploug et al., 2008; Iversen and Ploug, 2010), laboratorycontrolled experiments have demonstrated the potential for minerals to inhibit the microbial degradation of OM (Arnarson and Keil, 2005; Engel et al., 2009a; Le Moigne et al., 2013). On the other hand, it has been demonstrated that minerals could increase the velocity at which aggregates sink (e.g., De La Rocha and Passow, 2007; De La Rocha et al., 2008; Thomalla et al., 2008; Engel et al., 2009b; Iversen and Ploug, 2010) and drive large POC flux events (e.g., Thunell et al., 2007; Lee et al., 2009; Sanders et al., 2010; Ternon et al., 2010). Such fast-sinking POC, negligible within the euphotic zone, is sufficient to explain deep-ocean POC fluxes (Honda and Watanabe, 2010; Riley et al., 2012); this emphasizes the importance of the mode by which carbon is transferred downward when considering carbon flux parameterization.

Atmospherically and river-derived lithogenic material could act as an important carrier phase for POC to the deep ocean (Dunne et al., 2007; Salter et al., 2010). Often characterized by extreme events (Mahowald et al., 2003), atmospheric dust-deposition could induce a sudden and drastic increase in particle concentration and pronounced differential settling rates within surface waters. Such physical perturbation could profoundly change collision and coagulation rates, and, therefore, act as a catalyst during the aggregation process (Hamm, 2002; Lee et al., 2009). Atmospheric desert dust particles also constitute an important external source of new nutrients in large oceanic areas (Duce et al., 1991; Mahowald et al., 2009) that may alleviate biological limitation (Boyd et al., 2010 and references therein) and enhance export production (e.g., Bishop et al., 2002). Therefore, such a fertilization effect coupled with lithogenic ballasting has major consequences on ocean biogeochemistry and particle flux. However, while these simultaneous POC fluxes are selectively removed with respect to the ballast hypothesis, the difficulty of rigorously quantifying lithogenic ballasting due to this coupling of biotic and abiotic contributions remains.

As implicitly suggested by the ballast hypothesis, it has been demonstrated that POC–ballast associations take place within the upper ocean (Lee et al., 2009; Iversen et al., 2010; Sanders et al., 2010; Le Moigne et al., 2012) (i.e., before aggregates reach the mesopelagic depths where processes are determinant in setting the fraction of OM that reaches the deep ocean; Boyd and Trull, 2007). Although our mechanistic understanding of such an association must be improved, it seems that most lithogenic particles are not directly attached to biogenic particles, but form clusters that are likely bound by extracellular organic material (Hamm, 2002). In light of this fact, an alternative explanation states that correlations between POC and mineral fluxes may result from OM, which acts as "glue" for aggregate minerals that would otherwise not sink (Passow, 2004; Passow and De la Rocha, 2006). Without determining the catalyst for aggregation, Ternon et al. (2010) concluded that the simultaneous presence of OM and lithogenic particles is required for the formation of large and fast-sinking particles in the western Mediterranean Sea. Considering the strong spatial and temporal variability of dust-deposition events (Jickells et al., 2005; Mahowald et al., 2009), the water column into which atmospheric particles are deposited is characterized by a wide-ranging OM composition and abundance. Therefore, the intensity of the aggregation process and subsequent particulate export could vary, limiting the relevance of a global scale approach for estimating the contribution of lithogenic ballasting to POC export (Ragueneau et al., 2006; Boyd and Trull, 2007; De La Rocha and Passow, 2007).

Since only a few studies have reported the enhancement of export production following dust-deposition pulses in lownutrient, low-chlorophyll (LNLC) areas (e.g., Ternon et al., 2010), we sought to improve our understanding of this mechanism through this work. As a result of the episodic nature of dust deposition, artificial seeding in mesocosms constitutes a suitable approach for following, on an environmentally relevant timescale, the surface POC export associated with dust-deposition events. In this study, particulate export was investigated using two complementary approaches, as follows: (1) optical measurements were used to assess the rapid transfer of suspended particulate matter occurring a few hours following seeding, and (2) daily sediment trap samples were used to discriminate organic and lithogenic fractions, and to provide a global view of the flux that occurs following seedings. While determining surface fluxes is subject to trapping artefacts (Gardner, 2000; Scholten et al., 2001; Yu et al., 2001), such an experimental approach enabled us to investigate particulate fluxes using vertical coherence and a higher time resolution than that feasible with sediment traps. The objectives of the study are to determine (1) the causality of the relationship between dust deposition and subsequent POC export increase, (2) the respective contribution of biotic (fertilization) and abiotic (lithogenic ballasting) processes to total POC export, and (3) the variability of these processes for different initial biogeochemical conditions.

#### 2 Materials and methods

#### 2.1 Experimental setup and sampling

In the framework of the DUNE (a DUst experiment in a low-Nutrient, low-chlorophyll Ecosystem) project, a first artificial seeding experiment was performed in 2008 (DUNE 1) and a second in 2010 (DUNE 2). Since no optical measurements were performed during the DUNE 1 experiment, only the results from the DUNE 2 experiment are reported here. The mesocosm's experimental design and the methodology used to produce dust analog are reported in Guieu et al. (2010). Seven large mesocosms (height: 14.7 m, diameter: 2.3 m, surface area:  $4.15 \text{ m}^2$ , volume:  $52 \text{ m}^3$ ) were deployed within a coastal area (Corsica;  $42.374^\circ \text{ N}$ ,  $8.554^\circ \text{ E}$ ) during the summer oligotrophic period. Three mesocosms kept without seeding (here referred to as "control mesocosms") and three seeded mesocosms (here referred to as "+dust mesocosms") were devoted to biogeochemical studies. A seventh mesocosm, also seeded, was utilized for the deployment of optical instruments.

The mesocosms ended with a sediment trap (a 250 mL high-density polyethylene bottle) from which divers collected material every 24 h. Optical measurements were performed at high spatial and temporal resolutions during the first  $\sim$  24 h of each experiment for  $\sim$  2 days following each seeding, whereas biogeochemical parameters were sampled daily during the week. Primary production measurements were performed every 24 h at a depth of 5 m (Ridame et al., 2014). The data were integrated over the water column (0–12.5 m in depth) by assuming a homogeneous profile.

#### 2.2 The dust analog and the seeding

The finest dust fraction (<  $20 \,\mu$ m) was separated from the bulk soil samples – dominated by quartz (40%), calcite (30%) and clays (25%) – by grinding and dry-sieving. Then the dust analogs were processed to simulate cloud evapoc-ondensation cycles. The physicochemical characteristics of the dust analog are reported in Desboeufs et al. (2014). The resulting dust population presented a volume median diameter around 6.5  $\mu$ m and a peak at ~ 10  $\mu$ m, while the particle number size distribution peaked at 0.1  $\mu$ m (Guieu et al., 2010a).

The DUNE 2 experiment lasted 14 days, from 26 June to 9 July 2010. Two artificial seedings were successively conducted seven days apart within the same mesocosm and consisted of mimicking realistic wet deposition events with a dust flux of  $10 \text{ g m}^{-2}$ . Such a flux corresponds to 41.5 g of evapocondensed dust diluted in 2 L of ultrapure water and sprayed onto the surface of each of the mesocosms for a total duration of ~ 40 min.

In the Mediterranean Basin, dust is mainly derived from the Sahara desert in the form of strong pulses (Loÿe-Pilot et al., 1986; Bergametti et al., 1989; Guerzoni et al., 1999). Between 1984 and 1994, Loÿe-Pilot and Martin (1996) reported a mean annual flux in Corsica of  $12.5 \text{ g m}^{-2} \text{ yr}^{-1}$ , mainly attributed to pulses  $> 1 \text{ g m}^{-2}$ . According to the same authors, this deposition is mainly wet deposition and may occur only with few drops of rain, meaning that a high amount of dust can be deposited in time scales of minutes. Similar strong and sudden (few-hour-long) events have been recorded over the past decade, with African dust-deposition fluxes as high as  $22 \text{ g m}^{-2}$  (Bonnet and Guieu, 2006; Guieu et al., 2010b; Ternon et al., 2010). Our simulation, which allowed us to seed all the + dust mesocosms in a quasi-synoptic way, therefore appears realistic in terms of flux magnitude and duration (Guieu et al., 2013a).

#### 2.3 Optical measurements

Optical instruments, measurements, data processing and corrections are described in detail in Bressac et al. (2012). Briefly, an optical profiling package, including an ECO-BB3 (WET Labs, Inc.) backscattering sensor and a LISST-100 type B device (Sequoia Scientific Inc.), was deployed inside and outside the "optical" mesocosm. To minimize perturbation inside the mesocosm when profiling, vertical profiles were performed from 0 to 10 m depth at a speed of  $0.1 \text{ m s}^{-1}$  at the center of the bag. Vertical profiles, performed every hour using both sensors, consisted of continuous measurements for approximately 20 min between the subsurface and 10 m depth. For the remaining 40 min, measurements were performed continuously at 0.5 m depth. Control profiles were performed outside the mesocosm in the same manner. The particulate backscattering coefficient  $(b_{bp}(\lambda), m^{-1})$ was obtained from the light backscattered at  $117^{\circ}$  ( $\beta$  (117°),  $m^{-1} sr^{-1}$ ), measured using the ECO-BB3 sensor. Because of a logistic constraint, two different backscattering sensor were employed: a visible (440, 532, and 660 nm) sensor and a near-infrared (720, 770, and 870 nm) sensor; these were deployed during the first and second seeding experiment, respectively. Only measurements performed at 660 (first seeding) and 720 nm (second seeding) are reported here and compared to their respective initial values (see Sect. 3.3.2); indeed, the results at other wavelengths were similar. The LISST-100 type B sensor measured the beam attenuation coefficient at 670 nm along a path length of 5 cm. The ratio of the intensity of transmitted light in seawater to the intensity of light transmitted in ultrapure water was used to calculate the attenuation coefficient due to suspended particles, removing the attenuation due to water (Hill et al., 2011). Since light absorption by colored dissolved substances is negligible in the red spectral domain (Bricaud et al., 1981; Boss et al., 2009), the measured attenuation was assumed to be the particulate beam attenuation ( $c_{\rm p}, {\rm m}^{-1}$ ).

### 2.4 Sediment traps: sample treatment, elemental analyses, and calculations

Samples from sediment traps were analyzed within the seven mesocosms. The means of POC and lithogenic fluxes measured in the "optical" and + dust mesocosms were not significantly different from each other (p values > 0.05). However, since primary production was not measured, data from the "optical" mesocosm are not reported here. Following collection, samples were preserved using a solution of 5 % buffered formaldehyde and stored in the dark at 4 °C until processed. Swimmers were carefully handpicked using a binocular microscope. The remaining sample was desalted using ultrapure water and freeze-dried. Mass flux was determined by weighing the entire freeze-dried sample five times, with the accuracy of the weighing being < 1% over the whole data series.

The total concentration of carbon (TC) was measured in duplicate using a CHN analyzer (PerkinElmer 2400) on aliquots of desiccated samples (5-10 mg). HNO<sub>3</sub>/HF aciddigestion was performed in 7 mL Teflon flasks at 150 °C on aliquots of desiccated samples ( $\sim 20 \text{ mg}$ ). Following complete evaporation, samples were diluted in 0.1 M HNO<sub>3</sub> and analyzed for their calcium (Ca), aluminium (Al), and sulfur (S) concentrations by ICP-AES (inductively coupled plasma-atomic emission spectrometry) using the procedures described by Desboeufs et al. (2014). The detection limit of the apparatus for Ca, Al, and S was well below the lowest concentration of the digested aliquot (Desboeufs et al., 2014). Aliquots of the blanks and the certified reference materials (GBW07313) were digested and analyzed for Al, Ca, and S under the same conditions. In dust analogs Ca is present both as calcium carbonate and as calcium sulfate (Guieu et al., 2010a; Desboeufs et al., 2014). The part of Ca associated with sulfate (% Ca<sub>CaSO4</sub>) was estimated from the particulate S concentration within sediment traps. Therefore, the particulate Ca concentration as carbonate (% Ca<sub>CaCO3</sub>) corresponded to the difference between total % Ca and %  $Ca_{CaSO_4}$ . Then the carbonate fraction (CaCO<sub>3</sub>) was determined from the %Ca<sub>CaCO3</sub> as follows:

$$% CaCO_3 = 100/40 \times % Ca_{CaCO_3}.$$
 (1)

Particulate inorganic carbon (PIC) was deduced from the CaCO<sub>3</sub> fraction by assuming that, as follows:

$$\text{\% PIC} = \frac{12}{100} \times \text{\% CaCO}_3.$$
 (2)

POC was determined by subtracting PIC from TC. The lithogenic fraction was determined from particulate Al concentrations measured in the + dust mesocosms by considering the Al composition of the dust analogs (3.32%; Desboeufs et al., 2014). The protocol for the elemental analyses, the calculations for the various fractions, and the complete sediment trap data set are fully described in Desboeufs et al. (2014).

#### 3 Results

*Calculation and nomenclature* – POC flux data are presented for all of the mesocosms (Figs. 5 and 6) or as the average of the three + dust and three control values (Figs. 1, 7, and 8). The difference between these averaged POC fluxes is referred to as "net" (Fig. 1). " $T_x$ " represents the sampling time expressed in hours (x) since the first seeding. For samples from the second experiment, time (in hours) since the second seeding (y) is added in parentheses, as follows: " $T_{x(y)}$ ".

### **3.1** Evolution of the physical and biogeochemical parameters

Throughout the entire experiment, the seawater temperature ranged from 20.0 to  $27.3 \,^{\circ}$ C (Guieu et al., 2013a). The water



**Fig. 1.** Temporal evolution of the  $POC_{dust}$  flux (black line), the  $POC_{control}$  flux (light-gray portion), and the  $POC_{net}$  flux (dark-gray portion). Data represents the average (+SD) of the three replicate mesocosms.

column was well mixed during the first seeding experiment  $(T_0-T_{166(0)})$ . Toward the last day of the first experiment, solar heating within the surface layer led to the onset of stratification. The vertical temperature gradient was rapidly disrupted on the night after the second seeding ( $\sim T_{190(24)}-T_{210(44)}$ ). The water column remained well mixed until the last day of the experiment, when a strong thermal stratification appeared. Stratification was higher during the second seeding experiment but interrupted by wind-driven mixing events.

Considering the biological response to the first seeding (Giovagnetti et al., 2013; Ridame et al., 2014), the biogeochemical status of the water column changed between the experiments. Prior to the first seeding, the chlorophyll *a* concentration (Chl *a*) was characteristic of oligotrophic conditions  $(0.07 \pm 0.02 \text{ mg m}^{-3})$ . Seven days after the first seeding (i.e., prior to the second seeding), Chl *a* was ~1.7 times higher  $(0.12 \pm 0.04 \text{ mg m}^{-3})$ . In the control mesocosms Chl *a* remained constant for the entire experiment  $(0.06 \pm 0.01 \text{ mg m}^{-3})$ , indicating that changes in the + dust mesocosms were directly linked to the input of new nutrients provided by the dust.

Initial conditions indicate (1) very low dissolved inorganic phosphorus concentrations ([DIP]) ( $5 \pm 3$  nM; Pulido-Villena et al., 2014), (2) dissolved inorganic nitrogen concentrations ([DIN]) below the detection limit (< 30 nM; Ridame et al., 2013), and (3) dissolved iron concentrations ([DFe]) typical of coastal area ( $3.3 \pm 0.8$  nM; Wuttig et al., 2013). Following the first seeding, a decrease in [DFe] due to scavenging by sinking dust (Wuttig et al., 2013) and a transient increase in [DIP] (Pulido-Villena et al., 2014) were observed. The second addition of dust induced a significant increase in [DFe] (Wuttig et al., 2013) and [DIP] (Pulido-Villena et al., 2014). Furthermore, significant increases in [DIN] were observed following both seedings (Ridame et al., 2013). By increasing [DIP] and [DIN], the seedings relieved the ambient nutrient limitation and strongly stimulated primary production (see Sect. 4.1). Based on estimation of the new production, Ridame et al. (2013) observed a switch from a regenerated-production-based system to a new-production-based system 24 h after the seeding.

#### **3.2** Evolution of the POC flux

The temporal evolution of the averaged POC fluxes in the dust-seeded (POC<sub>dust</sub>) and control (POC<sub>control</sub>) mesocosms, as well as the difference between both fluxes (POC<sub>net</sub>), are reported in Fig. 1. It should be noted that the organic carbon contained in dust (0.31 %) constituted a very small fraction of the POC flux and is not discussed here since its fate remains unknown (Desboeufs et al., 2014). The first seeding resulted in a POC<sub>dust</sub> flux (111 mg m<sup>-2</sup>) 2.3 times higher than the POC<sub>control</sub> flux (47.4 mg m<sup>-2</sup>). The POC<sub>dust</sub> flux remained high during the first 96 h (> 18.5 mg m<sup>-2</sup> d<sup>-1</sup>), and then decreased by ~ 16 mg m<sup>-2</sup> d<sup>-1</sup> from  $T_{96}$  (day 5). Until the end of the first experiment ( $T_{166}$ ), the POC<sub>dust</sub> flux remained higher than the POC<sub>control</sub> flux.

The second seeding strongly reactivated the downward POC export, while the POC<sub>control</sub> flux (on average, ~6.5 times lower) continually decreased, reaching a near-zero value at the end of the experiment (~0.8 mg m<sup>-2</sup> d<sup>-1</sup>). The increase in POC<sub>dust</sub> flux was particularly significant during the first 24 h ( $T_{166(0)}-T_{190(24)}$ ) following the second seeding (+22.6 mg m<sup>-2</sup> d<sup>-1</sup>), and was characterized by a larger variation (45 %) between + dust mesocosms than the variation observed 24 h following the first seeding (20 %).

During the first 24 h following the second seeding, the POC<sub>net</sub> flux was 1.8 times higher than that 24 h after the first seeding. On days 2 and 3, POC<sub>net</sub> fluxes were similar between both experiments ( $\sim 10-15 \text{ mg m}^{-2} \text{ d}^{-1}$ ) but differed widely on day 4, as follows: 15.4 and 5.5 mg m<sup>-2</sup> d<sup>-1</sup> for the first and the second experiments, respectively. The POC<sub>net</sub> fraction collected over six days was slightly higher following the first seeding (62 mg m<sup>-2</sup>) than after the second seeding (54 mg m<sup>-2</sup>). Although these POC<sub>net</sub> fluxes were similar, export dynamics differed in magnitude and duration. The POC export that stretched over time following the first seeding was intense, but was shorter following the second seeding.

#### 3.3 Evolution of the optical parameters

 $b_{bp}$  and  $c_p$  measurements are representative of the various fractions of particulate matter stock (e.g., Loisel et al., 2011).  $b_{bp}$  is highly sensitive to changes in the abundance of small non-living particles, such as dust (Stramski and Kiefer, 1991; Ulloa et al., 1994), while the  $c_p$  (670) is dependent on the entire particle assemblage, excluding dissolved matter (Bricaud et al., 1981; Boss et al., 2009).

#### **3.3.1 Depth integration**

Thanks to the high-resolution vertical coverage of the water column using optical measurements, the integration of  $c_p$  (670) from the surface to 10 m depth ( $c_p$  (0–10 m)) provides a record of particulate matter (of both biological and lithogenic origin) accumulation or depletion over time (Fig. 2). At  $T_0$ , the  $c_p$  (0–10 m) was very low (0.46 m<sup>-1</sup>), consistent with the oligotrophic status of the water column, and immediately increased by  $3.2 \text{ m}^{-1}$  following the first seeding; subsequently it continually increased, reaching  $8.7 \pm 1.4 \text{ m}^{-1}$  ( $T_{48}$ – $T_{53}$ ). At  $T_{166(0)}$ ,  $c_p$  (0–10 m) was ~ 10 times higher (4.93 m<sup>-1</sup>) than the  $T_0$  value. Immediately following the second seeding,  $c_p$  (0–10 m) increased by  $4.28 \text{ m}^{-1}$  (9.21 m<sup>-1</sup>), and then continually decreased, reaching a lower value (2.62 m<sup>-1</sup>) than the initial value.

#### **3.3.2** The vertical particle profiles

The different initial values and the opposite evolution of  $c_{\rm p}$  (0–10 m) during the two seeding experiments suggest a different behavior of the dust and/or biological communities. The calculation of  $\Delta c_p$  (670) ( $c_p$  (670)– $c_p$  (670)<sub>initial</sub>;  $m^{-1}$ ) allowed us to investigate the net effect of seedings over time and depth (Fig. 3). Following the first seeding, the decrease (~1 m<sup>-1</sup>) of  $\Delta c_{\rm p}(670)$  within surface waters was rapidly (T<sub>6</sub>) followed by an increase ( $\sim 0.2 \,\mathrm{m}^{-1}$ ) at depth (i.e., by an accumulation of particulate matter in comparison to the initial inventory of the water column). Between  $T_{30}$  and  $T_{48}$ , high and homogeneous  $\Delta c_{\rm p}$  (670) values were observed over the entire water column. The second seeding induced a higher accumulation of particulate matter within the first two meters of the water column ( $\Delta c_{\rm p}$  (670) up to  $2.4 \text{ m}^{-1}$  at 0.5 m depth). As demonstrated by the dust profile issued from particulate Al measurements within the water column (Desboeufs et al., 2014), the accumulation of particulate matter is mainly associated with the addition of dust. Negative  $\Delta c_p$  (670) values were rapidly observed ( $T_3$ – $T_8$ ) at depth, demonstrating the rapid removal of particulate matter present before the second seeding. Between  $T_8$  and  $T_{30}$  the decrease of  $\Delta c_{\rm p}$  (670) at the surface was followed by an increase at depth, indicating that an important transfer of matter toward depth occurred. Two days following the second seeding,  $\Delta c_{\rm p}$  (670) was homogeneous but negative along the 0-10 m water column, indicating that most of the particulate export occurred during this period, in agreement with the POC<sub>dust</sub> flux trend (Fig. 1).

The evolution of  $\Delta b_{\rm bp}(\lambda)$   $(b_{\rm bp}(\lambda)-b_{\rm bp}(\lambda)_{\rm initial}$  in m<sup>-1</sup>; Fig. 4) was also investigated in order to focus on the fate of the added particles. The wavelength used differed between experiments (see Sect. 2.3); however, we assumed that this calculation allowed us to compare the respective evolution of  $\Delta b_{\rm bp}$ . Differences in the  $\Delta b_{\rm bp}(\lambda)$  profiles between experiments were less marked as compared to the  $\Delta c_{\rm p}$  (670) profiles. Decreases in the  $\Delta b_{\rm bp}(\lambda)$  within surface waters were not associated with a proportional increase at depth, and could indicate that the "cleaning effect" of the water column highlighted using  $\Delta c_{\rm p}$  (670) profiles could result from the sinking of larger particles, such as mixed aggregates, that



**Fig. 2.** Temporal evolution of the vertically integrated (0-10 m) particle attenuation coefficient  $(c_p (0-10 \text{ m}))$  during both seeding experiments.

backscatter light less efficiently than small lithogenic particles (Twardowsky et al., 2001), as supported by particulate Al profiles showing a depletion of dust particles in the first 10 m between  $T_{24}$  and  $T_{48}$  (Desboeufs et al., 2014). Two days following both seedings, the  $\Delta b_{\rm bp}(\lambda)$  profiles were homogeneous along the water column and still positive, indicating that a portion of the lithogenic particle pool was still in suspension as suggested by Bressac et al. (2012).

#### 4 Discussion

### 4.1 A comparison between primary production, POC, and lithogenic fluxes

The isolation of the water mass by the mesocosm could be conceptually compared to the effect of the strong stratification within the upper water column that typically occurs during the summer period. Within the sediment traps of control mesocosms, the observed very high POC to lithogenic ratio (not shown) indicated that lithogenic ballasting was negligible. In this context, POC<sub>control</sub> export can be considered as representative of the surface POC export that occurs in natural systems during oligotrophic periods without external forcing.

Since optical data was limited to the first 48 h of experiments, sediment trap data and primary production (PP) were used to determine the biological and lithogenic contributions in POC export over experiments. Similar increases in PP (by a factor of 2.3–2.4) were observed within + dust mesocosms following both seedings, while PP remained constant within control mesocosms (Ridame et al., 2014). Both data sets remained significantly different (p values < 0.05) throughout the experiment. The strong stimulation of PP went along with a rapid enhancement of the pigment content per cell and a rapid photoprotective response (Giovagnetti et al., 2013).



**Fig. 3.** The vertical profiles of  $\Delta c_p$  (670) ( $c_p$  (670)– $c_p$  (670)<sub>initial</sub> in m<sup>-1</sup>) for different times following both seedings. Negative values correspond to a decrease compared to the initial  $c_p$  (670) coefficient.



**Fig. 4.** Vertical profiles of  $\Delta b_{\rm bp}(\lambda)$  ( $b_{\rm bp}(\lambda) - b_{\rm bp}(\lambda)_{\rm initial}$  in m<sup>-1</sup>) at different times following both seedings. The wavelength used for this calculation differed between experiments (see Sect. 2.3). The insets (the  $T_{30}$  and  $T_{48}$  panels) show a close-up of the 0–0.006 m<sup>-1</sup> values.

Therefore, the increase in PP seems to have been favored by such physiological acclimation in relation to the high light environment (Guieu et al., 2013a).

The continual decrease in POC<sub>control</sub> flux during the experiment (Fig. 1) indicates that the constant primary production observed within control mesocosms was not sufficient to sustain the longevity of POC export throughout the

experiment, in agreement with the critical particle concentration concept which proposes that the (re)activation of the aggregation process and subsequent particulate export requires a particle concentration above a threshold value (Jackson and Lochmann, 1992; Boyd et al., 2005; Burd and Jackson, 2009). Atmospheric deposition could help to reach this threshold via the direct input of lithogenic particles, and indirectly through the supply of new nutrients, leading to an increase in algal biomass.

In the + dust mesocosms, a comparison between integrated PP data and POC fluxes indicated that no statistically significant relationship existed between these two parameters (p values > 0.05; Fig. 5). Introducing a time lag of 12 or 24 h between production and flux did not improve this relationship. The apparent decoupling between primary production and POC flux indicated that production alone could not be used to accurately predict POC fluxes.

Furthermore, following both seedings, the biomass of siliceous phytoplankton remained very low and a major response was observed for non-siliceous phytoplanktonic organisms (Giovagnetti et al., 2013). Although calcified organisms likely responded to the seedings, the fraction of biogenic CaCO<sub>3</sub> collected in sediment traps of + dust mesocosms remained negligible  $(3 \pm 0.5\%)$  in mass; Desboeufs et al., 2014). Therefore, biomineral ballasting was considered to be negligible and lithogenic particles were considered to be the main carrier phase of POC in + dust mesocosms, as confirmed by the scatter plot of the POC<sub>dust</sub> flux versus the lithogenic flux, which indicated a statistically significant correlation (p value, < 0.0001; Fig. 6). For these analyses we assumed that the regressions did not pass through the origin (i.e., that an additional POC fraction not directly associated with the lithogenic flux may be a significant component of the fluxes (Honda and Watanabe, 2010)), as follows:

$$POC_{dust}$$
 flux =  $a \times lithogenic$  flux +  $b$ , (3)

where *a* is defined as the carrying coefficient (Klaas and Archer, 2002) and *b* corresponds to the unassociated POC flux. A simple linear regression analysis revealed that the lithogenic fluxes explained more than 85% of the variance in POC<sub>dust</sub> fluxes. The ratios of the POC<sub>dust</sub> flux to the lithogenic flux (i.e., the carrying coefficient *a*) ranged from 0.013 (during the second seeding) to 0.016 (during the first seeding). A possible explanation for the decrease in the carrying coefficient following the second seeding is discussed later (see Sect. 4.3.2).

#### 4.2 The dynamics and timing of particulate export

The  $c_p$  (670) coefficient, usually associated with the particle concentration (< 20 µm) (Chung et al., 1998), is widely used to follow particulate export (e.g., Bishop et al., 2002; Smetacek et al., 2012). As with the POC<sub>dust</sub> fluxes and optical parameters, the phytoplankton community responded dif-



**Fig. 5.** POC flux for all of the mesocosms versus integrated primary production (PP). No significant correlation existed between PP and the POC flux in the + dust mesocosms (Fisher test, p values > 0.05).

ferently to both seedings with a distinct magnitude, timing, and composition change (Giovagnetti et al., 2013).

Although optical measurements revealed rapid particulate export following the first seeding (Bressac et al., 2012),  $c_{\rm p}$  (0–10 m) continually increased during the  $T_0-T_{53}$  period (Fig. 2). This increase, particularly marked between  $T_{48}$ and  $T_{53}$ , was consistent with an increase in picophytoplankton biomass ( $< 3 \mu m$ ) that occurred two days following the first seeding, while larger cell-sized phytoplankton (>  $3 \mu m$ ) needed three days to increase in terms of biomass (Giovagnetti et al., 2013). The  $c_p$  (0–10 m) evolution indicated that biological production exceeded and masked particulate export, explaining the longer duration of POC<sub>dust</sub> export during the first experiment (Fig. 1). The long time lag between the first seeding and biological response (2-3 days) and immediate POC<sub>dust</sub> export activation (Fig. 1) highlighted the decoupling between POC export and biological production. On the basis of these observations, we suggest that the immediate twofold increase in POC<sub>dust</sub> flux resulted from abiotic processes such as lithogenic ballasting (i.e., by aggregating cells and increasing particle sinking velocity), in agreement with the strong correlation observed between the POC and lithogenic fluxes during the entire first experiment (Fig. 6).

The opposite evolution of  $c_p (0-10 \text{ m})$  following the second seeding (Fig. 2) indicates that particulate export exceeded biological production  $(T_{166(0)}-T_{214(48)})$ . Even if a stronger biological response was observed (a threefold Chl *a* increase), dominated by the nano- and microphytoplankton communities (> 3 µm), it only occurred 24–48 h after the second seeding (Giovagnetti et al., 2013). Therefore, the negative  $\Delta c_p$  (670) values rapidly observed  $(T_3-T_8)$  at depth following the second seeding suggest that this immediate "cleaning effect" resulted from lithogenic ballasting. On a longer timescale, we can hypothesize that the preferential response of large cells, more prone to be exported, may have



**Fig. 6.** POC<sub>dust</sub> versus lithogenic fluxes. Solid lines represent linear regressions with the following slopes, intercepts, determination coefficients, and probabilities (Fisher test). First seeding: N = 21, slope = 0.016 ± 0.002, intercept =  $3.11 \pm 1.48$ ,  $R^2 = 0.85$ , p < 0.0001. Second seeding: N = 17, slope =  $0.013 \pm 0.001$ , intercept =  $2.41 \pm 0.80$ ,  $R^2 = 0.88$ , p < 0.0001. The highlighted outlier was excluded from the linear regression for second seeding data. Uncertainties correspond to the standard deviation of one sample measured two times.

increased the budget of exported carbon and changed the "biotic contribution" to POC export.

These observations suggest that a sequence of mechanisms controlled the particulate export. While the biological response was delayed, particulate export driven by algal aggregation during the first 48 h following both seedings is unlikely (e.g., Prieto et al., 2002). On the other hand, the input of 41.5 g of dust (with a grain size distribution from < 0.1 to ~ 20 µm) likely catalyzed the aggregation and adsorption processes on a short timescale (e.g., Lee et al., 2009) and activated particulate export. Thus, abiotic processes were likely dominant during the first 24–48 h until the biological contribution became significant and stretched over the time of particulate export. The decline could finally result from the imbalance between production and export, leading to a lower particle concentration within the water column than that required for the aggregation process.

#### 4.3 The relationship between POC and lithogenic fluxes

The first general assessment for the carrying coefficients of ballast minerals was performed on a global scale (Klaas and Archer, 2002). Others studies that have generally focused on biominerals allowed these coefficients to vary in space (Ragueneau et al., 2006; Le Moigne et al., 2012; Wilson et al., 2012), or assessed them from individual sites (Wong et al., 1999; Conte et al., 2001; Honda and Watanabe, 2010). All of these analyses, performed at an annual timescale, reflect average fluxes during both productive and oligotrophic periods missing important temporal variability within the flux relationships. Therefore, we focused on specific dust-deposition events on timescales on the order of a week, representative of the residence time for dust within surface waters estimated from days to month (Buat-Ménard et al., 1989; Bory and Newton, 2000; Croot et al., 2004; Frew et al., 2006). During this artificial seeding experiment in LNLC ecosystems, the lithogenic fluxes were high and the algal biomass was likely too low to drive large POC flux events. Therefore, the low carrying coefficients estimated in this study (0.013–0.016; Fig. 6) did not correspond to a low absolute associated POC flux but rather suggested a low lithogenic normalized POC flux. Despite the strong and statistically robust relationships observed in the + dust mesocosms between POC and lithogenic fluxes (Fig. 6), the mechanistic basis for these OM-lithogenic particle interactions remains uncertain. The correlations could result from (1) the lithogenic ballasting of organic matter (Armstrong et al., 2002; Francois et al., 2002; Klaas and Archer, 2002), (2) the inclusion of lithogenic particles within organic matter (freshly produced) (Passow and De La Rocha, 2006); or (3) the coupling of both processes. To determine the respective contribution of these biotic and abiotic processes to POC export, two comparative approaches were successively employed.

#### 4.3.1 The effect of biogeochemical conditions

As a result of the high variability in physical conditions during the course of this experiment (Sect. 3.1), a direct comparison of POC<sub>dust</sub> fluxes between both experiments could bias our interpretation. Therefore, the POC<sub>dust</sub> fluxes were compared to the respective POC<sub>control</sub> fluxes, first highlighting the global effect (the biotic plus abiotic processes) of both seedings (Fig. 7). For the same initial biogeochemical conditions (i.e., the first seeding), the POC<sub>dust</sub> flux was, on average, 2.33 ( $\pm$  0.12) times higher than the POC<sub>control</sub> flux. Under different initial biogeochemical conditions (i.e., the second seeding), the POC<sub>dust</sub> flux was, on average,  $6.68 (\pm 0.51)$ times higher than the POC<sub>control</sub> flux. On the basis of these observations, we conclude that the net effect of a dustdeposition event on POC export depends, in part, on the biogeochemical properties of the water column (the organic matter pool, the phytoplankton community) at the time of deposition. Among these biogeochemical parameters, the stronger increase in large cell (>  $3 \mu m$ ) biomass (Giovagnetti et al., 2013) could partly explain the larger difference observed during the second experiment.

Sorption of dissolved organic matter (DOM) from solution onto mineral surfaces could reach equilibrium very rapidly (within ~1 h; Arnarson and Keil, 2000, 2005). This process likely contributed to the downward POC export in the first 24 h following the seedings. By using the same dust analog and flux in 0.2  $\mu$ m filtered seawater, Bressac and Guieu (2013) quantified this abiotically driven organic carbon export and its variation as a function of the composition and abundance of DOM. During this abiotic experiment, the



**Fig. 7.** POC<sub>dust</sub> versus POC<sub>control</sub> fluxes (mean fluxes and standard deviations of the mesocosm triplicates). The best linear regressions and the 1:1 lines are shown. First seeding: N = 7, slope =  $2.33 \pm 0.12$ ,  $R^2 = 0.94$ , p < 0.0001 (Fisher test). Second seeding: N = 6, slope =  $6.68 \pm 0.51$ ,  $R^2 = 0.93$ , p < 0.0001 (Fisher test).

amount of organic carbon adsorbed onto dust particles and exported downward was of the same order of magnitude as the POC fluxes observed in the + dust mesocosms. Furthermore, sticky particles such as transparent exopolymer particles (TEP), known for promoting the aggregation process (Chin et al., 1998; Passow, 2002; Engel et al., 2004; Verdugo et al., 2004), could also have indirectly influenced the particulate export. Although DOM and TEP concentrations were not measured during this experiment, the abundance of such organic material prior to the second seeding was likely higher within the + dust than within the control mesocosms, contributing to the larger difference in POC fluxes observed following the second seeding.

Although lithogenic ballasting cannot be strictly considered as a pump (as surface primary production is required to sustain it), Bressac and Guieu (2013) introduced the concept of the "lithogenic carbon pump" in order to highlight the fact that aggregation and adsorption processes could trigger a dust-induced POC export event. As with the fertilization effect (e.g., Marañón et al., 2010), POC export abiotically triggered by dust deposition could vary according to the biogeochemical state of sea surface waters at the time of deposition (Bressac and Guieu, 2013).

#### 4.3.2 Biotic and abiotic contributions to POC export

With the objective of determining the strength of this "lithogenic carbon pump", the second comparative approach consisted of a deconvolution of the "ballasted" and "unballasted" POC fluxes. For this purpose, the  $POC_{dust}$  flux could be partitioned into two pools, a fraction associated with lithogenic particles (POC<sub>lith</sub>) and a "freshly" produced one induced by the input of new nutrients (POC<sub>ferti</sub>), as follows:

 $POC_{dust} = POC_{lith} + POC_{ferti}$ . Also known as the particle export efficiency (e.g., Buesseler, 1998; Lutz et al., 2002; Henson et al., 2012), POC<sub>lith</sub> was estimated by normalizing the POC<sub>dust</sub> fluxes to the respective integrated PP value (Fig. 8). As biomineral ballasting was negligible during this experiment (Sect. 4.1), the difference in this parameter between the control and + dust mesocosms was assumed to result from lithogenic ballasting. POC<sub>lith</sub> was  $1.17 (\pm 0.19)$  (during the first seeding) and 2.84 ( $\pm$  0.19) (during the second seeding) times higher than the control's POClith. According to the relationships obtained using both comparative approaches (Figs. 7 and 8), POC<sub>lith</sub> represented  $50 \pm 8$  and  $42 \pm 3\%$  of the POC<sub>dust</sub> fluxes after the first and the second seeding, respectively; consistent with a higher biological contribution to POC export following the second seeding (Sect. 4.2). A possible explanation of the relative decrease in the POClith after the second seeding is the scavenging of most of the biogenic particles from the water column following the first seeding. Such a decrease was observed by Lee et al. (2009) following successive dust-deposition events and suggests that POC<sub>lith</sub> following the second seeding was allowed by the OM production induced by the first pulse of new nutrients, reflecting a highly dynamic system. A second key finding is that POC<sub>ferti</sub> largely exceeded the variance in POC fluxes unexplained by lithogenic fluxes ( $\sim 15\%$ ; Fig. 6), demonstrating that the carrying coefficient should be carefully interpreted. Such a finding calls into question whether or not a simple fertilization mechanism exists or whether the lithogenic ballasting of POC<sub>ferti</sub> could occur and favor its downward export.

## 4.4 One dust-deposition event, several particle settling populations

Both the coupling (e.g., Jickells et al., 1998; Migon et al., 2002; Ternon et al., 2010; Brust et al., 2011) and decoupling (e.g., Buat-Ménard et al., 1989; Migon et al., 2002; Ternon et al., 2010) of dust deposition and particulate export have been reported. During our experiment, only  $\sim 57$ and 41 % (by mass) of the lithogenic particles initially added were recovered in the sediment traps six days after the first and second seedings, respectively (Desboeufs et al., 2014). Neuer et al. (2004) collected samples at  $\sim$  500 m depth with between 13 and 20% of lithogenic material deposited, and concluded that a large fraction of lithogenic matter was not removed by particle sedimentation. Based on Stokes' law, we estimated that approximately half (in mass) of the dust population was large enough  $(>5 \,\mu\text{m})$  to settle without being included in aggregates, and would reach the sediment traps of mesocosms before the end of the experiment. The smaller lithogenic particles ( $< 5 \,\mu m$ ) that were numerically dominant likely remained in suspension, in agreement with  $\Delta b_{\rm bp}$  (660), still positive at  $T_{166(0)}$  (not shown). The result could have been influenced by ephemeral dust-induced phytoplankton blooms, which did not supply a sufficient amount



**Fig. 8.** POC<sub>dust</sub> versus POC<sub>control</sub> fluxes, both normalized with respect to integrated primary production values (PP). The best linear regressions and the 1 : 1 lines are shown. First seeding: N = 6, slope =  $1.17 \pm 0.19$ ,  $R^2 = 0.56$ , p < 0.0001 (Fisher test). Second seeding: N = 6, slope =  $2.84 \pm 0.19$ ,  $R^2 = 0.96$ , p < 0.0001 (Fisher test).

of OM "glue", and by the rapid scavenging of most of the biogenic particles following both seedings.

In the open ocean, this pool of small lithogenic particles remains stored within surface layers during the stratification period until either the onset of vertical convection during winter (e.g., Migon et al., 2002), or their inclusion into faecal pellets (e.g., Buat-Ménard et al., 1989) or larger aggregates (e.g., Burd and Jackson, 2009). For this last case, POC fluxes could scavenge these slow-settling lithogenic particles from the water column and determine their flux (Passow and De La Rocha, 2006). The particle population could also act as a nucleation point for aggregate formation, and, therefore, contribute to a second POC export event (e.g., Ternon et al., 2010). Actually, a decoupling between the deposition and export of this particle population will occur, preventing a direct temporal comparison. Finally, the long residence time within surface waters for such highly refractive submicron lithogenic particles has been advanced as an explanation of the color anomaly of the ultra-oligotrophic waters of the Mediterranean Sea (Claustre et al., 2002; Loisel et al., 2011). Our observations demonstrated the possible lasting impact of atmospheric lithogenic particles on seawater optical properties.

#### 5 Conclusions

By combining the use of sediment traps and high-resolution vertical optical profiles inside large mesocosms deployed within a LNLC system, we report surface particulate export following two successive dust-deposition events. Although the mechanism driving the relationship between POC and ballast material remains uncertain (De La Rocha and Passow, 2007), we have demonstrated in the present study that (1) dust deposition catalyzes particle-sinking events (e.g., Lee et al., 2009; Ternon et al., 2010), and (2) the intensity of POC–lithogenic particle association depends on the biogeochemical conditions of surface seawater at the time of deposition. The results obtained could be used to parameterize surface POC export in oligotrophic systems receiving high rates of dust deposition (e.g., the Mediterranean Sea) and to better predict the effects of future changes in atmospheric dust fluxes (Mahowald et al., 2009).

The lack of a significant relationship between POC flux and primary production combined with the high degree of covariance between POC and lithogenic fluxes adds further support to the ballast hypothesis. The probable composition of sinking aggregates (the high content of lithogenic material and the low amount of relatively buoyant OM), as well as their sinking velocity ( $\sim 24-86 \text{ m d}^{-1}$ ; Bressac et al., 2012) suggest low specific carbon remineralization (Riley et al., 2012). Furthermore, the inherent sporadic character of the lithogenic carbon pump may play a fundamental role in carbon sequestration (Ragueneau et al., 2006) since episodic events may lead to the rapid sinking of particles that will not be fully exploited by the deep sea bacterial community (Hansell and Ducklow, 2003: Nagata et al., 2000, 2010). As a result, the major fraction of the deep sea POC flux following dust-deposition events likely corresponds to the remnants of surface ballasted POC stocks (e.g., Thunell et al., 2007; Honda and Watanabe, 2010).

Impacts on heterotrophic activity must be taken into account in order to achieve connections between dustdeposition and ocean carbon cycling (Pulido-Villena et al., 2008). Both seedings maintained the strong net heterotrophic character of the ecosystem, indicating that bacteria are key players in the ecosystem response to dust depositions (Pulido-Villena et al., 2014), which leads to the remineralization of a fraction of fixed algal carbon, and, therefore, to a decrease in the amount of exported organic carbon (Guieu et al., 2013b). While dust deposition can have a fertilization effect on the phytoplankton compartment by bringing new nutrients, the labile OM produced could escape being remineralized within the upper ocean through its physical associations (i.e., aggregation and adsorption processes) with dust particles and subsequent lithogenic ballasting.

Acknowledgements. This work was funded by the ANR DUNE project under ANR-07-BLAN-0126-01. contract DUNE was endorsed by the international SOLAS (Surface Ocean - Lower Atmosphere) program in February 2009 (http://www.solas-int.org/activities/project-endorsement.html). F. Louis and J.-M. Grisoni are thanked for their help in the conception of the "optical" mesocosm. This field campaign could not have taken place without the logistical support and facilities of the Parc Naturel Régional de Corse, and the diving expertise of J.-M. Grisoni, D. Luquet, and C. Rouvière from the Observatoire Océanologique de Villefranche. A grant provided by the ACRI-ST company and the French National Association for Research and Technology (ANRT) supported M. Bressac. The authors wish to thank the two anonymous reviewers for their helpful comments in the discussion version of the paper.

Edited by: F. Dulac



The publication of this article is financed by CNRS-INSU.

#### References

- Armstrong, R. A., Lee, C., Hedges, J. I., Honjo, S., and Wakeham, S. G.: A new, mechanistic model for organic carbon fluxes in the ocean: based on the quantitative association of POC with ballast minerals, Deep-Sea Res. Pt. II, 49, 219–236, 2002.
- Arnarson, T. S. and Keil, R. G.: Mechanisms of pore water organic matter adsorption to montmorillonite, Mar. Chem., 71, 309–320, 2000.
- Arnarson, T. S. and Keil, R. G.: Influence of organic-mineral aggregates on microbial degradation of the dinoflagellate Scrippsiella trochoidea, Geochim. Cosmochim. Ac., 69, 2111–2117, 2005.
- Bergametti, G., Gomes, L., Remoudaki, E., Desbois, M., Martin, D., and Buat-Ménard, P.: Present, transport and deposition patterns of African dusts to the north-western Mediterranean, in: Paleoclimatology and Paleometeorology: Modern and Past Patterns of Global Atmospheric Transport, edited by: Leinen, M. and Sarnthein, M., Kluwer, 227–252, 1989.
- Bishop, J. K. B., Davis, E. R., and Sherman, J. T.: Robotic Observations of Dust Storm Enhancement of Carbon Biomass in the North Pacific, Science, 298, 817–821, doi:10.1126/science.1074961, 2002.
- Bonnet, S. and Guieu, C.: Atmospheric forcing on the annual iron cycle in the Mediterranean Sea, a one-year survey, J. Geophys. Res., 111, C09010, doi:10.1029/2005JC003213, 2006.
- Bory, A. J.-M. and Newton, P. P.: Transport of airborne lithogenic material down through the water column in two contrasting regions of eastern subtropical North Atlantic Ocean, Global Biogeochem. Cy., 14, 297–315, 2000.
- Boss, E., Slade, W., and Hill, P.: Effect of particulate aggregation in aquatic environments on the beam attenuation and its utility as a proxy for particulate mass, Opt. Express., 17, 9408–9420, doi:10.1364/OE.17.009408, 2009.
- Boyd, P. W. and Trull, T. W.: Understanding the export of biogenic particles in oceanic waters: Is there consensus?, Prog. Oceanogr., 72, 276–312, doi:10.1016/j.pocean.2006.10.007, 2007.
- Boyd, P. W, Strzpek, R., Takeda, S., Jackson, G. A., Wong, C. S., McKay, R. M., Law, C., Kiyosawa, H., Saito, H., Sherry, N., Johnson, K., Gower, J., and Ramaiah, N.: The evolution and termination of an iron-induced mesoscale bloom in the North East subarctic Pacific, Limnol. Oceanogr., 50, 1872–1886, 2005.
- Boyd, P. W., Jickells, T., Law, C. S., Blain, S., Boyle, E. A., Buesseler, K. O., Coale, K. H., Cullen, J. J., de Baar, H. J. W., Follows, M., Harvey, M., Lancelot, C., Levasseur, M., Owens, N. P. J., Pollard, R., Rivkin, R. B., Sarmiento, J., Schoemann, V., Smetacek, V., Takeda, S., Tsuda, A., Turner, S., and Watson, A.

J.: Mesoscale iron enrichment experiments 1993–2005: synthesis and future directions, Science, 315, 612–617, 2007.

- Boyd, P. W., Mackie, D. S., and Hunter, K. A.: Aerosol iron deposition to the surface ocean: modes of iron supply and biological responses, Mar. Chem., 120, 128–143, 2010.
- Bressac, M. and Guieu, C.: Post-depositional processes: What really happens to new atmospheric iron in the ocean's surface? Global Biogeochem. Cy., 27, 859–870, doi:10.1002/gbc.20076, 2013.
- Bressac, M., Guieu, C., Doxaran, D., Bourrin, F., Obolensky, G., and Grisoni J.-M.: A mesocosm experiment coupled with optical measurements to assess the fate and sinking of atmospheric particles in clear oligotrophic waters, Geo-Mar. Lett., 32, 153–164, doi:10.1007/s00367-011-0269-4, 2012.
- Bricaud, A., Morel, A., and Prieur, L.: Absorption by dissolved organic matter of the sea (yellow substance) in the UV and visible domains, Limnol. Oceanogr., 26, 43–53, 1981.
- Brust, J., Schulz-Bull, D. E., Leipe, T., Chavagnac, V., and Waniek, J. J.: Descending particles: From the atmosphere to the deep ocean – A time series study in the subtropical NE Atlantic, Geophys. Res. Lett., 38, L06603, doi:10.1029/2010GL045399, 2011.
- Buat-Ménard, P., Davies, P. J., Remoudaki, E., Miquel, J. C., Bergametti, G., Lamber, C. E., Ezat, E., Quétel, C. R., La Rosa J., and Fowler, S. W.: Non-steady-state biological removal of atmospheric particles from Mediterranean surface waters, Nature, 340, 131–133, doi:10.1038/340131a0, 1989.
- Buesseler, K. O.: The decoupling of production and particulate export in the surface ocean, Global Biogeochem. Cy., 12, 297–310, 1998.
- Burd, A. B. and Jackson, G. A.: Particle Aggregation, Annu. Rev. Mar. Sci., 1, 65–90, 2009.
- Chin, W. C., Orellana, M. V., and Verdugo, P.: Spontaneous assembly of marine dissolved organic matter into polymer gels, Nature, 391, 568–572, doi:10.1038/35345, 1998.
- Chung, S. P., Gardner, W. D., Landry, M. R., Richardson, M. J., and Walsh, I. D.: Beam attenuation by microorganisms and detrital particles in the equatorial Pacific, J. Geophys. Res., 103, 12669– 12681, doi:10.1029/98JC00608, 1998.
- Claustre, H., Morel, A., Hooker, S. B., Babin, M., Antoine, D., Oubelkheir, K., Bricaud, A., Leblanc, K., Quéguiner, B., and Maritorena, S.: Is desert dust making oligotrophic waters greener?, Geophys. Res. Lett., 29, 1469, doi:10.1029/2001GL014056, 2002.
- Conte, M., Ralph, R., and Ross, E.: Seasonal and interannual variability in deep ocean particle fluxes at the Ocean Flux Program (OFP)/ Bermuda Atlantic Time Series (BATS) site in the western Sargasso Sea near Bermuda, Deep-Sea Res. Pt. II, 48, 1471–1505, doi:10.1016/S0967-0645(00)00150-8, 2001.
- Croot, P. L., Streu, P., and Baker, A. R.: Short residence time for iron in surface seawater impacted by atmospheric dry deposition from Saharan dust events, Geophys. Res. Lett., 31, L23S08, doi:10.1029/2004GL020153, 2004.
- De La Rocha, C. L. and Passow, U.: Factors influencing the sinking of POC and the efficiency of the biological carbon pump, Deep-Sea Res. Pt. II, 54, 639–658, doi:10.1016/j.dsr2.2007.01.004, 2007.
- De La Rocha, C. L., Nowald, N., and Passow, U.: Interactions between diatom aggregates, minerals, particulate organic carbon, and dissolved organic matter: Further implications for

the ballast hypothesis, Global Biogeochem. Cy., 22, GB4005, doi:10.1029/2007GB003156, 2008.

- Desboeufs, K. V., Leblond, N., Wagener, T., Bon Nguyen, E., and Guieu, C.: Chemical fate and settling of mineral dust in surface seawater after atmospheric deposition observed from dust seeding experiments in large mesocosms, Biogeosciences, submitted, 2014.
- Deuser, W. G., Ross, E. H., and Anderson, R. F.: Seasonality in the supply of sediment to the deep Sargasso Sea and implications for the rapid transfer of matter to the deep ocean, Deep-Sea Res. Pt. I, 28, 495–505, 1981.
- Deuser, W. G., Brewer, P. G., Jickells, T. D., and Commeau, R. F.: Biological control of the removal of abiogenic particles from the surface ocean, Science, 219, 388–391, 1983.
- Duce, R. A., Liss, P. S., Merrill, J. T., Atlas, E. L., Buat-Ménard, P., Hicks, B. B., Miller, J. M., Prospero, J. M., Arimoto, R., Church, T. M., Ellis, W., Galloway, J. N., Hansen, L., Jickells, T. D., Knap, A. H., Reinhardt, K. H., Schneider, B., Soudine, A., Tokos, J. J., Tsunogai, S., Wollast, R., and Zhou, M.: The atmospheric input of trace species to the world ocean, Global Biogeochem. Cy., 5, 193–259, doi:10.1029/91GB01778, 1991.
- Dunne, J. P., Sarmiento, J. L., and Gnanadesikan, A.: A synthesis of global particle export from the surface ocean and cycling through the ocean interior and on the seafloor, Global Biogeochem. Cy., 21, GB4006, doi:10.1029/2006GB002907, 2007.
- Engel, A., Abramson, L., Szlosek, J., Liu, Z., Stewart, G., Hirschberg, D., and Lee, C.: Investigating the effect of ballasting by CaCO<sub>3</sub> in Emiliana huxleyi, II: Decomposition of particulate organic matter, Deep-Sea Res. Pt. II, 56, 1408–1419, doi:10.1016/j.dsr2.2008.11.028, 2009a.
- Engel, A., Szlosek, J., Abramson, L., Liu, Z., and Lee, C.: Investigating the effect of ballasting by CaCO<sub>3</sub> in Emiliania huxleyi:
  I. Formation, settling velocities and physical properties of aggregates, Deep-Sea Res. Pt. II, 56, 1396–1407, 2009b.
- Engel, A., Thoms, S., Riebesell, U., Rochelle-Newall, E., and Zondervan, I.: Polysaccharide aggregation as a potential sink of marine dissolved organic carbon, Nature, 428, 929–932, doi:10.1038/nature02453, 2004.
- Francois, R., Honjo, S., Krishfiels, R., and Manganini, S.: Factors controlling the flux of organic carbon to the bathypelagic zone of the ocean, Global Biogeochem. Cy., 16, 1087, doi:10.1029/2001GB001722, 2002.
- Frew, R. D., Hutchins, D. A., Nodder, S., Sanudo-Wilhelmy, S., Tovar-Sanchez, A., Leblanc, K., Hare, C. E., and Boyd P. W.: Particulate iron dynamics during FeCycle in subantarctic waters southeast of New Zealand, Glob. Biogeochem. Cy., 20, GB1S93, doi:10.1029/2005GB002558, 2006.
- Gardner, W. D.: Sediment trap sampling in surface waters, in: The Changing Ocean Carbon Cycle: a Midterm Synthesis of the Joint Global Ocean Flux Study, vol. 5, edited by: Hanson, R. B., Ducklow, H. W., and Field, J. G., Cambridge University Press, Cambridge, 240–281, 2000.
- Giovagnetti, V., Brunet, C., Conversano, F., Tramontano, F., Obernosterer, I., Ridame, C., and Guieu, C.: Assessing the role of dust deposition on phytoplankton ecophysiology and succession in a low-nutrient low-chlorophyll ecosystem: a mesocosm experiment in the Mediterranean Sea, Biogeosciences, 10, 2973–2991, doi:10.5194/bg-10-2973-2013, 2013.

- Guerzoni, S., Chester, R., Dulac, F., Herut, B., Loÿe-Pilot, M.D., Measures, C., Migon, C., Molinaroli, E., Moulin, C., Rossini, P., Saydam, C., Soudine, A., and Ziveri, P.: The role of atmospheric deposition in the biogeochemistry of the Mediterranean Sea, Progr. Oceanog., 44, 147–190, 1999.
- Guieu, C., Dulac, F., Desboeufs, K., Wagener, T., Pulido-Villena, E., Grisoni, J.-M., Louis, F., Ridame, C., Blain, S., Brunet, C., Bon Nguyen, E., Tran, S., Labiadh, M., and Dominici, J. M.: Large clean mesocosms and simulated dust deposition: a new methodology to investigate responses of marine oligotrophic ecosystems to atmospheric inputs, Biogeosciences, 7, 2765– 2784, doi:10.5194/bg-7-2765-2010, 2010a.
- Guieu, C., Loÿe-Pilot, M.-D., Benyaya, L., and Dufour, A.: Spatial and temporal variability of atmospheric fluxes of metals (Al, Fe, Cd, Zn and Pb) and phosphorus over the whole Mediterranean from a one-year monitoring experiment, biogeochemical implications, Mar. Chem., 120, 164–178, 2010b.
- Guieu, C., Dulac, F., Ridame, C., and Pondaven, P.: Introduction to the project DUNE, a DUst experiment in a low Nutrient, low chlorophyll Ecosystem, Biogeosciences Discuss., 10, 12491– 12527, doi:10.5194/bgd-10-12491-2013, 2013a.
- Guieu, C., Ridame, C., Pulido-Villena, E., Bressac, M., Desboeufs, K., and Dulac, F.: Dust deposition in an oligotrophic marine environment: impact on the carbon budget, Biogeosciences Discuss., 11, 1707–1738, doi:10.5194/bgd-11-1707-2014, 2014.
- Hamm, C. E.: Interactive aggregation and sedimentation of diatoms and clay-sized lithogenic material, Limnol. Oceanogr. 47, 1790– 1795, doi:10.4319/lo.2002.47.6.1790, 2002.
- Hansell, D. A. and Ducklow, H. W.: Bacterioplankton distribution and production in the bathypelagic ocean: Directly coupled to particulate organic carbon export?, Limnol. Oceanogr., 48, 150– 156, 2003.
- Hedges, J. I., Baldock, J. A., Gélinas, Y., Lee, C., Peterson, M., and Wakeham S. G.: Evidence for non-selective preservation of organic matter in sinking marine particles, Nature, 409, 801–804, doi:10.1038/35057247, 2000.
- Henson, S. A., Sanders, R., and Madsen, E.: Global patterns in efficiency of particulate organic carbon export and transfer to the deep ocean, Global Biogeochem. Cy., 26, GB1028, doi:10.1029/2011GB004099, 2012.
- Hill, P. S., Boss, E., Newgard, J. P., Law, B. A., and Milligan T. G.: Observations of the sensitivity of beam attenuation to particle size in a coastal bottom boundary layer, J. Geophys. Res., 116, C02023, doi:10.1029/2010JC006539, 2011.
- Honda, M. C. and Watanabe, S.: Importance of biogenic opal as ballast of particulate organic carbon (POC) transport and existence of mineral ballast-associated and residual POC in the Western Pacific Subarctic Gyre, Geophys. Res. Lett., 37, L02605, doi:10.1029/2009GL041521, 2010.
- Ingalls, A. E., Liu, Z., and Lee, C.: Seasonal trends in the pigment and amino acid compositions of sinking particles in biogenic CaCO<sub>3</sub> and SiO<sub>2</sub> dominated regions of the Pacific sector of the Southern Ocean along  $170^{\circ}$  W, Deep-Sea Res. Pt. I, 53, 836–859, 2006.
- Ittekkot, V.: The abiotically driven biological pump in the ocean and short-term fluctuations in atmospheric CO<sub>2</sub> contents, Global Planet. Change, 8, 17–25, 1993.
- Iversen, M. H. and Ploug, H.: Ballast minerals and the sinking carbon flux in the ocean: carbon-specific respiration rates and

#### M. Bressac et al.: Quantification of the lithogenic carbon pump

sinking velocity of marine snow aggregates, Biogeosciences, 7, 2613–2624, doi:10.5194/bg-7-2613-2010, 2010.

- Iversen, M. H., Nowald, N., Ploug, H., Jackson, G. A., and Fischer, G.: High resolution profiles of vertical particulate matter export off Cape Blanc, Mauritania: degradation processes and ballasting effects, Deep-Sea Res. Pt. I, 57, 711–784, 2010.
- Jackson, G. A. and Lochman, S. E.: Effect of coagulation on nutrient and light limitation of an algal bloom, Limnol. Oceanogr., 37, 77–89, 1992.
- Jickells, T. D., Dorling, S., Deuser, W. G., Church, T. M., Arimoto, R., and Prospero, J. M.: Air–borne dust fluxes to a deep water sediment trap in the Sargasso Sea, Global Biogeochem. Cy., 12, 311–320, 1998.
- Jickells, T. D., An, Z. S., Andersen, K. K., Baker, A. R., Bergametti, G., Brooks, N., Cao, J. J., Boyd, P. W., Duce, R. A., Hunter, K. A., Kawahata, H., Kubilay, N., laRoche, J., Liss, P. S., Mahowald, N., Prospero, J. M., Ridgwell, A. J., Tegen, I., and Torres, R.: Global Iron Connections Between Desert Dust, Ocean Biogeochemistry, and Climate, Science, 308, 67– 71, doi:10.1126/science.1105959, 2005.
- Klaas, C. and Archer, D. E.: Association of sinking organic matter with various types of mineral ballast in the deep sea: Implications for the rain ratio, Global Biogeochem. Cy., 16, 1116, doi:10.1029/2001GB001765, 2002.
- Lee, C., Peterson, M. L., Wakeham, S. G., Armstrong, R. A., Cochran, J. K., Miquel, J. C., Fowler, S. W., Hirschberg, D., Beck, A., and Xue, J.: Particulate organic matter and ballast fluxes measured using time-series and settling velocity sediment traps in the northwestern Mediterranean Sea, Deep-Sea Res. Pt. II, 56, 1420–1436, 2009.
- Le Moigne, F. A. C., Sanders, R. J., Villa-Alfageme, M., Martin, A. P., Pabortsava, K., Planquette, H., Morris, P. J., and Thomalla, S. J.: On the proportion of ballast versus non-ballast associated carbon export in the surface ocean, Geophys. Res. Lett., 39, L15610, doi:10.1029/2012GL052980, 2012.
- Le Moigne, F. A. C., Gallinari, M., Laurenceau, E., and De La Rocha, C. L.: Enhanced rates of particulate organic matter remineralization by microzooplankton are diminished by added ballast minerals, Biogeosciences, 10, 5755–5765, doi:10.5194/bg-10-5755-2013, 2013.
- Loisel, H., Vantrepotte, V., Norkvist, K., Mériaux, X., Kheireddine, M., Ras, J., Pujo-Pay, M., Combet, Y., Leblanc, K., Dall'Olmo, G., Mauriac, R., Dessailly, D., and Moutin, T.: Characterization of the bio-optical anomaly and diurnal variability of particulate matter, as seen from scattering and backscattering coefficients, in ultra-oligotrophic eddies of the Mediterranean Sea, Biogeosciences, 8, 3295–3317, doi:10.5194/bg-8-3295-2011, 2011.
- Loÿe-Pilot, M.-D. and Martin, J. M.: Saharan dust input to the western Mediterranean: an eleven years record in Corsica, in: The Impact of Desert Dust Across the Mediterranean, edited by: Guerzoni, S. and Chester, R., Kluwer, 191–199, 1996.
- Loÿe-Pilot, M.-D., Martin, J. M., and Morelli, J.: Influence of Saharan dust on the rain acidity and atmospheric input to the Mediterranean, Nature, 321, 427–428, doi:10.1038/321427a0, 1986.
- Lutz, M., Dunbar, R., and Caldeira, K.: Regional variability in the vertical flux of particulate organic carbon in the ocean interior, Global Biogeochem. Cy., 16, doi:10.1029/2000GB001383, 2002.

- Mahowald, N., Luo, C., del Corral, J., and Zender, C. S.: Interannual variability in atmospheric mineral aerosols from a 22-year model simulation and observational data, J. Geophys. Res., 108, 4352, doi:10.1029/2002JD002821, 2003.
- Mahowald, N. M., Engelstaedter, S., Luo, C., Sealy, A., Artaxo, P., Benitez-Nelson, C., Bonnet, S., Chen, Y., Chuang, P. Y., Cohen, D. D., Dulac, F., Herut, B., Johansen, A. M., Kubilay, N., Losno, R., Maenhaut, W., Paytan, A., Prospero, J. M., Shank, L. M., and Siefert, R. L.: Atmospheric Iron Deposition: Global Distribution, Variability, and Human Perturbations, Annu. Rev. Mar. Sci., 1, 245–278, doi:10.1146/annurev.marine.010908.163727, 2009.
- Marañón, E., Fernández, A., Mouriño-Carballido, B., Martínez-García, S., Teira, E., Cermeño, P., Chouciño, P., Huete-Ortega, M., Fernández, E., Calvo-Díaz, A., Morán, X. A. G., Bode, A., Moreno-Ostos, E., Varela, M. M., Patey, M. D., and Achterberg, E. P.: Degree of oligotrophy controls the response of microbial plankton to Saharan dust, Limnol. Oceanogr., 55, 2339–2352, 2010.
- Martin, J. H.: Glacial-interglacial CO<sub>2</sub> change: the iron hypothesis, Paleoceanography, 5, 1–13, 1990.
- Migon, C., Sandroni, V., Marty, J. C., Gasser, B., and Miquel, J. C.: Transfer of atmospheric matter through the euphotic layer in the northwestern Mediterranean: seasonal pattern and driving forces, Deep-Sea Res. Pt. II, 49, 2125–2141, 2002.
- Nagata, T., Fukuda, H., Fukuda, R., and Koike, I.: Bacterioplankton distribution and production in deep Pacific waters: Large-scale geographic variations and possible coupling with sinking particle fluxes, Limnol. Oceanogr., 45, 426–435, 2000.
- Nagata, T., Tamburini, C., Arístegui, J., Baltar, F., Bochdansky, A. B., Fonda-Umani, S., Fukuda, H., Gogou, A., Hansell, D. A., Hansman, R. L., Herndl, G. J., Panagiotopoulos, C., Reinthaler, T., Sohrin, R., Verdugo, P., Yamada, N., Yamashita, Y., Yokokawa, T., and Bartlett, D. H.: Emerging concepts on microbial processes in the bathypelagic ocean: ecology, biogeochemistry, and genomics, Deep-Sea Res. Pt. II, 57, 1519–1536, 2010.
- Neuer, S., Torres-Padrón, M. E., Gelado-Caballero, M. D., Rueda, M. J., Hernández-Brito, J., Davenport, R., and Wefer, G.: Dust deposition pulses to the eastern subtropical North Atlantic gyre: does ocean's biogeochemistry respond? Global Biogeochem. Cy., 18, GB4020, doi:10.1029/2004GB002228, 2004.
- Passow, U.: Transparent exopolymer particles (TEP) in aquatic environments, Prog. Oceanogr., 55, 287–333, 2002.
- Passow, U.: Switching perspectives: Do mineral fluxes determine particulate organic carbon fluxes or vice versa?, Geochem. Geophys. Geosyst., 5, Q04002, doi:10.1029/2003GC000670, 2004.
- Passow, U. and De La Rocha, C. L.: Accumulation of mineral ballast on organic aggregates, Global Biogeochem. Cy., 20, GB1013, doi:10.1029/2005GB002579, 2006.
- Ploug, H., Iversen, M. H., and Fischer, G.: Ballast, sinking velocity, and apparent diffusivity within marine snow and zooplankton fecal pellets: Implications for substrate turnover by attached bacteria, Limnol. Oceanogr., 53, 1878–1886, 2008.
- Prieto, L., Ruiz, J., Echevarría, F., García, C. M., Bartual, A., Gálvez, J. A., Corzo, A., and Macías, D.: Scales and processes in the aggregation of diatom blooms: high time resolution and wide size range records in a mesocosm study, Deep-Sea Res. Pt. I, 49, 1233–1253, 2002.

#### M. Bressac et al.: Quantification of the lithogenic carbon pump

- Pulido-Villena, E., Wagener, T., and Guieu, C.: Bacterial response to dust pulses in the western Mediterranean: Implications for carbon cycling in the oligotrophic ocean, Global Biogeochem. Cy., 22, GB1020, doi:10.1029/2007GB003091, 2008.
- Pulido-Villena, E., Baudoux, A.-C., Obernosterer, I., Landa, M., Caparros, J., Catala, P., Georges, C., Harmand, J., and Guieu, C.: Microbial food web dynamics in response to a Saharan dust event: results from a mesocosm study in the oligotrophic Mediterranean Sea, Biogeosciences Discuss., 11, 337– 371, doi:10.5194/bgd-11-337-2014, 2014.
- Ragueneau, O., Schultes, S., Bidle, K., Claquin, P., and Moriceau, B.: Si and C interactions in the world ocean: Importance of ecological processes and implications for the role of diatoms in the biological pump, Global Biogeochem. Cy., 20, GB4S02, doi:10.1029/2006GB002688, 2006.
- Ridame, C., Guieu, C., and L'Helguen, S.: Strong stimulation of N<sub>2</sub> fixation in oligotrophic Mediterranean Sea: results from dust addition in large in situ mesocosms, Biogeosciences, 10, 7333– 7346, doi:10.5194/bg-10-7333-2013, 2013.
- Ridame, C., Dekaezemacker, J., Guieu, C., Bonnet, S., L'Helguen, S., and Malien, F.: Phytoplanktonic response to contrasted Saharan dust deposition events during mesocosm experiments in LNLC environment, Biogeosciences Discuss., 11, 753-796, doi:10.5194/bgd-11-753-2014, 2014.
- Riley, J. S., Sanders, R., Marsay, C., Le Moigne, F. A. C., Achterberg, E. P., and Poulton, A. J.: The relative contribution of fast and slow sinking particles to ocean carbon export, Global Biogeochem. Cy., 26, GB1026, doi:10.1029/2011GB004085, 2012.
- Salter, I., Kemp, A. E. S., Lampitt, R. S., and Gledhill, M.: The association between biogenic and inorganic minerals and the amino acid composition of settling particles, Limnol. Oceanogr., 55, 2207–2218, 2010.
- Sanders, R., Morris, P. J., Poulton, A. J., Stinchcombe, M. C., Charalampopoulou, A., Lucas, M. I., and Thomalla, S. J.: Does a ballast effect occur in the surface ocean?, Geophys. Res. Lett., 37, L08602, doi:10.1029/2010GL042574, 2010.
- Scholten, J. C., Fietzke, F., Vogler, S., Rutgers van der Loeff, M., Mangini, A., Koeve, W., Waniek, J., Stoffers, P., Antia, A., and Kuss, J.: Trapping efficiency of sediment traps from the deep eastern North Atlantic: <sup>230</sup>Th calibration, Deep-Sea Res. Pt. II, 48, 243–268, 2001.
- Smetacek, V., Klaas, C., Strass, V. H., Assmy, P., Montresor, M., Cisewski, B., Savoye, N., Webb, A., d'Ovidio, F., Arrieta, J. M., Bathmann, U., Bellerby, R., Berg, G. M., Croot, P., Gonzalez, S., Henjes, J., Herndl, G. J., Hoffmann, L. J., Leach, H., Losch, M., Mills, M. M., Neill, C., Peeken, I., Röttgers, R., Sachs, O., Sauter, E., Schmidt, M. M., Schwarz, J., Terbrüggen, A., and Wolf-Gladrow, D.: Deep carbon export from a Southern Ocean iron-fertilized diatom bloom, Nature, 487, 313, doi:10.1038/nature11229, 2012.
- Stramski, D. and Kiefer, D. A.: Light scattering by microorganisms in the open ocean, Progr. Oceanogr., 28, 343–383, doi:10.1016/0079-6611(91)90032-H, 1991.

- Ternon, E., Guieu, C., Loye-Pilot, M.-D., Leblond, N., Bosc, E., Gasser, B., Miquel, J.-C., and Martin, J.: The impact of Saharan dust on the particulate export in the water column of the North Western Mediterranean Sea, Biogeosciences, 7, 809–826, doi:10.5194/bg-7-809-2010, 2010.
- Thomalla, S. J., Poulton, A. J., Sanders, R., Turnewitsch, R., Holligan, P. M., and Lucas, M. I.: Variable export fluxes and efficiencies for calcite, opal, and organic carbon in the Atlantic Ocean: A ballast effect in action?, Global Biogeochem. Cy., 22, GB1010, doi:10.1029/2007GB002982, 2008.
- Thunell, R., Benitez-Nelson, C., Varela, R., Astor, Y., and Muller-Karger, F.: Particulate organic carbon fluxes along upwellingdominated continental margins: Rates and mechanisms, Global Biogeochem. Cy., 21, GB1022, doi:10.1029/2006GB002793, 2007.
- Twardowsky, M. S., Boss, E., Macdonald, J. B., Pegau, W. S., Barnard, A. H., and Zaneveld J. R. V.: A model for estimating bulk refractive index from the optical backscattering ratio and the implications for understanding particle composition in case I and case II waters, J. Geophys. Res., 106, 14129–14142, doi:10.1029/2000JC000404, 2001.
- Ulloa, O., Sathyendranath, S., and Platt, T.: Effect of the particlesize distribution on the backscattering ratio in seawater, Appl Optics, 33, 7070–7077, doi:10.1364/AO.33.007070, 1994.
- Verdugo, P., Alldredge, A. L., Azam, F., Kirchman, D. L., Passow, U., and Santschi, P. H.: The oceanic gel phase: a bridge in the DOM–POM continuum, Mar. Chem., 92, 67–85, doi:10.1016/j.marchem.2004.06.017, 2004.
- Wedepohl, K. H.: The composition of the continental crust, Geochim. Cosmochim. Ac., 59, 1217–1232, 1995.
- Wilson, J. D., Barker, S., and Ridgwell, A.: Assessment of the spatial variability in particulate organic matter and mineral sinking fluxes in the ocean interior: Implications for the ballast hypothesis, Global Biogeochem. Cy., 26, GB4011, doi:10.1029/2012GB004398, 2012.
- Wong, C., Whitney, F., Crawford, K., Iseki, K., Matear, R., Johnson, W., Page, J., and Timothy, D.: Seasonal and interannual variability in particle fluxes of carbon, nitrogen and silicon from time series of sediment traps at Ocean Station P, 1982–1993: Relationship to changes in subarctic primary productivity, Deep-Sea Res., Pt. II, 46, 2735–2760, doi:10.1016/S0967-0645(99)00082-X, 1999.
- Wuttig, K., Wagener, T., Bressac, M., Dammshäuser, A., Streu, P., Guieu, C., and Croot, P. L.: Impacts of dust deposition on dissolved trace metal concentrations (Mn, Al and Fe) during a mesocosm experiment, Biogeosciences, 10, 2583–2600, doi:10.5194/bg-10-2583-2013, 2013.
- Yu, E. F., Francois, R., Bacon, M. P., Honjo, S., Fleer, A. P., Manganini, S. J., Rutgers van der Loeff, M. M., and Ittekot, V.: Trapping efficiency of bottom-tethered sediment traps estimated from the intercepted fluxes of <sup>230</sup>Th and <sup>231</sup>Pa, Deep-Sea Res. Pt. I, 48, 865–889, 2001.