



HAL
open science

Estimation of population parameters in stochastic differential equations with random effects in the diffusion coefficient

Maud Delattre, Valentine Genon-Catalot, Adeline Samson

► **To cite this version:**

Maud Delattre, Valentine Genon-Catalot, Adeline Samson. Estimation of population parameters in stochastic differential equations with random effects in the diffusion coefficient. 2014. hal-00954438

HAL Id: hal-00954438

<https://hal.science/hal-00954438>

Submitted on 3 Mar 2014

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Estimation of population parameters in stochastic differential equations with random effects in the diffusion coefficient.

Maud Delattre¹, Valentine Genon-Catalot² and Adeline Samson^{3*}

¹ AgroParisTech, UMR518, 75005 Paris, France

² UMR CNRS 8145, Laboratoire MAP5, Université Paris Descartes, Sorbonne Paris Cité, France

³ Laboratoire Jean Kuntzmann, UMR CNRS 5224, Univ Grenoble-Alpes, France

March 2, 2014

Abstract

We consider N independent stochastic processes $(X_i(t), t \in [0, T_i])$, $i = 1, \dots, N$, defined by a stochastic differential equation with diffusion coefficients depending on a random variable ϕ_i . The distribution of the random effect ϕ_i depends on unknown population parameters which are to be estimated from a discrete observation of the processes (X_i) . The likelihood generally does not have any closed form expression. Two estimation methods are proposed: one based on the Euler approximation of the likelihood and another based on estimations of the random effects. When the distribution of the random effects is Gamma, the asymptotic properties of the estimators are derived when both N and the number of observations per subject tend to infinity. The estimators are computed on simulated data for several models and show good performances.

Key Words: Approximate maximum likelihood estimator, asymptotic normality, consistency, estimating equations, random effects models, stochastic differential equations.

*Corresponding author

Email addresses: maud.delattre@agroparistech.fr (Maud Delattre), valentine.genon-catalot@parisdescartes.fr (Valentine Genon-Catalot), adeline.leclercq-samson@imag.fr (Adeline Samson)

1 Introduction

Stochastic differential equations (SDEs) with random effects have been the subject of several recent contributions, with various applications such as pharmacokinetic/pharmacodynamic, neuronal modeling (Picchini *et al.*, 2010; Delattre and Lavielle, 2013; Donnet and Samson, 2013). Several estimation methods have been proposed to provide estimators in these complex models. The exact maximum likelihood can be studied theoretically (Nie, 2006) but the likelihood has no explicit expression except in some special cases. In Delattre *et al.* (2013), the case of a linear random effect in the drift together with a specific distribution for the random effects is investigated. In this case, the exact maximum likelihood estimator is explicit and studied. In the general case, Picchini *et al.* (2010); Picchini and Ditlevsen (2011) propose approximations of the likelihood based on Hermite expansion and Gaussian quadrature. All these references work with random effects in the drift, and not in the diffusion coefficient, except Delattre and Lavielle (2013) who incorporate measurement error and propose an approximation of the likelihood with the extended Kalman filter.

Here, we focus on discretely observed SDEs with a random effect in the diffusion coefficient. The distribution of the random effect depends on unknown parameters to be estimated. For simplicity, we assume that the drift is zero and that there is a linear random effect in the diffusion coefficient. Extensions are discussed in Sections 4 and 5, in particular the case of non null drift and of a more general diffusion coefficient.

Statistical inference for discretely observed SDEs with no random effects has been widely studied (see Kessler *et al.*, 2012, and references therein). In Genon-Catalot and Jacod (1993) the estimation of unknown fixed parameters in the diffusion coefficient is studied with discrete observations of a single trajectory when the sampling interval tends to zero. The likelihood of these observations is not explicit, therefore estimating equations are built based on the Euler approximation of the SDE with drift set to zero. One of the strategies described below follows the same idea, but here the parameters are random. This complicates the definition and the theoretical study of the estimator.

More precisely, we consider N real valued stochastic processes $(X_i(t), t \geq 0)$, $i = 1, \dots, N$, with dynamics ruled by the following SDEs:

$$dX_i(t) = \phi_i \sigma(X_i(t)) dW_i(t), \quad X_i(0) = x_i^0, \quad i = 1, \dots, N, \quad (1)$$

where $(W_i)_{1 \leq i \leq N}$ are N independent Wiener processes, $(\phi_i)_{1 \leq i \leq N}$ are N *i.i.d.* random variables taking values in $(0, +\infty)$, $(\phi_i)_{1 \leq i \leq N}$ and $(W_i)_{1 \leq i \leq N}$ are independent. The function $\sigma(x)$ is known and real-valued. Each process $(X_i(t))$ represents an individual, the variable ϕ_i represents the random effect of individual i . The variables $(\phi_i)_{1 \leq i \leq N}$ have a common distribution $g(\varphi, \theta)d\nu(\varphi)$ on $(0, +\infty)$ where ν is a dominating measure and θ is a vector of unknown parameters called population parameters, belonging to a set $\Theta \subset \mathbb{R}^p$.

Our aim is to estimate θ from discrete observations $\{X_i(t_{i,j}), j = 1, \dots, n, i = 1, \dots, N\}$. In the case of a linear random effect in the diffusion coefficient (1), choosing an inverse Gamma distribution leads to explicit estimators. Therefore, we consider the specific case

$$\phi_i = \frac{1}{\Gamma_i^{1/2}} \quad \text{with} \quad \Gamma_i \sim G(a, \lambda), \quad a > 0, \lambda > 0, \quad \theta = (a, \lambda). \quad (2)$$

We study the exact maximum likelihood estimator in the case $\sigma(\cdot) \equiv 1$. When $\sigma(\cdot) \not\equiv 1$, we build estimating equations based on the Euler approximation of the fixed effect diffusion model. The difficulty of these estimating equations is that the Euler approximation has to be integrated out with respect to the distribution of the random effects. Moreover, we build another type of estimating equations, corresponding to the ideal likelihood of directly observed random effects where estimators of the random effects are plugged in. This second approach has the advantage to be easily generalized to any distribution for the random effects.

The paper is organized as follows. Section 2 introduces some assumptions and gives the exact likelihood and its approximation obtained by Euler scheme. Our asymptotic framework is when the number N of subjects tends to infinity. In Section 3, we study the asymptotic properties of the estimators. When $\sigma(\cdot) \equiv 1$, the exact maximum likelihood estimator of θ is asymptotically Gaussian with rate \sqrt{N} both for fixed number of measurements per subject n and for n tending to infinity. When $\sigma(\cdot) \not\equiv 1$, we must assume that n depends on N and satisfies the constraint $N/n \rightarrow 0$ for the first method, $\sqrt{N}/n \rightarrow 0$ for the second. Our estimators are asymptotically Gaussian with rate \sqrt{N} . Simulations illustrate the behavior of the estimators and results are presented in Section 4. Section 5 concludes the paper with some extensions. Proofs are gathered in Appendix.

2 Exact and approximate likelihoods

Consider N real valued stochastic processes $(X_i(t), t \geq 0)$, $i = 1, \dots, N$, with dynamics ruled by (1). The processes $(W_i)_{1 \leq i \leq N}$ and the r.v.'s $(\phi_i)_{1 \leq i \leq N}$ are defined on a common probability space $(\Omega, \mathcal{F}, \mathbb{P})$. Consider the filtration $(\mathcal{F}_t, t \geq 0)$ defined by $\mathcal{F}_t = \sigma(\phi_i, W_i(s), s \leq t, i = 1, \dots, N)$. As $\mathcal{F}_t = \sigma(W_i(s), s \leq t) \vee \mathcal{F}_t^i$, with $\mathcal{F}_t^i = \sigma(\phi_i, \phi_j, W_j(s), s \leq t, j \neq i)$ independent of W_i , each process W_i is a $(\mathcal{F}_t, t \geq 0)$ -Brownian motion. Moreover, the random variables ϕ_i are \mathcal{F}_0 -measurable. In what follows, we assume that

(H1) The function σ belongs to $C^2(\mathbb{R})$ and for all $x \in \mathbb{R}$, $0 < \sigma_0^2 \leq \sigma^2(x) \leq \sigma_1^2$,
 $|\sigma'(x)| + |\sigma''(x)| \leq K$.

Under (H1), the process $(X_i(t))$ is well-defined and $(\phi_i, X_i(t))$ is strong Markov adapted to the filtration $(\mathcal{F}_t, t \geq 0)$. The N processes $(\phi_i, X_i(\cdot))_{1 \leq i \leq N}$ are independent. For all φ , and all $x_i^0 \in \mathbb{R}$, the fixed effect SDE

$$dX_i^{\varphi, x_i^0}(t) = \varphi \sigma(X_i^{\varphi, x_i^0}(t)) dW_i(t), \quad X_i^{\varphi, x_i^0}(0) = x_i^0 \quad (3)$$

admits a unique strong solution process $(X_i^{\varphi, x_i^0}(t), t \geq 0)$ adapted to the filtration $(\mathcal{F}_t, t \geq 0)$. From the Markov property of $(\phi_i, X_i(t))$, we deduce that the conditional distribution of X_i given $\phi_i = \varphi$ is identical to the distribution of X_i^{φ, x_i^0} (for more details, see Delattre *et al.*, 2013).

For $i = 1, \dots, N$, the process $(X_i(t), t \in [0, T_i])$ is discretely observed at times $t_{i,j} = jT_i/n, j = 0, \dots, n$ and we set

$$\Delta_i = \frac{T_i}{n}, \quad X_i = (X_i(t_{i,j}), j = 1, \dots, n), \quad \text{with } T_i \leq T, i = 1, \dots, N, \quad (4)$$

where T_1, \dots, T_N, T are fixed. The number of observations per subject is the same for all subjects but the sampling intervals may be distinct.

We start by the exact likelihood of (4). The distribution of the observations $(X_i)_{1 \leq i \leq N}$ on $\prod_{i=1}^N \mathbb{R}^n$ has the form $P_\theta = \otimes_{i=1}^N P_\theta^i$ where P_θ^i is the distribution of X_i on \mathbb{R}^n . If $Q_{\varphi, x_i^0}^i$ denotes the distribution of $X_i^\varphi = (X_i^{\varphi, x_i^0}(t_{i,j}), j = 1, \dots, n)$ and $p_t(x, y, \varphi)$ the transition density of (3), then $Q_{\varphi, x_i^0}^i$ admits the density $\prod_{j=1}^n p_{\Delta_i}(x_{i,j-1}, x_{i,j}, \varphi)$ w.r.t. the Lebesgue measure of \mathbb{R}^n (with $x_{i,0} = x_i^0$). Therefore, the density of P_θ^i w.r.t. the Lebesgue measure of \mathbb{R}^n is given by:

$$\lambda_i(\theta, x_i) = \int_0^{+\infty} g(\varphi, \theta) \prod_{j=1}^n p_{\Delta_i}(x_{i,j-1}, x_{i,j}, \varphi) d\nu(\varphi).$$

The exact likelihood is $\Lambda_N(\theta) = \prod_{i=1}^N \lambda_i(\theta, X_i)$. Here, we are faced with two problems. First, the transition densities of (3) are generally not explicit. Second, and this is specific to SDE with random effect, even if these transition densities were explicit, the density of (ϕ_i, X_i) would be explicit but it is generally not possible to get a closed-form expression for the marginal density of X_i , which corresponds to the integral $\lambda_i(\theta, x_i)$. Therefore the exact likelihood is not explicit and difficult to study theoretically and numerically.

Instead of using the exact transition densities of(3), it is now standard to use the approximation given by the transition densities of the corresponding Euler scheme, *i.e.* the one-step discretisation of (3) (see *e.g.* Genon-Catalot and Jacod, 1993; Donnet and Samson, 2008; Kessler *et al.*, 2012). Therefore, we introduce

$$\tilde{L}_i(X_i, \varphi) = \frac{1}{\varphi^n \prod_{j=1}^n \sigma(X_i(t_{i,j}))} \exp\left(-\frac{S_i}{2\varphi^2}\right) \propto \frac{1}{\varphi^n} \exp\left(-\frac{S_i}{2\varphi^2}\right),$$

with, for $i = 1, \dots, N$,

$$S_i = \frac{1}{\Delta_i} \sum_{j=1}^n \frac{(X_i(t_{i,j}) - X_i(t_{i,j-1}))^2}{\sigma^2(X_i(t_{i,j-1}))} \quad (5)$$

To estimate θ , instead of the exact likelihood, we introduce the approximate likelihood, corresponding to the Euler scheme integrated with respect to the random effects distribution:

$$\tilde{\Lambda}_N(\theta) = \prod_{i=1}^N \int_0^{+\infty} \varphi^{-n} \exp\left(-\frac{S_i}{2\varphi^2}\right) g(\varphi, \theta) d\nu(\varphi). \quad (6)$$

A theoretical study of the estimators based on $\tilde{\Lambda}_N(\theta)$ could be possible using the approach developed by Nie (2006) but his assumptions are generally difficult to verify. Below, as in Delattre *et al.* (2013), we rather introduce a specific distribution for the random effects allowing to obtain an explicit formula for (6). In Section 3, we are able to directly study the corresponding estimators.

Remark 1. *Except in the case $\sigma(\cdot) \equiv 1$ where (6) is the exact likelihood, our approach based on an approximate likelihood imposes a double asymptotic framework where both N and n tend to infinity. As $n \rightarrow \infty$, note that the statistic S_i based on the i -th trajectory provides an estimator of the random effect ϕ_i .*

Indeed, let $M_i(t) = \int_0^t \sigma(X_i(s))dW_i(s)$, and

$$R_i = \sum_{j=1}^n (M_i(t_{i,j}) - M_i(t_{i,j-1}))^2 / \sigma^2(X_i(t_{i,j-1})). \quad (7)$$

By standard properties of quadratic variations, $R_i/T_i \rightarrow 1$ in probability as $n \rightarrow \infty$. Thus, $S_i/n = \phi_i^2 R_i/T_i$ tends to ϕ_i^2 .

3 A specific distribution for the random effect

For a general distribution $g(\varphi, \theta)d\nu(\varphi)$ of the random effect ϕ_i , the integral in (6) has no explicit expression. However, for the conjugate distribution, namely the inverse Gamma (2), an explicit expression is obtained. The unknown parameter is then $\theta = (a, \lambda) \in \Theta = \mathbb{R}^+ \times \mathbb{R}^+$. The true value is denoted by θ_0 .

Let us start with the ideal case of directly observed random effects ϕ_i (or Γ_i). Then, the exact log-likelihood of $(\Gamma_1, \dots, \Gamma_N)$ is given by:

$$\ell_N(\theta) = Na \log \lambda - N \log \Gamma(a) + (a-1) \sum_{i=1}^N \log \Gamma_i - \lambda \sum_{i=1}^N \Gamma_i \quad (8)$$

with associated score function $\mathcal{S}_N(\theta) = \left(\frac{\partial}{\partial \lambda} \ell_N(\theta) \quad \frac{\partial}{\partial a} \ell_N(\theta) \right)'$ where

$$\frac{\partial}{\partial \lambda} \ell_N(\theta) = \sum_{i=1}^N \left(\frac{a}{\lambda} - \Gamma_i \right), \quad \frac{\partial}{\partial a} \ell_N(\theta) = \sum_{i=1}^N (-\psi(a) + \log \lambda + \log \Gamma_i),$$

where $\psi(z) = \frac{\Gamma'(z)}{\Gamma(z)}$ is the di-gamma function. By standard properties of Gamma distributions, we have, under the true value θ_0 , $(1/\sqrt{N})\mathcal{S}_N(\theta_0) \rightarrow_{\mathcal{D}} \mathcal{N}_2(0, \mathcal{I}(\theta_0))$, where $\mathcal{I}(\theta)$ is

$$\mathcal{I}(\theta) = \begin{pmatrix} \frac{a}{\lambda^2} & -\frac{1}{\lambda} \\ -\frac{1}{\lambda} & \psi'(a) \end{pmatrix}. \quad (9)$$

Note that using properties of the di-gamma function (see Section 7), $\mathcal{I}(\theta)$ is invertible for all $\theta \in (0, +\infty)^2$. The maximum likelihood estimator based on the observation of $\Gamma_1, \dots, \Gamma_N$, denoted $\theta_N = \theta_N(\Gamma_1, \dots, \Gamma_N)$ is consistent and satisfies $\sqrt{N}(\theta_N - \theta_0) \rightarrow_{\mathcal{D}} \mathcal{N}_2(0, \mathcal{I}^{-1}(\theta_0))$ as N tends to infinity.

But the Γ_i 's are not observed. Two different strategies are studied. Following

Remark 1, a natural idea consists in plugging in $\ell_N(\theta)$ the estimator n/S_i of Γ_i . This reveals to be more complex than expected (Section 3.2) and we will need to truncate the estimator n/S_i . The other strategy (Section 3.1) is based on (6). We provide asymptotic results when n is fixed and $N \rightarrow \infty$ in the case $\sigma(\cdot) \equiv 1$, and when both $n, N \rightarrow \infty$ for a general $\sigma(\cdot)$.

3.1 Estimation based on the Euler approximation of the likelihood.

Let $\tilde{\mathcal{L}}_N(\theta) = \log \tilde{\Lambda}_N(\theta)$ be the log contrast Euler (see (6)).

Proposition 1. *Under (H1) and (2), we have:*

$$\tilde{\mathcal{L}}_N(\theta) = \sum_{i=1}^N \log \left(\frac{\Gamma(a + n/2)}{\Gamma(a)} \right) + aN \log \lambda - \sum_{i=1}^N (a + n/2) \log \left(\lambda + \frac{1}{2} S_i \right). \quad (10)$$

The associated gradient vector (pseudo-score function)

$$G_N(\theta) = \left(\frac{\partial}{\partial \lambda} \tilde{\mathcal{L}}_N(\theta) \quad \frac{\partial}{\partial a} \tilde{\mathcal{L}}_N(\theta) \right)' \quad (11)$$

is given by $\frac{\partial}{\partial \lambda} \tilde{\mathcal{L}}_N(\theta) = \sum_{i=1}^N \left(\frac{a}{\lambda} - \frac{a+n/2}{\lambda+S_i/2} \right)$ and $\frac{\partial}{\partial a} \tilde{\mathcal{L}}_N(\theta) = \sum_{i=1}^N (\psi(a + n/2) - \psi(a)) + \sum_{i=1}^N \log \left(\frac{\lambda}{\lambda+S_i/2} \right)$. For the Hessian matrix (pseudo Fisher information matrix)

$$\tilde{\mathcal{I}}_N(\theta) = - \begin{pmatrix} \frac{\partial^2}{\partial \lambda^2} \tilde{\mathcal{L}}_N(\theta) & \frac{\partial^2}{\partial \lambda \partial a} \tilde{\mathcal{L}}_N(\theta) \\ \frac{\partial^2}{\partial \lambda \partial a} \tilde{\mathcal{L}}_N(\theta) & \frac{\partial^2}{\partial a^2} \tilde{\mathcal{L}}_N(\theta) \end{pmatrix}, \quad (12)$$

we get $\frac{\partial^2}{\partial \lambda^2} \tilde{\mathcal{L}}_N(\theta) = - \sum_{i=1}^N \left(\frac{a}{\lambda^2} - \frac{a+n/2}{(\lambda+S_i/2)^2} \right)$, $\frac{\partial^2}{\partial \lambda \partial a} \tilde{\mathcal{L}}_N(\theta) = \sum_{i=1}^N \left(\frac{1}{\lambda} - \frac{1}{\lambda+S_i/2} \right)$ and $\frac{\partial^2}{\partial a^2} \tilde{\mathcal{L}}_N(\theta) = - \sum_{i=1}^N (\psi'(a) - \psi'(a + n/2))$. We study the estimators defined by the estimating equation:

$$G_N(\tilde{\theta}_N) = 0. \quad (13)$$

We consider two asymptotics: n fixed (section 3.1.1) and $n \rightarrow \infty$ (section 3.1.2).

3.1.1 Fixed number of observations per subject

We assume that the number n of observations per subject is fixed and that the number of subjects N tends to infinity. The only model that enters this

asymptotic is the special case $\sigma(\cdot) \equiv 1$. We denote by the upper index 1 all the quantities associated to this model: $dX_i^1(t) = \phi_i dW_i(t)$, and the statistic is

$$S_i^1 = \sum_{j=1}^n (X_i^1(t_{i,j}) - X_i^1(t_{i,j-1}))^2 / \Delta_i. \quad (14)$$

The distribution of S_i^1 can be explicitly computed.

Proposition 2. *Under P_θ , the random variables $\beta_i^1(\lambda) = \frac{\lambda}{\lambda + S_i^1/2}$, $i = 1, \dots, N$, are independent and $\beta_i^1(\lambda)$ has distribution beta of the first kind on $(0, 1)$ with parameters $(a, n/2)$. The random variables $S_i^1/(2\lambda)$ are independent with distribution on $(0, +\infty)$ beta of the second kind with parameters $(n/2, a)$.*

Then, $\tilde{\mathcal{L}}_N(\theta) = \tilde{\mathcal{L}}_N^1(\theta)$ where S_i is replaced by S_i^1 is the exact log-likelihood. Define the associated exact maximum likelihood estimator as any solution of:

$$\hat{\theta}_N^1 = \text{Argsup}_\theta \tilde{\mathcal{L}}_N^1(\theta). \quad (15)$$

The asymptotic study of $\hat{\theta}_N^1$ when n is fixed and N tends to infinity is standard: inference on θ is simply based of the *i.i.d.* sample $(S_i^1, i = 1, \dots, N)$.

Proposition 3. *Assume that n is fixed. Then, the maximum likelihood estimator $\hat{\theta}_N^1$ (15) is consistent. Let*

$$\mathcal{I}_n(\theta) = \begin{pmatrix} \frac{a(n/2)}{\lambda^2(a+1+n/2)} & -\frac{n/2}{\lambda(a+n/2)} \\ -\frac{n/2}{\lambda(a+n/2)} & \psi'(a) - \psi'(a+n/2) \end{pmatrix} \quad (16)$$

Then, the matrix $\mathcal{I}_n(\theta_0)$ is invertible and under P_{θ_0} $\sqrt{N}(\hat{\theta}_N^1 - \theta_0) \rightarrow_{\mathcal{D}} \mathcal{N}_2(0, \mathcal{I}_n^{-1}(\theta_0))$.

Remark that $\mathcal{I}_n(\theta) = \mathcal{I}(\theta) + O(\frac{1}{n})$.

3.1.2 Number of observations per subject goes to infinity

Now, we assume that both n and N tend to infinity with all T_i 's fixed and $T_i \leq T$ for some fixed T . The strategy consists in studying the case $\sigma(\cdot) \equiv 1$ where computations can all be done explicitly and then studying the difference between the general case and the case $\sigma(\cdot) \equiv 1$. Some preliminary results are needed. For these results, we do not assume that the ϕ_i^2 's have inverse Gamma distribution. As already said before, $S_i/n = \phi_i^2 R_i/T_i$ tends to ϕ_i^2 as n tends to infinity (see (7) for the definition of R_i). Let us define the equivalent of R_i for the model $\sigma(\cdot) \equiv 1$: $R_i^1 = \sum_{j=1}^n (W_i(t_{i,j}) - W_i(t_{i,j-1}))^2$.

We denote by $\mathbb{P}_\theta = \otimes_{i \geq 1} \mathbb{P}_\theta^i$ the distribution of the sequence of processes $(\phi_i, (X_i(t), t \in [0, T_i]), i \geq 1)$ on $(0, +\infty) \times \prod_{i \geq 1} C([0, T_i])$, by \mathbb{E}_θ the corresponding expectation. Note that P_θ is the marginal distribution of $(X_i, i = 1, \dots, N)$ under \mathbb{P}_θ . Both R_i/T_i and R_i^1/T_i tend to 1 in probability as $n \rightarrow \infty$. Furthermore:

Proposition 4. *Under (H1), for all θ , we have $\mathbb{E}_\theta \left(\frac{R_i^1}{T_i} - 1 | \phi_i \right) = 0$, $|\mathbb{E}_\theta \left(\frac{R_i}{T_i} - 1 | \phi_i \right)| \leq C \frac{T_i}{n} \phi_i^2$, and $|\mathbb{E}_\theta \left(\frac{R_i}{T_i} - \frac{R_i^1}{T_i} | \phi_i \right)| \leq C \frac{T_i}{n} \phi_i^2$, and for all $p \geq 1$, $\mathbb{E}_\theta \left(\left(\frac{R_i}{T_i} - 1 \right)^{2p} | \phi_i \right) \leq C \left(\frac{T_i^p}{n^p} + \frac{T_i^{2p}}{n^{2p}} (\phi_i^{2p} + \phi_i^{4p}) \right)$ and $\mathbb{E}_\theta \left(\left(\frac{R_i}{T_i} - \frac{R_i^1}{T_i} \right)^{2p} | \phi_i \right) \leq C \frac{T_i^{2p}}{n^{2p}} (\phi_i^{2p} + \phi_i^{4p})$.*

We now study the score function (11) and the Fisher information matrix (12).

Proposition 5. *Recall $G_N(\theta_0)$ defined by (11) and $\mathcal{I}(\theta_0)$ given in (9).*

For $\sigma(\cdot) \equiv 1$, as N, n tend to infinity, under P_{θ_0} , $G_N(\theta_0)/\sqrt{N}$ converges in distribution to $\mathcal{N}_2(0, \mathcal{I}(\theta_0))$.

In the general case, if $\mathbb{E}_{\theta_0} \phi_i^8 < +\infty$, i.e. if $a_0 > 4$, $n > 8$ and N, n tend to infinity in such a way that N/n tends to 0, the same result holds.

The convergence of the Fisher information matrix is as follows:

Proposition 6. *In the case $\sigma(\cdot) \equiv 1$ and the general case, the Fisher information matrix given in (12), $\tilde{\mathcal{I}}_N(\theta_0)/N$, converges in probability to $\mathcal{I}(\theta_0)$ as N, n tend to infinity, under P_{θ_0} .*

Now we study the estimator $\tilde{\theta}_N$ defined by (13).

Proposition 7. *Assume that $n, N \rightarrow +\infty$ in such a way that N/n tends to 0.*

Then, an estimator $\tilde{\theta}_N$ which solves (13) exists with probability tending to one as N tends to infinity under P_{θ_0} and is weakly consistent. The matrix $\mathcal{I}(\theta_0)$ is invertible and under P_{θ_0} , $\sqrt{N}(\tilde{\theta}_N - \theta_0) \rightarrow_{\mathcal{D}} \mathcal{N}_2(0, \mathcal{I}^{-1}(\theta_0))$.

Moreover, the estimator $\tilde{\theta}_N$ is asymptotically equivalent to the MLE $\theta_N = \theta_N(\Gamma_1, \dots, \Gamma_N)$ based on the direct observation of $(\Gamma_1, \dots, \Gamma_N)$.

The constraint $N/n = o(1)$ is the same than what would be required with N observed trajectories of a fixed effect SDE but the rate of convergence would be \sqrt{Nn} while it is only \sqrt{N} with a random effect SDE. This rate is equivalent to the one obtained when the random effects Γ_i are directly observed, but in the latter case, the constraint $N/n = o(1)$ is not needed.

3.2 Approach based on estimators of the random effects.

In this section, we exploit directly the fact that the random effect $\phi_i^2 = \Gamma_i^{-1}$ can be estimated using the trajectory $X_i(t), t \leq T_i$ by S_i/n . The idea is simply to

replace the random variables Γ_i by their estimator n/S_i in the likelihood (8) of $(\Gamma_1, \dots, \Gamma_N)$. But this works only when $\sigma(\cdot) \equiv 1$, otherwise we need to truncate the estimators. More precisely, let us set, in the case $\sigma(\cdot) \equiv 1$:

$$U_N(\theta) = Na \log \lambda - N \log \Gamma(a) + (a-1) \sum_{i=1}^N \log(n/S_i^1) - \lambda \sum_{i=1}^N (n/S_i^1).$$

and consider the estimators θ_N^* given by

$$\nabla U_N(\theta_N^*) = 0. \quad (17)$$

Otherwise, we define truncated estimators of $\log \Gamma_i$ and Γ_i as follows:

$$\widetilde{\log \Gamma}_i = \log(n/S_i) \mathbb{1}_{(S_i/n \geq k/\sqrt{n})}, \quad \widetilde{\Gamma}_i = (n/S_i) \mathbb{1}_{(S_i/n \geq k/\sqrt{n})}.$$

where k is a constant. Note that, by the above definitions, $\widetilde{\log \Gamma}_i$ and $\widetilde{\Gamma}_i$ are set to 0 outside the set $(S_i/n \geq k/\sqrt{n})$ where S_i/n is not bounded from below. Then we consider the function

$$V_N(\theta) = Na \log \lambda - N \log \Gamma(a) + (a-1) \sum_{i=1}^N \widetilde{\log \Gamma}_i - \lambda \sum_{i=1}^N \widetilde{\Gamma}_i,$$

and the associated estimator θ_N^{**} defined by the estimating equation:

$$\nabla V_N(\theta_N^{**}) = 0. \quad (18)$$

Proposition 8. *Assume that $\sigma(\cdot) \equiv 1$. If N, n tend to infinity in such a way that \sqrt{N}/n tends to 0, then an estimator θ_N^* which solves (17) exists with probability tending to 1 under \mathbb{P}_{θ_0} and is weakly consistent. Moreover, $\sqrt{N}(\theta_N^* - \theta_0)$ converges in distribution to $\mathcal{N}_2(0, \mathcal{I}^{-1}(\theta_0))$ and θ_N^* is asymptotically equivalent to the exact MLE θ_N associated to $(\Gamma_1, \dots, \Gamma_N)$, i.e. $\sqrt{N}(\theta_N^* - \theta_N) = o_{P_{\theta_0}}(1)$. When $\sigma(\cdot)$ is not equal to 1, the same result holds for θ_N^{**} under the condition $\mathbb{E}_{\theta_0} \phi_i^8 < +\infty$, i.e. $a_0 > 4$.*

Note that in this approach, even when $\sigma(\cdot) \equiv 1$, the constraint $\sqrt{N}/n \rightarrow 0$ is required.

4 Numerical simulation results.

We compare the performances of both estimation methods on simulated data for several models. Two sets of population parameters θ_0 are used: $(a_0 = 6, \lambda_0 = 1)$ and $(a_0 = 5, \lambda_0 = 3)$. In each case, 100 datasets are generated with an Euler scheme with sampling interval $\delta = 10^{-4}T$ on time interval $[0, T]$, with $T = 5$, and $N = 50, 100$ subjects, $n = 500, 1000, 10000$. The parameter θ_0 is estimated via $\tilde{\theta}_N$ (method 1) and via either θ_N^* or θ_N^{**} (method 2). The empirical mean and standard deviation are computed from the 100 datasets. We consider:

Example 1. $dX_i(t) = \phi_i dW_i(t)$, $X_i(0) = 0$.

Example 2. $dX_i(t) = \phi_i \sqrt{1 + X_i^2(t)} dW_i(t)$, $X_i(0) = 0$.

For *Example 1*, estimation method 1 leads to the exact MLE of θ_0 . For *Example 2*, the volatility function does not match the whole conditions from Section 2 ($\sigma(x)$ not bounded above). Nevertheless, considering models where the volatility is a sub-linear function does not change the results stated above.

A SDE with drift $dX_i(t) = b(X_i(t))dt + \phi_i \sigma(X_i(t))dW_i(t)$, $X_i(0) = 0$, can also be considered as if $b(\cdot) \equiv 0$ using the same inferential strategies (this is done in the case of fixed-effects in Genon-Catalot and Jacod (1993)). As the statistic S_i only depends on the volatility function, the estimation methods 1 and 2 do not require to know the expression of the drift function. We consider:

Example 3. $dX_i(t) = -\rho X_i(t)dt + \phi_i dW_i(t)$, $X_i(0) = 0$.

Example 4. $dX_i(t) = -\rho X_i(t)dt + \phi_i \sqrt{1 + X_i^2(t)} dW_i(t)$, $X_i(0) = 0$.

Note that for method 1, numerical difficulties may appear due to the term $\Gamma(a + n/2)$ in (10). To avoid these, we use the approximation for all (a, a') : $\log \Gamma(a + \frac{n}{2}) - \log \Gamma(a' + \frac{n}{2}) = (a - a') \log \frac{n}{2} + O(\frac{1}{n})$. Thus, we have

$$\begin{aligned} \frac{1}{N} (\tilde{\mathcal{L}}_N(\theta) - \tilde{\mathcal{L}}_N((1, 1))) &= a \log \lambda - (a + \frac{n}{2}) \frac{1}{N} \sum_{i=1}^N \log \frac{n^{-1} \lambda + (n^{-1}/2) S_i}{n^{-1} + (n^{-1}/2) S_i} \\ &\quad - (a - 1) \frac{1}{N} \sum_{i=1}^N \log (2n^{-1} + n^{-1} S_i) - \log \Gamma(a) + O(\frac{1}{n}). \end{aligned}$$

The results for *Examples 1* to *4* are displayed in Tables 1 to 4 respectively. The results are satisfactory overall and similar for the 4 models, even when the model includes a drift. Method 1 estimators are biased for $n = 500, 1000$. When $\sigma(\cdot) \neq 1$, this is expected due to the Euler approximation of the likelihood. Nevertheless, for fixed N , we observe the convergence of the estimators to the true value when n increases. Even though method 1 estimators seem

	$N = 50$			$N = 100$		
	$n = 500$	$n = 1000$	$n = 10000$	$n = 500$	$n = 1000$	$n = 10000$
	$(a_0 = 5, \lambda_0 = 3)$					
\tilde{a}	4.38 (0.60)	4.67 (0.74)	5.04 (0.97)	4.38 (0.46)	4.67 (0.56)	5.03 (0.70)
$\tilde{\lambda}$	2.63 (0.39)	2.81 (0.47)	3.03 (0.61)	2.64 (0.31)	2.82 (0.37)	3.04 (0.45)
a^*	5.00 (0.94)	5.03 (0.96)	5.09 (1.00)	4.97 (0.70)	5.02 (0.71)	5.08 (0.72)
λ^*	3.00 (0.59)	3.02 (0.60)	3.06 (0.63)	2.99 (0.46)	3.02 (0.46)	3.06 (0.47)
	$(a_0 = 6, \lambda_0 = 1)$					
\tilde{a}	5.12 (0.74)	5.61 (0.94)	6.32 (1.34)	5.02 (0.45)	5.50 (0.59)	6.08 (0.80)
$\tilde{\lambda}$	0.86 (0.13)	0.94 (0.16)	1.06 (0.23)	0.84 (0.08)	0.91 (0.11)	1.01 (0.14)
a^*	6.26 (1.40)	6.32 (1.38)	6.41 (1.41)	6.00 (0.79)	6.10 (0.83)	6.16 (0.84)
λ^*	1.04 (0.23)	1.06 (0.23)	1.07 (0.24)	1.00 (0.14)	1.01 (0.14)	1.02 (0.15)

Table 1: Example 1: $\sigma(\cdot) \equiv 1$. Empirical mean and standard deviation (in brackets) of $\tilde{\theta}$ (method 1) and θ^* (method 2) computed from 100 datasets.

slightly overestimated when $n = 10000$, the bias tends to vanish when N increases. This clearly illustrates consistency of the estimators when both n and N tend to infinity. The bias of method 2 estimators is much less important than with method 1, especially for $n = 500$ but the precision is generally lower (bigger standard deviation). As for method 1, we observe the convergence of the estimators to the true values when both N and n increase. For both methods, the precision of the estimators improves when N becomes larger. Contrary to method 1, the precision of the method 2 estimators does not depend on n . Method 1 is numerically more difficult to implement, this may explain why the standard deviations slightly increase with n . Finally, the implementation of the method 2 in *Examples 2, 3, 4* requires to choose a value for the threshold k . The results are displayed for $k = 0.5$. Simulations with various values of k have not shown any significant impact of k on the estimators performances.

5 Extensions and concluding remarks.

In this paper, we study the estimation of population parameters in a SDE with a linear random effect in the diffusion coefficient from discrete observations of N *i.i.d.* trajectories on a fixed length time interval (1). We especially study the case of a null drift and of $\phi_i = 1/\Gamma_i^{1/2}$ with $\Gamma_i \sim G(a, \lambda)$. This leads to estimators using two different approaches. The first method is based on an approximation of the exact likelihood relying on the Euler scheme of the SDE.

	$N = 50$			$N = 100$		
	$n = 500$	$n = 1000$	$n = 10000$	$n = 500$	$n = 1000$	$n = 10000$
	$(a_0 = 5, \lambda_0 = 3)$					
\tilde{a}	4.52 (0.69)	4.89 (0.85)	5.33 (1.14)	4.38 (0.51)	4.70 (0.60)	5.07 (0.75)
$\tilde{\lambda}$	2.75 (0.43)	2.97 (0.52)	3.23 (0.70)	2.65 (0.34)	2.84 (0.40)	3.06 (0.49)
a^{**}	5.24 (1.15)	5.33 (1.15)	5.38 (1.19)	4.99 (0.78)	5.05 (0.77)	5.12 (0.78)
λ^{**}	3.18 (0.70)	3.23 (0.71)	3.26 (0.73)	3.01 (0.51)	3.05 (0.50)	3.09 (0.51)
	$(a_0 = 6, \lambda_0 = 1)$					
\tilde{a}	5.10 (0.69)	5.57 (0.87)	6.23 (1.19)	4.98 (0.47)	5.43 (0.59)	6.03 (0.81)
$\tilde{\lambda}$	0.86 (0.12)	0.93 (0.15)	1.04 (0.20)	0.84 (0.09)	0.91 (0.10)	1.01 (0.14)
a^{**}	6.19 (1.24)	6.25 (1.25)	6.32 (1.24)	5.94 (0.81)	6.02 (0.82)	6.10 (0.84)
λ^{**}	1.03 (0.21)	1.05 (0.21)	1.06 (0.21)	0.99 (0.14)	1.01 (0.14)	1.02 (0.15)

Table 2: Example 2: $\sigma^2(x) = 1 + x^2$. Empirical mean and standard deviation (in brackets) of $\tilde{\theta}$ (method 1) and θ^{**} (method 2) computed from 100 datasets.

The second method uses a plug-in of estimators of the random effects in the likelihood of (ϕ_1, \dots, ϕ_N) .

Several extensions are possible. First, assumption (H1) on the function $\sigma(x)$ can be weakened. For instance, it is enough to assume that $\sigma, \sigma', \sigma''$ have linear growth (instead of bounded). But, this would lengthen considerably proofs. Second, we can consider a SDE with drift (known or unknown): $dX_i(t) = b(X_i(t))dt + \phi_i \sigma(X_i(t))dW_i(t)$, and use exactly the same estimators, ignoring the drift. This was done in Genon-Catalot and Jacod (1993) in the case of non random effects. Here again, this would considerably lengthen proofs, as we would have to deal with error terms, including the drift. In simulations, models with drifts have been implemented and the results are quite satisfactory.

Another direction for extensions is to look at other distributions for the random effects. In particular, the plug-in method applies for any distribution provided that we introduce appropriate truncations as is done here.

For a more general model for the diffusion coefficient including a non linear random effect, the two approaches studied here could be extended, in particular the second method.

References

Barndorff-Nielsen, O. and Sorensen, M. (1991). Information quantities in non-classical settings. *Computational Statistics & Data Analysis*, **12**, 143–158.

	$N = 50$			$N = 100$		
	$n = 500$	$n = 1000$	$n = 10000$	$n = 500$	$n = 1000$	$n = 10000$
	$(a_0 = 5, \lambda_0 = 3)$					
\tilde{a}	4.50 (0.63)	4.86 (0.81)	5.26 (1.02)	4.43 (0.47)	4.75 (0.57)	5.12 (0.72)
$\tilde{\lambda}$	2.68 (0.41)	2.89 (0.51)	3.14 (0.64)	2.64 (0.31)	2.85 (0.37)	3.08 (0.46)
a^{**}	5.20 (1.05)	5.28 (1.09)	5.31 (1.06)	5.06 (0.73)	5.12 (0.74)	5.16 (0.75)
λ^{**}	3.08 (0.65)	3.14 (0.67)	3.17 (0.66)	3.01 (0.46)	3.06 (0.47)	3.10 (0.48)
	$(a_0 = 6, \lambda_0 = 1)$					
\tilde{a}	5.14 (0.64)	5.64 (0.81)	6.32 (1.13)	5.06 (0.53)	5.51 (0.68)	6.16 (0.92)
$\tilde{\lambda}$	0.85 (0.11)	0.93 (0.14)	1.05 (0.19)	0.84 (0.09)	0.92 (0.12)	1.03 (0.16)
a^{**}	6.26 (1.18)	6.34 (1.18)	6.41 (1.19)	6.08 (0.94)	6.14 (0.95)	6.24 (0.96)
λ^{**}	1.03 (0.20)	1.05 (0.20)	1.06 (0.20)	1.00 (0.16)	1.02 (0.16)	1.04 (0.16)

Table 3: Example 3 ($\rho = 1$): Empirical mean and standard deviation (in brackets) of $\tilde{\theta}$ (method 1) and θ^{**} (method 2) computed from 100 datasets.

Delattre, M. and Lavielle, M. (2013). Coupling the SAEM algorithm and the extended kalman filter for maximum likelihood estimation in mixed-effects diffusion models. *Statistics and Its Interface*, **6**, 519–532.

Delattre, M., Genon-Catalot, V., and Samson, A. (2013). Maximum likelihood estimation for stochastic differential equations with random effects. *Scandinavian Journal of Statistics*, **40**, 322–343.

Donnet, S. and Samson, A. (2008). Parametric inference for mixed models defined by stochastic differential equations. *ESAIM P&S*, **12**, 196–218.

Donnet, S. and Samson, A. (2013). A review on estimation of stochastic differential equations for pharmacokinetic - pharmacodynamic models. *Advanced Drug Delivery Reviews*, **65**, 929D939.

Genon-Catalot, V. and Jacod, J. (1993). On the estimation of the diffusion coefficient for multi-dimensional diffusion processes. *Ann. Inst. H. Poincaré Probab. Statist.*, **29**(1), 119–151.

Genon-Catalot, V., Jeantheau, T., and Laredo, C. (1999). Parameter estimation for discretely observed stochastic volatility models. *Bernoulli*, **5**, 855–872.

Kessler, M., Lindner, A., and Sorensen, M., editors (2012). *Statistical Methods for Stochastic Differential Equations*. CRC Press.

	$N = 50$			$N = 100$		
	$n = 500$	$n = 1000$	$n = 10000$	$n = 500$	$n = 1000$	$n = 10000$
	$(a_0 = 5, \lambda_0 = 3)$					
\tilde{a}	4.48 (0.67)	4.82 (0.83)	5.24 (1.09)	4.30 (0.42)	4.59 (0.49)	4.92 (0.61)
$\tilde{\lambda}$	2.67 (0.46)	2.88 (0.56)	3.15 (0.72)	2.56 (0.29)	2.74 (0.32)	2.94 (0.39)
a^{**}	5.17 (1.07)	5.23 (1.10)	5.29 (1.13)	4.85 (0.63)	4.91 (0.62)	4.95 (0.63)
λ^{**}	3.07 (0.70)	3.13 (0.72)	3.18 (0.74)	2.88 (0.40)	2.92 (0.39)	2.96 (0.40)
	$(a_0 = 6, \lambda_0 = 1)$					
\tilde{a}	5.08 (0.72)	5.57 (0.92)	6.25 (1.29)	5.06 (0.42)	5.53 (0.54)	6.13 (0.75)
$\tilde{\lambda}$	0.85 (0.14)	0.93 (0.17)	1.05 (0.23)	0.85 (0.08)	0.93 (0.10)	1.03 (0.13)
a^{**}	6.17 (1.32)	6.26 (1.34)	6.34 (1.36)	6.06 (0.76)	6.15 (0.77)	6.20 (0.78)
λ^{**}	1.03 (0.23)	1.05 (0.24)	1.06 (0.24)	1.01 (0.13)	1.03 (0.13)	1.04 (0.14)

Table 4: Example 4. ($\rho = 1$) Empirical mean and standard deviation (in brackets) of $\tilde{\theta}$ (method 1) and θ^{**} (method 2) computed from 100 datasets.

Nie, L. (2006). Strong consistency of the maximum likelihood estimator in generalized linear and nonlinear mixed-effects models. *Metrika*, **63**(2), 123–143.

Picchini, U. and Ditlevsen, S. (2011). Practicle estimation of high dimensional stochastic differential mixed-effects models. *Computational Statistics & Data Analysis*, **55**, 1426–1444.

Picchini, U., De Gaetano, A., and Ditlevsen, S. (2010). Stochastic differential mixed-effects models. *Scandinavian Journal of Statistics*, **37**, 67–90.

Sweeting, T. (1980). Uniform asymptotic normality of the maximum likelihood estimator. *Annals of Statistics*, **8**, 1179–1403.

6 Appendix: proofs

Proof of Proposition 1 Using the fact that ϕ_i^{-2} has Gamma distribution $G(a, \lambda)$, we get the result as:

$$\tilde{\lambda}_i(\theta, X_i) = \int_{(0, +\infty)} \frac{\lambda^a \gamma^{a-1+n/2}}{\Gamma(a)} \exp[-\gamma(\lambda + \frac{1}{2}S_i)] d\gamma = \frac{\lambda^a \Gamma(a + n/2)}{\Gamma(a) (\lambda + \frac{1}{2}S_i)^{a+n/2}}. \square$$

Proof of Proposition 2 Let $\chi_i = \sum_{j=1}^n (W_i(t_{i,j}) - W_i(t_{i,j-1}))^2 / \Delta_i = R_i^1 / \Delta_i$. As

$S_i^1 = \Gamma_i^{-1} \chi_i$, (see 14), $\beta_i^1(\lambda) = \frac{\lambda \Gamma_i}{\lambda \Gamma_i + \frac{1}{2} \chi_i}$, $\frac{S_i^1}{2\lambda} = \frac{\chi_i/2}{\lambda \Gamma_i}$ Under P_θ , Γ_i and χ_i are independent, Γ_i is $G(a, \lambda)$ and χ_i is $\chi^2(n) = G(n/2, 1/2)$. Hence the results using Proposition 10. \square

Proof of Proposition 3 The proof is elementary using Propositions 2 and 10. \square

Proof of Proposition 4 We need the following Lemma and Proposition:

Lemma 1. For all θ , $\mathbb{E}_\theta((X_i(t) - X_i(s))^{2p} | \phi_i) \leq C(2p) \sigma_1^{2p} \phi_i^{2p} |t - s|^p$ where $C(2p)$ is a numerical constant.

Proposition 9. $(R_i^1/T_i) - 1 = T_i^{-1} \int_0^{T_i} H_{i,1}^n(s) dW_i(s)$ and

$$(R_i/T_i) - 1 = T_i^{-1} \left(\int_0^{T_i} H_i^n(s) dW_i(s) + \int_0^{T_i} K_i^n(s) dW_i(s) + \int_0^{T_i} L_i^n(s) ds \right),$$

where, for $j = 1, \dots, n$ and $s \in]t_{i,j-1}, t_{i,j}]$, $H_{i,1}^n(s) = 2(W_i(s) - W_i(t_{i,j-1}))$,

$$H_i^n(s) = 2 \frac{(M_i(s) - M_i(t_{i,j-1})) \sigma(X_i(s))}{\sigma^2(X_i(t_{i,j-1}))}, \quad K_i^n(s) = 2\phi_i(t_{i,j} - s) \frac{\sigma^2(X_i(s)) \sigma'(X_i(s))}{\sigma^2(X_i(t_{i,j-1}))},$$

$$L_i^n(s) = \phi_i^2 \kappa(X_i(s)) (t_{i,j} - s) \frac{\sigma^2(X_i(s))}{\sigma^2(X_i(t_{i,j-1}))}, \quad \text{with } \kappa = \sigma \sigma'' + (\sigma')^2.$$

Lemma 1 and Proposition 9 yield $\mathbb{E}_\theta(\frac{R_i^1}{T_i} - 1 | \mathcal{F}_0) = 0$ and

$$\mathbb{E}_\theta\left(\frac{R_i}{T_i} - 1 | \mathcal{F}_0\right) = \mathbb{E}_\theta\left(\frac{R_i}{T_i} - \frac{R_i^1}{T_i} | \mathcal{F}_0\right) = \frac{1}{T_i} \int_0^{T_i} \mathbb{E}_\theta(L_i^n(s) | \mathcal{F}_0) ds.$$

Using (H1), we get $|L_i^n(s)| \leq C \phi_i^2 \sum_{j=1}^n \mathbb{1}_{]t_{i,j-1}, t_{i,j}]}(s) (t_{i,j} - s) \leq C \phi_i^2 \Delta_i \mathbb{1}_{]0, T_i]}(s)$, for C depending on σ_0, σ_1, K . Thus, the first inequality of Proposition 4.

As $(\frac{R_i}{T_i} - 1)^{2p} = (A_1 + A_2 + A_3)^{2p} \leq 3^{2p-1} \sum_{i=1}^3 A_i^{2p}$, we study separately the three terms A_i^{2p} . We have $A_3^{2p} = \left(\frac{1}{T_i} \int_0^{T_i} L_i^n(s) ds\right)^{2p} \leq (C \phi_i^2 \Delta_i)^{2p}$. Next, we use the Burkholder-Davies-Gundy (BDG), the Hölder inequalities and (H1):

$$T_i^{2p} \mathbb{E}_\theta(A_2^{2p} | \mathcal{F}_0) \leq C(2p) \mathbb{E}_\theta\left(\left(\int_0^{T_i} (K_i^n(s))^2 ds\right)^p | \mathcal{F}_0\right) \leq C(2p) T_i^{p-1} \mathbb{E}_\theta\left(\int_0^{T_i} \mathbb{E}_\theta((K_i^n(s))^{2p} | \mathcal{F}_0) ds\right)$$

where $(K_i^n(s))^{2p} \leq C \phi_i^{2p} \Delta_i^{2p} \mathbb{1}_{]0, T_i]}(s)$. Finally, for $T_i^{2p} \mathbb{E}_\theta(A_1^{2p} | \mathcal{F}_0)$ we study

$\left(\int_0^{T_i} (H_i^n(s))^2 ds\right)^p$. By the Holder inequality, we have,

$$\left(\int_0^{T_i} (H_i^n(s))^2 ds\right)^p \leq C^p n^{p-1} \sum_{j=1}^n \Delta_i^{p-1} \int_{t_{i,j-1}}^{t_{i,j}} (M_i(s) - M_i(t_{i,j-1}))^{2p} ds.$$

Consequently, for constants C depending on σ_0, σ_1, K ,

$$\mathbb{E}_\theta\left(\left(\int_0^{T_i} (H_i^n(s))^2 ds\right)^p \mid \mathcal{F}_0\right) \leq C \sum_{j=1}^n \int_{t_{i,j-1}}^{t_{i,j}} \mathbb{E}_\theta\left(\left(\int_{t_{i,j-1}}^s \sigma^2(X_i(u)) du\right)^p \mid \mathcal{F}_0\right) ds \leq C \Delta_i^p.$$

Finally, to study the difference $R_i - R_i^1$, we only need to study the term: $\int_0^{T_i} (H_i^n(s) - H_{i,1}^n(s)) dW_i(s) = \int_0^{T_i} 2 \sum_{k=1}^3 \sum_{j=1}^n \mathbf{1}_{]t_{i,j-1}, t_{i,j}[}(s) Z_{i,j}^k(s) dW_i(s)$ with

$$\begin{aligned} Z_{i,j}^1(s) &= \frac{\sigma(X_i(s)) - \sigma(X_i(t_{i,j-1}))}{\sigma^2(X_i(t_{i,j-1}))} \int_{t_{i,j-1}}^s (\sigma(X_i(u)) - \sigma(X_i(t_{i,j-1}))) dW_i(u), \\ Z_{i,j}^2(s) &= \frac{\sigma(X_i(s)) - \sigma(X_i(t_{i,j-1}))}{\sigma(X_i(t_{i,j-1}))} \int_{t_{i,j-1}}^s dW_i(u), \\ Z_{i,j}^3(s) &= \frac{1}{\sigma(X_i(t_{i,j-1}))} \int_{t_{i,j-1}}^s (\sigma(X_i(u)) - \sigma(X_i(t_{i,j-1}))) dW_i(u), \end{aligned}$$

These terms are studied analogously using the BDG and Cauchy-Schwarz inequalities. \square

Proof of Lemma 1 Recall that ϕ_i is \mathcal{F}_0 -measurable and when dealing with the process $(X_i(t))$, conditioning on ϕ_i is equal to conditioning on \mathcal{F}_0 . We have $\mathbb{E}_\theta((X_i(t) - X_i(s))^{2p} \mid \mathcal{F}_0) = \phi_i^{2p} \mathbb{E}_\theta((M_i(t) - M_i(s))^{2p} \mid \mathcal{F}_0)$. By the BDG inequality and (H1), for $s \leq t$,

$$\mathbb{E}_\theta((M_i(t) - M_i(s))^{2p} \mid \mathcal{F}_0) \leq C(2p) \mathbb{E}\left(\left(\int_s^t \sigma^2(X_i(u)) du\right)^p \mid \mathcal{F}_0\right) \leq C(2p) \sigma_1^{2p} (t-s)^{2p}. \square$$

Proof of Proposition 9 By the Ito formula, we have:

$$(M_i(t_{i,j}) - M_i(t_{i,j-1}))^2 = 2 \int_{t_{i,j-1}}^{t_{i,j}} (M_i(s) - M_i(t_{i,j-1})) \sigma(X_i(s)) dW_i(s) + \int_{t_{i,j-1}}^{t_{i,j}} \sigma^2(X_i(s)) ds.$$

We split: $\sigma^2(X_i(s)) = \sigma^2(X_i(t_{i,j-1})) + \sigma^2(X_i(s)) - \sigma^2(X_i(t_{i,j-1}))$ and use the Ito formula: $\sigma^2(X_i(s)) - \sigma^2(X_i(t_{i,j-1})) = \phi_i \int_{t_{i,j-1}}^s (\sigma^2)'(X_i(u)) \sigma(X_i(u)) dW_i(u) +$

$\frac{1}{2}\phi_i^2 \int_{t_{i,j-1}}^s (\sigma^2)''(X_i(u))\sigma^2(X_i(u))du$. Integrating, Fubini formula yields:

$$\int_{t_{i,j-1}}^{t_{i,j}} (\sigma^2(X_i(s)) - \sigma^2(X_i(t_{i,j-1})))ds = \phi_i \int_{t_{i,j-1}}^{t_{i,j}} (t_{i,j} - u)(\sigma^2)'(X_i(u))\sigma(X_i(u))dW_i(u) \\ + \frac{1}{2}\phi_i^2 \int_{t_{i,j-1}}^{t_{i,j}} (t_{i,j} - u)(\sigma^2)''(X_i(u))\sigma^2(X_i(u))du.$$

The result for R_i^1 corresponds to $\sigma(\cdot) \equiv 1$. Hence the results. \square

Proof of Proposition 5 Recall that $S_i = n\Gamma_i^{-1}R_i/T_i$. We have by (11) $G_N(\theta_0) = S_N(\theta_0) + \left(\sum_{i=1}^N Y_i(\theta_0) \quad \sum_{i=1}^N Z_i(\theta_0) \right)'$ where $Y_i(\theta_0) = \Gamma_i - \frac{(a_0 + \frac{n}{2})\Gamma_i}{\lambda_0\Gamma_i + C_i}$ and $Z_i(\theta_0) = \psi(a_0 + \frac{n}{2}) - \log(\lambda_0\Gamma_i + C_i)$.

Therefore, we have to prove that $\frac{1}{\sqrt{N}} \left(\sum_{i=1}^N Y_i(\theta_0) \quad \sum_{i=1}^N Z_i(\theta_0) \right)'$ tends to 0 in \mathbb{P}_{θ_0} -probability as n, N tend to infinity. To distinguish the two cases $\sigma(\cdot) \equiv 1$ and $\sigma(\cdot) \neq 1$, we introduce the random variables $Y_i^1(\theta_0), Z_i^1(\theta_0)$ where we replaced S_i by S_i^1 . We proceed on two steps:

- (1) $\frac{1}{\sqrt{N}} \left(\sum_{i=1}^N Y_i^1(\theta_0) \quad \sum_{i=1}^N Z_i^1(\theta_0) \right)' = o_{\mathbb{P}_{\theta_0}}(1)$ as $N, n \rightarrow \infty$.
- (2) $\frac{1}{\sqrt{N}} \left(\sum_{i=1}^N (Y_i(\theta_0) - Y_i^1(\theta_0)) \quad \sum_{i=1}^N (Z_i(\theta_0) - Z_i^1(\theta_0)) \right)' = o_{\mathbb{P}_{\theta_0}}(1)$ as $N, n \rightarrow \infty$ under the constraints $N/n \rightarrow 0$ and $\mathbb{E}_{\theta_0}\phi_i^8 < +\infty$.

Proof of (1): Let $C_i^1 = nR_i^1/(2T_i) = \Gamma_i S_i^1/2$ and $G_i^0 = \lambda_0\Gamma_i$. We have $Y_i^1(\theta_0) = \Gamma_i - \frac{a_0 + n/2}{\lambda_0} \frac{G_i^0}{G_i^0 + C_i^1}$. In what follows, we use repeatedly the fact that G_i^0 and C_i^1 are independent, that $G_i^0 \sim G(a_0, 1)$ and $C_i^1 \sim G(n/2, 1)$. Hence, $G_i^0 + C_i^1$ and $G_i^0/(G_i^0 + C_i^1)$ are independent, the latter with distribution $\beta^{(1)}(a_0, (n/2))$, the former with distribution $G(a_0 + (n/2), 1)$.

The r.v. $Y_i^1(\theta_0), i = 1, \dots, N$ are *i.i.d.* with $\mathbb{E}_{\theta_0}(Y_i^1(\theta_0)) = 0$, $\mathbb{E}_{\theta_0}(Y_i^1(\theta_0)^2) = \frac{a_0(a_0+1)}{\lambda_0^2(a_0+1+n/2)}$. Therefore, $N^{-1/2} \sum_{i=1}^N Y_i^1(\theta_0) = o_{\mathbb{P}_{\theta_0}}(1)$.

Analogously, $Z_i^1(\theta_0) = \psi(a_0 + n/2) - \log(G_i^0 + C_i^1)$ satisfies $\mathbb{E}_{\theta_0}(Z_i^1(\theta_0)) = 0$, $\mathbb{E}_{\theta_0}(Z_i^1(\theta_0))^2 = 1/(a_0 + n/2) + o(1/n)$. Thus, $N^{-1/2} \sum_{i=1}^N Z_i^1(\theta_0) = o_{\mathbb{P}_{\theta_0}}(1)$.

Proof of (2): We introduce $C_i = nR_i/(2T_i)$. We have $Y_i(\theta_0) - Y_i^1(\theta_0) = \frac{(a_0 + \frac{n}{2})\Gamma_i}{G_i^0 + C_i^1} - \frac{(a_0 + \frac{n}{2})\Gamma_i}{G_i^0 + C_i}$, and $Z_i(\theta_0) - Z_i^1(\theta_0) = \log(G_i^0 + C_i^1) - \log(G_i^0 + C_i)$. Thus,

$$Y_i(\theta_0) - Y_i^1(\theta_0) = \frac{a_0 + \frac{n}{2}}{\lambda_0} \frac{G_i^0}{(G_i^0 + C_i)(G_i^0 + C_i^1)} \frac{n}{2} \left(\frac{R_i}{T_i} - \frac{R_i^1}{T_i} \right)$$

We introduce the set $\Omega_i = \{|(R_i/T_i) - 1| \leq 1/2\}$. On Ω_i , we use $G_i^0 + C_i \geq (n/4)$.

So

$$\mathbb{E}_{\theta_0} |Y_i(\theta_0) - Y_i^1(\theta_0)| \mathbf{1}_{\Omega_i} \leq 2 \frac{a_0 + \frac{n}{2}}{\lambda_0} \mathbb{E}_{\theta_0} \left(\frac{G_i^0}{G_i^0 + C_i^1} \left| \frac{R_i}{T_i} - \frac{R_i^1}{T_i} \right| \right).$$

Then, if $\mathbb{E}_{\theta_0} \phi_i^4 < +\infty$, we have by proposition 4

$$\mathbb{E}_{\theta_0} \left(\frac{G_i^0}{G_i^0 + C_i^1} \right)^2 = O\left(\frac{1}{n^2}\right) \quad \text{and} \quad \mathbb{E}_{\theta_0} \left(\frac{R_i}{T_i} - \frac{R_i^1}{T_i} \right)^2 \leq (T_i/n)^2 \mathbb{E}_{\theta_0} \phi_i^4.$$

Therefore, using the Cauchy-Schwarz inequality, $\mathbb{E}_{\theta_0} |Y_i(\theta_0) - Y_i^1(\theta_0)| \mathbf{1}_{\Omega_i} \leq Cn \left(\frac{C}{n^4}\right)^{1/2} \leq \frac{C}{n}$.

On Ω_i^c , we use $G_i^0/(G_i^0 + C_i) \leq 1$. Therefore,

$$|Y_i(\theta_0) - Y_i^1(\theta_0)| \mathbf{1}_{\Omega_i^c} \leq \frac{\frac{n}{2}(a_0 + \frac{n}{2})}{\lambda_0} \frac{1}{G_i^0 + C_i^1} \left| \frac{R_i}{T_i} - \frac{R_i^1}{T_i} \right| \mathbf{1}_{\Omega_i^c}.$$

We have if $\mathbb{E}_{\theta_0} \phi_i^8 < +\infty$, by Proposition 4, $\mathbb{E}_{\theta_0} \left(\frac{1}{G_i^0 + C_i^1} \right)^4 = O\left(\frac{1}{n^4}\right)$, $\mathbb{E}_{\theta_0} \left(\frac{R_i}{T_i} - \frac{R_i^1}{T_i} \right)^4 \leq (T_i/n)^4 \mathbb{E}_{\theta_0} (\phi_i^4 + \phi_i^8)$, and $\mathbb{P}_{\theta_0}(\Omega_i^c) \leq 2^{2p} \mathbb{E}_{\theta_0} \left| \frac{R_i}{T_i} - 1 \right|^{2p}$. Using the Cauchy-Schwarz inequality twice, the above inequality with $p = 2$ and Proposition 4 with the condition $\mathbb{E}_{\theta_0} \phi_i^8 < +\infty$, we get:

$$\mathbb{E}_{\theta_0} |Y_i(\theta_0) - Y_i^1(\theta_0)| \mathbf{1}_{\Omega_i^c} \leq Cn^2 (C/n^4)^{1/4} (C/n^4)^{1/4} (\mathbb{P}_{\theta_0}(\Omega_i^c))^{1/2} \leq C/n.$$

We can conclude that under the condition $\mathbb{E}_{\theta_0} \phi_i^8 < +\infty$,

$$\mathbb{E}_{\theta_0} \left| \frac{1}{\sqrt{N}} \sum_{i=1}^N (Y_i(\theta_0) - Y_i^1(\theta_0)) \right| \leq C\sqrt{N}/n. \quad (19)$$

We now turn to the other difference. We have by the Taylor formula

$$Z_i(\theta_0) - Z_i^1(\theta_0) = \frac{n}{2} \frac{1}{G_i^0 + n/2} \left(\frac{R_i^1}{T_i} - \frac{R_i}{T_i} \right) + \frac{n}{2} \left(\frac{R_i^1}{T_i} - \frac{R_i}{T_i} \right) \int_0^1 f_i(s) ds,$$

where $f_i(s) = -\frac{s \frac{n}{2} \left(\frac{R_i^1}{T_i} - 1 \right) + (1-s) \frac{n}{2} \left(\frac{R_i}{T_i} - 1 \right)}{(G_i^0 + n/2) \left(G_i^0 + s \frac{n}{2} \frac{R_i^1}{T_i} + (1-s) \frac{n}{2} \frac{R_i}{T_i} \right)}$. We use that:

$$\left| \mathbb{E}_{\theta_0} \frac{n}{2} \frac{1}{G_i^0 + n/2} \left(\frac{R_i^1}{T_i} - \frac{R_i}{T_i} \right) \middle| \mathcal{F}_0 \right| \leq \frac{nC}{2} \frac{1}{G_i^0 + n/2} \phi_i^2 = C \frac{1}{\Gamma_i^2(G_i^0 + n/2)}$$

and if $\mathbb{E}_{\theta_0} \phi_i^4 < +\infty$, $\mathbb{E}_{\theta_0} \frac{1}{\Gamma_i^2(G_i^0 + n/2)} \leq C \left(\frac{1}{(a_0 + n/2 - 1)(a_0 + n/2 - 2)} \right)^{1/2}$, to obtain

that ,

$$\frac{1}{\sqrt{N}} \sum_{i=1}^N \mathbb{E}_{\theta_0} \left| \mathbb{E}_{\theta_0} \frac{n}{2} \frac{1}{G_i^0 + n/2} \left(\frac{R_i^1}{T_i} - \frac{R_i}{T_i} \right) | \mathcal{F}_0 \right| \leq \frac{C\sqrt{N}}{n}$$

On the other hand, noticing that for $s \in [0, 1]$: $|f_i(s)| \leq \frac{n/2}{(G_i^0 + n/2)G_i^0} \left(\left| \frac{R_i^1}{T_i} - 1 \right| + \left| \frac{R_i}{T_i} - 1 \right| \right)$, if $\mathbb{E}_{\theta_0} \phi_i^8 < +\infty$,

$$\mathbb{E}_{\theta_0} \left(\int_0^1 f_i(s) ds \right)^2 \leq \left(\mathbb{E}_{\theta_0} \frac{1}{(G_i^0)^4} \mathbb{E}_{\theta_0} \left(\left| \frac{R_i^1}{T_i} - 1 \right|^4 + \left| \frac{R_i}{T_i} - 1 \right|^4 \right) \right)^{1/2} \leq C \left(\frac{T_i}{n} \right).$$

Finally, we have $\mathbb{E}_{\theta_0} \frac{n}{2} \left(\frac{R_i^1}{T_i} - \frac{R_i}{T_i} \right) \int_0^1 f_i(s) ds \leq \left(\frac{C}{n} \right)^{1/2}$. Therefore, $\mathbb{E}_{\theta_0} \left| \frac{1}{\sqrt{N}} \sum_{i=1}^N (Z_i(\theta_0) - Z_i^1(\theta_0)) \right| \leq C\sqrt{N}(1/n + 1/n^{1/2})$. \square

Proof of Proposition 6: To obtain $\tilde{\mathcal{I}}_N(\theta_0)/N = \mathcal{I}(\theta_0) + o_{\mathbb{P}_{\theta_0}}(1)$, we have to prove that $\frac{1}{N} \sum_{i=1}^N A_i(\theta_0) \rightarrow 0$, and $\frac{1}{N} \sum_{i=1}^N B_i(\theta_0) \rightarrow 0$, in \mathbb{P}_{θ_0} - probability where

$$A_i(\theta_0) = \frac{a_0 + n/2}{(\lambda_0 + S_i/2)^2} = \frac{a_0 + n/2}{\lambda_0^2} \frac{(G_i^0)^2}{(G_i^0 + C_i)^2}, \quad B_i(\theta_0) = \frac{1}{\lambda_0 + S_i/2} = \frac{\Gamma_i}{G_i^0 + C_i}. \quad (20)$$

As in the previous proposition, we separate the cases $\sigma(\cdot) \equiv 1$ and $\sigma(\cdot) \neq 1$ and define the random variables $A_i^1(\theta_0), B_i^1(\theta_0)$ where S_i is replaced by S_i^1 :

$$A_i^1(\theta_0) = \frac{a_0 + n/2}{\lambda_0^2} \frac{(G_i^0)^2}{(G_i^0 + C_i^1)^2}, \quad B_i^1(\theta_0) = \frac{\Gamma_i}{G_i^0 + C_i^1}.$$

Recall that $C_i^1 \sim G(n/2, 1)$ and is independent of Γ_i . Thus,

$$\mathbb{E}_{\theta_0} A_i^1(\theta_0) = \frac{a_0(a_0 + 1)}{\lambda_0^2(a_0 + n/2 + 1)} = O\left(\frac{1}{n}\right), \quad \mathbb{E}_{\theta_0} B_i^1(\theta_0) = \frac{a_0}{\lambda_0(a_0 + n/2)} = O\left(\frac{1}{n}\right).$$

This implies $\frac{1}{N} \sum_{i=1}^N \mathbb{E}_{\theta_0} A_i^1(\theta_0) = O\left(\frac{1}{n}\right)$, $\frac{1}{N} \sum_{i=1}^N \mathbb{E}_{\theta_0} B_i^1(\theta_0) = O\left(\frac{1}{n}\right)$. Next, we study the differences $A_i(\theta_0) - A_i^1(\theta_0), B_i(\theta_0) - B_i^1(\theta_0)$.

$$A_i(\theta_0) - A_i^1(\theta_0) = \frac{a_0 + n/2}{\lambda_0^2} (C_i - C_i^1) \left(\frac{(G_i^0)^2}{(G_i^0 + C_i)(G_i^0 + C_i^1)^2} + \frac{(G_i^0)^2}{(G_i^0 + C_i)^2(G_i^0 + C_i^1)} \right)$$

Thus: $|A_i(\theta_0) - A_i^1(\theta_0)| \leq \frac{a_0 + n/2}{\lambda_0^2} |C_i^1 - C_i| \frac{2G_i^0}{(G_i^0 + C_i)(G_i^0 + C_i^1)}$. We introduce again the set $\Omega_i = \{ |(R_i/T_i) - 1| \leq 1/2 \}$. On Ω_i , using $G_i^0 + C_i \geq (n/4)$, we have

$$\mathbb{E}_{\theta_0} |A_i(\theta_0) - A_i^1(\theta_0)| \mathbf{1}_{\Omega_i} \leq 2 \frac{a_0 + n/2}{\lambda_0^2(n/4)} \left[\mathbb{E}_{\theta_0} \left(\frac{G_i^0}{G_i^0 + C_i^1} \right)^2 \mathbb{E}_{\theta_0} (C_i^1 - C_i)^2 \right]^{1/2} \leq C/n.$$

Next, using $G_i^0/(G_i^0+C_i) \leq 1$ on Ω_i^c , we have $|A_i(\theta_0) - A_i^1(\theta_0)|\mathbb{1}_{\Omega_i^c} \leq \frac{a_0+n/2}{\lambda_0^2}|C_i^1 - C_i| \frac{2}{(G_i^0+C_i^1)}$. Thus, using the same arguments than in proof of proposition 5

$$\mathbb{E}_{\theta_0}(|A_i(\theta_0) - A_i^1(\theta_0)|\mathbb{1}_{\Omega_i^c}) \leq C (\mathbb{P}_{\theta_0}(\Omega_i^c))^{1/2} \leq \frac{C}{n}.$$

Analogously, $B_i(\theta_0) - B_i^1(\theta_0) = \frac{1}{\lambda_0}(C_i^1 - C_i) \frac{G_i^0}{(G_i^0+C_i)(G_i^0+C_i^1)}$. Introducing the set Ω_i again, we obtain $|B_i(\theta_0) - B_i^1(\theta_0)| \leq C \left(\frac{G_i^0}{G_i^0+C_i^1} + \mathbb{1}_{\Omega_i^c} \frac{n}{G_i^0+C_i^1} \right) |R_i/T_i - R_i^1/T_i^1|$. We conclude $\mathbb{E}_{\theta_0}|B_i(\theta_0) - B_i^1(\theta_0)| \leq \frac{C}{n} + (\mathbb{P}_{\theta_0}(\Omega_i^c))^{1/2}$. So the proof is complete. \square

Proof of Proposition 7: To obtain the weak consistency of $\tilde{\theta}_N$ and its asymptotic normality, we follow the scheme described in Barndorff-Nielsen and Sorensen (1991) (Theorem 3.4 and Lemma 3.5) and Genon-Catalot *et al.* (1999) (Theorem 4.1), see also Sweeting (1980). We must prove that:

- (1) Under P_{θ_0} , $G_N(\theta_0)/\sqrt{N} \rightarrow_{\mathcal{D}} \mathcal{N}_2(0, \mathcal{I}(\theta_0))$ as $N \rightarrow \infty$.
- (2) $\tilde{\mathcal{I}}_N(\theta_0)/N \rightarrow \mathcal{I}(\theta_0)$ in P_{θ_0} -probability.
- (3) $\sup_{\theta \in M_{c,N}} |\tilde{\mathcal{I}}_N(\theta)/N - \mathcal{I}(\theta_0)| \rightarrow 0$ in P_{θ_0} -probability, where $M_{c,N} = \{\theta \in (0, +\infty)^2, \|\theta - \theta_0\| \leq c/\sqrt{N}\}$. (Uniformity condition)

Points (1) and (2) are directly implied by Propositions 5 and 6. It remains to prove (3). We will prove

- (a) $\mathbb{E}_{\theta_0}(\sup_{\theta \in M_{c,N}} |\tilde{\mathcal{I}}_N^1(\theta)/N - \tilde{\mathcal{I}}_N^1(\theta_0)/N|) \rightarrow 0$.
- (b) $\mathbb{E}_{\theta_0}(\sup_{\theta \in M_{c,N}} |\tilde{\mathcal{I}}_N^1(\theta)/N - \tilde{\mathcal{I}}_N(\theta)/N|) \rightarrow 0$.

Point (a) Let $\varepsilon > 0$ be such that $a_0 - \varepsilon > 0, \lambda_0 - \varepsilon > 0$. Choose N large enough to ensure that $M_{c,N} = \{(a, \lambda) \in (0, +\infty)^2, |a - a_0| \leq c/\sqrt{N}, |\lambda - \lambda_0| \leq c/\sqrt{N}\} \subset [a_0 - \varepsilon, a_0 + c] \times [\lambda_0 - \varepsilon, \lambda_0 + c]$ and $n > 8$. We have

$$\tilde{\mathcal{I}}_N^1(\theta)/N - \tilde{\mathcal{I}}_N^1(\theta_0)/N = \mathcal{I}(\theta) - \mathcal{I}(\theta_0) + \begin{pmatrix} D_N^{11}(\theta, \theta_0) & D_N^{12}(\theta, \theta_0) \\ D_N^{12}(\theta, \theta_0) & D_N^{22}(\theta, \theta_0) \end{pmatrix}$$

where $D_N^{11}(\theta, \theta_0) = \frac{1}{N} \sum_{i=1}^N (A_i^1(\theta) - A_i^1(\theta_0))$, $D_N^{12}(\theta, \theta_0) = \frac{1}{N} \sum_{i=1}^N (B_i^1(\theta) - B_i^1(\theta_0))$, $D_N^{22}(\theta, \theta_0) = -(\psi'(a + n/2) - \psi'(a_0 + n/2))$. We only study D^{11} and D^{12} which are the most difficult. We can write $D_N^{11}(\theta, \theta_0) = c_N + d_N$, with

$G_i = \lambda \Gamma_i$ and

$$c_N = (\lambda - \lambda_0)(a_0 + n/2) \frac{1}{N} \sum_{i=1}^N \frac{\Gamma_i(G_i + G_i^0 + 2C_i^1)}{(G_i + C_i^1)^2 (G_i^0 + C_i^1)^2}, \quad d_N = (a_0 - a) \frac{1}{N} \sum_{i=1}^N \frac{\Gamma_i}{(G_i + C_i^1)^2}.$$

For $\theta \in M_{c,N}$, we have the bounds $|d_N| \leq \frac{c}{(\lambda_0 - \varepsilon)^2 \sqrt{N}}$, and

$$|c_N| \leq \frac{c(a_0 + n/2)}{\sqrt{N}} \left[(2\lambda_0 + c) \left(\frac{\Gamma_i}{C_i^1} \right)^4 + \left(2 \frac{\Gamma_i}{C_i^1} \right)^3 \right]$$

We have, for $n > 2k$, $\mathbb{E}_{\theta_0} \left(\frac{\Gamma_i}{2C_i^1} \right)^k = (2\lambda_0)^{-k} \frac{(a_0 + k - 1)(a_0 + k - 2) \dots a_0}{(n/2 - 1)(n/2 - 2) \dots (n/2 - k)}$.

Thus, $\mathbb{E}_{\theta_0} \sup_{\theta \in M_{c,N}} |c_N| \leq \frac{C}{\sqrt{N} n^2}$. Thus $\mathbb{E}_{\theta_0} \sup_{\theta \in M_{c,N}} |D_N^{11}(\theta, \theta_0)| = O(1/\sqrt{N})$.

For the other term, (a) is proved as we have

$$\sup_{\theta \in M_{c,N}} |D_N^{12}(\theta, \theta_0)| \leq |\lambda - \lambda_0| \frac{1}{\lambda_0(\lambda_0 - \varepsilon)} = O(1/\sqrt{N}).$$

We now prove (b). For this, we prove the convergence to 0 of (see (20):

$\mathbb{E}_{\theta_0} \sup_{\theta \in M_{c,N}} |A_i(\theta) - A_i^1(\theta)|$ and $\mathbb{E}_{\theta_0} \sup_{\theta \in M_{c,N}} |B_i(\theta) - B_i^1(\theta)|$. We have $|A_i(\theta) - A_i^1(\theta)| \leq \frac{a_0 + n/2 + c}{(\lambda_0 - \varepsilon)^2} |C_i^1 - C_i| \frac{2G_i}{(G_i + C_i)(G_i + C_i^1)}$. On the set $\Omega_i = \{|(R_i/T_i) - 1| \leq 1/2\}$, $R_i/T_i \geq 1/2$ and $G_i + C_i \geq n/4$. Thus, as $G_i + C_i^1 > C_i^1$ and G_i and C_i^1 are independent, for $n > 4$,

$$\begin{aligned} & \mathbb{E}_{\theta_0} \sup_{\theta \in M_{c,N}} |A_i(\theta) - A_i^1(\theta)| \mathbf{1}_{\Omega_i} \\ & \leq 2 \frac{a_0 + n/2 + c}{(\lambda_0 - \varepsilon)^2 (n/4)} (n/2) (\mathbb{E}_{\theta_0} (R_i^1/T_i - R_i/T_i)^2 \mathbb{E}_{\theta_0} ((\lambda_0 + c)\Gamma_i)^2 \mathbb{E}_{\theta_0} (1/C_i^1)^2)^{1/2} \\ & \leq C \frac{T_i}{n} (\mathbb{E}_{\theta_0} (\phi_i^2 + \phi_i^4))^{1/2} = O\left(\frac{1}{n}\right). \end{aligned}$$

Next ($n > 8$, $\mathbb{E}_{\theta_0} \phi_i^8 < +\infty$)

$$\begin{aligned} & \mathbb{E}_{\theta_0} \sup_{\theta \in M_{c,N}} |A_i(\theta) - A_i^1(\theta)| \mathbf{1}_{\Omega_i^c} \\ & \leq \frac{2(a_0 + n/2 + c)n/2}{(\lambda_0 - \varepsilon)^2} (\mathbb{P}_{\theta_0}(\Omega_i^c))^{1/2} \left(\mathbb{E}_{\theta_0} (R_i^1/T_i - R_i/T_i)^4 \mathbb{E}_{\theta_0} \frac{1}{(nR_i^1/T_i)^4} \right)^{1/4} = o(1) \end{aligned}$$

We have $B_i(\theta) - B_i^1(\theta) = \frac{\Gamma_i}{(\lambda\Gamma_i + \frac{nR_i^1}{2T_i})(\lambda\Gamma_i + \frac{nR_i^1}{2T_i})} (\frac{nR_i^1}{2T_i} - \frac{nR_i}{2T_i})$. Using that $\lambda \geq \lambda_0 - \varepsilon$,

$$\begin{aligned} \mathbb{E}_{\theta_0} \sup_{\theta \in M_{\varepsilon, N}} |B_i(\theta) - B_i^1(\theta)| &\leq C \frac{n}{2(\lambda_0 - \varepsilon)} \frac{T_i}{n} \left(\frac{1}{(n/2 - 1)(n/2 - 2)} \mathbb{E}_{\theta_0}(\phi_i^2 + \phi_i^4) \right)^{1/2} \\ &= O\left(\frac{1}{n}\right). \end{aligned}$$

Therefore, the proof of the first part of Proposition 7 is complete.

The fact that $\sqrt{N}(\tilde{\theta}_n - \theta_N) = o_{\mathbb{P}_{\theta_0}}(1)$ can be deduced from the above proof. \square

Proof of Proposition 8: We first consider the case $\sigma(\cdot) \equiv 1$ and the estimating function $\nabla U_N(\theta)$. To get the result, it is enough to prove that:

$$\frac{1}{\sqrt{N}} \sum_{i=1}^N \left(\frac{n}{S_i^1} - \Gamma_i \right) = o_{\mathbb{P}_{\theta_0}}(1), \quad \frac{1}{\sqrt{N}} \sum_{i=1}^N \left(\log \frac{n}{S_i^1} - \log \Gamma_i \right) = o_{\mathbb{P}_{\theta_0}}(1), \quad (21)$$

where we recall that $S_i^1/n = \Gamma_i^{-1} R_i^1/T_i$ and nR_i^1/T_i is independent of Γ_i and has distribution $\chi^2(n)$. Using results recalled in Section 7, for $n > 2$, we have $\mathbb{E}_{\theta_0}(\frac{n}{S_i^1} - \Gamma_i) = \mathbb{E}_{\theta_0} \Gamma_i \mathbb{E}_{\theta_0}(\frac{n/2}{C_i^1} - 1) = \frac{a_0}{\lambda_0} O(n^{-1})$. Analogously, for $n > 4$, $\mathbb{E}_{\theta_0}(\frac{n}{S_i^1} - \Gamma_i)^2 = \mathbb{E}_{\theta_0} \Gamma_i^2 \mathbb{E}_{\theta_0}(\frac{n/2}{C_i^1} - 1)^2 = O(n^{-1})$. This implies: $\mathbb{E}_{\theta_0}(\frac{1}{\sqrt{N}} \sum_{i=1}^N (\frac{n}{S_i^1} - \Gamma_i))^2 = O(n^{-1}) + \frac{N-1}{2} (O(n^{-1}))^2$.

Hence, the first part of (21) holds provided that $\sqrt{N}/n = o(1)$.

For the second assertion, we compute $\mathbb{E}_{\theta_0}(\log \frac{n}{S_i^1} - \log \Gamma_i) = -\psi(n/2) + \log(n/2) = O(n^{-1})$, and $\text{Var}_{\theta_0}(\log \frac{n}{S_i^1} - \log \Gamma_i) = \mathbb{E}_{\theta_0}(-\log C_i^1 + \psi(n/2))^2 = O(n^{-1})$. Therefore, the second part of (21) also holds for $\sqrt{N}/n = o(1)$.

Next, the result on θ_N^* will follow analogously from the fact that $T_N = (\sqrt{N})^{-1} \sum_{i=1}^N (\tilde{\log} \Gamma_i - \log \Gamma_i)$ and $\tau_N = (\sqrt{N})^{-1} \sum_{i=1}^N (\tilde{\Gamma}_i - \Gamma_i)$ both tend to 0 in probability. The result follows from the first part of the proof and the following lemma.

Lemma 2. *Assume that $\mathbb{E}_{\theta_0} \phi_i^8 < +\infty$, i.e. $a_0 > 4$ and that $n > 4$. Then,*

$$\mathbb{E}_{\theta_0} \left(\frac{n}{S_i} \mathbf{1}_{(S_i/n \geq k/\sqrt{n})} - \frac{n}{S_i^1} \right)^2 + \mathbb{E}_{\theta_0} \left(\log \frac{S_i}{n} \mathbf{1}_{(S_i/n \geq k/\sqrt{n})} - \log \frac{S_i^1}{n} \right)^2 \leq \frac{C}{n^2}$$

Proof of Lemma 2 We write: $\tilde{\Gamma}_i - \frac{n}{S_i^1} = \frac{n}{S_i} \mathbf{1}_{(S_i/n \geq k/\sqrt{n})} - \frac{n}{S_i^1} = \nu_i + \nu'_i$ with

$$\nu_i = \left(\frac{n}{S_i} - \frac{n}{S_i^1} \right) \mathbf{1}_{(S_i/n \geq k/\sqrt{n})}, \quad \nu'_i = -\frac{n}{S_i^1} \mathbf{1}_{(S_i/n < k/\sqrt{n})}. \quad (22)$$

And analogously $\log \frac{S_i}{n} \mathbb{1}_{(S_i/n \geq k/\sqrt{n})} - \log \frac{S_i^1}{n} = \tau_i + \tau'_i$ with

$$\tau_i = \left(\log \frac{S_i}{n} - \log \frac{S_i^1}{n} \right) \mathbb{1}_{(S_i/n \geq k/\sqrt{n})}, \quad \tau'_i = -\log \frac{S_i^1}{n} \mathbb{1}_{(S_i/n < k/\sqrt{n})}. \quad (23)$$

For $n > 4$ and $\mathbb{E}_{\theta_0} \phi_i^8 < +\infty$, using explicit computations, $\mathbb{E}_{\theta_0} \left(\frac{n}{S_i^1} \right)^2 = O(1)$, $\mathbb{E}_{\theta_0} \log^2 \frac{S_i^1}{n} = O(1)$. To obtain that $\mathbb{E}_{\theta_0} (\nu'_i + \tau'_i)^2 = O(n^{-2})$, we now prove that:

$$\mathbb{P}_{\theta_0}(S_i/n < k/\sqrt{n}) \leq C/n^2. \quad (24)$$

Proof of (24): We remark:

$$\begin{aligned} \left(\frac{S_i}{n} < \frac{k}{\sqrt{n}} \right) &\subset \left(\left| \phi_i^2 - \frac{S_i}{n} \right| > \phi_i^2 - \frac{k}{\sqrt{n}}, \phi_i^2 \geq 2 \frac{k}{\sqrt{n}} \right) \cup \left(\phi_i^2 < 2 \frac{k}{\sqrt{n}} \right) \\ &\subset \left(\left| \phi_i^2 - \frac{S_i}{n} \right| > \frac{\phi_i^2}{2} \right) \cup \left(\phi_i^{-2} > \frac{\sqrt{n}}{2k} \right) = \left(\left| 1 - \frac{R_i}{T_i} \right| > \frac{1}{2} \right) \cup \left(\phi_i^{-2} > \frac{\sqrt{n}}{2k} \right). \end{aligned}$$

Consequently, using the Markov inequality and Proposition 4 yields:

$$\mathbb{P}_{\theta_0}(S_i/n < k/\sqrt{n}) \leq C \left(2^4 \left(\frac{T_i}{n} \right)^2 (1 + \mathbb{E}_{\theta_0}(\phi_i^4 + \phi_i^8)) + \frac{(2k)^4}{n^2} \mathbb{E}_{\theta_0} \phi_i^{-4} \right) \leq c'/n^2.$$

So the proof of (24) is complete. \square

It remains to study the terms ν_i, τ_i . We have on $(S_i/n \geq k/\sqrt{n})$:

$$\begin{aligned} |\nu_i| &= \left| \frac{1}{S_i^1/n} (S_i^1/n - S_i/n) \left(\frac{1}{S_i/n} - \Gamma_i + \Gamma_i \right) \right| \\ &\leq \frac{1}{R_i^1/T_i} \left| \left(\frac{R_i^1}{T_i} - \frac{R_i}{T_i} \right) \left(1 - \frac{R_i}{T_i} \right) \right| \Gamma_i^{-1} \frac{\sqrt{n}}{k} + \frac{\Gamma_i}{R_i^1/T_i} \left| \frac{R_i^1}{T_i} - \frac{R_i}{T_i} \right|. \end{aligned}$$

We use Proposition 4, the Cauchy-Schwarz inequality and the exact distribution of Γ_i and $\frac{1}{R_i^1/T_i}$ to obtain, for $n > 4$ and $\mathbb{E}_{\theta_0} \phi_i^8 < +\infty$:

$$\mathbb{E}_{\theta_0} \nu_i^2 \leq C \left[\frac{n}{k^2} \left(\mathbb{E}_{\theta_0} \left(\frac{R_i^1}{T_i} - \frac{R_i}{T_i} \right)^4 \mathbb{E}_{\theta_0} \left(1 - \frac{R_i}{T_i} \right)^4 \right)^{1/2} + \left(\mathbb{E}_{\theta_0} \left(\frac{R_i^1}{T_i} - \frac{R_i}{T_i} \right)^2 \right) \right] \leq \frac{C'}{n^2}.$$

For the term τ_i , we use the Taylor formula and get:

$$\tau_i = (S_i/n - S_i^1/n) \int_0^1 \frac{ds}{s(S_i/n) + (1-s)(S_i^1/n)}.$$

Then, we split the integral:

$$\int_0^1 \frac{ds}{s(S_i/n) + (1-s)(S_i^1/n)} = \frac{1}{S_i^1/n} + \int_0^1 \frac{s(S_i^1/n - S_i/n)}{(s(S_i/n) + (1-s)(S_i^1/n))S_i^1/n} ds.$$

Thus, on $(S_i/n \geq k/\sqrt{n})$, we obtain, after simplifications:

$$|\tau_i| \leq \frac{1}{R_i^1/T_i} |R_i/T_i - R_i^1/T_i| + \frac{\sqrt{n}}{k} \frac{1}{R_i^1/T_i} \Gamma_i^{-1} (R_i/T_i - R_i^1/T_i)^2.$$

This yields $\mathbb{E}_{\theta_0} \tau_i^2 \leq C/n^2$. The proof of Lemma 2 is now complete. \square

Applying Lemma 2, we obtain the result of Proposition 8. \square

7 Auxiliary results

We recall some properties of Gamma and related distributions. The Gamma distribution with parameters (a, λ) ($a > 0, \lambda > 0$), $G(a, \lambda)$, has density $\gamma_{a,\lambda}(x) = (\lambda^a/\Gamma(a))x^{a-1}e^{-x}\mathbb{1}_{(0,+\infty)}(x)$, where $\Gamma(a)$ is the Gamma function. The digamma function $\psi(a) = \Gamma'(a)/\Gamma(a)$ admits the following integral representation: $\psi(z) = -\gamma + \int_0^1 (1-t^{z-1})/(1-t)dt$. (where $\gamma = \psi(1) = \Gamma'(1)$). For all positive a , we have $\psi'(a) = -\int_0^1 \frac{\log t}{1-t} t^{a-1} dt$. The following asymptotic expansions as a tends to infinity hold:

$$\log \Gamma(a) = (a - \frac{1}{2}) \log a - a + \frac{1}{2} \log 2\pi + O(\frac{1}{a}), \quad (25)$$

$$\psi(a) = \log a - \frac{1}{2a} + O(\frac{1}{a^2}), \quad \psi'(a) = \frac{1}{a} + O(\frac{1}{a^2}). \quad (26)$$

The following results are classical.

Proposition 10. *If X has distribution $G(a, \lambda)$, then λX has distribution $G(a, 1)$. For all integer k , $\mathbb{E}(\lambda X)^k = \frac{\Gamma(a+k)}{\Gamma(a)}$. For $a > k$, $\mathbb{E}(\lambda X)^{-k} = \frac{\Gamma(a-k)}{\Gamma(a)}$. Moreover, $\mathbb{E} \log(\lambda X) = \psi(a)$, $\text{Var}[\log(\lambda X)] = \psi'(a)$.*

If X, Y are independent, X having distribution $G(a, 1)$ and Y having distribution $G(b, 1)$ ($a, b > 0$), then, $X + Y$ and $X/(X + Y)$ are independent, $X + Y$ has distribution $G(a + b, 1)$, $T = \frac{X}{X+Y}$ has distribution beta of the first kind with parameters (a, b) , denoted by $\beta^{(1)}(a, b)$, and density

$$f_T(t) = \frac{1}{B(a, b)} t^{a-1} (1-t)^{b-1} \mathbb{1}_{(0,1)}(t),$$

with $B(a, b) = \frac{\Gamma(a)\Gamma(b)}{\Gamma(a+b)}$ and $Z = X/Y$ has distribution beta of the second kind with parameters (a, b) , denoted by $\beta^{(2)}(a, b)$, and density

$$f_Z(z) = \frac{1}{B(a, b)} \frac{z^{a-1}}{(1+z)^{a+b}} \mathbb{1}_{(0, +\infty)}(z).$$

We have $E(T) = \frac{a}{a+b}$, $E(T^2) = \frac{a(a+1)}{(a+b)(a+b+1)}$, $Var(T) = \frac{ab}{(a+b)^2(a+b+1)}$, $E \log T = \psi(a) - \psi(a+b)$, $Var(\log T) = \psi'(a) - \psi'(a+b)$, $Cov(T, \log T) = \frac{b}{(a+b)^2}$.