



HAL
open science

Completeness in approximation classes beyond APX

Bruno Escoffier, Vangelis Paschos

► **To cite this version:**

| Bruno Escoffier, Vangelis Paschos. Completeness in approximation classes beyond APX. 2005. <hal-00958036>

HAL Id: hal-00958036

<https://hal.science/hal-00958036v1>

Preprint submitted on 11 Mar 2014

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



HAL Authorization

CAHIER DU LAMSADE

225

Septembre 2005

Completeness in approximation classes beyond APX

Bruno Escoffier, Vangelis Th. Paschos

Completeness in approximation classes beyond APX

Bruno Escoffier Vangelis Th. Paschos

LAMSADE, Université Paris-Dauphine
Place du Maréchal De Lattre de Tassigny, 75775 Paris Cedex 16, France
{escoffier,paschos}@lamsade.dauphine.fr

September 13, 2005

Abstract

We present a reduction that allows us to establish completeness results for several approximation classes mainly beyond **APX**. Using it, we extend one of the basic results of S. Khanna, R. Motwani, M. Sudan, and U. Vazirani (*On syntactic versus computational views of approximability*, SIAM J. Comput., 28:164–191, 1998) by proving the existence of complete problems for the whole **Log-APX**, the class of problems approximable within ratios that are logarithms of the size of the instance. We also introduce a new approximability class, called **Poly-APX(Δ)**, dealing with graph-problems approximable with ratios functions of the maximum degree Δ of the input graph. For this class also, using the reduction propose, we establish complete problems.

1 Introduction and preliminaries

Consider an **NPO** problem¹ $\Pi = (\mathcal{I}_\Pi, \text{Sol}_\Pi, m_\Pi, \text{goal}_\Pi)$, where: \mathcal{I}_Π denotes the set of instances of Π ; for any instance $x \in \mathcal{I}_\Pi$, $\text{Sol}_\Pi(x)$ is the set of feasible solutions of x ; for any $x \in \mathcal{I}_\Pi$ and any $y \in \text{Sol}_\Pi(x)$, $m_\Pi(x, y)$ denotes the value of y ; finally, goal_Π is max, or min. For any $x \in \mathcal{I}_\Pi$, let $\text{opt}_\Pi(x)$ be the value of an optimal solution for x . Then, the approximation ratio of an algorithm **A** computing a feasible solution $\mathbf{A}(x) = y \in \text{Sol}_\Pi(x)$ is defined by $m_\Pi(x, y)/\text{opt}_\Pi(x)$. The objective of the polynomial approximation theory is double: on the one hand it aims at devising polynomial algorithms achieving good approximation ratios for **NP**-hard problems; on the other hand, it aims at building a hierarchy of these problems, elements of which correspond to strata of problems sharing common approximability properties (they notably are approximable within comparable – in some predefined sense – approximation ratios) and at investigating relations between problems in the same stratum (notably to exhibit problems that are harder from others). This second objective is the scientific are called in short structure in approximability classes.

Study of structure in approximability classes is in the heart of the research in polynomial approximation since the seminal papers of [10, 11, 6]. By using suitable approximation-preserving reductions, the existence of natural complete problems for almost all the known approximation classes has been established. For instance, **MAX WSAT** for **NPO** under **AP**-reduction ([3, 5]), or **PTAS**-reduction ([7]), **MAX WSAT-B** for **APX**, the class of the problems approximable within (fixed) constant ratios, under **P**-reduction ([6]), **MAX 3-SAT** for **APX** under **AP**-reduction ([3, 5]), or **PTAS**-reduction ([7]), **MAX PLANAR INDEPENDENT SET** for **PTAS**, the class of the problems solvable by polynomial time approximation schemata, under **FT**-reduction ([4]), **MAX INDEPENDENT SET** for **Poly-APX** under **PTAS**-reduction ([4]). Also, under **E**-reduction, completeness in **Log-APX-PB**, the subclass of **Log-APX** dealing with polynomially bounded problems (the

¹An **NPO** problem is an optimization problem, the decision version of which is in **NP**.

class of problems whose values are bounded by a polynomial in the size of the instance), has also been established. One problem that, to our knowledge, remains open from this ambitious but so successful research program, is the establishing of completeness for the whole **Log-APX**.

As one can see by the unravelling of the fascinating history of the approximation-preserving reductions, the one that allows achievement of the most of completeness results is the **PTAS-reduction**, originally introduced in [7]. Let Π and Π' be two maximization **NPO** problems (the case of minimization is completely analogous). We say that Π *PTAS-reduces to* Π' if and only if there exist three functions f , g and c such that:

- for any $x \in \mathcal{I}_\Pi$ and any $\varepsilon \in]0, 1[$, $f(x, \varepsilon) \in \mathcal{I}_{\Pi'}$; f is computable in time polynomial with $|x|$;
- for any $x \in \mathcal{I}_\Pi$, any $\varepsilon \in]0, 1[$ and any $y \in \text{Sol}(f(x, \varepsilon))$, $g(x, y, \varepsilon) \in \text{Sol}(x)$; g is computable in time polynomial with $|x|$ and $|y|$;
- $c :]0, 1[\rightarrow]0, 1[$;
- for any $x \in \mathcal{I}_\Pi$ and any $\varepsilon \in]0, 1[$, $r_{\Pi'}(f(x, \varepsilon), y) \geq 1 - c(\varepsilon)$ implies $r_\Pi(x, g(x, y, \varepsilon)) \geq 1 - \varepsilon$.

This is the successor of a number of powerful reductions as the **L-reduction** ([11]) or the **E-reduction** ([9]) that, even if they allowed achievement of completeness results in natural approximability sub-classes (e.g., **Max-SNP** \subset **APX**, for the former, or **Log-APX-PB** \subset **Log-APX**, for the latter) were not able to extend them to the whole of the classes dealt. In fact, as shown in [7], these reductions suffered from the fact that they map optimal solutions to optimal solutions and, in this sense, it is very unlikely that they can allow completeness of a polynomially bounded problem in the classes dealt (unless $\mathbf{P}^{\text{SAT}} = \mathbf{P}^{\text{SAT}[O(\log(n))]}$, where \mathbf{P}^{SAT} and $\mathbf{P}^{\text{SAT}[O(\log(n))]}$ are the classes of decision problems solvable by using, respectively, a polynomial and a logarithmic number of calls to an oracle solving SAT). On the contrary, **PTAS-reduction**, by allowing functions f and g to depend on ε , does not necessarily map optimal solutions between them and consequently, it allows that general problems are **PTAS-reducible** to polynomially bounded ones. Indeed, by means of **PTAS-reduction** it is proved in [7] that **MAX WSAT-B** reduces **MAX WSAT-B** with weights bounded by a polynomial to the size of the instance.

In [4] the power of **PTAS-reduction** has been confirmed since it has allowed the achievement of **Poly-APX-complete** problems. This was not possible with the **E-reduction** under which only **Poly-APX-PB** completeness have been obtained in [9]. Given a family \mathbf{F} of functions, denote by **F-APX** the subclass of **NPO** whose problems are approximable in polynomial time within ratio $g(n)$ (in the case of minimization), or $1/g(n)$ (for maximization), for a $g \in \mathbf{F}$. Here we further confirm its scope by generalizing the result of [4], providing a way to find complete problems, under **PTAS-reducibility**, for any approximation class **F-APX**, where \mathbf{F} is a class of polynomially bounded functions. This in particular allows us to establish the **Log-APX-completeness** of **MIN SET COVER**. Recall that, in [9], only the **Log-APX-PB** completeness of this problem has been established under the **E-reduction**.

Before continuing, we define some key-notions for what follows, namely, additivity and canonical hardness of a maximization **NPO** problem. They are introduced in [9]. A problem $\Pi \in \mathbf{NPO}$ is said *additive* if and only if there exist an operator \oplus and a function f , both computable in polynomial time, such that:

- \oplus associates with any pair $(x_1, x_2) \in \mathcal{I}_\Pi \times \mathcal{I}_\Pi$ an instance $x_1 \oplus x_2 \in \mathcal{I}_\Pi$ with $\text{opt}(x_1 \oplus x_2) = \text{opt}(x_1) + \text{opt}(x_2)$;
- with any solution $y \in \text{sol}_\Pi(x_1 \oplus x_2)$, f associates two solutions $y_1 \in \text{sol}_\Pi(x_1)$ and $y_2 \in \text{sol}_\Pi(x_2)$ such that $m_\Pi(x_1 \oplus x_2, y) = m_\Pi(x_1, y_1) + m_\Pi(x_2, y_2)$.

A set \mathbf{F} of functions from \mathbb{N} to \mathbb{N} will be called *downward close* if, for any function $g \in \mathbf{F}$ and any constant c , if $h(n) = O(g(n^c))$, then $h \in \mathbf{F}$. A function $g : \mathbb{N} \rightarrow \mathbb{N}$ is *hard* for \mathbf{F} if and only if, for any $h \in \mathbf{F}$, there exists a constant c such that $h(n) = O(g(n^c))$; if, in addition, $g \in \mathbf{F}$, then g is said *complete* for \mathbf{F} .

A maximization (resp., minimization) problem $\Pi \in \mathbf{NPO}$ is *canonically hard for $\mathbf{F-APX}$* , for a downward close family \mathbf{F} of functions, if and only if there exist a polynomially computable function f , two constants n_0 and c and a function F , hard for \mathbf{F} , such that:

- for any instance φ of 3SAT on $n \geq n_0$ variables and for any $N \geq n^c$, $f(\varphi, N)$ belongs to \mathcal{I}_Π ;
- if φ is satisfiable, then $\text{opt}(f(\varphi, N)) = N$;
- if x is not satisfiable, then $\text{opt}(f(\varphi, N)) = N/F(N)$ (resp., $NF(N)$);
- for any $y \in \text{sol}_\Pi(f(\varphi, N))$ such that $m(f(\varphi, N), y) > N/F(N)$ (resp., $m(f(\varphi, N), y) < NF(N)$), one can polynomially determine a truth assignment satisfying φ .

More generally, since 3SAT is \mathbf{NP} -complete, a maximization (resp., minimization) problem Π is canonically hard for $\mathbf{F-APX}$ if and only if, for any decision problem $\Pi' \in \mathbf{NP}$, given an instance x' of Π' , one can construct in polynomial time an instance x of Π such that:

- if x' is a positive instance, then $\text{opt}(x) = N$;
- if x' is a negative instance, then $\text{opt}(x) = N/F(N)$ (resp. $NF(N)$);
- given a solution $y \in \text{sol}_\Pi(x)$ such that $m(x, y) > N/F(N)$ (resp., $< NF(N)$), one can polynomially determine a certificate proving that x' is a positive instance.

In [9], the following major theorem is proved, that constitutes the starting point of the paper at hand.

Theorem 1. ([9]) *Let \mathbf{F} be any class of downward close polynomially bounded functions and Ω be an additive maximization problem canonically hard for \mathbf{F} . Then, any maximization problem $\Pi \in \mathbf{F-APX-PB}$ (the class of problems in $\mathbf{F-APX}$ whose values are bounded by a polynomial with the size of the instance) \mathbf{E} -reduces to Ω .*

In order to extend result of Theorem 1, and since, as we have already mentioned, it seems very unlikely that one could establish completeness for the whole $\mathbf{F-APX}$ under \mathbf{E} -reducibility (at least for polynomially bounded problems), we introduce a modification of PTAS-reducibility, called MPTAS-reduction, \mathbf{M} standing for multivalued, where function f is allowed to be multivalued. Formally, MPTAS-reduction can be defined as follows.

Definition 1. Let Π and Π' be two maximization \mathbf{NPO} problems (the case of minimization is completely analogous). Then, Π *MPTAS-reduces to* Π' , if and only if there exist three functions f and g , computable in polynomial time, and a function c such that:

- for any $x \in \mathcal{I}_\Pi$ and any $\varepsilon \in]0, 1[$, $f(x, \varepsilon) = (f_1(x, \varepsilon), f_2(x, \varepsilon), \dots)$ is a family of instances of Π' (this family is necessarily of size polynomial with x);
- for any $x \in \mathcal{I}_\Pi$, any $\varepsilon \in]0, 1[$ and any family of feasible solutions $y = (y_1, y_2, \dots)$, where y_i is a feasible solution of $f_i(x, \varepsilon)$, $g(x, y, \varepsilon) \in \text{Sol}(x)$;
- $c :]0, 1[\rightarrow]0, 1[$;
- there exists an index j such that, for any $x \in \mathcal{I}_\Pi$ and any $\varepsilon \in]0, 1[$, $r_{\Pi'}(f_j(x, \varepsilon), y_j) \geq 1 - c(\varepsilon)$ implies $r_\Pi(x, g(x, y, \varepsilon)) \geq 1 - \varepsilon$. ■

It is easy to see that MPTAS-reduction preserves membership in **PTAS**.

We show that using MPTAS-reducibility, **F-APX**-completeness can be extended for any class **F** of functions even if they are not polynomially bounded. Furthermore, the fact that function f in Definition 1 is allowed to be multivalued, relaxes the restriction to additive problems. As we will see, as a corollary, we have an alternative proof (without explicit use of the Cook-Levin theorem) for the **Exp-APX**-completeness of MIN TSP on general graphs, where **Exp** is the class of exponential functions (note that the **NPO**-completeness of MIN TSP immediately derives its **Exp-APX**-completeness).

The approximation classes beyond **APX** dealt until now in the literature are defined with respect ratios depending on the size of the instances and not on other parameters even natural. This can be considered as somewhat restrictive given that a lot of approximation (and of inapproximability results) are established with respect to other parameters of the instances. Such parameters can be, for example, the maximum or the average degree of the input-graph, when dealing with graph-problems, the maximum set-cardinality, when dealing with problems on set-systems as the MIN SET COVER, or the MAX SET PACKING, etc. So, the third and last issue of this paper is to consider the class **Poly-APX(Δ)**, of maximization (resp., minimization) graph-problems approximable within ratios which are inverse polynomials (resp., polynomials) of the maximum degree Δ of the input-graph. We prove that MAX INDEPENDENT SET is complete for this class, under MPTAS.

2 **F-APX**-completeness for any downward close function

We show in this section that, using MPTAS-reduction (Definition 1), we can apprehend completeness for any class **F-APX** even containing exponential functions. In a first time we shall deal with maximization problems. Extension to minimization problems will be performed after the proof of Theorem 2 that follows.

Theorem 2. *Let \mathbf{F} be a class of downward close functions, and $\Omega \in \mathbf{NPO}$ be a maximization problem canonically hard for **F-APX**. Then, any maximization **NPO** problem in **F-APX** MPTAS-reduces to Ω .*

Proof. Let Π be a maximization problem of **F-APX** and let \mathbf{A} be an algorithm achieving approximation ratio $1/r(\cdot)$, $r \in \mathbf{F}$. Since $\Pi \in \mathbf{NPO}$, the value of the solutions for any instance $x \in \mathcal{I}_\Pi$ is bounded above by $2^{p(|x|)}$ for some polynomial p . Let Ω be as assumed in theorem's statement. Let F be a function hard for **F**, and k , n_0 and c' be constants such that, for $n \geq n_0$ $r(n) - 1 \leq k(F(n^{c'}) - 1)$. Finally, consider $x \in \mathcal{I}_\Pi$ and $\varepsilon \in (0, 1)$.

In order to build function $f(x, \varepsilon)$, claimed by Definition 1, we first partition interval $[1, 2^{p(|x|)}]$ of the possible values for $\text{opt}(x)$ into a polynomial number of subintervals $[(1/(1-\varepsilon))^{i-1}, (1/(1-\varepsilon))^i]$, for $i = 1, \dots, M = \lceil p(|x|) \ln 2 / \ln(1/(1-\varepsilon)) \rceil$ (i.e., $(1/(1-\varepsilon))^M \geq 2^{p(|x|)}$). Consider then, for $i = 1, \dots, M$, the languages $L_i = \{x \in \mathcal{I}_\Pi : \text{opt}_\Pi(x) \geq (1/(1-\varepsilon))^{i-1}\}$. Set $N = |x|^{c'}$. Since Ω is canonically hard for **F-APX**, we can build, for any $i = 1, \dots, M$, an instance $\omega_i \in \mathcal{I}_\Omega$ such that, if $x \in L_i$, then $\text{opt}_\Omega(\omega_i) = N$, otherwise, $\text{opt}_\Omega(\omega_i) = N/F(N)$. We set $f(x, \varepsilon) = \Upsilon = (\omega_i, 1 \leq i \leq M)$.

We now show how $g(x, y, \varepsilon)$ can be built. Consider a solution $y = (y_1, \dots, y_M)$ of Υ . Then, for any $i \in \{1, \dots, M\}$, if $m_\Omega(\omega_i, y_i) > N/F(N)$, one can find a polynomial certificate proving that $x \in L_i$, i.e., a solution $y'_i \in \text{Sol}(x)$ such that

$$m_\Pi(x, y'_i) \geq \left(\frac{1}{1-\varepsilon} \right)^{i-1} \quad (1)$$

otherwise (i.e., if $m_\Omega(\omega_i, y_i) \leq N/F(N)$), we consider solution $\mathbf{A}(x)$. Finally, $g(x, y, \varepsilon)$ is the best among the solutions so produced.

Let us now prove that we really deal with an MPTAS-reduction. Let j be an index verifying:

$$\left(\frac{1}{1-\varepsilon}\right)^{j-1} \leq \text{opt}_{\Pi}(x) \leq \left(\frac{1}{1-\varepsilon}\right)^j \quad (2)$$

and set $c(\varepsilon) = \varepsilon/(\varepsilon + k(1 - \varepsilon))$.

Assume first that $m_{\Omega}(\omega_j, y_j) > N/F(N)$. Then, using (1) and (2), we get:

$$\frac{m_{\Pi}(x, g(x, y, \varepsilon))}{\text{opt}_{\Pi}(x)} \geq 1 - \varepsilon$$

On the other hand, if $m_{\Omega}(\omega_j, y_j) \leq N/F(N)$, then $\rho(\omega_j, y_j) \geq 1 - c(\varepsilon)$ implies:

$$\frac{1}{F(N)} \geq \frac{k(1 - \varepsilon)}{\varepsilon + k(1 - \varepsilon)} \quad (3)$$

By the assumptions made above, we have:

$$F(N) \geq 1 + \frac{r(|x|) - 1}{k} \quad (4)$$

$$\rho(x, g(x, y, \varepsilon)) \geq \frac{1}{r(|x|)} \quad (5)$$

Using (3), (4) and (5), we get the following implications:

$$\rho(\omega_j, y_j) \geq 1 - c(\varepsilon) \implies 1 + \frac{r(|x|) - 1}{k} \geq \frac{\varepsilon + k(1 - \varepsilon)}{k(1 - \varepsilon)} \implies r(|x|) \geq 1 + k \left(\frac{\varepsilon}{k(1 - \varepsilon)} \right) = \frac{1}{1 - \varepsilon}$$

i.e., $\rho(x, g(x, y, \varepsilon)) \geq 1 - \varepsilon$. So, the reduction exhibited is indeed a MPTAS-reduction and the proof of the theorem is completed. ■

In order to extend the result of Theorem 2 to minimization problems, we are based upon an analogous result in [9]. Let Π be a maximisation problem in **F-APX** approximable within ratio $1/f(\cdot)$. One can define a minimization problem Π' , identical to Π up to its objective function defined as: $m_{\Pi'}(x, y) = \lfloor 2M^2(x)/m_{\Pi}(x, y) \rfloor$, where $M(x)$ is an upper bound for the values in $\text{Sol}(x)$. Then, Π E-reduces (hence, MPTAS-reduces also) to Π' and, furthermore, Π' is approximable within ratio $f(\cdot)(1 + 2/M(x))$. In completely analogous way, one can MPTAS-reduce a minimization problem, approximable within ratio $f(\cdot)$, to a maximization problem approximable within ratio $1/(f(\cdot)(1 + 2/M(x)))$. This allows the existence of additive canonically hard *minimization* problems also to be **F-APX**-complete, since the above transformation derives, from such a problem Π , an additive canonically hard maximization problem Π' (notice that $f(\cdot)$ and $f(\cdot)(1 + 2/M(x))$ have the same hardness status with respect to **F**). Then, it suffices that one applies Theorem 2 as stated. The remark above applies for any family **F**. As a consequence, throughout this paper we restrict ourselves to maximization problems.

Theorem 3. *Let **F** be a class of downward close functions, and $\Omega \in \mathbf{NPO}$ be a problem canonically hard for **F**. Then, any **NPO** problem in **F-APX** MPTAS-reduces to Ω .*

From the discussion above, two are the main differences between Theorems 1 and 3. The first one is, as it has been already announced, that the latter one applies to the whole of the problems of any class **F-APX** and not only to the bounded ones. The second difference is that, the use of a multivalued version of PTAS-reduction allows us to relax additivity from the conditions of Theorem 1 enlarging so the scope of the applicability of Theorem 3.

Obviously, the negative result for MIN TSP of [13] (see also [8]) can immediately be extended for showing that MIN TSP is canonically hard for **Exp-APX**. Consequently, the **Exp-APX**-completeness of MIN TSP can be derived as an immediate corollary of Theorem 3.

Corollary 1. *MIN TSP is **Exp-APX**-complete under MPTAS-reducibility.*

3 Log-APX-completeness

We deal in this section with **Log-APX**. For this class, as well as for any class of polynomially bounded downward close functions, we can obviously apply Theorem 3 to get completeness sufficient conditions. However, in what follows, we will use the fact that functions dealt are polynomially bounded in order to prove an equivalent of Theorem 3 but with respect to PTAS-reducibility. As noticed above, we can restrict ourselves to maximization problems.

Theorem 4. *Let \mathbf{F} be any class of downward close polynomially bounded functions and Ω be an additive maximization problem canonically hard for $\mathbf{F-APX}$. Then, any maximization problem $\Pi \in \mathbf{F-APX}$ PTAS-reduces to Ω .*

Proof. The proof goes along the lines of the corresponding proof of [4] for **Poly-APX**. Let Π be a maximization problem of $\mathbf{F-APX}$ and let \mathbf{A} be an algorithm achieving approximation ratio $1/r(\cdot)$, $r \in \mathbf{F}$. Let Ω be as assumed in theorem's statement. Let F be a function hard for \mathbf{F} , and k, n_0 and c' be constants such that, for $n \geq n_0$, $r(n) \leq k(F(n^{c'}) - 1)$. Finally, consider $x \in \mathcal{I}_\Pi$, $\varepsilon \in (0, 1)$ and set $|x| = n$.

We first construct function $f(x, \varepsilon)$. Set $m(x, \mathbf{A}(x)) = m \geq \text{opt}_\Pi(x)/r(n)$. Partition the interval $[0, mr(n)]$ of the possible values for $\text{opt}_\Pi(x)$ into $q(n)$ regular subintervals, where $q(n)$ is bounded by polynomial of n . More precisely, we set $q(n) = r(n)/\varepsilon$. Consider now, for $i \in \{1, \dots, q(n)\}$, the sets of languages $L_i = \{x \in \mathcal{I}_\Pi : \text{opt}_\Pi(x) \geq imr(n)/q(n)\}$ and set $N = n^{c'}$. Since Ω is canonically hard for $\mathbf{F-APX}$, one can build, for any i , an instance $\omega_i \in \mathcal{I}_\Omega$ such that, if $x \in L_i$, then $\text{opt}_\Omega(\omega_i) = N$ and if $x \notin L_i$, then $\text{opt}_\Omega(\omega_i) = N/F(N)$. We define $f(x, \varepsilon) = \Upsilon = \bigoplus_{1 \leq i \leq q(n)} \omega_i$. Observing that $r(n)/q(n) = \varepsilon$, we get:

$$\text{opt}_\Omega(\Upsilon) = N |\{i \leq q(n) : \text{opt}_\Pi(x) \geq im\varepsilon\}| + \frac{N}{F(N)} (q(n) - |\{i \leq q(n) : \text{opt}_\Pi(x) \geq im\varepsilon\}|) \quad (6)$$

We now construct function $g(x, y, \varepsilon)$. Consider a solution $y \in \text{Sol}(\Upsilon)$. By the additivity of Ω , one can compute, for any ω_i , a solution y_i in such a way that $m_\Omega(\Upsilon, y) = \sum_i m_\Omega(\omega_i, y_i)$. Let j be the largest among the indices i such that $m_\Omega(\omega_i, y_i) > N/F(N)$ ($j = 0$, if no i verifies the inequality). Then, one can find a polynomial certificate proving that $x \in L_j$, i.e., a solution $y'_1 \in \text{Sol}(x)$ verifying: $m_\Pi = (x, y'_1) \geq jm\varepsilon$. Furthermore, $m_\Omega(\Upsilon, y) \leq Nj + (q(n) - j)N/F(N)$.

We then define $g(x, y, \varepsilon) = y'$ as the best (the largest value one) among y'_1 and $y'_2 = \mathbf{A}(x)$. Obviously, $m_\Pi(x, y') \geq \max\{m, jm\varepsilon\}$.

Let us now prove that the reduction described is indeed a PTAS-reduction. We first notice that, using expressions for m_Π and m_Ω , the following inequality is derived (after some algebra) for the approximation ratio $\rho_\Omega(\Upsilon, y)$, using that $q(n) = r(n)/\varepsilon \leq k(F(N) - 1)/\varepsilon$:

$$\rho_\Omega(\Upsilon, y) \leq \frac{j + \frac{k}{\varepsilon}}{\frac{\text{opt}_\Pi(x)}{\varepsilon m} - 1 + \frac{k}{\varepsilon}}$$

We now consider two cases, namely, $j \leq 1/\varepsilon$ and $j \geq 1/\varepsilon$. For the first one, $j \leq 1/\varepsilon$, we get from the expression for ρ_Ω (after some algebra and using also that $\rho_\Pi(x, y') \geq m/\text{opt}_\Pi(x)$):

$$\rho_\Omega(\Upsilon, y) \leq \frac{\rho_\Pi(x, y') (1 + k)}{1 + \rho_\Pi(x, y') (k - \varepsilon)}$$

On the other hand, if $j \geq 1/\varepsilon$, noticing that, using expression for m_Π , $\rho_\Pi(x, y') \geq jm\varepsilon/\text{opt}_\Pi(x)$, we also get:

$$\rho_\Omega(\Upsilon, y) \leq \frac{\rho_\Pi(x, y') (1 + k)}{1 + \rho_\Pi(x, y') (k - \varepsilon)}$$

Assume $c(\varepsilon) = \varepsilon^2 / (1 + (1 - \varepsilon)(k - \varepsilon))$. Then, after some algebra one gets: $\rho_{\Pi}(x, y') \geq 1 - \varepsilon$, proving so that the reduction just devised is indeed a PTAS-reduction and completing the proof of the theorem. ■

Consider now **Log-APX**, the class of **NPO** problems approximable with logarithmic (resp., inversely logarithmic when goal = max) ratios, i.e., within ratios $O(\log(\cdot))$ (resp., $O(1/\log(\cdot))$). Consider also MIN SET COVER that is approximable within ratio $O(\log n)$ ([14]), where n is the size of the ground set in the instance, but canonically hard to approximate within ratios $O(\log n)$ ([12]). Moreover, one can easily see that MIN SET COVER is additive. So, immediate application of Theorem 4 definitely settles the status of MIN SET COVER with respect to the class **Log-APX**.

Theorem 5. MIN SET COVER is **Log-APX**-complete under PTAS-reducibility.

4 Completeness in **F-APX**(Δ)

We now deal with a new approximation class, **F-APX**(Δ), namely, the class of graph-problems approximable with ratio $f(\Delta)$, where Δ is the degree of the input-graph, i.e., the maximum degree of its vertices, **F** a downward close class of functions and $f \in \mathbf{F}$.

Definition 2. Let **F** be a downward close class of functions and $\Pi \in \mathbf{NPO}$ be a maximization (resp., minimization) graph-problem. Then, Π is said *canonically hard* for **F-APX**(Δ), if there exist three functions f , α and β , computable in polynomial time, a constant Δ_0 and a function F , hard for **F** such that:

- for any instance φ of 3-SAT and any $\Delta \geq \Delta_0$, $G_{\varphi} = f(\varphi, \Delta)$ is a graph (instance of Π) of maximum degree Δ ;
- if φ is satisfiable, then $\text{opt}_{\Pi}(G_{\varphi}) \geq \alpha(G_{\varphi})$ (resp., $\text{opt}_{\Pi}(G_{\varphi}) \leq \alpha(G_{\varphi})$);
- if φ is not satisfiable, then $\text{opt}_{\Pi}(G_{\varphi}) \leq \beta(G_{\varphi})$ (resp., $\text{opt}_{\Pi}(G_{\varphi}) \geq \beta(G_{\varphi})$);
- $\alpha(G_{\varphi})/\beta(G_{\varphi}) \geq f(\Delta)$ (resp., $\beta(G_{\varphi})/\alpha(G_{\varphi}) \geq f(\Delta)$);
- given a solution $y \in \text{sol}(G_{\varphi})$ of value strictly greater (resp., smaller) than $\beta(G_{\varphi})$, one can determine, in polynomial time a truth assignment satisfying φ . ■

For the same reasons as previously, we will restrict ourselves to maximization problems and are going to prove the following theorem.

Theorem 6. If **F** is a downward close family of functions and Ω is a maximization problem canonically hard for **F-APX**(Δ), then any problem in **F-APX**(Δ) MPTAS-reduces to Ω .

Proof. Consider a maximization problem $\Pi \in \mathbf{F-APX}(\Delta)$ a graph $G \in \mathcal{I}_{\Pi}$ of order n and an algorithm **A** for Π achieving approximation ratio $1/r(\Delta(G))$, where $r \in \mathbf{F}$. The proof goes along the same line as the one of Theorem 2. For constructing function $f(G, \varepsilon)$ claimed by Definition 1 we partition the interval $[1, 2^{p(n)}]$ in the same way as in the proof of Theorem 2, and we consider, for any i , the analogous sets L_i of languages. We set $\Delta = \Delta^c(G)$, we build, for any i an instance H_i of Ω , of maximum degree Δ such that if $G \in L_i$, then $\text{opt}_{\Omega}(H_i) \geq \alpha(H_i)$, otherwise $\text{opt}_{\Omega}(H_i) \leq \beta(H_i)$, with $\alpha(H_i)/\beta(H_i) \geq F(\Delta)$ and, finally we specify $f(G, \varepsilon) = H = (H_i, 1 \leq i \leq M)$.

Function $g(G, y, \varepsilon)$ is defined as in the proof of Theorem 2, setting $\beta(H_i)$ instead of $N/F(N)$. The proof for the transfer of the approximation ratios is also done in the same way as there (always with $\beta(H_i)$ instead of $N/F(N)$). ■

Denote by **Poly-APX**(Δ) the subclass of the graph-problems in **NPO** which are approximable within polynomials of Δ^{-1} (of Δ when dealing with minimization problems). We are going now to establish an interesting completeness result for this class by showing that one of the most paradigmatic problems for the polynomial approximation theory and the combinatorial optimization, the **MAX INDEPENDENT SET**, is complete for **Poly-APX**(Δ). For this we will use the following theorem ([1]).

Theorem 7. ([1]) *Let $(a, b) \in [0, 1]^2$, $a > b$. There exists $\varepsilon_0 > 0$ such that, for any $\Delta \geq 3$, there exists a function f that transforms an instance G of **MAX INDEPENDENT SET** into an instance of **MAX INDEPENDENT SET- Δ** , i.e., into a graph with degree bounded by Δ , and two positive quantities α and β such that:*

- if $\text{opt}(G) \geq an$, then $\text{opt}(f(G)) \geq \alpha(f(G))$;
- if $\text{opt}(G) \leq bn$, then $\text{opt}(f(G)) \leq \beta(f(G))$;
- $\alpha(f(G))/\beta(f(G)) \geq \Delta^{\varepsilon_0}$.

Thanks to the **PCP** theorem, stating that there exist a and b such that it is **NP**-complete to distinguish graphs with maximum independent set with size at least a times their order, from graphs having maximum independent set at most b times their order, the following corollary can be derived ([1]).

Corollary 2. ([1]) *There exists an ε_0 such that, for any $\Delta \geq 3$, **MAX INDEPENDENT SET- Δ** is not approximable within ratio $\Delta^{-\varepsilon_0}$.*

We will also use the following proposition, derived from an immediate application of the **PCP** theorem.

Proposition 1. *There exists $(a, b) \in [0, 1]^2$, $a > b$, such that, for any $\Pi \in \mathbf{NP}$ there exists a function f which transforms an instance x of Π into a graph $G = f(x)$, instance of **MAX INDEPENDENT SET**, that verifies:*

1. if x is positive, then $\text{opt}(G) \geq an$;
2. if x is negative, then $\text{opt}(G) \leq bn$;
3. given an independent set S in G with size $|S| > bn$, one can determine, in polynomial time, a witness proving that x is positive.

Indeed, Proposition 1, is an approximation preserving transfer, by say an L-reduction ([11]), of an analogous result dealing with **MAX 3-SAT** (appearing in [2]) to **MAX INDEPENDENT SET**.

Theorem 8. *There exists an $\varepsilon_0 > 0$ such that, for any $\Delta \geq 3$, given $\Pi \in \mathbf{NP}$, there exists a function f computable in polynomial time, that transforms an instance x of Π into a graph $G = f(x)$, instance of **MAX INDEPENDENT SET**, and two positive quantities α and β verifying:*

1. if x is positive, then $\text{opt}(G) \geq \alpha(G)$;
2. if x is negative, then $\text{opt}(G) \leq \beta(G)$;
3. the maximum degree of G is Δ ;
4. $\alpha(G)/\beta(G) \geq \Delta^{\varepsilon_0}$;

5. given an independent set S of G such that $|S| > \beta$, one can build, in polynomial time, a certificate proving that x is positive.

Proof. Items 1 to 4 are direct applications of Theorem 7 and Proposition 1. Item 5 is derived by the proof of Theorem 7 in [1]. Indeed, following the relations between the independent sets in G and those in its derandomized product $f(G)$, from an independent set of $f(G)$ of size strictly greater than β , one can easily derive an independent set of G of size b times the order of G . Then, using item 3 of Proposition 1, one can specify a polynomial certificate proving that x is positive. ■

Consider now **Poly-APX(Δ)** and MAX INDEPENDENT SET- Δ . In order to apply Theorem 6, one has just to show that the complexity of the function f stated in Theorem 8 is polynomial in Δ . This is derived after a careful reading of the proof of Theorem 7 ([1]). So, the following theorem concludes the paper.

Theorem 9. MAX INDEPENDENT SET- Δ is **Poly-APX(Δ)**-complete under MPTAS-reducibility.

References

- [1] N. Alon, U. Feige, A. Wigderson, and D. Zuckerman. Derandomized graph products. *Computational Complexity*, 5:60–75, 1995.
- [2] S. Arora, C. Lund, R. Motwani, M. Sudan, and M. Szegedy. Proof verification and intractability of approximation problems. In *Proc. FOCS'92*, pages 14–23, 1992.
- [3] G. Ausiello, P. Crescenzi, G. Gambosi, V. Kann, A. Marchetti-Spaccamela, and M. Protasi. *Complexity and approximation. Combinatorial optimization problems and their approximability properties*. Springer, Berlin, 1999.
- [4] C. Bazgan, B. Escoffier, and V. Th. Paschos. Completeness in standard and differential approximation classes: Poly-(D)APX- and (D)PTAS-completeness. *Theoret. Comput. Sci.*, 339:272–292, 2005.
- [5] P. Crescenzi, V. Kann, R. Silvestri, and L. Trevisan. Structure in approximation classes. *SIAM J. Comput.*, 28(5):1759–1782, 1999.
- [6] P. Crescenzi and A. Panconesi. Completeness in approximation classes. *Information and Computation*, 93(2):241–262, 1991.
- [7] P. Crescenzi and L. Trevisan. On approximation scheme preserving reducibility and its applications. *Theory of Computing Systems*, 33(1):1–16, 2000.
- [8] M. R. Garey and D. S. Johnson. *Computers and intractability. A guide to the theory of NP-completeness*. W. H. Freeman, San Francisco, 1979.
- [9] S. Khanna, R. Motwani, M. Sudan, and U. Vazirani. On syntactic versus computational views of approximability. *SIAM J. Comput.*, 28:164–191, 1998.
- [10] P. Orponen and H. Mannila. On approximation preserving reductions: complete problems and robust measures. Technical Report C-1987-28, Dept. of Computer Science, University of Helsinki, Finland, 1987.
- [11] C. H. Papadimitriou and M. Yannakakis. Optimization, approximation and complexity classes. *J. Comput. System Sci.*, 43:425–440, 1991.

- [12] R. Raz and S. Safra. A sub-constant error probability low-degree test and a sub-constant error probability PCP characterization of NP. In *Proc. STOC'97*, pages 475–484, 1997.
- [13] S. Sahni and T. Gonzalez. P-complete approximation problems. *J. Assoc. Comput. Mach.*, 23:555–565, 1976.
- [14] P. Slavík. A tight analysis of the greedy algorithm for set cover. In *Proc. STOC'96*, pages 435–441, 1996.