



HAL
open science

Abelian and metabelian quotients of surface braid groups

Paolo Bellingeri, Eddy Godelle, John Guaschi

► **To cite this version:**

Paolo Bellingeri, Eddy Godelle, John Guaschi. Abelian and metabelian quotients of surface braid groups. 2014. hal-00965389

HAL Id: hal-00965389

<https://hal.science/hal-00965389>

Submitted on 25 Mar 2014

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Abelian and metabelian quotients of surface braid groups

Paolo Bellingeri, Eddy Godelle and John Guaschi

Abstract

In this paper we study Abelian and metabelian quotients of braid groups on oriented surfaces with boundary components. We provide group presentations and we prove rigidity results for these quotients arising from exact sequences related to (generalised) Fadell-Neuwirth fibrations.

1 Introduction

For $n \in \mathbb{N}$, let B_n denote the Artin braid group on n strings. The *Burau representation*, which is known not to be faithful if $n \geq 5$, and the *Bigelow-Krammer-Lawrence representation*, which is faithful [7, 25], play an important rôle in the theory of B_n [9, 24]. They arise as representatives of a larger family of representations of B_n defined via the action on a regular covering space of the k^{th} permuted configuration space of the n -punctured disc \mathbb{D}_n [29]. As we will explain in Section 5, these regular coverings are obtained from a pair (G_k, p_k) , $k \geq 1$, where $p_k : B_k(\mathbb{D}_n) \rightarrow G_k$ is the surjective homomorphism from the k -string braid group of \mathbb{D}_n onto a group G_k that is defined by the following commutative diagram of short exact sequences:

$$\begin{array}{ccccccccc}
 1 & \longrightarrow & B_k(\mathbb{D}_n) & \longrightarrow & B_{k,n} & \longrightarrow & B_n & \longrightarrow & 1 \\
 & & \downarrow p_k & & \downarrow r_{k,n} & & \downarrow r_n & & \\
 1 & \longrightarrow & G_k & \longrightarrow & B_{k,n}/\Gamma_2(B_{k,n}) & \longrightarrow & B_n/\Gamma_2(B_n) & \longrightarrow & 1,
 \end{array} \tag{1}$$

where $B_{k,n}$ is the mixed Artin braid group on (k, n) strings and $\Gamma_2(B_{k,n})$ is the second term in the lower central series of $B_{k,n}$ (recall that the *lower central series* of G is the filtration $G = \Gamma_1(G) \supseteq \Gamma_2(G) \supseteq \dots$, where $\Gamma_i(G) = [G, \Gamma_{i-1}(G)]$ for $i \geq 2$), and the vertical maps $r_{k,n}$ and r_n are the Abelianisation homomorphisms (see Section 2 for precise definitions and Section 4 for more details about this construction). Although similar constructions have been carried out for other related groups, such as Artin-Tits groups of spherical type [10, 11], their generalisation in a more topological direction, to braid and mapping class groups of surfaces for example, remains a largely open problem, with the exception of a few results [3, 8, 21].

Let $\widehat{\Sigma}_g$ be an orientable, compact surface of positive genus g and with one boundary component, and let $\widehat{\Sigma}_{g,n} = \widehat{\Sigma}_g \setminus \{x_1, \dots, x_n\}$, where $\{x_1, \dots, x_n\}$ is an n -point subset in the interior of $\widehat{\Sigma}_g$. In [1], An and Ko described an extension of the Bigelow-Krammer-Lawrence representations of B_n to surface braid groups $B_n(\widehat{\Sigma}_g)$ based on the regular covering associated to a projection map $\Phi_\Sigma : B_k(\widehat{\Sigma}_{g,n}) \rightarrow G_\Sigma$ of the braid group $B_k(\widehat{\Sigma}_{g,n})$ onto a specific group G_Σ , which can be seen as a kind of generalised Heisenberg group and that is constructed as a subgroup of a group H_Σ . The group H_Σ is defined abstractly in terms of its group presentation, and is chosen to satisfy certain technical homological constraints (Section 3.1 of [1]). An and Ko prove a rigidity result for H_Σ , which states intuitively that it is the ‘best possible’ group that satisfies the constraints. However the choices of H_Σ and G_Σ seem to be based on *ad hoc* technical arguments.

2000 AMS Mathematics Subject Classification: 20F14, 20F36. *Keywords:* surface braid groups, lower central series

Our first objective is to show that G_Σ defined in [1] may be constructed using short exact sequences of surface braid groups emanating from Fadell-Neuwirth fibrations, in which the lower central series $(\Gamma_i)_{i \in \mathbb{N}}$ of G_Σ plays a prominent rôle. These sequences are similar to those for G_k given by equation (1), but as we shall see in Lemma 5.1, there is a marked difference with the case of the Artin braid groups, since the short exact sequence on the Γ_2 -level does not yield the expected group G_Σ and homomorphism Φ_Σ . However, we prove that at the following stage, at the Γ_3 -level, the construction does indeed give rise to G_Σ . Consider the following commutative diagram of short exact sequences (the first line is the short exact sequence (SMB) that we shall recall in Section 2):

$$\begin{array}{ccccccc}
1 & \longrightarrow & B_k(\widehat{\Sigma}_{g,n}) & \longrightarrow & B_{k,n}(\widehat{\Sigma}_g) & \xrightarrow{\psi_k} & B_n(\widehat{\Sigma}_g) \longrightarrow 1 \\
& & \downarrow \Phi_k & & \downarrow \rho_{k,n} & & \downarrow \rho_n \\
1 & \longrightarrow & G_k(\widehat{\Sigma}_g) & \longrightarrow & B_{k,n}(\widehat{\Sigma}_g)/\Gamma_3(B_{k,n}(\widehat{\Sigma}_g)) & \xrightarrow{\bar{\psi}_k} & B_n(\widehat{\Sigma}_g)/\Gamma_3(B_n(\widehat{\Sigma}_g)) \longrightarrow 1,
\end{array} \tag{2}$$

where $\rho_{k,n}$ and ρ_n denote the two canonical projections, Φ_k is the restriction of $\rho_{k,n}$ to $B_k(\widehat{\Sigma}_{g,n})$, $\psi_k : B_{k,n}(\widehat{\Sigma}_g) \rightarrow B_n(\widehat{\Sigma}_g)$ is obtained geometrically by forgetting the first k strings, $\bar{\psi}_k$ is the map induced by ψ_k and $G_k(\widehat{\Sigma}_g)$ is the kernel of $\bar{\psi}_k$. We shall prove that $G_\Sigma = G_k(\widehat{\Sigma}_g)$ and $\Phi_\Sigma = \Phi_k$. More precisely:

Theorem 1.1. *Let $k, n \geq 3$. There is a canonical isomorphism of groups $\iota : G_\Sigma \rightarrow G_k(\widehat{\Sigma}_g)$. Moreover one has $\iota \circ \Phi_\Sigma = \Phi_k$.*

Our second objective is to obtain rigidity results within a completely algebraic framework for some of the groups appearing in equation (2), thus extending those of [1] mentioned above.

Theorem 1.2. *Let $k, n \geq 3$.*

(i) *Denote by $\iota_{k,n} : B_{k,n} \rightarrow B_{k,n}(\widehat{\Sigma}_g)$, $\iota_k : B_k(\mathbb{D}_n) \rightarrow B_k(\widehat{\Sigma}_{g,n})$ and $\iota_n : B_n \rightarrow B_n(\widehat{\Sigma}_g)$ the natural inclusions (see [28]). There exist injective homomorphisms $\gamma_k : G_k \rightarrow G_k(\widehat{\Sigma}_g)$, $\alpha_{k,n} : B_{k,n}/\Gamma_2(B_{k,n}) \rightarrow B_{k,n}(\widehat{\Sigma}_g)/\Gamma_3(B_{k,n}(\widehat{\Sigma}_g))$ and $\alpha_n : B_n/\Gamma_2(B_n) \rightarrow B_n(\widehat{\Sigma}_g)/\Gamma_3(B_n(\widehat{\Sigma}_g))$ so that following diagram of horizontal exact sequences is commutative:*

$$\begin{array}{ccccccc}
1 & \longrightarrow & G_k & \longrightarrow & B_{k,n}/\Gamma_2(B_{k,n}) & \longrightarrow & B_n/\Gamma_2(B_n) \longrightarrow 1 \\
& & \uparrow p_k & & \uparrow r_{k,n} & & \uparrow r_n \\
1 & \longrightarrow & B_k(\mathbb{D}_n) & \longrightarrow & B_{k,n} & \longrightarrow & B_n \longrightarrow 1 \\
& & \downarrow \iota_k & \swarrow \gamma_k & \downarrow \iota_{k,n} & \swarrow \alpha_{k,n} & \downarrow \iota_n \swarrow \alpha_n \\
1 & \longrightarrow & B_k(\widehat{\Sigma}_{g,n}) & \longrightarrow & B_{k,n}(\widehat{\Sigma}_g) & \longrightarrow & B_n(\widehat{\Sigma}_g) \longrightarrow 1 \\
& & \downarrow \Phi_k & & \downarrow \rho_{k,n} & & \downarrow \rho_n \\
1 & \longrightarrow & G_k(\widehat{\Sigma}_g) & \longrightarrow & B_{k,n}(\widehat{\Sigma}_g)/\Gamma_3(B_{k,n}(\widehat{\Sigma}_g)) & \longrightarrow & B_n(\widehat{\Sigma}_g)/\Gamma_3(B_n(\widehat{\Sigma}_g)) \longrightarrow 1,
\end{array} \tag{3}$$

where the rows are the short exact sequences of the commutative diagrams (1) and (2).

(ii) *Let G be a group, and $\Phi_G : B_k(\widehat{\Sigma}_{g,n}) \rightarrow G$ be a surjective homomorphism whose restriction to the group $B_k(\mathbb{D}_n)$ induces an injective homomorphism from G_k to G . Then $G = G_k(\widehat{\Sigma}_g)$ up to isomorphism; more precisely, $\Phi_G = \theta_G \circ \Phi_k$ where θ_G is an isomorphism.*

- (iii) Let H be a group, and $\rho_H : B_{k,n}(\widehat{\Sigma}_g) \rightarrow H$ be a surjective homomorphism whose restriction to the group $B_{k,n}$ induces an injective homomorphism from $B_{k,n}/\Gamma_2(B_{k,n})$ to H . Then $H = B_{k,n}(\widehat{\Sigma}_g)/\Gamma_3(B_{k,n}(\widehat{\Sigma}_g))$ up to isomorphism; more precisely, $\rho_H = \theta_H \circ \rho_{k,n}$ where θ_H is an isomorphism.
- (iv) Let K be a group, and $\rho_K : B_n(\widehat{\Sigma}_g) \rightarrow K$ be a surjective homomorphism whose restriction to the group B_n induces an injective homomorphism from $B_n/\Gamma_2(B_n)$ to K . Then $K = B_n(\widehat{\Sigma}_g)/\Gamma_3(B_n(\widehat{\Sigma}_g))$ up to isomorphism; more precisely, $\rho_K = \theta_K \circ \rho_n$ where θ_K is an isomorphism.

We will in fact obtain a stronger result, by proving that some of the above results remain true when the assumptions on Φ_G , ρ_K and ρ_H are relaxed. Remarking that the group H_Σ of [1] is a quotient of $B_{k,n}(\widehat{\Sigma}_g)/\Gamma_3(B_{k,n}(\widehat{\Sigma}_g))$, in Proposition 4.5, we exhibit an alternative proof of [1, Theorem 4.3].

This paper is organised as follows. In Section 2 we recall the definitions of (mixed) surface braid groups and their associated short exact sequences. The first part of Section 3 is devoted to obtaining presentations of mixed surface braid groups. In Section 3.2 we describe the Abelianisations of (mixed) surface braid groups (Propositions 3.3 and 3.4) and we show in particular that it is not possible to embed $B_{k,n}/\Gamma_2(B_{k,n})$ in any Abelian quotient of $B_{k,n}(\widehat{\Sigma}_g)$. In Section 3.3, we obtain similar results but at the level of quotients of surface mixed braid groups by Γ_3 rather than by Γ_2 . We give a presentation for $B_{k,n}(\widehat{\Sigma}_g)/\Gamma_3(B_{k,n}(\widehat{\Sigma}_g))$ (Proposition 3.7) and a normal form for elements of this quotient (Corollary 3.9), and we prove several rigidity results for (metabelian) quotients of $B_{k,n}(\widehat{\Sigma}_g)$ (Corollary 3.11). In particular, we show that $B_{k,n}/\Gamma_2(B_{k,n})$ embeds in $B_{k,n}(\widehat{\Sigma}_g)/\Gamma_3(B_{k,n}(\widehat{\Sigma}_g))$ and we deduce Theorem 1.2(iii). Similar results are given in Section 3.4 for surface braid groups, the main result being Corollary 3.11, which implies Theorem 1.2(i) and (iv).

In Section 4, we prove Theorem 1.1 and Theorem 1.2(ii), and we show that it is not possible to extend the length function $\lambda : B_n \rightarrow \mathbb{Z}$ to braid groups of closed oriented surfaces (Proposition 4.5). Finally in Section 5, we describe an algebraic approach to the Burau and Bigelow-Krammer-Lawrence representations that is based on the lower central series, and we explain why it is not possible to extend them to representations of surface braid groups. This latter remark was made in [1] under certain conditions of a homological nature. Within a purely algebraic framework, we prove this non-existence result with fewer conditions than those given in [1].

2 Preliminaries on configuration spaces

Surface braid groups are a natural generalisation of both the classical braid groups and the fundamental group of surfaces. We recall Fox's definition in terms of fundamental groups of configuration spaces [15]. Let Σ be a connected surface, and let $\mathbb{F}_n(\Sigma) = \Sigma^n \setminus \Delta$, where Δ is the set of n -tuples $(x_1, \dots, x_n) \in \Sigma^n$ for which $x_i = x_j$ for some $i \neq j$. The fundamental group $\pi_1(\mathbb{F}_n(\Sigma))$ is called the *pure braid group* on n strings of the surface Σ and shall be denoted by $P_n(\Sigma)$. The symmetric group S_n acts freely on $\mathbb{F}_n(\Sigma)$ by permutation of coordinates, and the fundamental group $\pi_1(\mathbb{F}_n(\Sigma)/S_n)$ of the resulting quotient space, denoted by $B_n(\Sigma)$, is the *braid group* on n strings of the surface Σ . Further, $\mathbb{F}_n(\Sigma)$ is a regular $n!$ -fold covering of $\mathbb{F}_n(\Sigma)/S_n$, from which we obtain the following short exact sequence:

$$1 \rightarrow P_n(\Sigma) \rightarrow B_n(\Sigma) \rightarrow S_n \rightarrow 1. \quad (4)$$

In the case of the disc \mathbb{D}^2 , it is well known that $B_n(\mathbb{D}^2) \cong B_n$ and $P_n(\mathbb{D}^2) \cong P_n$.

Let $k, n \in \mathbb{N}$. Regarded as a subgroup of S_{k+n} , the group $S_k \times S_n$ acts on $\mathbb{F}_{k+n}(\Sigma)$. The fundamental group $\pi_1(\mathbb{F}_{k+n}(\Sigma)/(S_k \times S_n))$ will be called the *mixed braid group of Σ on (k, n) strings*, and shall be denoted by $B_{k,n}(\Sigma)$. We shall denote $B_{k,n}(\mathbb{D}^2)$ simply by $B_{k,n}$. In an obvious manner, we have $P_{k+n}(\Sigma) \subset B_{k,n}(\Sigma) \subset B_{k+n}(\Sigma)$. Mixed braid groups, which play an important rôle in [1], were defined previously in [17, 27, 28], and were studied in more detail in [18] in the case where Σ is the 2-sphere \mathbb{S}^2 .

Consider the Fadell-Neuwirth fibration $\mathbb{F}_{k+n}(\Sigma) \rightarrow \mathbb{F}_n(\Sigma)$ given by forgetting the first k coordinates. Its long exact sequence in homotopy yields the *surface pure braid group* short exact sequence [12]:

$$1 \rightarrow P_k(\Sigma \setminus \{x_1, \dots, x_n\}) \rightarrow P_{k+n}(\Sigma) \rightarrow P_n(\Sigma) \rightarrow 1, \quad (\text{SPB})$$

where $n \geq 3$ (resp. $n \geq 2$) if $\Sigma = \mathbb{S}^2$ (resp. Σ is the projective plane $\mathbb{R}P^2$). In a similar manner, the map $\mathbb{F}_{k+n}(\Sigma)/(S_k \times S_n) \rightarrow \mathbb{F}_n(\Sigma)/S_n$ defined by forgetting the first k coordinates, is a locally-trivial fibration whose fibre may be identified with $\mathbb{F}_k(\Sigma \setminus \{x_1, \dots, x_n\})/S_k$. With the same constraints on n if $\Sigma = \mathbb{S}^2$ or $\mathbb{R}P^2$, this fibration gives rise to the *surface mixed braid group* short exact sequence:

$$1 \rightarrow B_k(\Sigma \setminus \{x_1, \dots, x_n\}) \rightarrow B_{k,n}(\Sigma) \xrightarrow{\psi_k} B_n(\Sigma) \rightarrow 1, \quad (\text{SMB})$$

where ψ_k is the epimorphism given in diagram 2 that may be interpreted geometrically by forgetting the first k strings. Note that (SPB) is the restriction of (SMB) to the corresponding pure braid groups. From now on, we denote $\mathbb{D}^2 \setminus \{x_1, \dots, x_n\}$ by \mathbb{D}_n .

The group $B_k(\mathbb{D}_n)$ (resp. $P_k(\mathbb{D}_n)$) may be seen to be isomorphic to the subgroup of B_{k+n} (resp. P_{k+n}) consisting of braids whose last n strings are trivial (vertical). As we shall see in Section 4, one important ingredient in the construction of representations of the Artin braid groups is the splitting of the short exact sequences (SMB) and (SPB) when $\Sigma = \mathbb{D}^2$; in both cases, a section, which we refer to henceforth as the *standard section*, is given by adding k vertical strings (see for instance [1, 4]). With the aim of obtaining representations of the braid groups of Σ , it is thus natural to ask in which cases these sequences split. Note that there is a commutative diagram of short exact sequences, where the first line is (SPB), the second line is (SMB), and the third line is:

$$1 \rightarrow S_k \rightarrow S_k \times S_n \rightarrow S_n \rightarrow 1.$$

The question of the splitting of (SPB) has been solved completely (see [20] for a summary). In particular, if Σ is a compact surface without boundary and different from \mathbb{S}^2 and $\mathbb{R}P^2$ then (SPB) only splits if $n = 1$, and one may show that this implies the splitting of (SMB) in this case. Using the methods of [16], it follows that both (SPB) and (SMB) split if Σ has non-empty boundary. Some partial results for the splitting of (SMB) are known if Σ has empty boundary (see for example [13, 18] for the case of \mathbb{S}^2), but in general the question remains unanswered.

3 Surface braid groups and lower central series

As in the Introduction, let $\widehat{\Sigma}_g$ be a compact, connected orientable surface of genus $g \geq 0$ with a single boundary component, and let $k \geq 1$ and $n \geq 0$. We will make use of the notation introduced in the commutative diagrams (2) and (3). We focus on $\widehat{\Sigma}_g$ essentially for two reasons; the first is that as we mentioned in Section 2, the short exact sequence (SMB) for $\widehat{\Sigma}_g$ splits and therefore $B_n(\widehat{\Sigma}_g)$ acts by conjugation on $B_k(\widehat{\Sigma}_{g,n})$. The second reason is that Theorem 1.2(iv) is not valid if we replace $\widehat{\Sigma}_g$ by a compact surface without boundary (this fact will be a straightforward consequence of Proposition 4.5).

In this section, we shall prove parts (i), (iii) and (iv) of Theorem 1.2. Taking into account the commutative diagrams (1) and (2) as well as the following commutative diagram:

$$\begin{array}{ccccccc} 1 & \longrightarrow & B_k(\mathbb{D}_n) & \longrightarrow & B_{k,n} & \longrightarrow & B_n \longrightarrow 1 \\ & & \downarrow \iota_k & & \downarrow \iota_{k,n} & & \downarrow \iota_n \\ 1 & \longrightarrow & B_k(\widehat{\Sigma}_{g,n}) & \longrightarrow & B_{k,n}(\widehat{\Sigma}_g) & \longrightarrow & B_n(\widehat{\Sigma}_g) \longrightarrow 1, \end{array} \quad (5)$$

to prove Theorem 1.2(i), it will suffice to show the existence of the homomorphisms $\gamma_k, \alpha_k, \alpha_{k,n}$ and α_n , and to verify commutativity in the vertical parts of the commutative diagram (3). The main result of

this section is Proposition 3.6, which is a stronger version of Theorem 1.2(iii). In Section 3.1, we start by exhibiting a presentation for $B_{k,n}(\widehat{\Sigma}_g)$. In what follows, we will consider the following disjoint sets:

$$\begin{aligned} S &= \{\sigma_1, \dots, \sigma_{k-1}\}, & \widetilde{S} &= \{\widetilde{\sigma}_1, \dots, \widetilde{\sigma}_{n-1}\}, \\ AB &= \{a_1, b_1, \dots, a_g, b_g\}, & \widetilde{AB} &= \{\widetilde{a}_1, \widetilde{b}_1, \dots, \widetilde{a}_g, \widetilde{b}_g\}, \\ Z &= \{\zeta_1, \dots, \zeta_n\}. \end{aligned}$$

If $k = 1$ (resp. $n = 0, n = 1, g = 0$) then S (resp. $\widetilde{S} \cup Z, \widetilde{S}, AB \cup \widetilde{AB}$) is taken to be empty. For $c, d \in AB$ we write $c < d$ if $c \in \{a_i, b_i\}$ and $d \in \{a_j, b_j\}$ with $i < j$. Similarly, for $c, d \in \widetilde{AB}$ we write $c < d$ if $c \in \{\widetilde{a}_i, \widetilde{b}_i\}$ and $d \in \{\widetilde{a}_j, \widetilde{b}_j\}$ with $i < j$. If x and y are elements of a group then we set $x^y = y^{-1}xy$ and $[x, y] = xyx^{-1}y^{-1}$.

3.1 Presentations of surface mixed braid groups

In order to prove Theorem 1.2, we will need to understand the structure of surface (mixed) braid groups and some of their quotients. With this in mind, in this section, we recall presentations of $B_k(\widehat{\Sigma}_{g,n})$ and $B_n(\widehat{\Sigma}_g)$, and we derive a presentation of $B_{k,n}(\widehat{\Sigma}_g)$.

If $g = 0$, part (i) of the following result is proved in [22]. If $g \geq 1$, the presentations of parts (i) and (ii) may be found in [4]. From hereon, if $g = 0$ then the elements of $AB \cup \widetilde{AB}$ and all relations containing these elements should be suppressed.

Proposition 3.1. *Let $k, n \geq 1$, and let $g \geq 0$.*

(i) *The group $B_k(\widehat{\Sigma}_{g,n})$ admits the following group presentation:*

Generating set: $S \cup AB \cup Z$;

Relations:

$$\begin{aligned} (a.1) \quad & \sigma_i \sigma_j = \sigma_j \sigma_i, & |i - j| \geq 2; \\ (a.2) \quad & \sigma_i \sigma_{i+1} \sigma_i = \sigma_{i+1} \sigma_i \sigma_{i+1}, & 1 \leq i \leq k - 2; \\ (a.3) \quad & c \sigma_i = \sigma_i c, & i \neq 1, c \in AB \cup Z; \\ (a.4) \quad & c \sigma_1 c \sigma_1 = \sigma_1 c \sigma_1 c, & c \in AB \cup Z; \\ (a.5) \quad & a_i \sigma_1 b_i = \sigma_1 b_i \sigma_1 a_i \sigma_1, & i \in \{1, \dots, g\}; \\ (a.6) \quad & (\sigma_1^{-1} c \sigma_1) d = d (\sigma_1^{-1} c \sigma_1), & c, d \in AB, c < d; \\ (a.7) \quad & (\sigma_1^{-1} \zeta_i \sigma_1) c = c (\sigma_1^{-1} \zeta_i \sigma_1), & c \in AB, \zeta_i \in Z; \\ (a.8) \quad & (\sigma_1^{-1} \zeta_i \sigma_1) \zeta_j = \zeta_j (\sigma_1^{-1} \zeta_i \sigma_1), & i < j. \end{aligned}$$

(ii) *The group $B_n(\widehat{\Sigma}_g)$ admits the following group presentation:*

Generating set: $\widetilde{S} \cup \widetilde{AB}$;

Relations:

$$\begin{aligned} (b.1) \quad & \widetilde{\sigma}_i \widetilde{\sigma}_j = \widetilde{\sigma}_j \widetilde{\sigma}_i, & |i - j| \geq 2; \\ (b.2) \quad & \widetilde{\sigma}_i \widetilde{\sigma}_{i+1} \widetilde{\sigma}_i = \widetilde{\sigma}_{i+1} \widetilde{\sigma}_i \widetilde{\sigma}_{i+1}, & 1 \leq i \leq n - 2; \\ (b.3) \quad & \widetilde{c} \widetilde{\sigma}_i = \widetilde{\sigma}_i \widetilde{c}, & i \neq 1, \widetilde{c} \in \widetilde{AB}; \\ (b.4) \quad & \widetilde{c} \widetilde{\sigma}_1 \widetilde{c} \widetilde{\sigma}_1 = \widetilde{\sigma}_1 \widetilde{c} \widetilde{\sigma}_1 \widetilde{c}, & \widetilde{c} \in \widetilde{AB}; \\ (b.5) \quad & \widetilde{a}_i \widetilde{\sigma}_1 \widetilde{b}_i = \widetilde{\sigma}_1 \widetilde{b}_i \widetilde{\sigma}_1 \widetilde{a}_i \widetilde{\sigma}_1, & i \in \{1, \dots, g\}; \\ (b.6) \quad & (\widetilde{\sigma}_1^{-1} \widetilde{c} \widetilde{\sigma}_1) \widetilde{d} = \widetilde{d} (\widetilde{\sigma}_1^{-1} \widetilde{c} \widetilde{\sigma}_1), & \widetilde{c}, \widetilde{d} \in \widetilde{AB}, \widetilde{c} < \widetilde{d}. \end{aligned}$$

The presentation of part (i) may be adapted to the case $n = 0$ by suppressing Z in the presentation of part (ii). However, to obtain a presentation of $B_{k,n}(\widehat{\Sigma}_g)$ from Proposition 3.1, it will be convenient to have both presentations at our disposal. If $g = 0$ then $B_{k,n}(\widehat{\Sigma}_g)$ is equal to $B_{k,n}$. An alternative presentation of $B_{k,n}(\widehat{\Sigma}_g)$ may be found in [1]; in the special case of $B_{k,n}$, see [14].

Proposition 3.2. *Let $k, n \geq 1$, and let $g \geq 0$. The group $B_{k,n}(\widehat{\Sigma}_g)$ admits the following group presentation:*

Generating set: $\Omega_{k,n} = S \cup \widetilde{S} \cup AB \cup \widetilde{A}\widetilde{B} \cup Z$;

Relations:

(a) the relations (a.1)–(a.8) given in Proposition 3.1(i).

(b) the relations (b.1)–(b.6) given in Proposition 3.1(ii).

(c) the relations that describe the action of $B_n(\widehat{\Sigma}_g)$ on $B_k(\widehat{\Sigma}_{g,n})$:

$$(c.1) \quad \widetilde{\sigma}_i \sigma_j \widetilde{\sigma}_i^{-1} = \widetilde{a}_i \sigma_j \widetilde{a}_i^{-1} = \widetilde{b}_i \sigma_j \widetilde{b}_i^{-1} = \sigma_j;$$

$$(c.2) \quad \widetilde{\sigma}_i a_j \widetilde{\sigma}_i^{-1} = a_j, \quad \widetilde{\sigma}_i b_j \widetilde{\sigma}_i^{-1} = b_j;$$

$$(c.3) \quad \left\{ \begin{array}{l} (c.3.1) \quad \widetilde{\sigma}_i \zeta_{i+1} \widetilde{\sigma}_i^{-1} = \zeta_i; \\ (c.3.2) \quad \widetilde{\sigma}_i \zeta_i \widetilde{\sigma}_i^{-1} = \zeta_i^{-1} \zeta_{i+1} \zeta_i; \\ (c.3.3) \quad \widetilde{\sigma}_i \zeta_j \widetilde{\sigma}_i^{-1} = \zeta_j, \end{array} \right. \quad j \neq i, i+1; \quad (c.4) \quad \left\{ \begin{array}{l} (c.4.1) \quad \widetilde{a}_i \zeta_1 \widetilde{a}_i^{-1} = \zeta_1^{a_i \zeta_1}; \\ (c.4.2) \quad \widetilde{b}_i \zeta_1 \widetilde{b}_i^{-1} = \zeta_1^{b_i \zeta_1}; \\ (c.4.3) \quad \widetilde{a}_i \zeta_j \widetilde{a}_i^{-1} = \zeta_j^{[a_i^{-1}, \zeta_1^{-1}]}, \quad j \neq 1; \\ (c.4.4) \quad \widetilde{b}_i \zeta_j \widetilde{b}_i^{-1} = \zeta_j^{[b_i^{-1}, \zeta_1^{-1}]}, \quad j \neq 1; \end{array} \right.$$

$$(c.5) \quad \left\{ \begin{array}{l} (c.5.1) \quad \widetilde{a}_i a_i \widetilde{a}_i^{-1} = \zeta_1^{-1} a_i \zeta_1; \\ (c.5.2) \quad \widetilde{a}_i a_j \widetilde{a}_i^{-1} = a_j^{[a_i^{-1}, \zeta_1^{-1}]}, \quad i > j; \\ (c.5.3) \quad \widetilde{a}_i a_j \widetilde{a}_i^{-1} = a_j, \quad j > i; \end{array} \right. \quad (c.6) \quad \left\{ \begin{array}{l} (c.6.1) \quad \widetilde{b}_i b_i \widetilde{b}_i^{-1} = \zeta_1^{-1} b_i \zeta_1; \\ (c.6.2) \quad \widetilde{b}_i b_j \widetilde{b}_i^{-1} = b_j^{[b_i^{-1}, \zeta_1^{-1}]}, \quad i > j; \\ (c.6.3) \quad \widetilde{b}_i b_j \widetilde{b}_i^{-1} = b_j, \quad j > i; \end{array} \right.$$



$$(c.7) \quad \left\{ \begin{array}{l} (c.7.1) \quad \widetilde{a}_i b_i \widetilde{a}_i^{-1} = b_i \zeta_1; \\ (c.7.2) \quad \widetilde{a}_i b_j \widetilde{a}_i^{-1} = b_j^{[a_i^{-1}, \zeta_1^{-1}]}, \quad i > j; \\ (c.7.3) \quad \widetilde{a}_i b_j \widetilde{a}_i^{-1} = b_j, \quad j > i \end{array} \right. \quad (c.8) \quad \left\{ \begin{array}{l} (c.8.1) \quad \widetilde{b}_i a_i \widetilde{b}_i^{-1} = \zeta_1^{-1} a_i [b_i^{-1}, \zeta_1^{-1}]; \\ (c.8.2) \quad \widetilde{b}_i a_j \widetilde{b}_i^{-1} = a_j^{[b_i^{-1}, \zeta_1^{-1}]}, \quad i > j; \\ (c.8.3) \quad \widetilde{b}_i a_j \widetilde{b}_i^{-1} = a_j, \quad j > i. \end{array} \right.$$

Proof. If $g \geq 1$, we first give a geometric interpretation of the generators of $B_{k,n}(\widehat{\Sigma}_g)$. We represent $\widehat{\Sigma}_g$ as a regular polygon with $4g$ edges, equipped with the standard identification of edges, and one boundary component. We consider braids to be paths on the polygon, which we draw with the usual over- and under-crossings, and we interpret the braids depicted in Figure 1 as geometric representatives of the generators of $B_k(\widehat{\Sigma}_{g,n})$, and those depicted in Figure 2 as the coset representatives of generators of $B_n(\widehat{\Sigma}_g)$ in $B_{k,n}(\widehat{\Sigma}_g)$. For example, for the braid a_r (respectively b_r), the only non-trivial string is the first one, which passes through the wall α_r (respectively the wall β_r). If $g \geq 0$, one can therefore check that relations hold for corresponding geometric braids, see Figure 3 for example.

From Section 2, the short exact sequence (SMB) splits. The group $B_{k,n}(\widehat{\Sigma}_g)$ is thus isomorphic to a semi-direct product $B_n(\widehat{\Sigma}_g) \ltimes B_k(\widehat{\Sigma}_{g,n})$. Proposition 3.1 implies that the set of relations (a) and (b) provide a complete set of relations for $B_k(\widehat{\Sigma}_{g,n})$ and $B_n(\widehat{\Sigma}_g)$ respectively. The set of relations (c) describes the action of the generators of $B_k(\widehat{\Sigma}_{g,n})$ on those of $B_n(\widehat{\Sigma}_g)$. The set of relations (a)–(c) therefore form a complete set of relations for $B_{k,n}(\widehat{\Sigma}_g)$ by [23, Chap. 10, Proposition 1]. \square

3.2 Abelian quotients of surface mixed braid groups

In this section, we use Propositions 3.1 and 3.2 to describe the Abelianisations of the (mixed) surface braid groups that arise in our study. We start with the case of mixed surface braid groups. We believe that the case $g = 0$ is well known to the experts in the field, but since we were not able to find a reference in the literature, we provide a short proof.


 Figure 1: The generators $\sigma_1, \dots, \sigma_{k-1}, a_1, b_1, \dots, a_g, b_g, \zeta_1, \dots, \zeta_n$

 Figure 2: The generators $\tilde{\sigma}_1, \dots, \tilde{\sigma}_{n-1}, \tilde{a}_1, \tilde{b}_1, \dots, \tilde{a}_g, \tilde{b}_g$

Proposition 3.3. Let $n, k \geq 1$ and $g \geq 0$, let $\hat{r}_{k,n} : B_{k,n}(\hat{\Sigma}_g) \rightarrow B_{k,n}(\hat{\Sigma}_g) / \Gamma_2(B_{k,n}(\hat{\Sigma}_g))$ denote the canonical projection (if $g = 0$ then $\hat{r}_{k,n}$ is the homomorphism $r_{k,n}$), and let


$$\hat{S} = \begin{cases} \emptyset & \text{if } n = k = 1 \\ \{\sigma\} & \text{if } k \geq 2 \text{ and } n = 1 \\ \{\tilde{\sigma}\} & \text{if } k = 1 \text{ and } n \geq 2 \\ \{\sigma, \tilde{\sigma}\} & \text{if } k, n \geq 2 \end{cases} \quad \text{and} \quad \hat{Z} = \begin{cases} \{\zeta\} & \text{if } g = 0 \\ \emptyset & \text{if } g \geq 1. \end{cases}$$

Then the group $B_{k,n}(\hat{\Sigma}_g) / \Gamma_2(B_{k,n}(\hat{\Sigma}_g))$ admits the following group presentation:

Generating set: $\hat{r}_{k,n}(S \cup \tilde{S} \cup Z \cup AB \cup \bar{A}\bar{B}) = \hat{S} \cup \hat{Z} \cup AB \cup \bar{A}\bar{B}$;

Relations:

- $xy = yx$ for $x, y \in \hat{S} \cup \hat{Z} \cup AB \cup \bar{A}\bar{B}$, $x \neq y$;
- if $g \geq 1$ then $s^2 = 1$ for all $s \in \hat{S}$.


 Figure 3: The braids $\tilde{a}_i b_i \tilde{a}_i^{-1}$ and $b_i \zeta_i$ are isotopic.

In particular, $\widehat{r}_{k,n}(S) = \{\sigma\}$ if $k \geq 2$, $\widehat{r}_{k,n}(\widetilde{S}) = \{\widetilde{\sigma}\}$ if $n \geq 2$, $\widehat{r}_{k,n}(Z) = \{\zeta\}$ if $g = 0$, and the group $B_{k,n}(\widehat{\Sigma}_g)/\Gamma_2(B_{k,n}(\widehat{\Sigma}_g))$ is isomorphic to $\mathbb{Z}^{|\widehat{S}|+1}$ if $g = 0$, and to $\mathbb{Z}^{4g} \times \mathbb{Z}_2^{|\widehat{S}|}$ if $g \geq 1$, where $|\widehat{S}|$ denotes the cardinal of \widehat{S} .

Proof. Consider the presentation of $B_{k,n}(\widehat{\Sigma}_g)$ given by Proposition 3.2. To obtain a presentation of the quotient $B_{k,n}(\widehat{\Sigma}_g)/\Gamma_2(B_{k,n}(\widehat{\Sigma}_g))$, we must add the relations of the form $xy = yx$ for all x, y in $\Omega_{k,n}$ to the presentation of $B_{k,n}(\widehat{\Sigma}_g)$. First suppose that $g = 0$. Then none of the relations (c) of Proposition 3.2 exist, with the exception of relation (c.3). The group $B_{k,n} = B_{k,n}(\widehat{\Sigma}_0)$ is generated by $S \cup \widetilde{S} \cup Z$. Under Abelianisation, if $k \geq 2$ (resp. $n \geq 2$), the elements of S (resp. \widetilde{S} , Z) are identified to a single element σ (resp. $\widetilde{\sigma}$, ζ) by relation (a.2) (resp. relation (b.2), relation (c.3)), so the given generating set of $B_{k,n}$ reduces to the generating set $\widehat{S} \cup \widehat{Z}$ of $B_{k,n}/\Gamma_2(B_{k,n})$, and the only defining relations are commutation relations. So the statement holds in the case $g = 0$. Assume now that $g \geq 1$. For the elements of $\widehat{S} \cup \widehat{Z}$, the same analysis holds as in the case $g = 0$. Additionally, relation (c.7.1) implies that $\widehat{r}_{k,n}(\zeta_1)$ is trivial, so $\widehat{Z} = \emptyset$, and thus the relations of type (c) do not give any extra information in $B_{k,n}(\widehat{\Sigma}_g)/\Gamma_2(B_{k,n}(\widehat{\Sigma}_g))$. Under $\widehat{r}_{k,n}$, if $k \geq 2$ (resp. $n \geq 2$), relations (a.1)–(a.8) (resp. relations (b.1)–(b.6)) do not yield any new relations, with the exception of relation (a.5) (resp. relation (b.5)) that reduces to $\sigma^2 = 1$ (resp. $\widetilde{\sigma}^2 = 1$) in $B_{k,n}(\widehat{\Sigma}_g)/\Gamma_2(B_{k,n}(\widehat{\Sigma}_g))$, and the result follows. If $k = 1$ (resp. $n = 1$) then relations (a.1)–(a.8) (resp. relations (b.1)–(b.6)) do not exist, and again we see that the statement holds, which completes the proof of the case $g \geq 1$. \square

For $k, g \geq 1$ and $n \geq 0$, let $\widehat{r}_k : B_k(\widehat{\Sigma}_{g,n}) \rightarrow B_k(\widehat{\Sigma}_{g,n})/\Gamma_2(B_k(\widehat{\Sigma}_{g,n}))$ denote the canonical projection. It will be convenient to denote the disc \mathbb{D}^2 by \mathbb{D}_0 , so $B_k(\mathbb{D}_0) = B_k$. Note once more that if $k = 1$ (resp. $n = 0, g = 0$) then all references to the element σ (resp. to the set Z , to the set AB) in the generators and relations should be suppressed.

Proposition 3.4. *Let $k \geq 1$ and $g, n \geq 0$. The group $B_k(\widehat{\Sigma}_{g,n})/\Gamma_2(B_k(\widehat{\Sigma}_{g,n}))$ admits the following presentation:*

Generating set: $\widehat{r}_k(S \cup AB \cup Z) = \{\sigma\} \cup AB \cup Z$;

Relations:

- $xy = yx$ for all $x, y \in \{\sigma\} \cup AB \cup Z$, $x \neq y$;
- if $g \geq 1$ then $\sigma^2 = 1$.

Consequently, $B_k(\mathbb{D}_n)/\Gamma_2(B_k(\mathbb{D}_n))$ is free Abelian on the set $\widehat{r}_k(S \cup Z) = \{\sigma\} \cup Z$, and if $g \geq 1$, the group $B_k(\widehat{\Sigma}_{g,n})/\Gamma_2(B_k(\widehat{\Sigma}_{g,n}))$ is isomorphic to $\mathbb{Z}^{2g+n} \times \mathbb{Z}_2$ if $k \geq 2$ and to \mathbb{Z}^{2g+n} if $k = 1$.

If $n = 0$, the result may be found in [5]. The proof of Proposition 3.4 follows in a similar manner to that of Proposition 3.3 using the presentation of $B_k(\widehat{\Sigma}_{g,n})$ given by Proposition 3.1.

For $k \geq 1$ and $n \geq 0$, let $r_k : B_k(\mathbb{D}_n) \rightarrow B_k(\mathbb{D}_n)/\Gamma_2(B_k(\mathbb{D}_n))$ denote the canonical projection. We now apply Propositions 3.3 and 3.4 to analyse the homomorphisms $\bar{\iota}_{k,n} : B_{k,n}/\Gamma_2(B_{k,n}) \rightarrow B_{k,n}(\widehat{\Sigma}_g)/\Gamma_2(B_{k,n}(\widehat{\Sigma}_g))$ and $\bar{\iota}_k : B_k(\mathbb{D}_n)/\Gamma_2(B_k(\mathbb{D}_n)) \rightarrow B_k(\widehat{\Sigma}_{g,n})/\Gamma_2(B_k(\widehat{\Sigma}_{g,n}))$ induced by the homomorphisms $\iota_{k,n} : B_{k,n} \rightarrow B_{k,n}(\widehat{\Sigma}_g)$ and $\iota_k : B_k(\mathbb{D}_n) \rightarrow B_k(\widehat{\Sigma}_{g,n})$ on the corresponding Abelianisations. These homomorphisms thus satisfy the relations $\widehat{r}_{k,n} \circ \iota_{k,n} = \bar{\iota}_{k,n} \circ r_{k,n}$ and $\widehat{r}_k \circ \iota_k = \bar{\iota}_k \circ r_k$.

Corollary 3.5. *Let $g, k \geq 1$, and let G be an Abelian group.*

- (i) *Let $n \geq 1$, and let $P_G : B_{k,n}(\widehat{\Sigma}_g) \rightarrow G$ and $\overline{P}_G : B_{k,n}/\Gamma_2(B_{k,n}) \rightarrow G$ be homomorphisms for which $P_G \circ \iota_{k,n} = \overline{P}_G \circ r_{k,n}$. Then \overline{P}_G is not injective. In particular, the homomorphism $\bar{\iota}_{k,n}$ is not injective.*
- (ii) *Let $n \geq 0$.*

- (a) If $k = 1$ then the homomorphism $\bar{\iota}_k$ is injective.
- (b) Suppose that $k \geq 2$. Let $P_G : B_k(\widehat{\Sigma}_{g,n}) \rightarrow G$ and $\overline{P}_G : B_k(\mathbb{D}_n)/\Gamma_2(B_k(\mathbb{D}_n)) \rightarrow G$ be homomorphisms for which $P_G \circ \iota_k = \overline{P}_G \circ r_k$. Then \overline{P}_G is not injective. In particular, the homomorphism $\bar{\iota}_k$ is not injective.

Proof.

- (i) We start by proving the non-injectivity of $\bar{\iota}_{k,n}$. Suppose that $G = B_{k,n}(\widehat{\Sigma}_g)/\Gamma_2(B_{k,n}(\widehat{\Sigma}_g))$ and $P_G = \widehat{r}_{k,n}$, and let $\overline{P}_G = \bar{\iota}_{k,n}$. Thus $\widehat{r}_{k,n} \circ \iota_{k,n} = \bar{\iota}_{k,n} \circ r_{k,n}$, and so $\bar{\iota}_{k,n}$ cannot be injective because $r_{k,n}(\zeta_1) \neq 1$ and $\widehat{r}_{k,n}(\zeta_1) = 1$ by Proposition 3.3. We now consider the general case. Let P_G and \overline{P}_G be as in the statement. Since G is Abelian, P_G factors through $\widehat{r}_{k,n}$, so as in the case of $\bar{\iota}_{k,n}$, we have $P_G(\zeta_1) = 1$ and $r_{k,n}(\zeta_1) \neq 1$, which implies the non-injectivity of \overline{P}_G using the relation $P_G \circ \iota_{k,n} = \overline{P}_G \circ r_{k,n}$.
- (ii) (a) If $k = 1$ then Proposition 3.4 implies that $B_k(\mathbb{D}_n)/\Gamma_2(B_k(\mathbb{D}_n))$ is a free Abelian group generated by Z (Z may be empty if $n = 0$), and that the homomorphism $\bar{\iota}_k$ identifies $B_k(\mathbb{D}_n)/\Gamma_2(B_k(\mathbb{D}_n))$ with the direct factor of $B_k(\widehat{\Sigma}_{g,n})/\Gamma_2(B_k(\widehat{\Sigma}_{g,n}))$ generated by Z , so $\bar{\iota}_k$ is injective.
- (b) If $k \geq 2$ then the argument is similar to that of part (i), where we replace ζ_1 by σ_1^2 . \square

3.3 Metabelian quotients of surface mixed braid groups

In this section, the aim is to obtain results similar to those of Section 3.2, but on the level of quotients by Γ_3 rather than by Γ_2 . With Section 4 in mind, our principal interest is in detecting the differences between these two types of quotient.

The following result will provide the crucial argument in the proof of Theorems 1.1 and 1.2.

Proposition 3.6. *Let $k, n \geq 3$, and let $g \geq 0$. Let H be a group, and let $\rho_H : B_{k,n}(\widehat{\Sigma}_g) \rightarrow H$ be a surjective homomorphism. Consider the presentation of $B_{k,n}(\widehat{\Sigma}_g)$ given in Proposition 3.2, and let R be a set of words on $\Omega_{k,n} \cup \Omega_{k,n}^{-1}$ whose normal closure in $B_{k,n}(\widehat{\Sigma}_g)$ is equal to $\ker(\rho_H)$. Assume that there exist $\sigma, \tilde{\sigma}$ in H such that*

$$\rho_H(S) = \{\sigma\} \text{ and } \rho_H(\tilde{S}) = \{\tilde{\sigma}\}.$$

Then the following assertions hold.

- (i) There exists $\zeta \in H$ such that $\rho_H(Z) = \{\zeta\}$.
- (ii) Let R_Ω denote the set of words obtained from R by replacing each of the letters $\sigma_i, \tilde{\sigma}_i$ and ζ_i by $\sigma, \tilde{\sigma}$ and ζ respectively (and their inverses by $\sigma^{-1}, \tilde{\sigma}^{-1}$ and ζ^{-1} respectively). Then the group H possesses the following presentation:
Generating set: $\Omega = \rho_H(\Omega_{k,n}) = \{\sigma, \tilde{\sigma}, \zeta\} \cup AB \cup \widetilde{AB}$;
Relations:

- $w = 1$ for all $w \in R_\Omega$;
- $xy = yx$ for all $x, y \in \Omega$, $x \neq y$, and $\{x, y\} \notin \left\{ \{a_i, b_i\}, \{\tilde{a}_i, \tilde{b}_i\}, \{b_i, \tilde{a}_i\}, \{\tilde{b}_i, a_i\}; i = 1, \dots, g \right\}$;
- $[a_i, b_i] = \sigma^2$; $[\tilde{a}_i, \tilde{b}_i] = \tilde{\sigma}^2$; $[a_i, \tilde{b}_i] = [\tilde{a}_i, b_i] = \zeta$ for $i = 1, \dots, g$.

In the above statement, we recall once more that if $g = 0$ then $\Omega = \{\sigma, \tilde{\sigma}, \zeta\}$, and that any relations involving the elements of the set $AB \cup \widetilde{AB}$ should be suppressed.

Proof.

- (i) We analyse the images under ρ_H of the relations of $B_{k,n}(\widehat{\Sigma}_g)$ given by the presentation of Proposition 3.2. Under ρ_H , for all $3 \leq j \leq n$, relation (c.3.3) with $i = j - 2$ becomes $\tilde{\sigma}\rho_H(\zeta_j)\tilde{\sigma}^{-1} = \rho_H(\zeta_j)$ and relation (c.3.1) with $i = j - 1$ becomes $\tilde{\sigma}\rho_H(\zeta_j)\tilde{\sigma}^{-1} = \rho_H(\zeta_{j-1})$, hence $\rho_H(\zeta_j) = \rho_H(\zeta_{j-1})$, and thus $\rho_H(\zeta_2) = \dots = \rho_H(\zeta_n)$. Relation (c.3.2) with $i = 1$ yields $\tilde{\sigma}\rho_H(\zeta_1)\tilde{\sigma}^{-1} = \rho_H(\zeta_1)^{-1}\rho_H(\zeta_2)\rho_H(\zeta_1)$, and relation (c.3.3) with $i = 2$ and $j = 1$ (recall that $n \geq 3$) becomes $\tilde{\sigma}\rho_H(\zeta_1)\tilde{\sigma}^{-1} = \rho_H(\zeta_1)$. Thus $\rho_H(\zeta_1) = \rho_H(\zeta_2)$, and all of the ζ_i have a common image ζ under ρ_H .
- (ii) By the assumption on the set R and the presentation of $B_{k,n}(\widehat{\Sigma}_g)$, the group H has a group presentation with $\Omega_{k,n}$ as a generating set, and whose defining relations are obtained by adding the relations of the form $w = 1$ for all w in R to those given in the presentation of Proposition 3.2. But by hypothesis, $\rho_H(S) = \{\sigma\}$ and $\rho_H(\tilde{S}) = \{\tilde{\sigma}\}$, so $\rho_H(Z) = \{\zeta\}$ by part (i). Hence we obtain a new presentation of H by replacing $\Omega_{k,n}$ by Ω and $\sigma_i, \tilde{\sigma}_i$ and ζ_i by $\sigma, \tilde{\sigma}$ and ζ respectively in all of the defining relations. Let us show that these relations reduce to those given in the statement. First, the relations obtained from (a.1), (a.2), (b.1) and (b.2) may be removed since they are satisfied trivially. The relations (c.3) become $[\tilde{\sigma}, \zeta] = 1$. Since $k, n \geq 3$, relations (a.3) and (b.3) yield $[\sigma, a_i] = [\sigma, b_i] = [\sigma, \zeta] = [\tilde{\sigma}, \tilde{a}_i] = [\tilde{\sigma}, \tilde{b}_i] = 1$, which implies that the relations (a.4), (a.8) and (b.4) may be removed. The relations (a.6) become $[a_i, a_j] = [a_i, b_j] = [b_i, b_j] = [b_i, a_j] = 1$ for all $i < j$, the relations (b.6) become $[\tilde{a}_i, \tilde{a}_j] = [\tilde{a}_i, \tilde{b}_j] = [\tilde{b}_i, \tilde{b}_j] = [\tilde{b}_i, \tilde{a}_j] = 1$ for all $i < j$, the relations (a.7) become $[\zeta, a_i] = [\zeta, b_i] = 1$, and the relations (a.5) and (b.5) become $[a_i, b_i] = \sigma^2$ and $[\tilde{a}_i, \tilde{b}_i] = \tilde{\sigma}^2$ respectively. The relations (c.1) and (c.2) reduce to $[\sigma, \tilde{a}_i] = [\sigma, \tilde{b}_i] = [\tilde{\sigma}, a_i] = [\tilde{\sigma}, b_i] = [\sigma, \tilde{\sigma}] = 1$, the relations (c.4) become $[\zeta, \tilde{a}_i] = [\zeta, \tilde{b}_i] = 1$, and for all $i < j$, the relations (c.5) reduce to $[\tilde{a}_i, a_j] = 1$, and the relations (c.6) become $[\tilde{b}_i, b_j] = 1$. The relations (c.7.1) and (c.8.1), which only exist if $g \geq 1$, yield $[\tilde{a}_i, b_i] = \zeta$ and $[\tilde{b}_i, a_i] = \zeta^{-1}$ respectively, the latter being equivalent to $[a_i, \tilde{b}_i] = \zeta$. The other relations of type (c.7) and (c.8) reduce to $[\tilde{a}_i, b_j] = [a_i, \tilde{b}_j] = 1$ for all $i \neq j$, and so we obtain the required presentation. \square

As a consequence of Proposition 3.6, we may exhibit a presentation of $B_{k,n}(\widehat{\Sigma}_g)/\Gamma_3(B_{k,n}(\widehat{\Sigma}_g))$, which will allow us later to decompose this quotient group as a semi-direct product.

Proposition 3.7. *Let $k, n \geq 3$ and $g \geq 0$. Let $\rho_{k,n} : B_{k,n}(\widehat{\Sigma}_g) \rightarrow B_{k,n}(\widehat{\Sigma}_g)/\Gamma_3(B_{k,n}(\widehat{\Sigma}_g))$ denote the canonical projection. The group $B_{k,n}(\widehat{\Sigma}_g)/\Gamma_3(B_{k,n}(\widehat{\Sigma}_g))$ admits the following group presentation:*

Generating set: $\Omega = \rho_{k,n}(\Omega_{k,n}) = \{\sigma, \tilde{\sigma}, \zeta\} \cup AB \cup \tilde{A}\tilde{B}$;

Relations:

(a) $xy = yx$ for all $x, y \in \Omega$, $x \neq y$, and $\{x, y\} \notin \{\{a_i, b_i\}, \{\tilde{a}_j, \tilde{b}_j\}, \{b_i, \tilde{a}_i\}, \{\tilde{b}_i, a_i\}; i = 1, \dots, g\}$;

(b) $[a_i, b_i] = \sigma^2$ and $[\tilde{a}_i, \tilde{b}_i] = \tilde{\sigma}^2$; $[a_i, \tilde{b}_i] = [\tilde{a}_i, b_i] = \zeta$ for all $i = 1, \dots, g$.

In particular, $\rho_{k,n}(S) = \{\sigma\}$, $\rho_{k,n}(\tilde{S}) = \{\tilde{\sigma}\}$ and $\rho_{k,n}(Z) = \{\zeta\}$.

Proof. We apply Proposition 3.6 with $H = B_{k,n}(\widehat{\Sigma}_g)/\Gamma_3(B_{k,n}(\widehat{\Sigma}_g))$ and $\rho_H = \rho_{k,n}$. We start by showing that the hypotheses of this proposition are indeed satisfied. By definition the normal subgroup $\Gamma_3(B_{k,n}(\widehat{\Sigma}_g))$ is generated by the infinite set of elements of the form $[g_1, [g_2, g_3]]$ where g_1, g_2, g_3 range over the elements of $B_{k,n}(\widehat{\Sigma}_g)$. We have $[\sigma_i, [\sigma_{i+1}, \sigma_i]] = \sigma_i\sigma_{i+1}\sigma_i\sigma_{i+1}^{-1}\sigma_i^{-1}\sigma_{i+1}^{-1}\sigma_i\sigma_{i+1}\sigma_i^{-1}\sigma_{i+1}^{-1} = \sigma_{i+1}\sigma_i\sigma_{i+1}\sigma_{i+1}^{-1}\sigma_i^{-1}\sigma_{i+1}\sigma_i^{-1}\sigma_{i+1}^{-1} = \sigma_{i+1}^2\sigma_i^{-1}\sigma_{i+1}^{-1}$, for all $i \in \{1, \dots, k-2\}$. Since $\rho_{k,n}([\sigma_i, [\sigma_{i+1}, \sigma_i]]) = 1$, it follows that $(\rho_{k,n}(\sigma_{i+1}))^2 = \rho_{k,n}(\sigma_i)\rho_{k,n}(\sigma_{i+1})$, so $\rho_{k,n}(\sigma_{i+1}) = \rho_{k,n}(\sigma_i)$, and hence all of the σ_i have a common image under $\rho_{k,n}$ that we denote by σ . Similarly, all of the $\tilde{\sigma}_i$ have a common image $\tilde{\sigma}$ under $\rho_{k,n}$. Applying Proposition 3.6, the group $B_{k,n}(\widehat{\Sigma}_g)/\Gamma_3(B_{k,n}(\widehat{\Sigma}_g))$ admits the presentation given in that proposition. The relations of the form $w = 1$ for all $w \in R_\Omega$ correspond to the relations $[x, [y, z]] = 1$,

where x, y, z range over all the words on $\Omega \cup \Omega^{-1}$. Comparing this presentation with that given in the statement of the corollary, it suffices to show that these relations are consequences of the relations (a) and (b) given in the statement of the corollary. Let \equiv denote the equivalence relation on words on $\Omega \cup \Omega^{-1}$ generated by the defining relations (a) and (b), and let x, y, z be words on $\Omega \cup \Omega^{-1}$. We wish to show that $[x, [y, z]] \equiv 1$, where 1 denotes the empty word. Applying the Witt-Hall commutator identities [26, Theorem 5.1(9) and (10)], induction on word length, and relations (a) and (b), we see that $[y, z]$ commutes with all words on $\Omega \cup \Omega^{-1}$, and so $[x, [y, z]] \equiv 1$ as required. \square

Corollary 3.8. *Let $k, n \geq 3$. The group $B_{k,n}/\Gamma_3(B_{k,n})$ coincides with $B_{k,n}/\Gamma_2(B_{k,n})$.*

Proof. Taking $g = 0$ in Proposition 3.7, we see that the group $B_{k,n}/\Gamma_3(B_{k,n})$ is isomorphic to \mathbb{Z}^3 , and is therefore Abelian. Thus $\Gamma_2(B_{k,n})$ is a subgroup of $\Gamma_3(B_{k,n})$, and since the converse inclusion holds by definition, we conclude that $\Gamma_3(B_{k,n}) = \Gamma_2(B_{k,n})$. \square

With respect to Corollary 3.8, one may ask whether $\Gamma_3(B_{k,n}) = \Gamma_2(B_{k,n})$ if $k \leq 2$ or if $n \leq 2$. If $k = 1$ (resp. $n = 1$), it can be checked easily using Proposition 3.2 (see also [19]) that the group $B_{1,n}$ (resp. $B_{k,1}$) is isomorphic to the m -string braid group $B_m(\widehat{\Sigma}_{0,1})$ of the annulus, where $m = n$ (resp. $m = k$). If $m = 1$ then $B_m(\widehat{\Sigma}_{0,1}) \cong \mathbb{Z}$, and $\Gamma_2(B_m(\widehat{\Sigma}_{0,1})) = \Gamma_3(B_m(\widehat{\Sigma}_{0,1}))$ trivially. If $m \geq 3$ then $\Gamma_2(B_m(\widehat{\Sigma}_{0,1})) = \Gamma_3(B_m(\widehat{\Sigma}_{0,1}))$ [6, 19]. So suppose that $m = 2$. Then $\Gamma_2(B_2(\widehat{\Sigma}_{0,1}))/\Gamma_3(B_2(\widehat{\Sigma}_{0,1})) \cong \mathbb{Z}_2$, and in fact $B_2(\widehat{\Sigma}_{0,1})$ is residually nilpotent [19]. This deals with the cases where one of k and n is equal to 1. The only remaining case is that of $k = n = 2$. We do not know whether the natural surjection $B_{2,2}/\Gamma_3(B_{2,2}) \rightarrow B_{2,2}/\Gamma_2(B_{2,2})$ is injective (in particular there is no reason for the images of ζ_1 and σ_1 in $B_{2,2}/\Gamma_3(B_{2,2})$ to commute). Note however that if $\Gamma_2(B_{2,2})/\Gamma_3(B_{2,2})$ were non trivial then it would be isomorphic to the direct sum of a finite (non-zero) number of copies of \mathbb{Z}_2 [19].

If $g \geq 1$, Proposition 3.7 allows us to give a more precise description of $B_{k,n}(\widehat{\Sigma}_g)/\Gamma_3(B_{k,n}(\widehat{\Sigma}_g))$.

Corollary 3.9. *Let $k, n \geq 3$, and let $g \geq 1$.*

- (i) *The group $B_{k,n}(\widehat{\Sigma}_g)/\Gamma_3(B_{k,n}(\widehat{\Sigma}_g))$ is isomorphic to a semi-direct product of the form $(\mathbb{Z}^3 \times \mathbb{Z}^{2g}) \rtimes \mathbb{Z}^{2g}$, the first factor \mathbb{Z}^3 being generated by $\{\sigma, \tilde{\sigma}, \zeta\}$, the second factor \mathbb{Z}^{2g} by $\{a_1, \dots, a_g, \tilde{a}_1, \dots, \tilde{a}_g\}$, and the third factor \mathbb{Z}^{2g} by $\{b_1, \dots, b_g, \tilde{b}_1, \dots, \tilde{b}_g\}$.*
- (ii) *Any element γ of $B_{k,n}(\widehat{\Sigma}_g)/\Gamma_3(B_{k,n}(\widehat{\Sigma}_g))$ may be written uniquely in the form:*

$$\gamma = \sigma^p \tilde{\sigma}^q \zeta^r \prod_{i=1}^g a_i^{m_i} \tilde{a}_i^{\tilde{m}_i} \prod_{i=1}^g b_i^{n_i} \tilde{b}_i^{\tilde{n}_i}, \quad \text{where } p, q, r, m_i, \tilde{m}_i, n_i, \tilde{n}_i \in \mathbb{Z}. \quad (6)$$

- (iii) *The centre of the group $B_{k,n}(\widehat{\Sigma}_g)/\Gamma_3(B_{k,n}(\widehat{\Sigma}_g))$ is the free Abelian group of rank three generated by $\sigma, \tilde{\sigma}$ and ζ .*

Proof.

- (i) For a set X , let $FA(X)$ denote the free Abelian group on X . Let $U = \{\sigma, \tilde{\sigma}, \zeta, a_1, \dots, a_g, \tilde{a}_1, \dots, \tilde{a}_g\}$ and let $V = \{b_1, \dots, b_g, \tilde{b}_1, \dots, \tilde{b}_g\}$. One may check that the map $\varphi : V \rightarrow \text{Aut}(FA(U))$ defined by:

$$\varphi(v)(u) = \begin{cases} \sigma^{-2}u & \text{if } v = b_i \text{ and } u = a_i, \text{ or if } v = \tilde{b}_i \text{ and } u = \tilde{a}_i \\ \zeta^{-1}u & \text{if } v = b_i \text{ and } u = \tilde{a}_i, \text{ or if } v = \tilde{b}_i \text{ and } u = a_i \\ u & \text{otherwise,} \end{cases}$$

is well defined, and that it extends to a homomorphism $\varphi : FA(V) \rightarrow \text{Aut}(FA(U))$. We may thus form the semi-direct product $FA(U) \rtimes_{\varphi} FA(V)$, and by standard results ([23] for example), this group admits a presentation that coincides with that of $B_{k,n}(\widehat{\Sigma}_g)/\Gamma_3(B_{k,n}(\widehat{\Sigma}_g))$ given in Proposition 3.7.

(ii) follows directly from (i).

(iii) The centrality of $\sigma, \tilde{\sigma}$ and ζ follows from Proposition 3.7, and the fact that the subgroup $\langle \sigma, \tilde{\sigma}, \zeta \rangle$ is free Abelian of rank three follows from (i). To see that this subgroup is indeed the centre of $B_{k,n}(\widehat{\Sigma}_g)/\Gamma_3(B_{k,n}(\widehat{\Sigma}_g))$, let γ be as in equation (6), and suppose that some m_j, \tilde{m}_j, n_j or \tilde{n}_j is non zero. If $m_j \neq 0$ say, then $b_j \gamma b_j^{-1} = (\sigma^{-2m_j}) \zeta^{-\tilde{m}_j} \gamma$, so $[\gamma, b_j] = \sigma^{-2m_j} \zeta^{-\tilde{m}_j} \neq 1$ since the set $\{\sigma, \zeta\}$ generates a free Abelian group of rank 2, and thus γ is non central. By replacing b_j by a_j, \tilde{b}_j and \tilde{a}_j respectively, we obtain the same conclusion in the cases $n_j, \tilde{m}_j, \tilde{n}_j \neq 0$. So if γ is central then $m_j = n_j = \tilde{m}_j = \tilde{n}_j = 0$ for all $j = 1, \dots, g$, and the result follows. \square

Before going any further, we state and prove the following lemma that shall be used at various points in the rest of the paper. If H is a group then we denote its centre by $Z(H)$.

Lemma 3.10. *Let K, H and \tilde{H} be groups, let $\gamma : K \rightarrow H$ and $\tau : H \rightarrow \tilde{H}$ be homomorphisms, and let $\tilde{\tau} : K \rightarrow \tilde{H}$ be defined by $\tilde{\tau} = \tau \circ \gamma$. Assume that $\Gamma_2(H) \subset \text{Im}(\gamma)$ and that $Z(H) \subset \text{Im}(\gamma)$. If $\tilde{\tau}$ is injective then τ is injective. In particular, if γ is injective, then $\tilde{\tau}$ is injective if and only if τ is injective.*

Proof. Suppose that $\Gamma_2(H) \subset \text{Im}(\gamma)$, $Z(H) \subset \text{Im}(\gamma)$ and that $\tilde{\tau}$ is injective, and let us prove that τ is injective. Let $g \in \ker \tau$. If $g' \in H$ then $[g, g'] \in \Gamma_2(H)$, so there exists $k \in K$ such that $\gamma(k) = [g, g']$. Then $\tilde{\tau}(k) = \tau \circ \gamma(k) = [\tau(g), \tau(g')] = 1$ since $g \in \ker \tau$. Thus $k = 1$ by injectivity of $\tilde{\tau}$, and hence $[g, g'] = 1$ for all $g' \in H$. It follows that $g \in Z(H)$, so there exists $k' \in K$ such that $\gamma(k') = g$. As in the previous sentence, we conclude that $k' = 1$, and so $g = 1$, which proves the injectivity of τ . The second assertion then follows easily. \square

The following result says that the inclusion of classical mixed braid groups in surface mixed braid groups induces an embedding on the level of metabelian quotients. As we saw in Corollary 3.5, there is no such embedding on the level of Abelian quotients.

Corollary 3.11. *Let $k, n \geq 3$, and let $g \geq 1$. Let H be a group, and let $\rho_H : B_{k,n}(\widehat{\Sigma}_g) \rightarrow H$ be a homomorphism.*

(i) *The following conditions are equivalent:*

- (a) *the homomorphism ρ_H factors through $\rho_{k,n} : B_{k,n}(\widehat{\Sigma}_g) \rightarrow B_{k,n}(\widehat{\Sigma}_g)/\Gamma_3(B_{k,n}(\widehat{\Sigma}_g))$.*
- (b) *the restriction of ρ_H to $B_{k,n}$ factors through $r_{k,n} : B_{k,n} \rightarrow B_{k,n}/\Gamma_2(B_{k,n})$.*
- (c) *there exist $\sigma, \tilde{\sigma} \in H$ such that $\rho_H(S) = \{\sigma\}$ and $\rho_H(\tilde{S}) = \{\tilde{\sigma}\}$.*

(ii) *There exists an injective homomorphism $\alpha_{k,n} : B_{k,n}/\Gamma_2(B_{k,n}) \rightarrow B_{k,n}(\widehat{\Sigma}_g)/\Gamma_3(B_{k,n}(\widehat{\Sigma}_g))$ for which $\alpha_{k,n} \circ r_{k,n} = \rho_{k,n} \circ \iota_{k,n}$ and whose image is the centre of $B_{k,n}(\widehat{\Sigma}_g)/\Gamma_3(B_{k,n}(\widehat{\Sigma}_g))$.*

(iii) *Let $\rho_{H,3} : B_{k,n}(\widehat{\Sigma}_g)/\Gamma_3(B_{k,n}(\widehat{\Sigma}_g)) \rightarrow H$ be a homomorphism, and let $\rho_{H,2} : B_{k,n}/\Gamma_2(B_{k,n}) \rightarrow H$ be the homomorphism defined by $\rho_{H,2} = \rho_{H,3} \circ \alpha_{k,n}$. Then $\rho_{H,3}$ is injective if and only if $\rho_{H,2}$ is injective.*

(iv) *Let $\rho_{H,2} : B_{k,n}/\Gamma_2(B_{k,n}) \rightarrow H$ be a homomorphism such that $\rho_H \circ \iota_{k,n} = \rho_{H,2} \circ r_{k,n}$. Then there exists a homomorphism $\rho_{H,3} : B_{k,n}(\widehat{\Sigma}_g)/\Gamma_3(B_{k,n}(\widehat{\Sigma}_g)) \rightarrow H$ such that $\rho_{H,3} \circ \rho_{k,n} = \rho_H$ and $\rho_{H,2} = \rho_{H,3} \circ \alpha_{k,n}$. Furthermore, $\rho_{H,3}$ is injective if and only if $\rho_{H,2}$ is injective. In particular, if $\rho_{H,2}$ is injective and ρ_H is surjective then $\rho_{H,3}$ is an isomorphism.*

Note that Theorem 1.2(iii) follows by taking $\theta_H = \rho_{H,3}$ in Corollary 3.11(iv).

Proof of Corollary 3.11. Replacing H by $\rho_H(B_{k,n}(\widehat{\Sigma}_g))$ if necessary, we may suppose that ρ_H is surjective.

- (i) The equivalence of (b) and (c) follows from Propositions 3.3 and 3.6 since $B_{k,n}$ is generated by $S \cup \widetilde{S} \cup Z$. The implication (a) \implies (c) also follows easily from Proposition 3.7. So it suffices to prove the implication (c) \implies (a). If (c) holds then Proposition 3.6 applies, and the comparison of the presentation of H given there with that of $B_{k,n}(\widehat{\Sigma}_g)/\Gamma_3(B_{k,n}(\widehat{\Sigma}_g))$ given in Proposition 3.7 implies that H is a quotient of $B_{k,n}(\widehat{\Sigma}_g)/\Gamma_3(B_{k,n}(\widehat{\Sigma}_g))$. Thus ρ_H factors through $\rho_{k,n}$ as required.
- (ii) Take $H = B_{k,n}(\widehat{\Sigma}_g)/\Gamma_3(B_{k,n}(\widehat{\Sigma}_g))$ and $\rho_H = \rho_{k,n}$ in part (i). Using Proposition 3.6, condition (a) is satisfied, and so applying condition (b), we deduce the existence of a homomorphism $\alpha_{k,n} : B_{k,n}/\Gamma_2(B_{k,n}) \rightarrow B_{k,n}(\widehat{\Sigma}_g)/\Gamma_3(B_{k,n}(\widehat{\Sigma}_g))$ that provides a factorisation of ρ_H through $r_{k,n}$, so that $\alpha_{k,n} \circ r_{k,n} = \rho_{k,n} \circ \iota_{k,n}$. This relation, the surjectivity of $r_{k,n}$ and Proposition 3.6 imply that the image of $\alpha_{k,n}$ is generated by $\{\sigma, \widetilde{\sigma}, \zeta\}$. By Corollary 3.9(iii), this image is a free Abelian group of rank 3 with basis $\{\sigma, \widetilde{\sigma}, \zeta\}$, and is equal to the centre of $B_{k,n}(\widehat{\Sigma}_g)/\Gamma_3(B_{k,n}(\widehat{\Sigma}_g))$. By Proposition 3.3, $B_{k,n}/\Gamma_2(B_{k,n})$ is also a free Abelian group of rank 3 with basis $\{\sigma, \widetilde{\sigma}, \zeta\}$. Since $\alpha_{k,n}$ sends this basis onto the given basis of the image of $\alpha_{k,n}$, we conclude that $\alpha_{k,n}$ is injective.
- (iii) We have $\rho_{H,2} = \rho_{H,3} \circ \alpha_{k,n}$, where $\alpha_{k,n}$ is injective. By standard commutator properties, we have $\Gamma_2(B_{k,n}(\widehat{\Sigma}_g)/\Gamma_3(B_{k,n}(\widehat{\Sigma}_g))) \subset Z(B_{k,n}(\widehat{\Sigma}_g)/\Gamma_3(B_{k,n}(\widehat{\Sigma}_g)))$. Further, $Z(B_{k,n}(\widehat{\Sigma}_g)/\Gamma_3(B_{k,n}(\widehat{\Sigma}_g)))$ is equal to the image of $\alpha_{k,n}$ by part (ii). The result then follows by applying Lemma 3.10.
- (iv) Since the restriction of ρ_H to $B_{k,n}$ factors through $r_{k,n}$, condition (i)(b) is satisfied, and so by condition (i)(a), there exists a homomorphism $\rho_{H,3} : B_{k,n}(\widehat{\Sigma}_g)/\Gamma_3(B_{k,n}(\widehat{\Sigma}_g)) \rightarrow H$ such that $\rho_{H,3} \circ \rho_{k,n} = \rho_H$. Using the homomorphism $\alpha_{k,n}$ of part (ii), we obtain the following diagram:

$$\begin{array}{ccc}
 B_{k,n} & \xrightarrow{r_{k,n}} & B_{k,n}/\Gamma_2(B_{k,n}) \\
 \downarrow \iota_{k,n} & & \downarrow \rho_{H,2} \\
 B_{k,n}(\widehat{\Sigma}_g) & \xrightarrow{\rho_H} & H \\
 & \searrow \rho_{k,n} & \swarrow \rho_{H,3} \\
 & & B_{k,n}(\widehat{\Sigma}_g)/\Gamma_3(B_{k,n}(\widehat{\Sigma}_g))
 \end{array}
 \quad \begin{array}{l}
 \curvearrowright \alpha_{k,n} \\
 \curvearrowleft \alpha_{k,n}
 \end{array}$$

which is commutative, except possibly for the relation $\rho_{H,3} \circ \alpha_{k,n} = \rho_{H,2}$. From the commutativity of the rest of the diagram, we see that $\rho_{H,3} \circ \alpha_{k,n} \circ r_{k,n} = \rho_{H,3} \circ \rho_{k,n} \circ \iota_{k,n} = \rho_H \circ \iota_{k,n} = \rho_{H,2} \circ r_{k,n}$. The surjectivity of $r_{k,n}$ implies that $\rho_{H,3} \circ \alpha_{k,n} = \rho_{H,2}$, which proves the existence of $\rho_{H,3}$ satisfying the given properties. The equivalence of the injectivity of $\rho_{H,3}$ and that of $\rho_{H,2}$ is given by part (iii), and the last part then follows easily. \square

3.4 Metabelian quotients of $B_k(\widehat{\Sigma}_{g,n})$ and $B_n(\widehat{\Sigma}_g)$

Starting with the presentations of Proposition 3.1 instead of that of $B_{k,n}(\widehat{\Sigma}_g)$ given in Proposition 3.2, many of the arguments of Section 3.3 may be repeated for $B_k(\widehat{\Sigma}_{g,n})$ and $B_n(\widehat{\Sigma}_g)$. As we already saw in Section 3.2, there are some minor differences in some of the proofs, for example, the ζ_i are not identified in quotients of $B_k(\widehat{\Sigma}_{g,n})$, and the case $k = 1$ gives rise to slightly different results. In what follows, $B_k(\widehat{\Sigma}_g)$ will also be denoted by $B_k(\widehat{\Sigma}_{g,0})$, and we shall consider its presentation given by Proposition 3.1(i) with $Z = \emptyset$ subject to the relations (a.1)–(a.6). We emphasise that in this section, we adopt the convention that if $g = 0$ (resp. $n = 0, k = 1$) then AB (resp. $Z, \{\sigma\}$) should be suppressed from the list of generators, and that any relations involving its elements should also be removed. Propositions 3.12 and 3.13 and

Corollaries 3.14 and 3.15 are the analogues for $B_k(\widehat{\Sigma}_{g,n})$ of Propositions 3.6 and 3.7 and Corollaries 3.9 and 3.11 respectively, and their proofs, which we leave to the reader, are similar.

Proposition 3.12. *Let $k \geq 3$, let $g, n \geq 0$, let G be a group, and let $\rho_G : B_k(\widehat{\Sigma}_{g,n}) \rightarrow G$ be a surjective homomorphism. Consider the presentation of $B_k(\widehat{\Sigma}_{g,n})$ given by Proposition 3.1, and let R be a set of words on $(S \cup AB \cap Z) \cup (S \cup AB \cup Z)^{-1}$ whose normal closure in $B_k(\widehat{\Sigma}_{g,n})$ generates $\ker(\rho_G)$. Suppose that there exists $\sigma \in G$ such that $\rho_G(S) = \{\sigma\}$. Let $R_{G,\sigma}$ be the set of words obtained from R by replacing all of the σ_i by σ (and σ_i^{-1} by σ^{-1}). Then the group G has the following presentation:*

Generating set: $\rho_G(S \cup AB \cup Z) = \{\sigma\} \cup AB \cup Z$;

Relations:

- $w = 1$ for all $w \in R_{G,\sigma}$;
- $xy = yx$ for all $x, y \in \{\sigma\} \cup AB \cup Z$, $x \neq y$, and $\{x, y\} \notin \{\{a_i, b_i\}; i = 1, \dots, g\}$;
- $[a_i, b_i] = \sigma^2$ for all $i = 1, \dots, g$.

Let $\rho_k : B_k(\widehat{\Sigma}_{g,n}) \rightarrow B_k(\widehat{\Sigma}_{g,n})/\Gamma_3(B_k(\widehat{\Sigma}_{g,n}))$ denote the canonical projection. Note that if $n = 0$ then this is the homomorphism ρ_k defined in the Introduction.

Proposition 3.13. *Let $k \geq 3$, and let $g, n \geq 0$. The group $B_k(\widehat{\Sigma}_{g,n})/\Gamma_3(B_k(\widehat{\Sigma}_{g,n}))$ admits the following group presentation:*

Generating set: $\rho_k(S \cup AB \cup Z) = \{\sigma\} \cup AB \cup Z$;

Relations:

- $xy = yx$ for all $x, y \in \{\sigma\} \cup AB \cup Z$, $x \neq y$, and $\{x, y\} \notin \{\{a_i, b_i\}; i = 1, \dots, g\}$;
- for all $i = 1, \dots, g$, $[a_i, b_i] = \sigma^2$.

As in Corollary 3.8, we deduce from Proposition 3.13 that $B_k(\mathbb{D}_n)/\Gamma_3(B_k(\mathbb{D}_n))$ coincides with its Abelianisation $B_k(\mathbb{D}_n)/\Gamma_2(B_k(\mathbb{D}_n))$.

Corollary 3.14. *Let $k \geq 3$, $g \geq 1$ and $n \geq 0$.*

(i) *The group $B_k(\widehat{\Sigma}_{g,n})/\Gamma_3(B_k(\widehat{\Sigma}_{g,n}))$ is isomorphic to a semi-direct product of the form $(\mathbb{Z}^{n+1} \times \mathbb{Z}^g) \rtimes \mathbb{Z}^g$. More precisely, the first factor \mathbb{Z}^{n+1} is generated by $\{\sigma\} \cup Z$, the second factor \mathbb{Z}^g is generated by $\{a_1, \dots, a_g\}$, and the third factor \mathbb{Z}^g is generated by $\{b_1, \dots, b_g\}$.*

(ii) *Every element $\gamma \in B_k(\widehat{\Sigma}_{g,n})/\Gamma_3(B_k(\widehat{\Sigma}_{g,n}))$ may be written uniquely in the form:*

$$\gamma = \sigma^p \prod_{i=1}^n \zeta_i^{q_i} \prod_{i=1}^g a_i^{m_i} \prod_{i=1}^g b_i^{n_i}, \text{ where } p, q_i, m_i, n_i \in \mathbb{Z}.$$

(iii) *The centre of the group $B_k(\widehat{\Sigma}_{g,n})/\Gamma_3(B_k(\widehat{\Sigma}_{g,n}))$ is isomorphic to \mathbb{Z}^{n+1} and is generated by $\{\sigma\} \cup Z$.*

Part of the following result contrasts with Corollary 3.5. More precisely the inclusion of classical braid groups in surface braid groups induces an embedding at the level of metabelian quotients, but as we saw in Corollary 3.5, there is no such embedding at the level of Abelian quotients.

Corollary 3.15. *Let $k \geq 3$, $g \geq 1$ and $n \geq 0$. Let G be a group, and let $\rho_G : B_k(\widehat{\Sigma}_{g,n}) \rightarrow G$ be a homomorphism.*

(i) *the following conditions are equivalent:*

- (a) *the homomorphism ρ_G factors through $\rho_k : B_k(\widehat{\Sigma}_{g,n}) \rightarrow B_k(\widehat{\Sigma}_{g,n})/\Gamma_3(B_k(\widehat{\Sigma}_{g,n}))$.*
- (b) *the restriction of ρ_G to $B_k(\mathbb{D}_n)$ factors through $r_k : B_k(\mathbb{D}_n) \rightarrow B_k(\mathbb{D}_n)/\Gamma_2(B_k(\mathbb{D}_n))$.*

- (c) There exists $\sigma \in G$ such that $\rho_G(S) = \{\sigma\}$.
- (ii) There exists an injective homomorphism $\alpha_k : B_k(\mathbb{D}_n)/\Gamma_2(B_k(\mathbb{D}_n)) \rightarrow B_k(\widehat{\Sigma}_{g,n})/\Gamma_3(B_k(\widehat{\Sigma}_{g,n}))$ that satisfies $\alpha_k \circ r_k = \rho_k \circ \iota_k$ and whose image is the centre of $B_k(\widehat{\Sigma}_{g,n})/\Gamma_3(B_k(\widehat{\Sigma}_{g,n}))$.
- (iii) Let $\rho_{G,3} : B_k(\widehat{\Sigma}_{g,n})/\Gamma_3(B_k(\widehat{\Sigma}_{g,n})) \rightarrow G$ be a homomorphism, and consider the homomorphism $\rho_{G,2} : B_k(\mathbb{D}_n)/\Gamma_2(B_k(\mathbb{D}_n)) \rightarrow G$ defined by $\rho_{G,2} = \rho_{G,3} \circ \alpha_k$. Then $\rho_{G,3}$ is injective if and only if $\rho_{G,2}$ is.
- (iv) Let $\rho_{G,2} : B_k(\mathbb{D}_n)/\Gamma_2(B_k(\mathbb{D}_n)) \rightarrow G$ be a homomorphism such that $\rho_G \circ \iota_k = \rho_{G,2} \circ r_k$. Then there exists a homomorphism $\rho_{G,3} : B_k(\widehat{\Sigma}_{g,n})/\Gamma_3(B_k(\widehat{\Sigma}_{g,n})) \rightarrow G$ such that $\rho_{G,3} \circ \rho_k = \rho_G$ and $\rho_{G,2} = \rho_{G,3} \circ \alpha_k$. Furthermore, $\rho_{G,3}$ is injective if and only if $\rho_{G,2}$ is injective. In particular, if $\rho_{G,2}$ is injective and ρ_G is surjective then $\rho_{G,3}$ is an isomorphism.

Corollary 3.15 may be proved in the same way as Corollary 3.11: in the proof, Propositions 3.3, 3.6 and 3.7 should be replaced by Propositions 3.4, 3.12 and 3.13 respectively.

We are now able to complete the proof of parts (i), (iii) and (iv) of Theorem 1.2. Part (ii) will be proved in Section 4.

Proof of parts (i), (iii) and (iv) of Theorem 1.2. Part (iii) was proved just after the statement of Corollary 3.11, and part (iv) follows in a similar way by Corollary 3.15(iv). It remains to prove part (i). Let $\alpha_{k,n} : B_{k,n}/\Gamma_2(B_{k,n}) \rightarrow B_{k,n}(\widehat{\Sigma}_g)/\Gamma_3(B_{k,n}(\widehat{\Sigma}_g))$ and $\alpha_n : B_n/\Gamma_2(B_n) \rightarrow B_n(\widehat{\Sigma}_g)/\Gamma_3(B_n(\widehat{\Sigma}_g))$ be the homomorphisms defined in Corollaries 3.11(ii) and 3.15(ii) respectively. Together with the commutative diagrams (1), (2) and (5), these two corollaries entail the existence and the commutativity of the diagram (3), with the exception, *a priori*, of the existence of γ_k and the commutativity of the first column, which we now prove. Let $\gamma_k : G_k \rightarrow G_k(\widehat{\Sigma}_g)$ denote the restriction of $\alpha_{k,n}$ to G_k . The commutativity of the rest of the diagram (3) implies that γ_k is well defined, and the injectivity of γ_k is a consequence of that of $\alpha_{k,n}$. Restricting appropriately the relation $\alpha_{k,n} \circ r_{k,n} = \rho_{k,n} \circ \iota_{k,n}$, we obtain $\gamma_k \circ p_k = \Phi_k \circ \iota_k$, and this completes the proof of Theorem 1.2(i). \square

4 The group $G_k(\widehat{\Sigma}_g)$

In this section, we exhibit a presentation of $G_k(\widehat{\Sigma}_g)$, we prove Theorem 1.1 and we complete the proof of Theorem 1.2.

Let $k, n \geq 1$ (resp. $k, n \geq 3$). The group G_k (resp. $G_k(\widehat{\Sigma}_g)$), which is defined by the commutative diagram (1) (resp. (2)), is described in Lemma 4.1 (resp. Proposition 4.2). Notice in particular that these groups only depend on k and g , and do not depend on n , which justifies the absence of n in the notation.

Lemma 4.1. *Let $k, n \geq 1$. Then the group G_k is a free Abelian group and is a direct factor of $B_{k,n}/\Gamma_2(B_{k,n})$. If $k = 1$ then G_k is isomorphic to \mathbb{Z} and is generated by $\{\zeta\}$; if $k \geq 2$ then G_k is of rank 2, and is generated by $\{\sigma, \zeta\}$.*

Proof. Let $k, n \geq 1$. We make use of the notation of Propositions 3.1 and 3.2 for the groups $B_{k,n}$ and B_n . Applying Propositions 3.3 and 3.4, the group $B_{k,n}/\Gamma_2(B_{k,n})$ is a free Abelian group with basis $\widehat{S} \cup \{\zeta\}$, and by Proposition 3.4, $B_n/\Gamma_2(B_n)$ is a free Abelian group with basis $\widehat{S} \setminus \{\sigma\}$, where \widehat{S} is as defined in Proposition 3.3.

In terms of the notation of Proposition 3.3, the group $B_{k,n}/\Gamma_2(B_{k,n})$ is a free Abelian group with basis $\widehat{S} \cup \{\zeta\}$, and by Proposition 3.4, $B_n/\Gamma_2(B_n)$ is a free Abelian group with basis $\widehat{S} \setminus \{\sigma\}$. The homomorphism from $B_{k,n}$ to B_n sends $\tilde{\sigma}_1, \dots, \tilde{\sigma}_{n-1}$ to $\tilde{\sigma}_1, \dots, \tilde{\sigma}_{n-1}$ respectively (if $n \geq 2$), and $\sigma_1, \dots, \sigma_{k-1}$ (if $k \geq 2$) and ζ_1, \dots, ζ_n onto the trivial element. Since $r_{k,n}$ identifies the elements of S (resp. of \tilde{S} , of Z) to $\{\sigma\}$ (resp. to $\{\tilde{\sigma}\}$, to $\{\zeta\}$), it follows that G_k is the kernel of the homomorphism from $B_{k,n}/\Gamma_2(B_{k,n})$ to

$B_n/\Gamma_2(B_n)$ that sends $\tilde{\sigma}$ to $\tilde{\sigma}$ (if $n \geq 2$), and σ (if $k \geq 2$) and ζ onto the trivial element. Hence G_k is the free Abelian group with basis $(\widehat{S} \setminus \{\tilde{\sigma}\}) \cup \{\zeta\}$. In particular, $G_k \cong \mathbb{Z}$ if $k = 1$, and $G_k \cong \mathbb{Z}^2$ if $k \geq 2$. \square

Proposition 4.2. *Let $k, n \geq 3$ and $g \geq 1$. The group $G_k(\widehat{\Sigma}_g)$ admits the following presentation:*

Generating set: $\{\sigma, \zeta\} \cup AB$;

Relations:

- $xy = yx$ for all $x, y \in \{\sigma, \zeta\} \cup AB$, $x \neq y$, and $\{x, y\} \notin \{\{a_i, b_i\}; i = 1, \dots, g\}$;
- $[a_i, b_i] = \sigma^2$ for all $i = 1, \dots, g$.

Proof. From the commutative diagram (2) of short exact sequences, the homomorphism $\Phi_k : B_k(\widehat{\Sigma}_{g,n}) \longrightarrow G_k(\widehat{\Sigma}_g)$ is the restriction of $\rho_{k,n}$ to $B_k(\widehat{\Sigma}_{g,n})$. In terms of the presentations of Propositions 3.1 and 3.2, ψ_k is defined by $\psi_k(x) = x$ for all $x \in \widetilde{S} \cup \widetilde{AB}$, and $\psi_k(x) = 1$ for all $x \in S \cup AB \cup Z$. It follows from Proposition 3.7 and the commutativity of the diagram (2) that the subgroup of $B_{k,n}(\widehat{\Sigma}_g)/\Gamma_3(B_{k,n}(\widehat{\Sigma}_g))$ generated by $\{\sigma, \zeta\} \cup AB$ is included in $G_k(\widehat{\Sigma}_g)$. Conversely, let $\gamma \in B_{k,n}(\widehat{\Sigma}_g)/\Gamma_3(B_{k,n}(\widehat{\Sigma}_g))$, that we write in the form of equation (6). Since $\bar{\psi}_k$ is induced by ψ_k , we see that $\bar{\psi}_k(\gamma) = \tilde{\sigma}^q \prod_{i=1}^g \tilde{a}_i^{\tilde{m}_i} \prod_{i=1}^g \tilde{b}_i^{\tilde{n}_i}$. Corollary 3.14(ii) implies that $\gamma \in \ker(\bar{\psi}_k)$ if and only if $q = \tilde{m}_i = \tilde{n}_i = 0$ for all $i = 1, \dots, g$, and so $\gamma \in \langle \{\sigma, \zeta\} \cup AB \rangle$ by equation (6). Consequently, $G_k(\widehat{\Sigma}_g)$ is the subgroup of $B_{k,n}(\widehat{\Sigma}_g)/\Gamma_3(B_{k,n}(\widehat{\Sigma}_g))$ generated by $\{\sigma, \zeta\} \cup AB$. Since the relations given in the statement of the proposition hold in $B_{k,n}(\widehat{\Sigma}_g)/\Gamma_3(B_{k,n}(\widehat{\Sigma}_g))$, they also hold in $G_k(\widehat{\Sigma}_g)$. Moreover, starting from any word w on the elements of $\{\sigma, \zeta\} \cup AB$ and their inverses, it follows from just these relations (and the relations $x^{-1}x = xx^{-1} = 1$ for $x \in \{\sigma, \zeta\} \cup AB$) that w may be transformed into the unique word of the form $\sigma^p \zeta^r \prod_{i=1}^g a_i^{m_i} \prod_{i=1}^g b_i^{n_i}$, $p, r, m_i, n_i \in \mathbb{Z}$, that represents the same element in $B_{k,n}(\widehat{\Sigma}_g)/\Gamma_3(B_{k,n}(\widehat{\Sigma}_g))$. This proves that the relations given in the proposition are indeed a complete set of relations for $G_k(\widehat{\Sigma}_g)$ for the generating set $\{\sigma, \zeta\} \cup AB$. \square

Let $k, n \geq 3$ and $g \geq 1$. The homomorphism $\gamma_k : G_k \longrightarrow G_k(\widehat{\Sigma}_g)$, which is the restriction of $\alpha_{k,n}$ to G_k , was seen to be injective in Theorem 1.2(i).

Corollary 4.3. *Let $g \geq 1$.*

- (i) *Let $k \geq 1$ and $n \geq 1$. Then the homomorphism $\Phi_k : B_k(\widehat{\Sigma}_{g,n}) \longrightarrow G_k(\widehat{\Sigma}_g)$ factors through the homomorphism $\rho_k : B_k(\widehat{\Sigma}_{g,n}) \longrightarrow B_k(\widehat{\Sigma}_{g,n})/\Gamma_3(B_k(\widehat{\Sigma}_{g,n}))$.*
- (ii) *Let $k, n \geq 3$. The group $G_k(\widehat{\Sigma}_g)$ is isomorphic to a semi-direct product of the form $(\mathbb{Z}^2 \times \mathbb{Z}^g) \rtimes \mathbb{Z}^g$. Its centre is a free Abelian group with basis $\{\sigma, \zeta\}$ and is equal to $\gamma_k(G_k)$. Moreover, every element of $G_k(\widehat{\Sigma}_g)$ may be written uniquely in the form $\sigma^p \zeta^q \prod_{i=1}^g a_i^{m_i} \prod_{i=1}^g b_i^{n_i}$, where $p, q, m_i, n_i \in \mathbb{Z}$.*
- (iii) *Let $k, n \geq 3$. The groups $G_k(\widehat{\Sigma}_g)$ and $M_k(\widehat{\Sigma}_g) := (B_k(\widehat{\Sigma}_{g,n})/\Gamma_3(B_k(\widehat{\Sigma}_{g,n}))) /_{\zeta_1 = \dots = \zeta_n}$ are isomorphic. Moreover, if $q_k : B_k(\widehat{\Sigma}_{g,n})/\Gamma_3(B_k(\widehat{\Sigma}_{g,n})) \longrightarrow G_k(\widehat{\Sigma}_g)$ is a homomorphism for which $\Phi_k = q_k \circ \rho_k$ then q_k induces an isomorphism from $M_k(\widehat{\Sigma}_g)$ to $G_k(\widehat{\Sigma}_g)$.*

Proof.

- (i) From the commutative diagram (2) of short exact sequences, $G_k(\widehat{\Sigma}_g)$ is a subgroup of the quotient $B_{k,n}(\widehat{\Sigma}_g)/\Gamma_3(B_{k,n}(\widehat{\Sigma}_g))$, so $[x, [y, z]] = 1$ for all $x, y, z \in G_k(\widehat{\Sigma}_g)$, and hence Φ_k factors through $\rho_k : B_k(\widehat{\Sigma}_{g,n}) \longrightarrow B_k(\widehat{\Sigma}_{g,n})/\Gamma_3(B_k(\widehat{\Sigma}_{g,n}))$.
- (ii) The result follows from Proposition 4.2 and Lemma 4.1 using arguments similar to those given in the proof of Corollary 3.9.
- (iii) This is a consequence of the presentations of the groups $B_k(\widehat{\Sigma}_{g,n})/\Gamma_3(B_k(\widehat{\Sigma}_{g,n}))$ and $G_k(\widehat{\Sigma}_g)$ given in Propositions 3.13 and 4.2 respectively. \square

We are now in a position to prove Theorem 1.1 and to finish the proof of Theorem 1.2.

Proof of Theorem 1.1. The proof is a straightforward consequence of Proposition 4.2. Indeed, for $k \geq 3$, the group G_Σ introduced in [1, Section 3] was abstractly defined by the group presentation of Proposition 4.2. Hence there exists a canonical isomorphism of groups $\iota : G_\Sigma \rightarrow G_k(\widehat{\Sigma}_g)$. Moreover, the homomorphism $\Phi_k : B_k(\widehat{\Sigma}_{g,n}) \rightarrow G_k(\widehat{\Sigma}_g)$ is the restriction of $\rho_{k,n} : B_{k,n}(\widehat{\Sigma}_g) \rightarrow B_{k,n}(\widehat{\Sigma}_g)/\Gamma_3(B_{k,n}(\widehat{\Sigma}_g))$, and by Proposition 3.7, is defined by $\Phi_k(S) = \{\sigma\}$, $\Phi_k(Z) = \{\zeta\}$, and $\Phi_k(x) = x$ for all $x \in AB$. So considering the definition of the homomorphism Φ_Σ given in [1, page 266], we conclude that $\iota \circ \Phi_\Sigma = \Phi_k$ as required. \square

Corollary 4.4. *Let $k, n \geq 3$ and $g \geq 1$. Let G be a group, and $\Phi_G : B_k(\widehat{\Sigma}_{g,n}) \rightarrow G$ be a homomorphism.*

(i) *The following conditions are equivalent:*

- (a) *the homomorphism Φ_G factors through $\Phi_k : B_k(\widehat{\Sigma}_{g,n}) \rightarrow G_k(\widehat{\Sigma}_g)$.*
- (b) *the restriction of Φ_G to $B_k(\mathbb{D}_n)$ factors through $p_k : B_k(\mathbb{D}_n) \rightarrow G_k$.*
- (c) *there exist $\sigma, \zeta \in G$ such that $\Phi_G(S) = \{\sigma\}$ and $\Phi_G(Z) = \{\zeta\}$.*

(ii) *Let $\Phi_{G,2} : G_k \rightarrow G$ be a homomorphism such that $\Phi_{G,2} \circ p_k$ is the restriction of Φ_G to $B_k(\mathbb{D}_n)$. Then there exists a homomorphism $\Phi_{G,3} : G_k(\widehat{\Sigma}_g) \rightarrow G$ such that $\Phi_{G,3} \circ \Phi_k = \Phi_G$. Further, $\Phi_{G,3}$ is injective if and only if $\Phi_{G,2}$ is injective.*

Proof.

(i) Suppose first that Φ_G factors through Φ_k . By the proof of Theorem 1.1, $\Phi_k(S) = \{\sigma\}$ and $\Phi_k(Z) = \{\zeta\}$, so using the same notation for the corresponding elements of G , there exist $\sigma, \zeta \in G$ such that $\Phi_G(S) = \{\sigma\}$ and $\Phi_G(Z) = \{\zeta\}$, hence (a) implies (c). Since $\gamma_k \circ p_k = \Phi_k \circ \iota_k$ by Theorem 1.2(i), the restriction of Φ_G to $B_k(\mathbb{D}_n)$ factors through p_k , so (a) implies (b).

Now suppose that (c) holds. Then $\Phi_G(S) = \{\sigma\}$, and by Corollary 3.15(i), Φ_G factors through ρ_k , so there exists a homomorphism $\Phi'_G : B_k(\widehat{\Sigma}_{g,n})/\Gamma_3(B_k(\widehat{\Sigma}_{g,n})) \rightarrow G$ such that $\Phi_G = \Phi'_G \circ \rho_k$. Further, $\Phi_G(Z) = \{\zeta\}$, and letting Z also denote the usual subset of elements of $B_k(\widehat{\Sigma}_{g,n})$, we have that $\rho_k(Z) = Z$, and thus $\Phi'_G(Z) = \{\zeta\}$. So Φ'_G factors through the canonical surjection $B_k(\widehat{\Sigma}_{g,n})/\Gamma_3(B_k(\widehat{\Sigma}_{g,n})) \rightarrow M_k(\widehat{\Sigma}_g)$. Taking $q_k : B_k(\widehat{\Sigma}_{g,n})/\Gamma_3(B_k(\widehat{\Sigma}_{g,n})) \rightarrow G_k(\widehat{\Sigma}_g)$ to be a homomorphism as in the statement of Corollary 4.3(iii) that satisfies $\Phi_k = q_k \circ \rho_k$, and applying the resulting induced isomorphism between $G_k(\widehat{\Sigma}_g)$ and $M_k(\widehat{\Sigma}_g)$, we obtain a homomorphism $\Phi_{G,3} : G_k(\widehat{\Sigma}_g) \rightarrow G$ such that $\Phi'_G = \Phi_{G,3} \circ q_k$. Hence $\Phi_G = \Phi'_G \circ \rho_k = \Phi_{G,3} \circ q_k \circ \rho_k = \Phi_{G,3} \circ \Phi_k$, therefore Φ_G factors through Φ_k , and thus (c) implies (a).

Finally, suppose that (b) holds. Then $p_k(S) = r_{k,n}(S) = \{\sigma\}$ and $p_k(Z) = r_{k,n}(Z) = \{\zeta\}$ by the commutative diagram (1) and Proposition 3.3. Since $\Phi_G \circ \iota_k$ factors through p_k , it follows that there exist $\sigma, \zeta \in G$ such that $\Phi_G(S) = \{\sigma\}$ and $\Phi_G(Z) = \{\zeta\}$, hence (b) implies (c).

(ii) Since the restriction of Φ_G to $B_k(\mathbb{D}_n)$ factors through p_k , it follows from part (i) that Φ_G factors through Φ_k , and so there exists a homomorphism $\Phi_{G,3} : G_k(\widehat{\Sigma}_g) \rightarrow G$ such that $\Phi_{G,3} \circ \Phi_k = \Phi_G$. By hypothesis, we have $\Phi_G \circ \iota_k = \Phi_{G,2} \circ p_k$, so $\Phi_{G,2} \circ p_k = \Phi_G \circ \iota_k = \Phi_{G,3} \circ \Phi_k \circ \iota_k = \Phi_{G,3} \circ \gamma_k \circ p_k$. The surjectivity of p_k implies that $\Phi_{G,2} = \Phi_{G,3} \circ \gamma_k$. Now γ_k is injective by Theorem 1.2(i), and applying Proposition 4.2 and Corollary 4.3(iii), we see that $\Gamma_2(G_k(\widehat{\Sigma}_g)) \subset Z(G_k(\widehat{\Sigma}_g))$ and $Z(G_k(\widehat{\Sigma}_g))$ is equal to the image of γ_k . The result then follows from Lemma 3.10. \square

This enables us to prove Theorem 1.2(ii).

Proof of Theorem 1.2(ii). With the notation of the proof of Corollary 4.4(iv), the existence and injectivity of $\Phi_{G,2}$ imply those of $\Phi_{G,3} : G_k(\widehat{\Sigma}_g) \rightarrow G$. The surjectivity of $\Phi_{G,3}$ follows from that of Φ_G . We thus have $\Phi_{G,3} \circ \Phi_k = \Phi_G$, where $\Phi_{G,3}$ is an isomorphism. \square

Taking into account the proof of parts (i), (iii) and (iv) in Section 3.4, this concludes the proof of Theorem 1.2.

These results allow us also to do make some remarks on the extension of the length function from B_n to surface braid groups. Let $k \geq 3$. The projection $r_k : B_k \rightarrow B_k/\Gamma_2(B_k)$ coincides (up to isomorphism) with the usual length function $\lambda : B_k \rightarrow \mathbb{Z}$ on B_k . If $g \geq 1$, we claim that, up to isomorphism, the only surjective homomorphism extending λ from B_k to $B_k(\widehat{\Sigma}_g)$ is $\rho_k : B_k(\widehat{\Sigma}_g) \rightarrow B_k(\widehat{\Sigma}_g)/\Gamma_3(B_k(\widehat{\Sigma}_g))$. Indeed, let G be a group, and let $\lambda_\Sigma : B_k(\widehat{\Sigma}_g) \rightarrow G$ be a surjective homomorphism that extends λ . So there exists an injective homomorphism $\rho_{G,2} : B_k/\Gamma_2(B_k) \rightarrow G$ satisfying $\rho_{G,2} \circ \lambda = \lambda_\Sigma \circ \iota_k$. Applying Corollary 4.4(ii), there exists an isomorphism $\rho_{G,3} : B_k(\widehat{\Sigma}_g)/\Gamma_3(B_k(\widehat{\Sigma}_g)) \rightarrow G$ such that $\rho_{G,3} \circ \rho_k = \lambda_\Sigma$, which proves the claim. In the case of surfaces without boundary, we have the following negative result, which as we mentioned at the beginning of Section 3, provides another reason why we only consider surfaces with boundary in this paper.

Proposition 4.5. *Let $n \geq 3$, and let Σ be a compact orientable surface of positive genus without boundary. It is not possible to extend the length function $\lambda : B_n \rightarrow \mathbb{Z}$ to $B_n(\Sigma)$. In other words, there does not exist a surjective homomorphism λ_Σ of $B_n(\Sigma)$ onto a group F whose restriction B_n coincides with λ .*

Proof. Let F be a group, and let $\lambda_\Sigma : B_n(\Sigma) \rightarrow F$ be a homomorphism that extends λ . Then there exists $\sigma \in F$ such that $\sigma = \lambda_\Sigma(\sigma_1) = \dots = \lambda_\Sigma(\sigma_{n-1})$. The group presentation of $B_n(\Sigma)$ given in [4] implies that $\sigma^{2(n+g-1)} = 1$ (for further details, see the proof of [6, Theorem 1]). The result then follows because σ is of finite order but $\lambda(\sigma_1)$ is of infinite order. \square

In [1, Section 3], the authors consider also a group H_Σ that is defined by its group presentation. One may check using this presentation that H_Σ is isomorphic to the quotient $(B_{k,n}(\widehat{\Sigma}_g)/\Gamma_3(B_{k,n}(\widehat{\Sigma}_g)))/\tilde{\sigma}=1$. In [1, Theorem 4.3(ii)], a rigidity result is proved for this group in the case $k \geq 3$. We conclude this section with an alternative proof of this theorem, based mainly on the results and arguments of Section 3.

Proposition 4.6. *Let $k, n \geq 3$. Let H be a group, and let $\Phi_H : B_{k,n}(\widehat{\Sigma}_g) \rightarrow H$ be a homomorphism such that:*

- (i) *the restriction of Φ_H to $B_{k,n}$ factors through $r_{k,n}$, in other words, there exists a homomorphism $\Phi_{H,2} : B_{k,n}/\Gamma_2(B_{k,n}) \rightarrow H$ such that $\Phi_H \circ \iota_{k,n} = \Phi_{H,2} \circ r_{k,n}$.*
- (ii) *the kernel of $\Phi_{H,2}$ is generated by $\tilde{\sigma}$.*

Then Φ_H induces an injective homomorphism from H_Σ to H . In particular, if Φ_H is surjective then the induced homomorphism is an isomorphism.

Note that the assumptions (i) and (ii) of Proposition 4.6 are equivalent to the following two conditions:

- (a) *the restriction of Φ_H to $B_k(\mathbb{D}_n)$ induces an injective homomorphism from G_k to H .*
- (b) *$\Phi_H(\tilde{S}) = \{1\}$.*

We remark that conditions (a) and (b) correspond to Hypothesis (†) in [1, Theorem 4.3(ii)].

Proof. Since the restriction of Φ_H to $B_{k,n}$ factors through $r_{k,n}$ using hypothesis (i), Corollary 3.11(i) implies that Φ_H factors through $\rho_{k,n}$, so there exists a homomorphism $\Phi_{H,3} : B_{k,n}(\widehat{\Sigma}_g)/\Gamma_3(B_{k,n}(\widehat{\Sigma}_g)) \rightarrow H$ such that $\Phi_H = \Phi_{H,3} \circ \rho_{k,n}$. Now $\Phi_H(\tilde{S}) = \{1\}$ using hypothesis (ii) and $\rho_{k,n}(\tilde{S}) = \{\tilde{\sigma}\}$ by Proposition 3.7, so it follows that $\Phi_{H,3}(\tilde{\sigma}) = \{1\}$. Thus the homomorphism $\Phi_{H,3}$ factors through the projection of the

group $B_{k,n}(\widehat{\Sigma}_g)/\Gamma_3(B_{k,n}(\widehat{\Sigma}_g))$ onto its quotient $(B_{k,n}(\widehat{\Sigma}_g)/\Gamma_3(B_{k,n}(\widehat{\Sigma}_g)))/_{\tilde{\sigma}=1}$, which from the preceding remarks, we know to be isomorphic to H_Σ . It remains to prove that the induced homomorphism from this quotient (or equivalently from H_Σ) to H is injective. To do so, first note that $\Phi_{H,3} \circ \alpha_{k,n} = \Phi_{H,2}$, where $\alpha_{k,n} : B_{k,n}/\Gamma_2(B_{k,n}) \rightarrow B_{k,n}(\widehat{\Sigma}_g)/\Gamma_3(B_{k,n}(\widehat{\Sigma}_g))$ is the homomorphism given by Corollary 3.11(ii). Since $\alpha_{k,n} \circ r_{k,n} = \rho_{k,n} \circ \iota_{k,n}$, we have $\alpha_{k,n}(\tilde{\sigma}) = \tilde{\sigma}$, and so the subgroup of $B_{k,n}(\widehat{\Sigma}_g)/\Gamma_3(B_{k,n}(\widehat{\Sigma}_g))$ generated by $\tilde{\sigma}$ is contained in $\ker(\Phi_{H,3})$. The proof of the converse is similar in spirit to that of Lemma 3.10. Let $\gamma \in \ker(\Phi_{H,3})$, and let $\gamma' \in B_{k,n}(\widehat{\Sigma}_g)/\Gamma_3(B_{k,n}(\widehat{\Sigma}_g))$. The element $[\gamma, \gamma']$ belongs to the centre of $B_{k,n}(\widehat{\Sigma}_g)/\Gamma_3(B_{k,n}(\widehat{\Sigma}_g))$, and therefore to the image of $\alpha_{k,n}$ by Corollary 3.11(ii). Let $h \in B_{k,n}/\Gamma_2(B_{k,n})$ be such that $\alpha_{k,n}(h) = [\gamma, \gamma']$. Then $\Phi_{H,2}(h) = \Phi_{H,3}([\gamma, \gamma']) = [\Phi_{H,3}(\gamma), \Phi_{H,3}(\gamma')] = 1$. Therefore $h = \tilde{\sigma}^\ell$ for some $\ell \in \mathbb{Z}$ using hypothesis (ii), and the relation $\alpha_{k,n}(\tilde{\sigma}) = \tilde{\sigma}$ implies that for all $\gamma' \in B_{k,n}(\widehat{\Sigma}_g)/\Gamma_3(B_{k,n}(\widehat{\Sigma}_g))$, the commutator $[\gamma, \gamma']$ is equal to a power of $\tilde{\sigma}$. Write γ in the form of equation (6). Then for all $j = 1, \dots, g$, we have $[\gamma, b_j] = (\sigma^{2m_j})\zeta^{\tilde{m}_j}$ and $[\gamma, a_j] = (\sigma^{-2n_j})\zeta^{-\tilde{n}_j}$ by Proposition 3.7. Since these commutators are powers of $\tilde{\sigma}$, we deduce that $m_j = n_j = \tilde{m}_j = \tilde{n}_j = 0$, and so γ belongs to the centre of $B_{k,n}(\widehat{\Sigma}_g)/\Gamma_3(B_{k,n}(\widehat{\Sigma}_g))$ by Corollary 3.9(iii). Repeating the argument with γ in place of $[\gamma, \gamma']$, there exists $h' \in B_{k,n}/\Gamma_2(B_{k,n})$ such that $\alpha_{k,n}(h') = \gamma$, so $\Phi_{H,2}(h') = 1$ because $\Phi_{H,3} \circ \alpha_{k,n} = \Phi_{H,2}$, and thus $\gamma = \tilde{\sigma}^\ell$ for some $\ell \in \mathbb{Z}$ by hypothesis (ii) and the fact that $\alpha_{k,n}(\tilde{\sigma}) = \tilde{\sigma}$. The kernel of $\Phi_{H,3}$ is therefore the subgroup of $B_{k,n}(\widehat{\Sigma}_g)/\Gamma_3(B_{k,n}(\widehat{\Sigma}_g))$ generated by $\tilde{\sigma}$, and hence the induced homomorphism from H_Σ to H is injective. \square

5 Representations of surface braid groups

In this section, we first describe an algebraic approach to the Burau and Bigelow-Krammer-Lawrence representations that is based on the lower central series. To our knowledge, this description has not appeared in the literature, although it is well known to the experts in the field. We then show why it is not possible to extend the Burau and Bigelow-Krammer-Lawrence representations to representations of surface braid groups. This latter remark was made in [1] under certain conditions of a homological nature. Within a purely algebraic framework, we prove this non-existence result with fewer conditions than those of [1].

Let us start recalling that B_n may be seen as the mapping class group of \mathbb{D}_n , thus giving rise to an action of B_n on $\pi_1(\mathbb{D}_n)$, the latter being isomorphic to the free group F_n on n generators [9]. This action coincides with the action by conjugation of B_n on $B_1(\mathbb{D}_n)$ defined by the standard section of (SMB), and gives rise to Artin's (faithful) representation of B_n as a subgroup of the group of automorphisms of F_n . The (non-reduced) Burau representation of B_n is then obtained by composing the Artin representation with the Magnus representation associated with the length function $\ell : B_1(\mathbb{D}_n) \rightarrow \mathbb{Z}$ (see for instance [2]), which we identify with the homomorphism $p_1 : B_1(\mathbb{D}_n) \rightarrow G_1$ of the commutative diagram (1) (see Lemma 4.1 for more details).

This representation also has a homological interpretation (see for instance [24, Chapter 3]): the group B_n acts on the infinite cyclic covering $\widetilde{\mathbb{D}}_n$ of \mathbb{D}_n , and since the action of B_n on $B_1(\mathbb{D}_n)$ commutes with the length function p_1 (whose image is \mathbb{Z}), the induced action on the first homology group of $\widetilde{\mathbb{D}}_n$ is the (reduced) Burau representation of B_n .

More generally, if $k \geq 1$, B_n , regarded as the mapping class group of \mathbb{D}_n , acts on $\mathbb{F}_k(\mathbb{D}_n)/S_k$ and therefore on its fundamental group $B_k(\mathbb{D}_n)$. Once more, the induced action of B_n on $B_k(\mathbb{D}_n)$ coincides with the action by conjugation of B_n on $B_k(\mathbb{D}_n)$ defined by the standard section of (SMB). In analogy with the case $k = 1$, in order to seek (linear) representations of $B_k(\mathbb{D}_n)$, we consider regular coverings associated with its normal subgroups, and we try to see if the induced action on homology is well defined. In other words, we wish to study surjections of $B_k(\mathbb{D}_n)$ onto a group subject to certain constraints. Now suppose that $k > 1$, and that the group G_k is a free Abelian group of rank 2 (*cf.* Lemma 4.1), and consider the homomorphism p_k of the commutative diagram (1). It is easy to check that the action of

B_n on $B_k(\mathbb{D}_n)$ commutes with p_k , that B_n acts on the regular covering of $\mathbb{F}_k(\mathbb{D}_n)/S_k$ corresponding to p_k , and that the induced action on the Borel-Moore middle homology group of this covering space defines a homological representation of B_n . When $k = 2$, we obtain the famous Bigelow-Krammer-Lawrence representation that is a faithful linear representation (see [24] for a complete description of this construction). In [29], it was proved that the corresponding representations are also faithful in the general case $k \geq 2$. In what follows, we will call this family of representations *BKL representations* (the Burau and the usual Bigelow-Krammer-Lawrence representations correspond to the cases $k = 1$ and $k = 2$).

With this algebraic constructions of Section 3 in mind, we would like to extend the BKL representations from B_n to $B_n(\widehat{\Sigma}_g)$. To do so, one might seek a projection $\Phi_G : B_k(\widehat{\Sigma}_{g,n}) \rightarrow G$ onto an Abelian group G that induces an action on the homology of the corresponding covering and whose restriction to $B_k(\mathbb{D}_n) \subset B_k(\widehat{\Sigma}_{g,n})$ coincides with $p_k : B_k(\mathbb{D}_n) \rightarrow G_k$ (more precisely, there is an injective homomorphism j_k from G_k to G so that the restriction of Φ_G to $B_k(\mathbb{D}_n)$ coincides with $j_k \circ p_k$). However, if there existed Φ_G such that $\Phi_G \circ \beta_* = \Phi_G$ for all $\beta \in B_n(\widehat{\Sigma}_g)$, where β_* denotes the automorphism of $B_k(\widehat{\Sigma}_{g,n})$ induced by conjugation by β , then the homomorphism Φ_G would extend to a homomorphism from $B_{k,n}(\widehat{\Sigma}_g)$ to G because G is Abelian, and we would obtain a linear representation of $B_n(\widehat{\Sigma}_g)$ in $\text{Aut}_{\mathbb{Z}[G]}(H_k^{\text{BM}}(\widehat{E_k/S_k}))$, where $H_k^{\text{BM}}(\widehat{E_k/S_k})$ is the Borel-Moore middle homology group of the covering space of E_k/S_k and $E_k = \mathbb{F}_k(\widehat{\Sigma}_{g,n})$. Unfortunately, the results of Section 3 imply that this approach is not valid when $k \geq 2$, even if we do not impose the equalities $\Phi_G \circ \beta_* = \Phi_G$.

Lemma 5.1. *Let $k \geq 1$ and $n \geq 1$, and let G be an Abelian group.*

- (i) *Let $k \geq 2$. Suppose that there exist homomorphisms $\Phi_G : B_k(\widehat{\Sigma}_{g,n}) \rightarrow G$ and $j_k : G_k \rightarrow G$ that satisfy $\Phi_G \circ \iota_k = j_k \circ p_k$. Then j_k is not injective.*
- (ii) *Let $k = 1$. For every non-trivial homomorphism $\Phi_G : B_1(\widehat{\Sigma}_{g,n}) \rightarrow G$, there exists an injective homomorphism $j_1 : G_1 \rightarrow G$ that satisfies $\Phi_G \circ j_1 = j_1 \circ p_1$.*

Proof.

- (i) Let Φ_G and j_k be as in the statement. Since G is Abelian, the homomorphism Φ_G factors through $\widehat{r}_k : B_k(\widehat{\Sigma}_{g,n}) \rightarrow B_k(\widehat{\Sigma}_{g,n})/\Gamma_2(B_k(\widehat{\Sigma}_{g,n}))$. By Proposition 3.4, $(\widehat{r}_k(\sigma_1))^2 = \sigma^2 = 1$, and so $\Phi_G(\sigma_1^2) = 1$. On the other hand, $p_k(\sigma_1) = \sigma$, and this element is of infinite order in G_k . The relation $\Phi_G \circ j_k = j_k \circ p_k$ implies the non-injectivity of j_k .
- (ii) Assume $k = 1$. As in (i), the homomorphism Φ_G factors through \widehat{r}_1 . But G_1 is isomorphic to \mathbb{Z} and generated by $p_1(\zeta_1)$ by Lemma 4.1; on the other hand $\widehat{r}_1(\zeta_1)$ is torsion free by Proposition 3.4. Thus $\widehat{r}_1 \circ j_1$ factors through p_1 . \square

This means that we cannot construct a linear representation for surface braid groups whose restriction to B_n is the Bigelow-Krammer-Lawrence representation. Furthermore if we impose that $\Phi_G \circ \beta_* = \Phi_G$ for any $\beta \in B_n(\widehat{\Sigma}_g)$, this negative result can be extended to the case $k = 1$ and to any group G . The following proposition is a reformulation in our framework of [1, Lemma 2.6] and of related remarks on the homological constraints.

Proposition 5.2. *Let $k \geq 1$ and $n \geq 2$, and let G be a group. Suppose that there exist homomorphisms $\Phi_G : B_k(\widehat{\Sigma}_{g,n}) \rightarrow G$ and $j_k : G_k \rightarrow G$ that satisfy $\Phi_G \circ \iota_k = j_k \circ p_k$ and $\Phi_G \circ \beta_* = \Phi_G$ for any $\beta \in B_n(\widehat{\Sigma}_g)$. Then j_k is not injective.*

Proof. The action of $B_n(\widehat{\Sigma}_g)$ on $B_k(\widehat{\Sigma}_{g,n})$ is completely described in Proposition 3.1. As remarked in [1, Lemma 2.6], combining relation (c.7.1) (we need therefore to suppose $n \neq 1$) with the hypothesis $\Phi_G \circ \beta_* = \Phi_G$, we see that $\Phi_G(\zeta_1) = 1$. On the other hand $p_k(\zeta_1) = \zeta$ by Proposition 3.4. The relation $\Phi_G \circ j_k = j_k \circ p_k$ implies the non-injectivity of j_k . \square

Acknowledgements. The research of the first author was supported by the French grant ANR-11-JS01-002-01, and that of the second and third authors was supported by the French grant ANR-08-BLAN-0269-02.

References

- [1] B. H. An and K. H. Ko, A family of representations of braid groups on surfaces, *Pacific J. Math.* **247** (2010), 257–282.
- [2] V. G. Bardakov, Extending representations of braid groups to the automorphism groups of free groups, *J. Knot Theory Ramifications* **14** (2005), 1087–1098.
- [3] V. G. Bardakov, Linear representations of the braid groups of some manifolds, *Acta Appl. Math.* **85** (2005), 41–48.
- [4] P. Bellingeri, On presentations of surface braid groups, *J. Algebra* **274** (2004), 543–563.
- [5] P. Bellingeri and L. Funar, Braids on surfaces and finite type invariants, *C. R. Math. Acad. Sci. Paris* **338** (2004), 157–162.
- [6] P. Bellingeri, S. Gervais and J. Guaschi, Lower central series of Artin-Tits and surface braid groups, *J. Algebra* **319** (2008), 1409–1427.
- [7] S. Bigelow, Braid groups are linear, *J. Amer. Math. Soc.* **14** (2001), 471–486.
- [8] S. Bigelow and R. Budney, The mapping class group of a genus to surface is linear, *Algebr. Geom. Topol.* **1** (2001), 699–708.
- [9] J. S. Birman, Braids, Links, and Mapping Class Groups, *Ann. Math. Stud.*, Princeton Univ. Press, vol. 82, 1974.
- [10] A. Cohen and D. B. Wales, Linearity of Artin groups of finite type, *Israel J. Math.* **131** (2002), 101–123.
- [11] F. Digne, On the linearity of Artin braid groups, *J. Algebra* **268** (2003), 39–57.
- [12] E. Fadell and L. Neuwirth, Configuration spaces, *Math. Scandinavica* **10** (1962), 111–118.
- [13] E. Fadell and J. Van Buskirk, The braid groups of E^2 and S^2 , *Duke Math. J.* **29** (1962), 243–257.
- [14] R. Fenn, D. Rolfsen and J. Zhu, Centralisers in the braid group and singular braid monoid, *Enseign. Math.* **42** (1996), 75–96.
- [15] R. H. Fox and L. Neuwirth, The braid groups, *Math. Scandinavica* **10** (1962), 119–126.
- [16] D. L. Gonçalves and J. Guaschi, On the structure of surface pure braid groups, *J. Pure Appl. Algebra* **182** (2003), 33–64 (due to a printer’s error, this article was republished in its entirety with the reference **186** (2004), 187–218).
- [17] D. L. Gonçalves and J. Guaschi, The roots of the full twist for surface braid groups, *Math. Proc. Camb. Phil. Soc.* **137** (2004), 307–320.
- [18] D. L. Gonçalves and J. Guaschi, The braid group $B_{n,m}(\mathbb{S}^2)$ and a generalisation of the Fadell-Neuwirth short exact sequence, *J. Knot Theory Ramifications* **14** (2005), 375–403.
- [19] D. L. Gonçalves and J. Guaschi, The lower central and derived series of the braid groups of the finitely-punctured sphere, *J. Knot Theory Ramifications* **18** (2009), 651–704.
- [20] D. L. Gonçalves and J. Guaschi, The Fadell-Neuwirth short exact sequence for non-orientable surfaces, *J. Pure Appl. Algebra* **214** (2010), 667–677.
- [21] D. L. Gonçalves and J. Guaschi, Surface braid groups and coverings, *J. London Math. Soc.* **85** (2012), 855–868.
- [22] R. Häring-Oldenburg and S. Lambropoulou, Knot theory in handlebodies, in *Knots 2000 Korea*, Vol. 3 (Yongpyong), *J. Knot Theory Ramifications* **11** (2002), 921–943.
- [23] D. L. Johnson, Presentation of groups, *LMS Lectures Notes* 22 (1976), Cambridge University Press.

- [24] C. Kassel and V. Turaev, Braid groups, *Graduate Texts in Mathematics* **247** Springer, New York, 2008.
- [25] D. Krammer, Braid groups are linear, *Ann. Math.* **155** (2002), 131–156.
- [26] W. Magnus, A. Karrass and D. Solitar, Combinatorial group theory, 2nd edition, Dover Publications Inc., Mineola, NY, 2004.
- [27] S. Manfredini, Some subgroups of Artin’s braid group, *Topology Appl.* **78** (1997), 123–142.
- [28] L. Paris and D. Rolfsen, Geometric subgroups of surface braid groups, *Ann. Inst. Fourier* **49** (1999), 417–472.
- [29] H. Zheng, Faithfulness of the Lawrence representation of braid group, [arXiv:math/0509074](https://arxiv.org/abs/math/0509074).

Paolo BELLINGERI, Eddy GODELLE and John GUASCHI,
Université de Caen Basse-Normandie, Laboratoire de Mathématiques Nicolas Oresme CNRS UMR 6139,
BP 5186, F-14032 Caen (France).
Email: paolo.bellingeri@unicaen.fr, eddy.godelle@unicaen.fr, john.guaschi@unicaen.fr