



**HAL**  
open science

## When Is $L_r(\mathbb{R})$ Contained in $L_p(\mathbb{R}) + L_q(\mathbb{R})$ ?

Jean-Baptiste Hiriart-Urruty, Patrice Lassère

► **To cite this version:**

Jean-Baptiste Hiriart-Urruty, Patrice Lassère. When Is  $L_r(\mathbb{R})$  Contained in  $L_p(\mathbb{R}) + L_q(\mathbb{R})$  ?. The American Mathematical Monthly, 2013, 120 (1), p. 55-61. 10.4169/amer.math.monthly.120.01.055 . hal-00966086

**HAL Id: hal-00966086**

**<https://hal.science/hal-00966086>**

Submitted on 6 Jun 2023

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

When is  $L^r(\mathbf{R})$  contained in  $L^p(\mathbf{R}) + L^q(\mathbf{R})$  ?

Jean-Baptiste Hiriart-Urruty and Lassère Patrice

### Abstract

We prove a necessary and sufficient condition on the exponents  $p, q, r \geq 1$  such that  $L^r(\mathbf{R}) \subset L^p(\mathbf{R}) + L^q(\mathbf{R})$ . In doing so, we explore the structure of  $L^p(\mathbf{R}) + L^q(\mathbf{R})$  as a normed vector space.

## 1 Introduction.

In a recent mathematical note aimed at undergraduate students and their teachers ([3]), J.-B. Hiriart-Urruty and M. Pradel proposed a way to extend the Fourier transformation to all the spaces  $L^r(\mathbf{R})$  with  $1 \leq r \leq 2$  in the following manner.

– First, they classically define the Fourier transformation on  $L^1(\mathbf{R})$ . Then they define it on  $L^2(\mathbf{R})$  using in that case the much less known Wiener’s approach which relies on a specific Hilbertian basis made of the so-called Bernstein functions.

– After having checked the coherence of both definitions on  $L^1(\mathbf{R}) \cap L^2(\mathbf{R})$ , they extend the definition of the Fourier transformation to the space  $L^1(\mathbf{R}) + L^2(\mathbf{R})$ .

– Finally, and this is the key-point, they prove the inclusion  $L^r(\mathbf{R}) \subset L^1(\mathbf{R}) + L^2(\mathbf{R})$  for all  $1 \leq r \leq 2$ , so that the Fourier transformation can be extended to all the Lebesgue spaces  $L^r(\mathbf{R})$ .

An obvious question which arises is, what happens if  $r \notin [1, 2]$  ? For example, is the inclusion  $L^3(\mathbf{R}) \subset L^1(\mathbf{R}) + L^2(\mathbf{R})$  true or not ? The objective of the present mathematical note is to answer this question. We provide a necessary and sufficient condition for the inclusion  $L^r(\mathbf{R}) \subset L^p(\mathbf{R}) + L^q(\mathbf{R})$  to hold true. For that purpose, concentrate on the sum  $L^p(\mathbf{R}) + L^q(\mathbf{R})$  and see how it is structured as a normed vector space.

## 2 How to norm a sum of normed vector spaces ?

Let  $V$  and  $W$  be two vector subspaces of a "holdall" vector space,  $E$ . If  $V$  is equipped with a norm  $\|\cdot\|_1$  and  $W$  with a norm  $\|\cdot\|_2$ , is there a natural way to define a norm on the vector subspace  $V + W$  ? Of course, we do not assume

that  $V \cap W = \{O_E\}$ . If this was the case, a natural way to define a norm  $N$  on  $V + W$  would be

$$N(u) = \|v\|_1 + \|w\|_2,$$

whenever  $u \in V + W$  is (uniquely) decomposed as  $u = v + w$ , with  $v \in V$  and  $w \in W$ .

What we have in mind is indeed  $V = L^p(\mathbf{R})$ ,  $W = L^q(\mathbf{R})$  and  $E = L(\mathbf{R})$  as the “holdall” vector space ( $L(\mathbf{R})$  stands for the set of all Lebesgue classes of measurable functions on  $\mathbf{R}$ ).

The theorem below answers the question posed above. It does not seem to be well-known, except by people who have to deal with the interpolation of functional spaces (like in [1]).

**Theorem 1** *Let  $N$  be defined on  $V + W$  as follows*

$$N(u) := \inf\{\|v\|_1 + \|w\|_2 ; u = v + w \text{ with } v \in V \text{ and } w \in W\}. \quad (1)$$

*Then,  $N$  is a semi-norm on  $V + W$ . It is a norm under the following “compatibility” assumption :*

$$(T) \quad \left. \begin{array}{l} z_k \in V \cap W, \\ z_k \longrightarrow a \text{ in } (V, \|\cdot\|_1) \\ z_k \longrightarrow b \text{ in } (W, \|\cdot\|_2) \end{array} \right\} \implies a = b.$$

**Proof:** To check that  $N(O_E) = 0$ ,  $N(\lambda u) = |\lambda|N(u)$  for all  $\lambda \in \mathbf{R}$ ,  $u \in V + W$ , and  $N(u_1 + u_2) \leq N(u_1) + N(u_2)$  for all  $u_1$  and  $u_2$  in  $V + W$ , does not raise any difficulty. It suffices to use the definition of the lower bound (or infimum) of a set of real numbers.

To prove that  $N(u) = 0$  implies that  $u = O_E$  is a bit more tricky. Our experience with that question with undergraduate students shows that they usually fail to answer it correctly. Their common mistake is to deduce that a sequence  $(v_k + w_k)_k$  converges to  $O_E$  using the fact that  $(v_k)_k$  converges to  $O_E$  in  $V$  and  $(w_k)_k$  converges to  $O_E$  in  $W$ . We therefore provide a proof here.

We first begin by observing that

$$\nu : V \cap W \ni u \mapsto \nu(u) := \max(\|u\|_1, \|u\|_2) \quad (2)$$

is a norm on  $V \cap W$  ; this is an easy result to prove.

Consider therefore,  $u \in V + W$  such that  $N(u) = 0$ . We take for example

$$u = v + w, \quad \text{with } v \in V \quad \text{and } w \in W. \quad (3)$$

Due to the definition (1) of  $N(u)$ , for all positive integers  $k$ , there exists  $v_k \in V$  and  $w_k \in W$  such that

$$u = v_k + w_k, \quad \text{and} \quad (4)$$

$$\|v_k\|_1 + \|w_k\|_2 \leq N(u) + \frac{1}{k} = \frac{1}{k}. \quad (5)$$

Thus,  $(v_k)_k$  converges to  $O_E$  in  $V$  and  $(w_k)_k$  converges to  $O_E$  in  $W$ . But what about  $(v_k + w_k)_k$ ? Recall that no norm, hence, no topology, has yet been defined on  $V + W$ . We infer from (3) and (4) that

$$v + w = v_k + w_k,$$

and thus

$$v - v_k = w_k - w. \tag{6}$$

As a consequence, this common vector  $z_k := v - v_k = w_k - w$  lies in  $V \cap W$  and, since  $\nu(z_k) = \|v - v_k\|_1$  or  $\|w_k - w\|_2$ ,

$$z_k \rightarrow v \text{ in } (V, \|\cdot\|_1) \text{ and } z_k \rightarrow -w \text{ in } (W, \|\cdot\|_2).$$

The assumption  $(\mathcal{T})$  then ensures that

$$v = -w, \text{ that is } u = v + w = O_E.$$

■

Note that the technical “compatibility” assumption  $(\mathcal{T})$  is satisfied

- trivially if  $V \cap W = \{O_E\}$  (in that case it amounts to  $0 = 0$ ).
- in the cases where  $V = L^p(\mathbb{R})$ ,  $W = L^q(\mathbb{R})$  (indeed, convergence of  $(f_k)_k$  towards  $f$  in  $L^p(\mathbb{R})$  implies convergence almost everywhere of a subsequence of  $(f_k)_k$  towards  $f$ ).
- when the “holdall” vector space  $E$  is a Hausdorff topological vector space in which  $V$  and  $W$  are continuously imbedded.

We suppose that  $(\mathcal{T})$  is in force for the rest of the section.

The vector space  $V+W$ , equipped with the norm  $N$  as defined in (1), inherits some properties of  $(V, \|\cdot\|_1)$  and  $(W, \|\cdot\|_2)$ . Here is one.

**Theorem 2** *If  $(V, \|\cdot\|_1)$  and  $(W, \|\cdot\|_2)$  are Banach spaces, then so is  $(V+W, N)$ .*

**Proof :** The proof of this theorem offers the opportunity to use a characterization of completeness of normed vector spaces which is not well-known. Let  $(X, \|\cdot\|)$  be a normed space. We have indeed

$$((X, \|\cdot\|) \text{ is complete}) \iff \left( \begin{array}{l} \text{Every series in } X, \text{ of general term } a_k, \text{ for} \\ \text{which } \sum_{k=0}^{\infty} \|a_k\| < +\infty \text{ does converges in } X \\ \text{(toward a sum denoted as } \sum_{k=0}^{\infty} a_k) \end{array} \right). \tag{7}$$

The implication  $(\Rightarrow)$  is classical, it is the most often used. The converse implication  $(\Leftarrow)$  is not often used, but we have an opportunity to do that here. For a proof of the equivalence (7), see for example ([6], Theorems 2-XIV-2.1

and 2.2, pages 164-165), ([4], page 20) or ([7], pages 262 and 270); it is also sketched in ([1], page 24). The proof is not very difficult however, but readers are encouraged to work through it themselves. We now proceed in this manner to a proof of Theorem 2.

Consider a series in  $V + W$ , of general term  $u_k$  for which  $\sum_{k=0}^{\infty} N(u_k) < \infty$ . We have to prove that  $\sum_{k=0}^n u_k$  converges as  $n \rightarrow +\infty$ , to some element  $u \in V + W$ . In view of the definition (1) of  $N$ , for all positive integer  $k$ , there exist  $v_k \in V$  and  $w_k \in W$  satisfying

$$u_k = v_k + w_k, \text{ and} \quad (8)$$

$$\|v_k\|_1 + \|w_k\|_2 \leq N(u_k) + \frac{1}{k^2}. \quad (9)$$

Thus,  $\sum_{k=0}^{\infty} \|v_k\|_1 < +\infty$  and  $\sum_{k=0}^{\infty} \|w_k\|_2 < +\infty$ . Since both  $(V, \|\cdot\|_1)$  and  $(W, \|\cdot\|_2)$  have been assumed to be complete, there exist  $v \in V$  and  $w \in W$  such that

$$\sum_{k=0}^n v_k \xrightarrow[n \rightarrow +\infty]{} v \text{ in } V, \text{ and } \sum_{k=0}^n w_k \xrightarrow[n \rightarrow +\infty]{} w \text{ in } W. \quad (10)$$

Let  $u := v + w \in V + W$ . Let us check that, as expected,  $\sum_{k=0}^n u_k$  converges to  $u$  in  $(V + W, N)$ . Indeed,

$$N\left(u - \sum_{k=0}^n u_k\right) = N\left(u + v - \sum_{k=0}^n (v_k + w_k)\right) \leq \left\|v - \sum_{k=0}^n v_k\right\|_1 + \left\|w - \sum_{k=0}^n w_k\right\|_2.$$

It remains to apply (10) to get the desired result,  $N(u - \sum_{k=0}^n u_k) \xrightarrow[n \rightarrow +\infty]{} 0$ . ■

**Comments :** The construction of the norm  $\nu$  on  $V \cap W$  and  $N$  on  $V + W$  deserves some geometrical interpretation. Even if  $\|\cdot\|_1$  (resp.  $\|\cdot\|_2$ ) is only defined on  $V \subset E$  (resp. on  $W \subset E$ ), we can extend it to the whole of  $E$  by setting  $\|u\|_1 = +\infty$  if  $u \notin V$  (resp.  $\|u\|_2 = +\infty$  if  $u \notin W$ ). We still denote by  $\|\cdot\|_1$  and  $\|\cdot\|_2$  the extended functions.

Clearly,  $\|\cdot\|_1$  and  $\|\cdot\|_2$  are convex positively homogeneous functions on  $E$ . Modern convex analysis accepts and can handle convex functions possibly taking the value  $+\infty$  ([5]). An important geometrical object associated with a convex function  $f : E \rightarrow \mathbf{R} \cup \{+\infty\}$  is its so-called strict epigraph

$$\text{epi}_s f := \{(x, r) \in E \times \mathbf{R} : f(x) < r\}$$

(literally, what is strictly above the graph of  $f$ ). In our situation,  $K_1 := \text{epi}_s \|\cdot\|_1$  and  $K_2 := \text{epi}_s \|\cdot\|_2$  are open convex cones of  $E$ . So, what are the strict epigraphs of the norm functions  $\nu$  and  $N$ ? One easily checks the following

$$\begin{aligned} \text{epi}_s \nu &= (\text{epi}_s \|\cdot\|_1) \cap (\text{epi}_s \|\cdot\|_2), \text{ and} \\ \text{epi}_s N &= (\text{epi}_s \|\cdot\|_1) + (\text{epi}_s \|\cdot\|_2). \end{aligned} \quad (11)$$

The sets where  $\nu$  (resp.  $N$ ) is finite, called the domain of  $\nu$  (resp. of  $N$ ) in convex analysis, is just  $V \cap W$  (resp.  $V + W$ ).

The binary operation which builds a convex function  $f$  from two other ones  $f_1$  and  $f_2$ , via the geometric construction

$$\text{epi}_s f = \text{epi}_s f_1 + \text{epi}_s f_2$$

is called the *infimal convolution* of  $f_1$  and  $f_2$  ([2],[5]). This operation enjoys properties similar to the usual (integral) convolution in classical analysis.

In brief, the norm  $N$  has been designed as an infimal convolution of the norms  $\|\cdot\|_1$  and  $\|\cdot\|_2$ .

Returning to our particular setting  $E = L(\mathbf{R})$ ,  $V = L^p(\mathbf{R})$  and  $W = L^q(\mathbf{R})$  with  $1 \leq p, q < +\infty$ , the vector space  $L^p(\mathbf{R}) + L^q(\mathbf{R})$  can be equipped with a norm that we denote  $\|\cdot\|_{p,q}$  as follows,

$$\|f\|_{p,q} = \inf\{\|g\|_p + \|h\|_q ; f = g + h \text{ with } g \in L^p(\mathbf{R}) \text{ and } h \in L^q(\mathbf{R})\}. \quad (12)$$

As proved in Theorem 2,  $(L^p(\mathbf{R}) + L^q(\mathbf{R}), \|\cdot\|_{p,q})$  is a Banach space.

### 3 Comparing $L^r(\mathbf{R})$ with $L^p(\mathbf{R}) + L^q(\mathbf{R})$

We know that the Lebesgue spaces  $L^r(\mathbf{R})$  and  $L^s(\mathbf{R})$  (for  $1 \leq r \neq s < +\infty$ ) cannot be compared. Neither  $L^r(\mathbf{R})$  is contained in  $L^s(\mathbf{R})$  nor the converse. A direct comparison is however possible if we deal with the sum of these spaces. Here is the main result of this section.

**Theorem 3** *If  $1 \leq p < q < +\infty$ , then we have the following :*

1.  $L^r(\mathbf{R})$  is contained in  $L^p(\mathbf{R}) + L^q(\mathbf{R})$  whenever  $r \in [p, q]$ ,
2. if  $1 \leq r < +\infty$  does not lie in  $[p, q]$ , then  $L^r(\mathbf{R})$  is not contained in  $L^p(\mathbf{R}) + L^q(\mathbf{R})$ .

**Proof :** (1) Let  $f \in L^r(\mathbf{R})$  and consider  $X := \{x \in \mathbf{R} : |f(x)| > 1\}$  (a measurable set defined within a set of null measure) as well as  $X^c = \mathbf{R} \setminus X$  (the complementary set of  $X$  in  $\mathbf{R}$ ). We decompose  $f$  as follows :

$$f = f_1 + f_2, \text{ with } f_1 = f \cdot \mathbf{1}_X \text{ and } f_2 = f \cdot \mathbf{1}_{X^c}. \quad (13)$$

We claim that (13) provides an explicit decomposition of  $f$  in  $L^p(\mathbf{R}) + L^q(\mathbf{R})$ , that is to say  $f_1 \in L^p(\mathbf{R})$  and  $f_2 \in L^q(\mathbf{R})$ . We first prove that,

$$\int_{\mathbf{R}} |f_1(x)|^p d\lambda(x) = \int_X |f(x)|^p d\lambda(x) = \int_X |f(x)|^{p-r} \cdot |f(x)|^r d\lambda(x). \quad (14)$$

For  $x \in X$ ,  $|f(x)| > 1$  and, since the exponent  $p-r$  is nonpositive,  $|f(x)|^{p-r} \leq 1$ . Consequently, the last integral in the string of equalities (14) is bounded above by  $\int_X |f(x)|^r d\lambda(x)$ . Finally,

$$(14') \quad \int_{\mathbf{R}} |f_1(x)|^p d\lambda(x) \leq \int_X |f(x)|^r d\lambda(x) \leq \int_{\mathbf{R}} |f(x)|^r d\lambda(x) < +\infty.$$

We thus have proved that  $f_1 \in L^p(\mathbf{R})$ .

Second, we prove that  $f_2 \in L^q(\mathbf{R})$ . Indeed,

$$\int_{\mathbf{R}} |f_2(x)|^q d\lambda(x) = \int_{X^c} |f(x)|^q d\lambda(x) = \int_{X^c} |f(x)|^{q-r} \cdot |f(x)|^r d\lambda(x). \quad (15)$$

For  $x \in X^c$ ,  $|f(x)| \leq 1$  and, since the exponent  $q-r$  is nonnegative,  $|f(x)|^{q-r} \leq 1$ . Again, the last integral in the string of equalities (15) is bounded above by  $\int_{X^c} |f(x)|^r d\lambda(x)$ . As a result,

$$\int_{\mathbf{R}} |f_2(x)|^q d\lambda(x) = \int_{X^c} |f(x)|^r d\lambda(x) = \int_{\mathbf{R}} |f(x)|^r d\lambda(x) < +\infty. \quad (16)$$

We therefore, have proved that  $f_2 \in L^q(\mathbf{R})$ .

(2) The second part of Theorem 3 is a bit harder to prove (like most of the negative results in mathematics). We actually have to distinguish two cases for  $r$  in the segment  $[p, q]$  :  $r < p$  and  $r > q$ .

**Case 1 :**  $r < p$ . Choose  $\alpha$  satisfying  $1/p < \alpha < 1/r$  and let  $f$  be defined on  $\mathbf{R}$  by  $f(x) = x^{-\alpha} \mathbf{1}_{(0,1]}(x)$ .

Since  $|f(x)|^r = x^{-\alpha r}$  for  $x \in (0, 1]$  and 0 elsewhere, the choice of  $\alpha$  implies that  $f \in L^r(\mathbf{R})$  (since  $\alpha r < 1$ ). The same argument shows that  $f \notin L^p(\mathbf{R})$  (since  $\alpha p > 1$ ).

Suppose now that  $L^r(\mathbf{R}) \subset L^p(\mathbf{R}) + L^q(\mathbf{R})$ . Then

$$f \in L^r(\mathbf{R}) \subset L^p(\mathbf{R}) + L^q(\mathbf{R}) \subset L^p([0, 1]) + L^q([0, 1]). \quad (17)$$

But since  $[0, 1]$  is of Lebesgue finite measure and  $p < q$ ,  $L^q([0, 1])$  is contained in  $L^p([0, 1])$ , so that (17) yields that  $f \in L^p([0, 1])$ . This is not the case.

Thus we have proved that  $L^r(\mathbf{R})$  is not contained in  $L^p(\mathbf{R}) + L^q(\mathbf{R})$ .

**Case 2 :**  $r > q$ . Our proof in this case relies on a technical lemma that we present separately.

**Lemma 1** *Let  $1 \leq p < q < +\infty$ , let  $\Omega$  be a measurable subset of  $\mathbf{R}$ , and let  $f \in L^p(\Omega) + L^q(\Omega)$ . Then  $f \in L^q(\Omega)$  whenever it is essentially bounded on  $\Omega$ .*

**Proof of the Lemma :** Let  $f$  be decomposed as  $f = f_p + f_q$ , with  $f_p \in L^p(\Omega)$  and  $f_q \in L^q(\Omega)$ . So, to prove that  $f \in L^q(\Omega)$  amounts to proving that  $f_p \in L^q(\Omega)$ .

Let  $X := \{x \in \mathbf{R} : |f_p(x)| > 1\}$ . To show that  $\int_{\Omega} |f_p(x)|^q d\lambda(x)$  is finite, we cut it into two pieces:  $\int_{\Omega \cap X^c} |f_p(x)|^q d\lambda(x)$  and  $\int_{\Omega \cap X} |f_p(x)|^q d\lambda(x)$ .

Consider the first piece. Since  $f_p \in L^p(\mathbf{R})$ , the set  $X$  is of finite (Lebesgue) measure. Now with the definition of  $X$  and the fact that  $q - p > 0$ , we obtain

$$\begin{aligned} \int_{\Omega \cap X^c} |f_p(x)|^q d\lambda(x) &= \int_{\Omega \cap X^c} |f_p(x)|^{q-p} \cdot |f_p(x)|^p d\lambda(x) \\ &\leq \int_{\Omega \cap X^c} |f_p(x)|^p d\lambda(x) \leq \int_{\mathbf{R}} |f_p(x)|^p d\lambda(x) < +\infty. \end{aligned}$$

This concludes the argument for the first piece.

Consider now the second piece. Since  $f$  has been assumed essentially bounded on  $\Omega$ ,

$$|f_p(x)| \leq |f(x) - f_q(x)| \leq \|f\|_{\infty} + |f_q(x)| \quad \text{for almost all } x \text{ in } \Omega.$$

Consequently,

$$\int_{\Omega \cap X} |f_p(x)|^q d\lambda(x) \leq \int_{\Omega \cap X} (\|f\|_{\infty} + |f_q(x)|)^q d\lambda(x). \quad (18)$$

The convexity of the function  $t \mapsto t^q$  on  $[0, +\infty)$  implies that  $(\|f\|_{\infty} + |f_q(x)|)^q \leq 2^{q-1} (\|f\|_{\infty}^q + |f_q(x)|^q)$ . So, we pursue the string of inequalities (18) with

$$\begin{aligned} \int_{\Omega \cap X} |f_p(x)|^q d\lambda(x) &\leq 2^{q-1} \int_{\Omega \cap X} (\|f\|_{\infty}^q + |f_q(x)|^q) d\lambda(x) \\ &\leq 2^{q-1} [\lambda(X) \|f\|_{\infty}^q + (\|f_q\|_q)^q]. \end{aligned}$$

To summarize, we have proved that

$$\int_{\Omega} |f_p(x)|^q d\lambda(x) = \int_{\Omega \cap X^c} |f_p(x)|^q d\lambda(x) + \int_{\Omega \cap X} |f_p(x)|^q d\lambda(x) < +\infty.$$

Thus,  $f_p \in L^p(\Omega)$ , which was our objective. That concludes the proof of the technical lemma.  $\square$

Let us go back to the second part of the proof of Theorem 3, the case where  $r > q$ . Choose  $\alpha$  satisfying  $1/r < \alpha < 1/q$  and let  $f$  be defined on  $\mathbf{R}$  by  $f(x) = x^{-\alpha} \mathbf{1}_{[1, +\infty)}(x)$ .

Since  $|f(x)|^r = x^{-\alpha r}$  for  $x \in [1, +\infty)$  and 0 elsewhere, the choice of  $\alpha$  implies that  $f \in L^r(\Omega)$  (since  $\alpha r > 1$ ). But also  $f$  is essentially bounded on  $\mathbf{R}$ . If  $f$  were in  $L^p(\mathbf{R}) + L^q(\mathbf{R})$ , the technical lemma would imply that  $f \in L^q(\mathbf{R})$ . But, this is not the case since  $|f(x)|^q = x^{-\alpha q}$  for  $x \in [1, +\infty)$  and 0 elsewhere, the choice of  $\alpha$  implies that  $\alpha q < 1$ .

Thus, again in the case where  $r > q$ ,  $L^r(\mathbf{R})$  is not contained in  $L^p(\mathbf{R}) + L^q(\mathbf{R})$ .  $\blacksquare$

We end this note with the following observation, which links sections 2 and 3. In the first part of Theorem 3, we have proved that  $L^r(\mathbf{R}) \subset L^p(\mathbf{R}) + L^q(\mathbf{R})$

whenever  $r \in [p, q]$ . In the course of its proof, a simple explicit decomposition of  $f \in L^r(\mathbf{R})$  as  $f = f_1 + f_2$ , with  $f_1 \in L^p(\mathbf{R})$  and  $f_2 \in L^q(\mathbf{R})$  has been provided (see (13) and the upper bounds (14') and (16)). Indeed, as a consequence of (14) and (16),

$$\|f\|_{p,q} \leq \|f_1\|_p + \|f_2\|_q \leq \|f\|_r^{r/p} + \|f\|_r^{r/q}. \quad (19)$$

Hence, the injection of  $L^r(\mathbf{R})$  into  $L^p(\mathbf{R}) + L^q(\mathbf{R})$  is continuous ; therefore, there exists  $C > 0$  such that

$$\|f\|_{p,q} \leq C\|f\|_r. \quad (20)$$

Indeed, using the inequality (19), we can get an upper bound for the norm of this injection (a somewhat complicated expression in terms of  $p, q, r$ ). This result complements a more classical one which says that, when  $r \in [p, q]$ ,  $L^p(\mathbf{R}) \cap L^q(\mathbf{R})$  is contained in  $L^r(\mathbf{R})$ , then the injection is continuous.

## References

- [1] J. Bergh; J.-B. L ofstr om, *Interpolation Spaces*, Grundlehren der mathematischen Wissenschaften, **223**, Springer-Verlag, New-York, 1976.
- [2] J.-B. Hiriart-Urruty; C. Lemarechal, *Fundamentals of Convex Analysis*, Grundlehren TextEditions, Springer-Verlag, New-York, 2001.
- [3] J.-B. Hiriart-Urruty; M. Pradel, *La transformation de Fourier de fonctions : au-del a de  $L^1$  et  $L^2$* , Revue de Math ematiques Sp eciales, **113** no.2 (2003) 25-33.
- [4] R.E. Megginson, *An Introduction to Banach Space Theory*, Graduate Texts in Mathematics, **183**. Springer-Verlag, New-York, 1998.
- [5] R.T. Rockafellar, *Convex Analysis*, Princeton University Press, 1970.
- [6] L. Schwartz, *Topologie G en erale et Analyse Fonctionnelle*, Hermann, Paris, 1970.
- [7] Y. Sonntag, *Topologie et Analyse Fonctionnelle*, Ellipses, Paris, 1998.

**JEAN-BAPTISTE HIRIART-URRUTY** Has been a professor of Mathematics at the Paul Sabatier University since 1981. He recieved his Doctorat  es sciences math ematiques from the Blaise Pascal University (Clermont-Ferrand) in 1977. His current research interests include Variational analysis (convex, nonsmooth, applied) and all the aspects of Optimization.

*Institute of Mathematics, Paul Sabatier University, TOULOUSE, FRANCE*  
*jbhu@cict.fr.*

[www.math.univ-toulouse.fr/~jbhu/](http://www.math.univ-toulouse.fr/~jbhu/)

**PATRICE LASSÈRE** received his Doctorat ès sciences mathématiques from the University Paul Sabatier, Toulouse, France, in 1991. He currently teaches at the University Paul Sabatier. His research interests includes complex analysis in one or several variables and functional analysis.

*Institute of Mathematics, Paul Sabatier University, TOULOUSE, FRANCE*

*lassere@math.ups-tlse.fr*

<http://www.math.univ-toulouse.fr/~lassere/>