



**HAL**  
open science

## On the Malliavin differentiability of BSDEs

Thibaut Mastrolia, Dylan Possamaï, Anthony Réveillac

► **To cite this version:**

Thibaut Mastrolia, Dylan Possamaï, Anthony Réveillac. On the Malliavin differentiability of BSDEs. 2014. hal-00971728v1

**HAL Id: hal-00971728**

**<https://hal.science/hal-00971728v1>**

Preprint submitted on 3 Apr 2014 (v1), last revised 11 Jun 2015 (v2)

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# On the Malliavin differentiability of BSDEs

Thibaut Mastrolia<sup>\*</sup>     Dylan Possamai<sup>†</sup>     Anthony Réveillac<sup>‡</sup>

Université Paris-Dauphine  
CEREMADE UMR CNRS 7534  
Place du Maréchal De Lattre De Tassigny  
75775 Paris cedex 16 France

April 3, 2014

## Abstract

In this paper we provide new conditions for the Malliavin differentiability of solutions of Lipschitz or quadratic BSDEs. Our results rely on the interpretation of the Malliavin derivative as a Gâteaux derivative in the directions of the Cameron-Martin space. Incidentally, we provide a new formulation for the characterization of the Malliavin-Sobolev type spaces  $\mathbb{D}^{1,p}$ .

*Key words:* Malliavin's calculus; abstract Wiener space; BSDEs.

*AMS 2010 subject classification:* Primary: 60H10; Secondary: 60H07.

## 1 Introduction

Backward Stochastic Differential Equations (BSDEs) have been studied extensively in the last two decades as they naturally arise in the context of stochastic control problems (for instance in Finance see [5]), and as they provide a probabilistic representation for solution to semi-linear parabolic PDEs, via a non-linear Feynman-Kac formula (see [12]). Before going further let us recall that this class of equations has been introduced in [2, 11, 12] and that a BSDE can be formulated as:

$$Y_t = \xi + \int_t^T f(s, Y_s, Z_s) ds - \int_t^T Z_s dW_s, \quad t \in [0, T], \quad (1.1)$$

where  $T$  is a fixed positive number,  $W := (W_t)_{t \in [0, T]}$  is a one-dimensional Brownian motion defined on a probability space  $(\Omega, \mathcal{F}_T, \mathbb{P})$  with natural filtration  $(\mathcal{F}_t)_{t \in [0, T]}$ . The data of the equation are the  $\mathcal{F}_T$ -measurable r.v.  $\xi$  which is called the terminal condition (as  $Y_T = \xi$ ) and the mapping  $f : [0, T] \times \Omega \times \mathbb{R}^2 \rightarrow \mathbb{R}$  which is a progressively measurable process and where according to the notations used in the literature we write  $f(t, y, z)$  for  $f(t, \omega, y, z)$ . A solution

---

<sup>\*</sup>mastrolia@ceremade.dauphine.fr

<sup>†</sup>possamai@ceremade.dauphine.fr

<sup>‡</sup>anthony.reveillac@ceremade.dauphine.fr

to the BSDE (1.1) is then a pair of predictable processes  $(Y, Z)$  (with appropriate integrability properties) such that Relation (1.1) holds  $\mathbb{P}$ -a.s.

When dealing with applications, one needs to obtain regularity properties on the solution  $(Y, Z)$ , such as the Malliavin differentiability of the random variables  $Y_t, Z_t$  at a given time  $t$  in  $[0, T]$  (note that for the  $Z$  component this question needs to be clarified a little bit because of the definition of  $Z$ , cf. Theorem 5.1 for a precise statement). More precisely, one needs to answer the following question:

Which conditions on the data  $\xi$  and  $f$  in (1.1) ensure that  $Y_t, Z_t$  are Malliavin differentiable?

This question was first addressed in the paper [12] in a Markovian setting, that is when  $\xi := g(X_T)$  and  $f(t, \omega, y, z) := h(t, X_t(\omega), y, z)$  where  $g : \mathbb{R} \rightarrow \mathbb{R}$  and  $h : [0, T] \times \mathbb{R}^3 \rightarrow \mathbb{R}$  are regular enough deterministic functions and  $X := (X_t)_{t \in [0, T]}$  is the unique solution to a SDE of the form:

$$X_t = X_0 + \int_0^t \sigma(s, X_s) dW_s + \int_0^t b(s, X_s) ds, \quad t \in [0, T],$$

with regular enough coefficients  $\sigma, b : [0, T] \times \mathbb{R} \rightarrow \mathbb{R}$ . In that framework, Pardoux and Peng proved in [12, Proposition 2.2] that, under (essentially) the following conditions:

(PP1)  $g$  is continuously differentiable with bounded derivative.

(PP2)  $h$  is continuously differentiable in  $(x, y, z)$  with bounded derivatives (uniformly in time),

$Y_t$  is Malliavin differentiable at any time  $t$  (with a similar statement for  $Z$ ) and the Malliavin derivatives of  $Y$  and  $Z$  provide a solution to an explicit linear BSDE. To be more precise, in [12] the authors make one assumption for the whole paper which is stronger than (PP1)-(PP2) above, however a careful reading of the proof of [12, Proposition 2.2] enables one to conclude that Conditions (PP1)-(PP2) are sufficient to obtain the Malliavin differentiability of the solution. Assumptions (PP1)-(PP2) look pretty intuitive since they basically require the Malliavin differentiability of the terminal condition  $\xi$  and of the generator  $f$  once the component  $(y, z)$  are frozen, i.e., of the process  $(t, \omega) \mapsto f(t, \omega, y, z)$  for given  $(y, z)$ . Hence, it is natural to expect that the latter conditions can be easily generalized to the non-Markovian framework. Unfortunately, the first result in that direction which was given by El Karoui, Peng and Quenez in [5] requires more stringent conditions than the aforementioned intuitive ones. More explicitly, the main result in [5] concerning the Malliavin differentiability of the solution to the BSDE (1.1) (essentially) involves the following conditions (see [5, Proposition 5.3] for a precise statement):

(EPQ1)  $\xi$  is Malliavin differentiable<sup>1</sup> and  $\mathbb{E}[|\xi|^4] < +\infty$ .

(EPQ2) At any time  $t \in [0, T]$ , the r.v.  $\omega \mapsto f(t, \omega, Y_t, Z_t)$  is Malliavin differentiable<sup>2</sup> with Malliavin derivative denoted by  $D_\theta f(t, Y_t, Z_t)$  such that there exists a predictable process  $K^\theta := (K_t^\theta)_{t \in [0, T]}$  with  $\int_0^T \mathbb{E}[(\int_0^T |K_s^\theta|^2 ds)^2] d\theta < +\infty$ , and such that for any  $(y_1, y_2, z_1, z_2) \in \mathbb{R}^4$  it holds for a.e.  $\theta \in [0, T]$  that:

$$|D_\theta f(t, \omega, y_1, z_1) - D_\theta f(t, \omega, y_2, z_2)| \leq K_t^\theta(\omega)(|y_1 - y_2| + |z_1 - z_2|).$$

---

<sup>1</sup>is in  $\mathbb{D}^{1,2}$

<sup>2</sup>in fact as an adapted process it belongs to  $\mathbb{D}^{1,2}$ , we refer to the space  $\mathbb{L}_{1,2}^\theta$  whose precise definition is recalled in [5, p. 58]

Roughly speaking, this means that  $\xi$  and  $\omega \mapsto f(t, \omega, y, z)$  have to be Malliavin differentiable (which is intuitively the minimal expected requirement), but in order to prove that  $Y$  and  $Z$  are Malliavin differentiable, one needs to enforce an extra regularity conditions on each of the data: that is  $\xi$  has a finite moment of order 4, and that the Malliavin derivative of the driver  $f$  is Lipschitz continuous in  $(y, z)$  with a stochastic Lipschitz constant  $K$  which is sufficiently integrable. Note that a careful reading of the proof allows one to conclude that the moment conditions on  $\xi$  and  $Df$  can actually be relaxed to hold only in  $L^{2+\varepsilon}$  for some  $\varepsilon > 0$ . Besides, as noted in [5, Remark at the bottom of p. 59], if  $K$  is bounded then the proof can be modified so that the extra integrability condition on  $\xi$  (i.e.  $\mathbb{E}[|\xi|^4] < +\infty$ ) can be dropped. However, even in that case, one can check that in the Markovian framework, Conditions (EPQ1)-(EPQ2) are strictly stronger than Conditions (PP1)-(PP2).

The aim of this paper is to provide an alternative sufficient condition to (EPQ1)-(EPQ2) for the Malliavin differentiability of the solution to a BSDE of the form (1.1) in the general non-Markovian setting. Our main result in that direction is Theorem 5.1 below, which uses a fundamentally different approach from [5, 12], as well as different type of assumptions. Since they involve some notations concerning the analysis on the Wiener space, we refrain from detailing them immediately, and rather explain informally what are the main differences between our approach and the one of [5]. A natural way to solve a BSDE of the form (1.1) when the driver  $f$  is Lipschitz in  $(y, z)$  is to make use of a Picard iteration, that is to say a family  $(Y^n, Z^n)$  of solutions to BSDEs satisfying

$$Y_t^n = \xi + \int_t^T f(s, Y_s^{n-1}, Z_s^{n-1}) ds - \int_t^T Z_s^n dW_s, \quad t \in [0, T], \quad (1.2)$$

where  $Y^0 \equiv Z^0 \equiv 0$ . Then, a fixed point argument allows one to construct, in appropriate spaces, a solution  $(Y, Z)$  to Equation (1.1). If  $\xi$  and  $f(t, y, z)$  are Malliavin differentiable (that is in the domain of the Malliavin derivative  $\mathbb{D}^{1,2}$  which is a Banach space equipped with a Sobolev type norm), then so is  $(Y^n, Z^n)$ . Then, it just remains to prove that this property extends to  $Y, Z$  which are limits (in the appropriate spaces) of respectively  $Y^n$  and  $Z^n$ . More precisely this is done by a uniform (in  $n$ ) control of the Sobolev norms of  $Y^n, Z^n$  or equivalently by proving that the Malliavin derivatives  $(DY^n, DZ^n)$  of  $(Y^n, Z^n)$  converge to the solution of a linear BSDE whose solution will be the Malliavin derivatives  $(DY, DZ)$  of  $Y$  and  $Z$ . This last step is exactly where the extra regularity (EPQ1)-(EPQ2) is needed. It appears quite clearly that for this approach, the conditions of [5] cannot be optimized in the general case. Even though this idea seems pretty natural, it is based on a choice which is somehow arbitrary. Indeed, a necessary condition for  $DY_t$  to be well-defined (at a given time  $t$ ) is that there exists a sequence of random variables  $(F^n)_n$  which converges to  $Y_t$  in  $L^2$  such that each variable  $F^n$  is Malliavin differentiable with derivative  $DF^n$  and such that  $DF^n$  converges (with respect to a suitable norm) to  $DY_t$ . As a consequence, in a sense in the approach described above one believes that this sequence  $(F^n)_n$  can be chosen to be the Picard iteration  $(Y^n)_n$ . Once again, this idea looks very natural (according to the same type of proofs for SDEs) but then one sees that in the BSDE framework this intuitive idea leads to pretty heavy assumptions. We elaborate a lithe bit more on this point in Section 6.3.

Regarding the discussion above, one could think of trying to find a sequence of processes which are known to approximate the Malliavin derivative of  $Y$  (and  $Z$ ) when  $Y$  is Malliavin differen-

tiable. This approximation is provided by the well-known interpretation of the Malliavin derivative as a Gâteaux derivative in the directions of the Cameron-Martin space. More precisely, a necessary condition for  $Y_t$  to belong to  $\mathbb{D}^{1,2}$ , is that for any absolutely continuous function  $h$  starting from 0 at 0 with derivative denoted  $\dot{h}$ , the difference quotient  $\varepsilon^{-1}(Y_t(\omega + \varepsilon h) - Y_t(\omega))$  converges (in a sense to be made precise) as  $\varepsilon$  goes to 0 to  $\langle DY_t, \dot{h} \rangle_{L^2([0,T])}$ . This fact was initially given by Malliavin and then extended by Stroock, Shigekawa, Kusuoka and Sugita in a series of papers [9, 15, 14, 8, 16]. In addition, Sugita proved in [16] that a r.v.  $F$  is Malliavin differentiable if it is *ray absolutely continuous*<sup>1</sup> and if it is stochastically Gâteaux differentiable. Using the main ideas of [16] we provide incidentally a new formulation of the characterization of the Malliavin-Sobolev type spaces  $\mathbb{D}^{1,p}$  in Theorem 4.1. Since we did not find explicitly this characterization in the literature, we believe that this result is new and maybe interesting by itself. The main point is that this formulation is especially handy when dealing with stochastic equations like BSDEs. With this result at hand, we obtain new conditions (see Assumptions (D),  $(H_1)$  and  $(H_2)$  at the beginning of Section 5) for  $Y, Z$  to be Malliavin differentiable, see Theorem 5.1. Our assumptions refine those of [12, 5] in the Markovian case, and our approach is directly applicable to quadratic growth BSDEs since we do not rely on any approximation procedure. We refer the reader to Section 6 for some examples and a discussion on the differences between our approach and the one of [12, 5].

We proceed as follows, we start below with some preliminaries. Then we turn to some elements of analyze on the Wiener space in Section 3. Our characterization of the sets  $\mathbb{D}^{1,p}$  is given in Section 4, and the material on the Malliavin differentiability of BSDEs itself is contained in Section 5. We provide applications and a comparison of the results in Section 6. Finally, we extend our approach to quadratic growth BSDEs in Section 7.

## 2 Preliminaries

### 2.1 Notations

We fix throughout the paper a time horizon  $T > 0$ . Let  $\Omega := C_0([0, T], \mathbb{R})$  be the canonical Wiener space of continuous function  $\omega$  from  $[0, T]$  to  $\mathbb{R}$  such that  $\omega(0) = 0$ . Let  $W := (W_t)_{t \in [0, T]}$  be the canonical Wiener process, that is, for any time  $t$  in  $[0, T]$ ,  $W_t$  denotes the evaluation mapping:  $W_t(\omega) := \omega_t$  for any element  $\omega$  in  $\Omega$ . We set  $\mathbb{F}^o$  the natural filtration of  $W$ . Under the Wiener measure  $\mathbb{P}_0$ , the process  $W$  is a standard Brownian motion and we denote by  $\mathbb{F} := (\mathcal{F}_t)_{t \in [0, T]}$  the usual augmentation (which is right-continuous and complete) of  $\mathbb{F}^o$  under  $\mathbb{P}_0$ . Unless otherwise stated, all the expectations considered in this paper will have to be understood as expectations under  $\mathbb{P}_0$ , and all notions of measurability for elements of  $\Omega$  will be with respect to the filtration  $\mathbb{F}$  or the  $\sigma$ -field  $\mathcal{F}_T$ .

For any Hilbert space  $\mathcal{K}$ , for any  $p \geq 1$  and for any  $t \in [0, T]$ , we set  $L^p([t, T]; \mathcal{K})$  to be following space

$$L^p([t, T]; \mathcal{K}) := \left\{ f : [t, T] \longrightarrow \mathcal{K}, \text{ Borel-measurable, s.t. } \int_t^T \|f(s)\|_{\mathcal{K}}^p ds < +\infty \right\},$$

where the norm  $\|\cdot\|_{\mathcal{K}}$  is the one canonically induced by the inner product on  $\mathcal{K}$ . We denote, for

---

<sup>1</sup>we refer to Section 4 where this notion is defined

simplicity, by  $\mathfrak{H} := L^2([0, T]; \mathbb{R})$  and by  $\langle \cdot, \cdot \rangle_{\mathfrak{H}}$  its canonical inner product, that is to say

$$\langle f, g \rangle_{\mathfrak{H}} := \int_0^T f(s)g(s)ds, \quad (f, g) \in L^2([0, T]; \mathbb{R}) \times L^2([0, T]; \mathbb{R}).$$

Let now  $H$  be the Cameron-Martin space that is the space of functions in  $\Omega$  which are absolutely continuous with square-integrable derivative and which start from 0 at 0:

$$H := \left\{ h : [0, T] \longrightarrow \mathbb{R}, \exists \dot{h} \in \mathfrak{H}, h(t) = \int_0^t \dot{h}(x)dx, \forall t \in [0, T] \right\},$$

which is an Hilbert space equipped with the inner product  $\langle h_1, h_2 \rangle_H := \langle \dot{h}_1, \dot{h}_2 \rangle_{\mathfrak{H}}$ , for any  $(h_1, h_2) \in H \times H$ , and with associated norm  $\|h\|_H^2 := \langle \dot{h}, \dot{h} \rangle_{\mathfrak{H}}$ . For any  $h$  in  $H$ , we will always denote by  $\dot{h}$  a version of its Radon-Nykodym density with respect to the Lebesgue measure. Define next  $L^p(\mathcal{K})$  as the set of all  $\mathcal{F}_T$ -measurable random variables  $F$  which are valued in an Hilbert space  $\mathcal{K}$ , and such that  $\|F\|_{L^p(\mathcal{K})}^p < +\infty$ , where

$$\|F\|_{L^p(\mathcal{K})} := (\mathbb{E} [\|F\|_{\mathcal{K}}^p])^{1/p}.$$

Let  $\mathcal{S}$  be the set of polynomial cylindrical functionals, that is the set of random variables  $F$  of the form

$$F = f(W(h_1), \dots, W(h_n)), \quad (h_1, \dots, h_n) \in H^n, f \in \mathbb{R}^n[X], \text{ for some } n \geq 1, \quad (2.1)$$

where  $W(h) := \int_0^T \dot{h}_s dW_s$  for any  $h$  in  $H$ . For any  $F$  in  $\mathcal{S}$  of the form (2.1), the Malliavin derivative  $\nabla F$  of  $F$  is defined as the following  $H$ -valued random variable:

$$\nabla F := \sum_{i=1}^n f_{x_i}(W(h_1), \dots, W(h_n))h_i, \quad (2.2)$$

where  $f_{x_i} := \frac{df}{dx_i}$ . It is then customary to identify  $\nabla F$  with the stochastic process  $(\nabla_t F)_{t \in [0, T]}$ . Denote then by  $\mathbb{D}^{1,p}$  the closure of  $\mathcal{S}$  with respect to the Malliavin-Sobolev semi-norm  $\|\cdot\|_{1,p}$ , defined as:

$$\|F\|_{1,p} := (\mathbb{E} [|F|^p] + \mathbb{E} [\|\nabla F\|_H^p])^{1/p}.$$

We set  $\mathbb{D}^{1,\infty} := \bigcap_{p \geq 2} \mathbb{D}^{1,p}$ . In order to link our notations with the ones of the related papers [12, 5] we make use of the notation  $DF$  to represent the derivative of  $\nabla F$  as:

$$\nabla_t F = \int_0^t D_s F ds, \quad t \in [0, T].$$

We denote by  $\delta : L^p(H) \longrightarrow L^p(\mathbb{R})$  the adjoint operator of the operator  $\nabla$  by the following duality relationship:

$$\mathbb{E}[F\delta(u)] = \mathbb{E}[\langle \nabla F, u \rangle_H], \quad \forall u \in \text{dom}(\delta), \quad \text{where}$$

$$\text{dom}(\delta) := \{u \in L^p(H), \exists c_u > 0, |\mathbb{E}[\langle \nabla F, u \rangle_H]| \leq c_u \|F\|_{L^p(\mathbb{R})}, \forall F \in \mathbb{D}^{1,p}\}.$$

$\delta$  is also known under the name of Skorohod (or divergence) operator. Recall that any element  $u$  of the form  $u := Gh$  with  $G$  in  $\mathcal{S}$  and  $h$  in  $H$  belongs to  $\text{dom}(\delta)$  and that

$$\delta(Gh) = GW(h) - \langle \nabla G, h \rangle_H, \quad (2.3)$$

see for example [10, Relation (1.46)].

We conclude this section by introducing the following norms and spaces which are of interest when studying BSDEs. For any  $p \geq 1$ , we set  $\mathbb{S}^p$  the space of  $\mathbb{R}$ -valued, continuous and  $\mathbb{F}$ -progressively measurable processes  $Y$  s.t.

$$\|Y\|_{\mathbb{S}^p}^p := \mathbb{E} \left[ \sup_{0 \leq t \leq T} |Y_t|^p \right] < +\infty.$$

We denote by  $\mathbb{H}^p$  the space of  $\mathbb{R}$ -valued and  $\mathbb{F}$ -predictable processes  $Z$  such that

$$\|Z\|_{\mathbb{H}^p}^p := \mathbb{E} \left[ \left( \int_0^T |Z_t|^2 dt \right)^{\frac{p}{2}} \right] < +\infty.$$

### 3 Some elements of analysis on the Wiener space

One of the main tool that we will use throughout this paper is the shift operator along directions in the Cameron-Martin space. More precisely, for any  $h \in H$ , we define the following shift operator  $\tau_h : \Omega \rightarrow \Omega$  by

$$\tau_h(\omega) := \omega + h.$$

Note that the fact that  $h$  belongs to  $H$  ensures that  $\tau_h$  is a measurable shift on the Wiener space. In fact, one can be a bit more precise, since according to [17, Lemma B.2.1] for any  $\mathcal{F}_T$ -measurable r.v.  $F$  the mapping  $h \mapsto F \circ \tau_h$  is continuous in probability from  $H$  to  $L^0(\mathbb{R})$ , the space of real-valued and  $\mathcal{F}_T$ -measurable random variables, see Lemma 3.2 below. Taking  $F = \text{Id}$ , one gets that  $\tau_h$  is a continuous mapping on  $\Omega$  for any  $h$  in  $H$ . We list below some other properties of such shifts.

**Lemma 3.1** (Appendix B.2, [17]). *Let  $X$  and  $Y$  be two  $\mathcal{F}_T$ -measurable random variables. If  $X = Y$ ,  $\mathbb{P}_0$ -a.s., then for any  $h$  in  $H$ ,*

$$X \circ \tau_h = Y \circ \tau_h, \mathbb{P}_0 - a.s.$$

We recall, the quite surprising result that any r.v. is continuous in probability in the directions of the Cameron-Martin space. More precisely:

**Lemma 3.2** (Lemma B.2.1, [17]). *Let  $F$  be a  $\mathcal{F}_T$ -measurable random variable. The mapping  $h \mapsto F \circ \tau_h$  is continuous from  $H$  to  $L^0(\mathbb{R})$  where the convergence is in probability.*

One of the main technique when working with shifts on the path space is the famous Cameron-Martin formula.

**Proposition 3.1.** (Cameron-Martin Formula, see e.g. [17, Appendix B.1]) *Let  $F$  be a  $\mathcal{F}_T$ -measurable random variable and let  $h$  be in  $H$ . Then, when both sides are well-defined*

$$\mathbb{E}[F \circ \tau_h] = \mathbb{E} \left[ F \exp \left( \int_0^T \dot{h}(s) dW_s - \frac{1}{2} \int_0^T |\dot{h}(s)|^2 ds \right) \right].$$

For further reference, we also emphasize that for any  $h \in H$  and for any  $p \geq 1$ , the stochastic exponential  $\mathcal{E} \left( \int_0^\cdot \dot{h}(s) dW_s \right) := \exp \left( \int_0^\cdot \dot{h}(s) dW_s - \frac{1}{2} \int_0^\cdot |\dot{h}(s)|^2 ds \right)$  verifies

$$\mathcal{E} \left( \int_0^\cdot \dot{h}(s) dW_s \right) \in \mathbb{S}^p, \quad \forall p \geq 1. \quad (3.1)$$

**Lemma 3.3.** *Let  $t$  in  $[0, T]$  and let  $F$  be a  $\mathcal{F}_t$ -measurable random variable. For any  $h$  in  $H$ , it holds that*

$$F \circ \tau_h = F \circ \tau_{h\mathbf{1}_{[0,t]}}, \quad \mathbb{P}_0 - a.s.$$

In particular,  $F \circ \tau_h$  is  $\mathcal{F}_t$ -measurable.

**Proof.** It is well-known that by definition of  $\mathbb{P}_0$ , any  $\mathcal{F}_t$ -measurable r.v. admits a  $\mathcal{F}_t^o$ -measurable version. Therefore, there exists some measurable map  $\varphi : \Omega \rightarrow \mathbb{R}$ , such that

$$F = \varphi(W_{\cdot \wedge t}), \quad \mathbb{P}_0 - a.s.$$

Hence, we deduce by Lemma 3.1 that for  $\mathbb{P}_0$ -a.e.  $\omega \in \Omega$

$$F \circ \tau_h(\omega) = \varphi(W_{\cdot \wedge t}(\omega)) \circ \tau_h = \varphi(W_{\cdot \wedge t} \circ \tau_h(\omega)) = \varphi(\omega(\cdot \wedge t) + h(\cdot \wedge t)) = F \circ \tau_{h\mathbf{1}_{[0,t]}}(\omega). \quad \square$$

We conclude this section with the following lemma which might be known. However since we did not find it in the literature we provide a proof in order to make this paper self-contained.

**Lemma 3.4.** *Let  $Z \in \mathbb{H}^2$  and  $h$  in  $H$ . It holds that*

$$\int_0^T Z_s dW_s \circ \tau_h = \int_0^T Z_s \circ \tau_h dW_s + \int_0^T Z_s \circ \tau_h \dot{h}(s) ds, \quad \mathbb{P}_0 - a.s.$$

**Proof.** Let  $\mathfrak{S}$  be the class of simple processes  $X$  of the form

$$X_t = \sum_{i=0}^n \lambda_{t_i} \mathbf{1}_{(t_i, t_{i+1}]}(t),$$

where  $n \in \mathbb{N}^*$ ,  $t_0 = 0 < t_1 < \dots < t_n = T$  and where for any  $0 \leq i \leq n$ ,  $(\lambda_{t_i})_{i=1, \dots, n}$  are  $\mathcal{F}_{t_i}$ -measurable and in  $L^2(\mathbb{R})$ .

We start by proving the result for  $Z$  in  $\mathfrak{S}$  and then we prove the result for any element  $Z$  in  $\mathbb{H}^2$  using a density argument. Let  $Z \in \mathfrak{S}$  with the decomposition

$$Z_s = \sum_{i=0}^n \lambda_{t_i} \mathbf{1}_{(t_i, t_{i+1}]}(s), \quad s \in [0, T].$$

Then, for any  $h \in H$  and for every  $\omega \in \Omega$ ,

$$\begin{aligned} \left( \int_0^T Z_s dW_s \circ \tau_h \right) (\omega) &= \left( \sum_{i=0}^n \lambda_{t_i} (W_{t_{i+1}} - W_{t_i}) \right) \circ \tau_h(\omega) \\ &= \sum_{i=0}^n \lambda_{t_i}(\omega + h) (W_{t_{i+1}} - W_{t_i})(\omega + h) \\ &= \sum_{i=0}^n \lambda_{t_i} \circ \tau_h(\omega) (\omega(t_{i+1}) - \omega(t_i) + h(t_{i+1}) - h(t_i)) \\ &= \int_0^T Z_s \circ \tau_h dW_s(\omega) + \int_0^T Z_s \circ \tau_h(\omega) dh_s, \end{aligned}$$



which gives the desired result since  $h$  is absolutely continuous. We extend this result to processes  $Z$  in  $\mathbb{H}^2$ . Let  $Z \in \mathbb{H}^2$ , then there exists a sequence  $(Z^n)_{n \in \mathbb{N}}$  in  $\mathfrak{S}$  which converges to  $Z$  in  $\mathbb{H}^2$ . Hence,

$$\begin{aligned} & \mathbb{E} \left[ \left| \int_0^T Z_s dW_s \circ \tau_h - \int_0^T Z_s \circ \tau_h dW_s - \int_0^T Z_s \circ \tau_h dh_s \right| \right] \\ & \leq \mathbb{E} \left[ \left| \int_0^T Z_s dW_s \circ \tau_h - \int_0^T Z_s^n dW_s \circ \tau_h \right| \right] + \mathbb{E} \left[ \left| \int_0^T Z_s^n \circ \tau_h dW_s - \int_0^T Z_s \circ \tau_h dW_s \right| \right] \\ & \quad + \mathbb{E} \left[ \left| \int_0^T Z_s^n \circ \tau_h dh_s - \int_0^T Z_s \circ \tau_h dh_s \right| \right] \\ & \leq \underbrace{\mathbb{E} \left[ \left| \int_0^T (Z_s - Z_s^n) dW_s \right| \circ \tau_h \right]}_{=: A^n} + \underbrace{\mathbb{E} \left[ \left| \int_0^T (Z_s^n - Z_s) \circ \tau_h dW_s \right| \right]}_{=: B^n} + \underbrace{\mathbb{E} \left[ \left| \int_0^T (Z_s^n - Z_s) \circ \tau_h dh_s \right| \right]}_{=: C^n}. \end{aligned}$$

Let us estimate these three terms. First, using Proposition 3.1, Burkholder-Davis-Gundy Inequality, then Cauchy-Schwarz Inequality, we have

$$\begin{aligned} A^n &= \mathbb{E} \left[ \left| \int_0^T (Z_s - Z_s^n) dW_s \right| e^{\int_0^T \dot{h}(s) dW_s - \frac{1}{2} \int_0^T |\dot{h}(s)|^2 ds} \right] \leq \mathbb{E} \left[ \int_0^T |Z_s - Z_s^n|^2 ds \right]^{1/2} \\ & \quad \times \mathbb{E} \left[ \mathcal{E} \left( \int_0^T \dot{h}(s) dW_s \right)_T \right]^{1/2}. \end{aligned}$$

By (3.1), this clearly goes to 0 as  $n$  goes to infinity. Similarly, using Burkholder-Davis-Gundy Inequality, we have

$$B^n \leq \mathbb{E} \left[ \left( \int_0^T ((Z_s^n - Z_s) \circ \tau_h)^2 ds \right)^{\frac{1}{2}} \right] = \mathbb{E} \left[ \left( \int_0^T (Z_s^n - Z_s)^2 ds \right)^{\frac{1}{2}} \circ \tau_h \right].$$

Therefore, we can use Proposition 3.1 and Cauchy-Schwarz Inequality, to also deduce that  $B^n \xrightarrow{n \rightarrow +\infty} 0$ . Finally, we have

$$\begin{aligned} C^n &= \mathbb{E} \left[ \mathcal{E} \left( \int_0^T \dot{h}(s) dW_s \right) \left| \int_0^T (Z_s^n - Z_s) \dot{h}(s) ds \right| \right] \\ & \leq \mathbb{E} \left[ \mathcal{E} \left( \int_0^T \dot{h}(s) dW_s \right)^2 \right]^{1/2} \mathbb{E} \left[ \left( \int_0^T |Z_s^n - Z_s| |\dot{h}(s)| ds \right)^2 \right]^{1/2} \\ & \leq \mathbb{E} \left[ \mathcal{E} \left( \int_0^T \dot{h}(s) dW_s \right)^2 \right]^{1/2} \mathbb{E} \left[ \int_0^T |Z_s^n - Z_s|^2 ds \right]^{1/2} \left( \int_0^T |\dot{h}(s)|^2 ds \right)^{1/2}, \end{aligned}$$

which also goes to 0 as  $n$  goes to infinity. Therefore the proof is complete.  $\square$

This result entails the following useful consequence. Let  $t$  in  $(0, T]$  and  $h$  in  $H$  such that  $\dot{h}_s = 0$  for  $s \geq t$ . Then for any  $Z$  in  $\mathbb{H}^2$ , it holds that:

$$\int_t^T Z_s dW_s \circ \tau_h = \int_t^T Z_s \circ \tau_h dW_s, \quad \mathbb{P}_0 - a.s., \quad (3.2)$$

since  $\int_t^T Z_s \circ \tau_h \dot{h}(s) ds = 0$ .

## 4 A characterization of Malliavin differentiability

Before going further we would like to recall the main finding of [16] which is that any Malliavin-Sobolev type space  $\mathbb{D}^{1,p}$  as defined in Section 2 (originally defined by Malliavin [9] and Shigekawa [14]) agrees with the Sobolev space (due to Stroock [15] and Kusuoka [8])  $\tilde{\mathbb{D}}^{1,p}$  which consists in the set of r.v.  $F$  in  $L^p(\mathbb{R})$  which are *Ray Absolutely Continuous* (RAC) and *Stochastically Gâteaux Differentiable* (SGD) where these notions are defined as follows:

(RAC) For any  $h$  in  $H$ , there exists a r.v.  $\tilde{F}_h$  such that  $\tilde{F}_h = F$ ,  $\mathbb{P}_0$ -a.s., and such that for any  $\omega$  in  $\Omega$ ,  $t \mapsto \tilde{F}_h(\omega + th)$  is absolutely continuous.

(SGD) There exists  $\mathcal{D}F$  in  $L^p(H)$  such that for any  $h$  in  $H$ ,

$$\frac{F \circ \tau_{\varepsilon h} - F}{\varepsilon} \xrightarrow{\varepsilon \rightarrow 0} \langle \mathcal{D}F, h \rangle_H, \text{ in probability.} \quad (4.1)$$

In addition, for any  $F$  in  $\mathbb{D}^{1,p}$ ,  $\nabla F = \mathcal{D}F$ ,  $\mathbb{P}_0$ -a.s. Note that according to the statement of Step 1 in the proof of [16, Theorem 3.1], if  $F$  is (RAC) and (SGD) then for any  $h$  in  $H$  and any  $\varepsilon > 0$  it holds that

$$\varepsilon^{-1}(\tilde{F}_h \circ \tau_{\varepsilon h} - \tilde{F}_h) = \varepsilon^{-1} \int_0^\varepsilon \langle \nabla F \circ \tau_{sh}, h \rangle_H ds, \quad \mathbb{P}_0 - a.s.$$

Furthermore, by Lemma 3.1, we have for any  $\varepsilon$  there exists a set  $A^\varepsilon$  such that  $\mathbb{P}_0[A^\varepsilon] = 0$  and  $F \circ \tau_{\varepsilon h} = \tilde{F}_h \circ \tau_{\varepsilon h}$  and  $F = \tilde{F}_h$  outside  $A^\varepsilon$ . Hence, for any  $\varepsilon$  in  $(0, 1)$ , the relation above rewrites as:

$$\varepsilon^{-1}(F \circ \tau_{\varepsilon h} - F) = \varepsilon^{-1} \int_0^\varepsilon \langle \nabla F \circ \tau_{sh}, h \rangle_H ds, \quad \mathbb{P}_0 - a.s. \quad (4.2)$$

The main result of this section is the following theorem whose proof is postponed to the end of the section.

**Theorem 4.1.** *Let  $p > 1$ .  $F$  belongs to  $\mathbb{D}^{1,p}$  if and only if there exist  $\mathcal{D}F$  in  $L^p(H)$  and  $q \in (1, p)$  such that for any  $h$  in  $H$*

$$\lim_{\varepsilon \rightarrow 0} \mathbb{E} \left[ \left| \frac{F \circ \tau_{\varepsilon h} - F}{\varepsilon} - \langle \mathcal{D}F, h \rangle_H \right|^q \right] = 0.$$

We recall the following lemma which can be found e.g. in [16, Corollary 2.1] and which characterizes the Malliavin derivative using the duality formula involving the Skorohod operator (also called divergence operator).

**Lemma 4.1.** *Let  $\varepsilon > 0$  and  $1 < p < +\infty$ . Suppose that  $F \in L^{1+\varepsilon}(\mathbb{R})$  and assume that there exists  $\mathcal{D}F$  in  $L^p(H)$  such that:*

$$\mathbb{E} [F \delta(Gh)] = \mathbb{E} [G \langle \mathcal{D}F, h \rangle_H],$$

for every  $G \in \mathcal{S}$  and  $h \in H$ . Then, it holds that  $F \in \mathbb{D}^{1,p}$ ,  $\mathcal{D}F = \nabla F$ ,  $\mathbb{P}_0 - a.s.$

We now prove the following lemma for the Malliavin differentiability of a given random variable.

**Lemma 4.2.** *Let  $p > 1$ . Let  $F$  be in  $\mathbb{D}^{1,p}$ . Then, for any  $q$  in  $(1, p)$  and for any  $h$  in  $H$ ,*

$$\frac{F \circ \tau_{\varepsilon h} - F}{\varepsilon} \xrightarrow{\varepsilon \rightarrow 0} \langle \nabla F, h \rangle_H \text{ in } L^q(\mathbb{R}).$$

**Proof.** Fix  $q$  in  $(1, p)$ ,  $h$  in  $H$  and  $\eta > 0$  such that  $q + \eta < p$ . We know from [16, Theorem 3.1] that since  $F$  is in  $\mathbb{D}^{1,p}$ ,  $F$  is (SGD), (RAC), and Relation (4.2) holds true. We thus have by Jensen's inequality,

$$\begin{aligned} & \mathbb{E} \left[ \left| \varepsilon^{-1} (F \circ \tau_{\varepsilon h} - F) \right|^{q+\eta} \right] \\ &= \mathbb{E} \left[ \varepsilon^{-(q+\eta)} \left| \int_0^\varepsilon \langle \nabla F \circ \tau_{sh}, h \rangle_H ds \right|^{q+\eta} \right] \\ &\leq \varepsilon^{-1} \mathbb{E} \left[ \int_0^\varepsilon |\langle \nabla F \circ \tau_{sh}, h \rangle_H|^{q+\eta} ds \right] \\ &\leq \varepsilon^{-1} \int_0^\varepsilon \mathbb{E} [|\langle \nabla F, h \rangle_H|^{q+\eta} \circ \tau_{sh}] ds \\ &= \varepsilon^{-1} \int_0^\varepsilon \mathbb{E} \left[ |\langle \nabla F, h \rangle_H|^{q+\eta} \mathcal{E} \left( s \int_0^T \dot{h}_r dW_r \right) \right] ds \\ &\leq \mathbb{E} [|\langle \nabla F, h \rangle_H|^p]^{\frac{q+\eta}{p}} \sup_{t \in (0,1)} \mathbb{E} \left[ \left| \mathcal{E} \left( t \int_0^T \dot{h}_r dW_r \right) \right|^{\frac{p}{p-q-\eta}} \right]^{\frac{p-q-\eta}{p}} < +\infty. \end{aligned}$$

Hence by de La Vallée Poussin Criterion, we deduce that the family of random variables  $(|\varepsilon^{-1}(F \circ \tau_{\varepsilon h} - F)|^q)_{\varepsilon \in (0,1)}$  is uniformly integrable which together with the convergence in probability (4.1) gives the result.  $\square$

**Remark 4.1.** *Note that the conclusion of the previous Lemma may fail for  $q = p$ .*

We can now proceed with the proof of Theorem 4.1.

**Proof of Theorem 4.1.** The necessary part follows from Lemma 4.2. We turn to the converse implication. Let  $F$  be such that there exists  $\mathcal{D}F$  in  $L^p(\Omega; H)$  and  $q \in (1, p)$  such that

$$\lim_{\varepsilon \rightarrow 0} \mathbb{E} \left[ \left| \frac{F \circ \tau_{\varepsilon h} - F}{\varepsilon} - \langle \mathcal{D}F, h \rangle_H \right|^q \right] = 0.$$

The proof consists in applying Lemma 4.1 by proving the duality relationship

$$\mathbb{E}[F \delta(Gh)] = \mathbb{E}[G \langle \mathcal{D}F, h \rangle_H], \quad G \in \mathcal{S}, \quad h \in H \quad (4.3)$$

which, by taking  $\varepsilon = 0$ , is itself a consequence of the following relation

$$\mathbb{E}[F \circ \tau_{\varepsilon h} \delta(Gh)] = \mathbb{E}[\langle (\mathcal{D}F) \circ \tau_{\varepsilon h}, h \rangle_H G]. \quad (4.4)$$

By Lemma A.1 (in the Appendix) it holds that

$$\begin{aligned} \mathbb{E}[F \circ \tau_{\varepsilon h} \delta(Gh)] &= \frac{d}{d\varepsilon} \mathbb{E}[F \circ \tau_{\varepsilon h} G] \\ &= \lim_{\eta \rightarrow 0} \frac{1}{\eta} \mathbb{E}[(F \circ \tau_{(\varepsilon+\eta)h} - F \circ \tau_{\varepsilon h})G] \\ &= \lim_{\eta \rightarrow 0} \mathbb{E} \left[ \left( \frac{F \circ \tau_{(\varepsilon+\eta)h} - F \circ \tau_{\varepsilon h}}{\eta} - \langle (\mathcal{D}F) \circ \tau_{\varepsilon h}, h \rangle_H \right) G \right] \\ &\quad + \mathbb{E}[\langle (\mathcal{D}F) \circ \tau_{\varepsilon h}, h \rangle_H G] \\ &= \mathbb{E}[\langle (\mathcal{D}F) \circ \tau_{\varepsilon h}, h \rangle_H G], \end{aligned} \quad (4.5)$$

where the proof that the first term on the right-hand side goes to 0 is reported below.

Note that  $\mathbb{E}[\langle (\mathcal{D}F) \circ \tau_{\varepsilon h}, h \rangle_H G] < +\infty$  since  $\langle (\mathcal{D}F) \circ \tau_{\varepsilon h}, h \rangle_H = \langle \mathcal{D}F, h \rangle_H \circ \tau_{\varepsilon h}$ ,  $\mathbb{P}_0$ -a.s.,  $G$  belongs to all the spaces  $L^r(\mathbb{R})$  for  $r \geq 1$  and

$$\mathbb{E}[|\langle \mathcal{D}F, h \rangle_H|^p] \leq \|h\|_H^p \mathbb{E}[\|\mathcal{D}F\|_H^p] < +\infty.$$

The Equality (4.5) is justified by Hölder's inequality since

$$\begin{aligned} & \mathbb{E} \left[ \left| \left( \frac{F \circ \tau_{(\varepsilon+\eta)h} - F \circ \tau_{\varepsilon h}}{\eta} - \langle (\mathcal{D}F) \circ \tau_{\varepsilon h}, h \rangle_H \right) G \right| \right] \\ & \leq \mathbb{E} \left[ \left| \frac{F \circ \tau_{(\varepsilon+\eta)h} - F \circ \tau_{\varepsilon h}}{\eta} - \langle (\mathcal{D}F) \circ \tau_{\varepsilon h}, h \rangle_H \right|^r \right]^{\frac{1}{r}} \mathbb{E}[|G|^{\bar{r}}]^{\frac{1}{\bar{r}}} \\ & = \mathbb{E} \left[ \left| \frac{F \circ \tau_{\eta h} - F}{\eta} - \langle \mathcal{D}F, h \rangle_H \right|^r \circ \tau_{\varepsilon h} \right]^{\frac{1}{r}} \mathbb{E}[|G|^{\bar{r}}]^{\frac{1}{\bar{r}}} \\ & \leq \mathbb{E} \left[ \left| \frac{F \circ \tau_{\eta h} - F}{\eta} - \langle \mathcal{D}F, h \rangle_H \right|^{q\gamma} \right]^{\frac{1}{q}} \mathbb{E} \left[ \mathcal{E} \left( \varepsilon \int_0^T \dot{h}(s) dW_s \right)^{\bar{\alpha}} \right]^{\frac{1}{\bar{\alpha}}} \mathbb{E}[|G|^{\bar{r}}]^{\frac{1}{\bar{r}}} \end{aligned}$$

where  $1 < r < q$  and  $\alpha := \frac{q}{r}$  and where  $\bar{r}$  (resp.  $\bar{\alpha}$ ) is the Hölder conjugate of  $r$  (resp.  $\alpha$ ), which establishes (4.4).  $\square$

**Remark 4.2.** Note that according to Lemma 4.2 the necessary part of Theorem 4.1 can be strengthened since the convergence of the difference quotient holds in any space  $L^q(\mathbb{R})$  with  $q < p$ .

**Remark 4.3.** From (4.4) we prove that for any  $\varepsilon > 0$  and any  $h$  in  $H$ ,  $F \circ \tau_{\varepsilon h}$  belongs to  $\mathbb{D}^{1,p}$  and that  $\nabla(F \circ \tau_{\varepsilon h}) = (\nabla F) \circ \tau_{\varepsilon h}$ .

## 5 Malliavin's differentiability of BSDEs

In this section we derive a sufficient condition ensuring that the solution to a BSDE is Malliavin differentiable. To simplify the comparisons of the results with the companion papers [5, 12] we adopt the notations used in these papers concerning the Malliavin calculus. More precisely, for any  $F$  in  $\mathbb{D}^{1,p}$  (for  $p > 1$ ) we have defined the Malliavin derivative  $\nabla F$  as an  $H$ -valued random variable. Recall that denoting  $DF$  the derivative of  $\nabla F$  that is  $\nabla_t F = \int_0^t D_s F ds$ ,  $DF$  coincides with the Malliavin derivative introduced in [5, 12, 10]. In particular  $\langle \nabla F, h \rangle_H = \langle DF, \dot{h} \rangle_{\mathcal{H}}$  for any  $h$  in  $H$ .

We consider now the following BSDE:

$$Y_t = \xi + \int_t^T f(s, Y_s, Z_s) ds - \int_t^T Z_s dW_s, \quad t \in [0, T], \quad \mathbb{P} - a.s., \quad (5.1)$$

where  $\xi$  is a  $\mathcal{F}_T$ -measurable r.v. and  $f : [0, T] \times \Omega \times \mathbb{R}^2 \rightarrow \mathbb{R}$  is a  $\mathbb{F}$ -progressively measurable process where as usual the  $\omega$ -dependence is omitted.

The aim of this section is to show that for any  $t \in [0, T]$ , we can apply Theorem 4.1 under the following assumptions:

- (L) The map  $(y, z) \mapsto f(\cdot, y, z)$  is differentiable with uniformly bounded and continuous partial derivatives.

(D)  $\xi$  belongs to  $\mathbb{D}^{1,2}$ , for any  $(y, z) \in \mathbb{R}^2$ ,  $(t, \omega) \mapsto f(t, \omega, y, z)$  is in  $L^2([0, T]; \mathbb{D}^{1,2})$ ,  $f(\cdot, y, z)$  and  $Df(\cdot, y, z)$  are  $\mathbb{F}$ -progressively measurable, and

$$\mathbb{E} \left[ \int_0^T \|D.f(s, Y_s, Z_s)\|_{\mathfrak{H}}^2 ds \right] < +\infty.$$

(H<sub>1</sub>) There exists  $p \in (1, 2)$  such that for any  $h \in H$

$$\lim_{\varepsilon \rightarrow 0} \mathbb{E} \left[ \left( \int_0^T \left| \frac{f(s, \cdot + \varepsilon h, Y_s, Z_s) - f(s, \cdot, Y_s, Z_s)}{\varepsilon} - \langle Df(s, \cdot, Y_s, Z_s), \dot{h} \rangle_{\mathfrak{H}} \right| ds \right)^p \right] = 0,$$

(H<sub>2</sub>) Let  $(\varepsilon_n)_{n \in \mathbb{N}}$  be a sequence in  $(0, 1]$  such that  $\lim_{n \rightarrow +\infty} \varepsilon_n = 0$ , and let  $(Y^n, Z^n)_n$  be a sequence of random variables which converges in  $\mathbb{S}^p \times \mathbb{H}^p$  for any  $p \in [1, 2)$  to some  $(Y, Z)$ . Then for all  $h \in H$ , the following convergences hold in probability

$$\begin{aligned} \|f_y(\cdot, \omega + \varepsilon_n h, Y^n, Z^n) - f_y(\cdot, \omega, Y, Z)\|_{\mathfrak{H}} &\xrightarrow{n \rightarrow +\infty} 0 \\ \|f_z(\cdot, \omega + \varepsilon_n h, Y^n, Z^n) - f_z(\cdot, \omega, Y, Z)\|_{\mathfrak{H}} &\xrightarrow{n \rightarrow +\infty} 0, \end{aligned} \quad (5.2)$$

or

$$\begin{aligned} \|f_y(\cdot, \omega + \varepsilon_n h, Y^n, Z^n) - f_y(\cdot, \omega, Y, Z)\|_{\mathfrak{H}} &\xrightarrow{n \rightarrow +\infty} 0 \\ \|f_z(\cdot, \omega + \varepsilon_n h, Y, Z^n) - f_z(\cdot, \omega, Y, Z)\|_{\mathfrak{H}} &\xrightarrow{n \rightarrow +\infty} 0. \end{aligned} \quad (5.3)$$

Before turning to the main result of this section, we would like to comment on Assumption (H<sub>2</sub>). On the one hand, by Lemma 3.2, at given  $(s, y, z)$ ,  $f_y(s, \omega + \varepsilon_n h, y, z)$  converges in probability to  $f_y(s, \omega, y, z)$  as  $n$  goes to infinity. On the other hand,  $f_y(s, \omega, \cdot)$  is continuous by assumption. Thus, Condition (H<sub>2</sub>) is just requiring joint continuity of  $f_y$  in  $L^2([0, T], \mathbb{R})$ . The same comment holds for  $f_z$ . Note finally, that since  $f_y$  is assumed to be bounded, a sufficient condition for (H<sub>2</sub>) to hold true is that  $f_y(t, Y_t^n, Z_t)$  converges in probability to  $f_y(t, Y_t, Z_t)$  for  $dt$ -almost every  $t$  (and the same for  $f_z$ ).

**Theorem 5.1.** *Let  $t$  be in  $[0, T]$ . Under Assumptions (L), (D), (H<sub>1</sub>) and (H<sub>2</sub>),  $Y_t$  belongs to  $\mathbb{D}^{1,2}$  and  $Z \in L^2([t, T]; \mathbb{D}^{1,2})$ .*

**Proof.** We proceed in two steps and only consider the case where (5.2) holds in Assumption (H<sub>2</sub>), since the other one can be treated similarly.

**Step 1:** We prove that  $Y_t$  belongs to  $\mathbb{D}^{1,p}$  where  $p \in (1, 2)$  is the exponent appearing in Assumption (H<sub>1</sub>). To this end we aim at applying Theorem 4.1. Let  $h$  in  $H$ . Since  $Y$  is  $\mathbb{F}$ -progressively measurable, by Lemma 3.3, we can assume without loss of generality that  $\dot{h}_s = 0$  for  $s > t$ . Let  $\varepsilon > 0$ . By Lemmas 3.1 and 3.4, it holds that<sup>1</sup>

$$Y_s \circ \tau_{\varepsilon h} = \xi \circ \tau_{\varepsilon h} + \int_s^T f(r, Y_r, Z_r) \circ \tau_{\varepsilon h} dr - \int_s^T Z_r \circ \tau_{\varepsilon h} dW_r, \quad \forall s \in [t, T], \mathbb{P}_0 - a.s.$$

---

<sup>1</sup>note that by Cameron-Martin formula the process  $(Y_s \circ \tau_{\varepsilon h})_s$  is continuous which enables us to get that the  $\Omega$ -exception set does not depend on  $s$ .

As a consequence, setting for simplicity

$$Y_s^\varepsilon := \frac{1}{\varepsilon}(Y_s \circ \tau_{\varepsilon h} - Y_s), \quad Z_s^\varepsilon := \frac{1}{\varepsilon}(Z_s \circ \tau_{\varepsilon h} - Z_s), \quad \xi^\varepsilon := \frac{1}{\varepsilon}(\xi \circ \tau_{\varepsilon h} - \xi), \quad s \in [t, T],$$

we have that  $(Y^\varepsilon, Z^\varepsilon)$  solves the BSDE:

$$Y_s^\varepsilon = \xi^\varepsilon + \int_s^T \tilde{A}_r^\varepsilon + \tilde{A}_r^{y,\varepsilon} Y_r^\varepsilon + \tilde{A}_r^{z,\varepsilon} Z_r^\varepsilon dr - \int_s^T Z_r^\varepsilon dW_r, \quad (5.4)$$

with

$$\begin{aligned} \tilde{A}_r^\varepsilon &:= \frac{1}{\varepsilon}(f(r, \cdot + \varepsilon h, Y_r, Z_r) - f(r, \cdot, Y_r, Z_r)), \\ \tilde{A}_r^{y,\varepsilon} &:= f_y(r, \cdot + \varepsilon h, \bar{Y}_r^{\varepsilon,h}, Z_r), \\ \tilde{A}_r^{z,\varepsilon} &:= f_z(r, \cdot + \varepsilon h, Y_r \circ \tau_{\varepsilon h}, \bar{Z}_r^{\varepsilon,h}), \end{aligned}$$

where  $\bar{Y}_r^{\varepsilon,h}$  (resp.  $\bar{Z}_r^{\varepsilon,h}$ ) is some intermediary point between  $Y_r$  and  $Y_r \circ \tau_{\varepsilon h}$  (resp.  $Z_r$  and  $Z_r \circ \tau_{\varepsilon h}$ ).

Under Assumptions (D) and (L), the following linear BSDE on  $[t, T]$  is also well-posed

$$\tilde{Y}_s^h = \langle D\xi, \dot{h} \rangle_{\mathfrak{H}} + \int_s^T \langle Df(r, \cdot, Y_r, Z_r), \dot{h} \rangle_{\mathfrak{H}} + \tilde{Y}_r^h f_y(r, \cdot, Y_r, Z_r) + \tilde{Z}_r^h f_z(r, \cdot, Y_r, Z_r) dr - \int_s^T \tilde{Z}_r^h dW_r. \quad (5.5)$$

Besides, we have  $(\tilde{Y}^h, \tilde{Z}^h) \in \mathbb{S}^2 \times \mathbb{H}^2$ . Using a priori estimates (see Proposition 3.2 in [3]) in  $L^p$ , we have for some constant  $C_p$ , independent of  $\varepsilon$

$$\begin{aligned} &\mathbb{E} \left[ \sup_{s \in [t, T]} |Y_s^\varepsilon - \tilde{Y}_s^h|^p \right] + \mathbb{E} \left[ \left( \int_t^T |Z_s^\varepsilon - \tilde{Z}_s^h|^2 ds \right)^{p/2} \right] \\ &\leq C_p \left( \mathbb{E} \left[ |\xi^\varepsilon - \langle D\xi, \dot{h} \rangle_{\mathfrak{H}}|^p \right] + \mathbb{E} \left[ \left( \int_0^T |\tilde{A}_s^\varepsilon - \langle Df(s, \cdot, Y_s, Z_s), \dot{h} \rangle_{\mathfrak{H}}|^p ds \right)^p \right] \right) \\ &\quad + C_p \mathbb{E} \left[ \left( \int_0^T |\tilde{A}_s^{y,\varepsilon} - f_y(s, \cdot, Y_s, Z_s)| |\tilde{Y}_s^h|^p ds \right)^p \right] \\ &\quad + C_p \mathbb{E} \left[ \left( \int_0^T |\tilde{A}_s^{z,\varepsilon} - f_z(s, \cdot, Y_s, Z_s)| |\tilde{Z}_s^h|^p ds \right)^p \right]. \quad (5.6) \end{aligned}$$

Since  $\xi$  is in  $\mathbb{D}^{1,2}$ ,  $\lim_{\varepsilon \rightarrow 0} \mathbb{E} \left[ |\xi^\varepsilon - \langle D\xi, \dot{h} \rangle_{\mathfrak{H}}|^p \right] = 0$  by Lemma 4.2. By Assumption  $(H_1)$ , the second term in the right-hand side of (5.6) goes to 0 as  $\varepsilon$  goes to 0. For the last two terms, we will use Assumption  $(H_2)$ . First, the above estimate implies directly that  $(Y \circ \tau_{\varepsilon h} - Y, Z \circ \tau_{\varepsilon h} - Z)_\varepsilon$  goes to 0 in  $\mathbb{S}^q \times \mathbb{H}^q$  for any  $q \in (1, 2)$ . We can therefore conclude with Assumption  $(H_2)$  together with the fact that  $f_y$  is bounded that by the dominated convergence theorem:

$$\begin{aligned} &\mathbb{E} \left[ \left( \int_0^T |\tilde{A}_s^{y,\varepsilon} - f_y(s, \cdot, Y_s, Z_s)| |\tilde{Y}_s^h|^p ds \right)^p \right] \\ &\leq C \mathbb{E} \left[ \left( \int_0^T |\tilde{A}_s^{y,\varepsilon} - f_y(s, \cdot, Y_s, Z_s)|^2 ds \right)^{p/2} \right] \mathbb{E} \left[ \left( \int_0^T |\tilde{Y}_s^h|^2 ds \right)^{p/2} \right] \\ &\rightarrow_{\varepsilon \rightarrow 0} 0. \end{aligned}$$

We can show similarly that the last term on the right-hand side of (5.6) also goes to 0, by using the fact that  $\tilde{Z}^h \in \mathbb{H}^2$ . Hence by Theorem 4.1,  $Y_t$  belongs to  $\mathbb{D}^{1,p}$ .

Moreover, Burkholder-Davis-Gundy's inequality implies that there exists  $\tilde{C}_p > 0$  such that

$$\begin{aligned} & \mathbb{E} \left[ \left| \varepsilon^{-1} \left( \int_t^T Z_r dW_r \circ \tau_{\varepsilon h} - \int_t^T Z_r dW_r \right) - \int_t^T \tilde{Z}_r^h dW_r \right|^p \right] \\ & \leq \mathbb{E} \left[ \sup_{t \leq s \leq T} \left| \int_t^s \varepsilon^{-1} (Z_r \circ \tau_{\varepsilon h} - Z_r) - \tilde{Z}_r^h dW_r \right|^p \right] \\ & \leq \tilde{C}_p \mathbb{E} \left[ \left( \int_t^T |Z_r^\varepsilon - \tilde{Z}_r^h|^2 dr \right)^{p/2} \right], \end{aligned}$$

which tends to 0 as  $\varepsilon$  goes to 0. Thus  $\int_t^T Z_s dW_s$  is in  $\mathbb{D}^{1,p}$ .

**Step 2:** Let  $t$  in  $(0, T]$ . We now prove that  $(Y_t, Z \mathbf{1}_{[t, T]}) \in \mathbb{D}^{1,2} \times L^2([t, T]; \mathbb{D}^{1,2})$ . Let  $(h_n)_n$  be a complete orthonormal system of  $H$ . We have that

$$\begin{aligned} \mathbb{E} \left[ \|\nabla Y_t\|_H^2 + \|\nabla \int_t^T Z_s dW_s\|_H^2 \right] &= \sum_{n \geq 1} \mathbb{E} \left[ |\langle \nabla Y_t, h_n \rangle_H|^2 + \left| \langle \nabla \int_t^T Z_s dW_s, h_n \rangle_H \right|^2 \right] \\ &= \sum_{n \geq 1} \mathbb{E} \left[ |\tilde{Y}_t^{h_n}|^2 + \left| \int_t^T \tilde{Z}_s^{h_n} dW_s \right|^2 \right] \\ &= \sum_{n \geq 1} \mathbb{E} \left[ |\tilde{Y}_t^{h_n}|^2 + \int_t^T |\tilde{Z}_s^{h_n}|^2 ds \right] \\ &\leq C \sum_{n \geq 1} \mathbb{E} \left[ |\langle \nabla \xi, h_n \rangle_H|^2 + \int_0^T |\langle \nabla f(s, Y_s, Z_s), h_n \rangle_H|^2 ds \right] \\ &= C \mathbb{E} [\|\nabla \xi\|_H^2] + \int_0^T \mathbb{E} [\|\nabla f(s, Y_s, Z_s)\|_H^2] ds \\ &= C \mathbb{E} [\|D\xi\|_{\mathfrak{H}}^2] + \int_0^T \mathbb{E} [\|Df(s, Y_s, Z_s)\|_{\mathfrak{H}}^2] ds < +\infty, \end{aligned}$$

where we have used the fact that  $\langle \nabla Y_t, h_n \rangle_H = \langle DY_t, \dot{h}_n \rangle_{\mathfrak{H}} = \tilde{Y}_t^{h_n}$  (and the same for the  $Z$  component) from Step 1, and a priori estimates (see e.g. [5, Proposition 2.1]) in  $\mathbb{S}^2 \times \mathbb{H}^2$  (which involves a generic constant  $C > 0$ ) for BSDEs of the form (5.5) since the data are square integrable. Hence,  $Y_t$ , and  $\int_t^T Z_s dW_s$  belong to  $\mathbb{D}^{1,2}$ , and by [12, Lemma 2.3],  $Z$  belongs to  $L^2([t, T]; \mathbb{D}^{1,2})$ . Finally note that a version of  $(D_s Y_t, D_s Z_t)_{0 \leq s \leq t, 0 \leq t \leq T}$  is given as the solution (*a priori* in  $\mathbb{S}^p \times \mathbb{H}^p$ ) to the linear BSDE:

$$D_s Y_t = D_s \xi + \int_t^T D_s f(r, \cdot, Y_r, Z_r) + f_y(r, \cdot, Y_r, Z_r) D_s Y_r + f_z(r, \cdot, Y_r, Z_r) D_s Z_r dr - \int_t^T D_s Z_r dW_r, \quad (5.7)$$

which admits in fact a unique solution in  $\mathbb{S}^2 \times \mathbb{H}^2$  by our assumptions.  $\square$

**Remark 5.1.** We would like to point out that since the process  $Z$  is defined as a  $\mathfrak{H}$ -valued r.v., one may be careful not to study  $Z$  directly at a given time, as  $Z_t$  is not well defined for a given  $t$ . Hence, in the proof we rather study at any time  $t$  the random variable  $\int_t^T Z_s dW_s$  and prove that it belongs to  $\mathbb{D}^{1,2}$ . Then by [12, Lemma 2.3] the latter result is equivalent to the fact that  $Z$  belongs to  $L^2([t, T]; \mathbb{D}^{1,2})$ .

## 6 Applications and discussion of the results

### 6.1 Application to FBSDEs

We consider in this section a FBSDE of the form

$$\begin{cases} X_t = X_0 + \int_0^t b(s, X_s) ds + \int_0^t \sigma(s, X_s) dW_s, & t \in [0, T], \mathbb{P} - a.s. \\ Y_t = g(X_T) + \int_t^T f(s, X_s, Y_s, Z_s) ds - \int_t^T Z_s dW_s, & t \in [0, T], \mathbb{P} - a.s. \end{cases} \quad (6.1)$$

We make the following Assumptions:

(A<sub>1</sub>)  $b, \sigma : [0, T] \times \mathbb{R} \rightarrow \mathbb{R}$  are continuous in time and continuously differentiable in space for any fixed time  $t$  and such that there exist  $k_b, k_\sigma > 0$  with

$$|b_x(t, x)| \leq k_b, \quad |\sigma_x(t, x)| \leq k_\sigma, \quad \text{for all } x \in \mathbb{R}.$$

Besides  $b(t, 0), \sigma(t, 0)$  are bounded functions of  $t$ .

(A<sub>2</sub>) (i)  $g$  is continuously differentiable with polynomial growth.

(ii)  $f : [0, T] \times \mathbb{R}^3 \rightarrow \mathbb{R}$  is continuously differentiable in  $(x, y, z)$  with bounded first partial derivatives in  $y, z$  uniformly in  $t$ , such that  $\mathbb{E}[\int_0^T |f(s, 0, 0, 0)|^2 ds] < +\infty$  and satisfying for some  $C > 0$

$$\exists (q, \kappa) \in \mathbb{R}_+ \times [0, 2), \quad |f_x(t, x, y, z)| \leq C(1 + |y|^\kappa + |z|^\kappa + |x|^q), \quad \forall (t, x, y, z) \in [0, T] \times \mathbb{R}^3.$$

The well-known following lemma provides the existence of a Malliavin derivative for  $X_t$  for all  $t \in [0, T]$  under Assumption (A<sub>1</sub>) (see e.g. [10, Theorem 2.2.1]).

**Lemma 6.1.** *Under Assumption (A<sub>1</sub>), for any  $p \geq 1$ ,  $X_t \in \mathbb{D}^{1,p}$  for all  $t \in [0, T]$ , and  $X \in \mathbb{S}^p$ .*

The theorem below shows that in the Markovian case, Theorem 5.1 holds directly under Assumption (A<sub>1</sub>) and (A<sub>2</sub>).

**Theorem 6.1.** *Let  $t \in [0, T]$  and assume that (A<sub>1</sub>) and (A<sub>2</sub>) hold. Then,  $Y_t \in \mathbb{D}^{1,2}$  and  $Z \in L^2([t, T]; \mathbb{D}^{1,2})$ .*

**Proof.** We aim at applying Theorem 4.1. Property (D) holds by the chain rule formula and (L) follows from our assumptions. It remains to prove (H<sub>1</sub>) and (H<sub>2</sub>). We start with (H<sub>1</sub>). Let  $1 < p < 2$  and  $h$  in  $H$ . Below  $C$  denotes a positive constant which can differ from line to line. Recall that from our assumptions,

$$\mathbb{E} \left[ \sup_{t \in [0, T]} |Y_t|^r + \left( \int_0^T |Z_t|^2 dt \right)^{r/2} \right] < \infty, \quad \forall r \leq 2. \quad (6.2)$$



Denoting by  $\bar{X}_t$  a random point between  $X_t$  and  $X_t \circ \tau_{\varepsilon h}$ , where we suppressed the dependence on  $\varepsilon$  for notational simplicity. We have for any  $t$  in  $[0, T]$ , that

$$\begin{aligned}
& \mathbb{E} \left[ \left| \varepsilon^{-1} (f(t, X_t \circ \tau_{\varepsilon h}, Y_t, Z_t) - f(t, X_t, Y_t, Z_t)) - f_x(t, X_t, Y_t, Z_t) \langle DX_t, \dot{h} \rangle_{\mathfrak{H}} \right|^p \right] \\
&= \mathbb{E} \left[ \left| \frac{X_t \circ \tau_{\varepsilon h} - X_t}{\varepsilon} f_x(t, \bar{X}_t, Y_t, Z_t) - f_x(t, X_t, Y_t, Z_t) \langle DX_t, \dot{h} \rangle_{\mathfrak{H}} \right|^p \right] \\
&\leq C \mathbb{E} \left[ \left| \varepsilon^{-1} (X_t \circ \tau_{\varepsilon h} - X_t) - \langle DX_t, \dot{h} \rangle_{\mathfrak{H}} \right|^p (1 + |Y_t|^{\kappa p} + |Z_t|^{\kappa p} + |X_t|^{pq} + |X_t \circ \tau_{\varepsilon h}|^{pq}) \right] \\
&\quad + C \mathbb{E} \left[ |f_x(t, \bar{X}_t, Y_t, Z_t) - f_x(t, X_t, Y_t, Z_t)|^p \left| \langle DX_t, \dot{h} \rangle_{\mathfrak{H}} \right|^p \right] \\
&\leq C \mathbb{E} \left[ \left| \varepsilon^{-1} (X_t \circ \tau_{\varepsilon h} - X_t) - \langle DX_t, \dot{h} \rangle_{\mathfrak{H}} \right|^{pr} \right]^{\frac{1}{r}} \mathbb{E} \left[ (1 + |Y_t|^{\kappa p} + |Z_t|^{\kappa p} + |X_t|^{pq} + |X_t \circ \tau_{\varepsilon h}|^{pq})^{\bar{r}} \right]^{\frac{1}{\bar{r}}} \\
&\quad + C \mathbb{E} \left[ |f_x(t, \bar{X}_t, Y_t, Z_t) - f_x(t, X_t, Y_t, Z_t)|^p \left| \langle DX_t, \dot{h} \rangle_{\mathfrak{H}} \right|^p \right] \\
&=: A_t^{1, \varepsilon} + A_t^{2, \varepsilon},
\end{aligned}$$

where  $\bar{r} > 1$  and  $p$  are chosen so that  $p\kappa\bar{r} < 2$  and  $r$  denotes the Hölder conjugate of  $\bar{r}$ . Using the above estimates, we deduce

$$\begin{aligned}
& \mathbb{E} \left[ \left( \int_0^T \left| \varepsilon^{-1} (f(t, X_t \circ \tau_{\varepsilon h}, Y_t, Z_t) - f(t, X_t, Y_t, Z_t)) - f_x(t, X_t, Y_t, Z_t) \langle DX_t, \dot{h} \rangle_{\mathfrak{H}} \right| dt \right)^p \right] \\
&\leq \int_0^T \left( A_t^{1, \varepsilon} + A_t^{2, \varepsilon} \right) dt.
\end{aligned}$$

Then, we have

$$\begin{aligned}
\int_0^T A_t^{1, \varepsilon} dt &\leq C \left( \int_0^T \mathbb{E} \left[ \left| \varepsilon^{-1} (X_t \circ \tau_{\varepsilon h} - X_t) - \langle DX_t, \dot{h} \rangle_{\mathfrak{H}} \right|^{pr} \right]^{2/r} dt \right)^{1/2} \\
&\quad \times \left( \int_0^T \mathbb{E} \left[ (1 + |Y_t|^{\kappa p} + |Z_t|^{\kappa p} + |X_t|^{pq} + |X_t \circ \tau_{\varepsilon h}|^{pq})^{\bar{r}} \right]^{2/\bar{r}} dt \right)^{1/2}. \quad (6.3)
\end{aligned}$$

In addition by Lemma 3.4, we have that  $M^{\varepsilon, h} := X \circ \tau_{\varepsilon h} - X$  is solution to the linear SDE:

$$dM_t^{\varepsilon, h} = M_t^{\varepsilon, h} (b_x(t, \underline{X}_t) dt + \sigma_x(t, \underline{X}_t) dW_t) + \varepsilon \sigma(t, X_t \circ \tau_{\varepsilon h}) \dot{h}_t dt,$$

where  $\underline{X}_s$  denotes once again a random point between  $X_s$  and  $X_s \circ \tau_{\varepsilon h}$ . Hence using Assumption  $(A_1)$  and standard estimates for SDEs, we get that for any  $q \geq 1$ ,

$$\lim_{\varepsilon \rightarrow 0} \mathbb{E} \left[ \sup_{t \in [0, T]} |X_t \circ \tau_{\varepsilon h} - X_t|^q \right] = 0.$$

Following the same lines as above, and recalling that  $N^h := \langle DX_t, \dot{h} \rangle_{\mathfrak{H}}$  is solution to the SDE:

$$dN_t^h = N_t^h (b_x(t, X_t) dt + \sigma_x(t, X_t) dW_t) + \sigma(t, X_t) \dot{h}_t dt,$$

we get that the process  $P^{\varepsilon, h} := \varepsilon^{-1} (X \circ \tau_{\varepsilon h} - X) - \langle DX, \dot{h} \rangle_{\mathfrak{H}}$  is solution to the affine SDE:

$$dP_t^{\varepsilon, h} = dH_t^\varepsilon + P_t^{\varepsilon, h} (b_x(t, \underline{X}_t) dt + \sigma_x(t, \underline{X}_t) dW_t),$$

with

$$dH_t^\varepsilon := \left( \langle DX_t, \dot{h} \rangle_{\mathfrak{H}} (b_x(t, \underline{X}_t) - b_x(t, X_t)) + \dot{h}_t(\sigma(t, X_t \circ \tau_{\varepsilon h}) - \sigma(t, X_t)) \right) dt \\ + \langle DX_t, \dot{h} \rangle_{\mathfrak{H}} (\sigma_x(t, \underline{X}_t) - \sigma_x(t, X_t)) dW_t$$

Using the fact that  $\sigma_x, b_x$  are bounded,  $\sigma$  has linear growth and is continuous, we get by similar computations than those done several times in this paper that:

$$\lim_{\varepsilon \rightarrow 0} \mathbb{E} \left[ \sup_{t \in [0, T]} |H_t^\varepsilon|^q \right] = 0, \quad \forall q \geq 1,$$

from which we deduce using the explicit representation of solutions to affine SDEs (see e.g. [13, Theorem V.9.53]) that

$$\lim_{\varepsilon \rightarrow 0} \mathbb{E} \left[ \sup_{t \in [0, T]} |\varepsilon^{-1}(X \circ \tau_{\varepsilon h} - X) - \langle DX_t, \dot{h} \rangle_{\mathfrak{H}}|^q \right] = 0, \quad \forall q \geq 1.$$

As a consequence, combining this estimate with (6.3), we get that:

$$\int_0^T A_t^{1, \varepsilon} dt \leq C \left( \mathbb{E} \left[ \sup_{t \in [0, T]} |\varepsilon^{-1}(X_t \circ \tau_{\varepsilon h} - X_t) - \langle DX_t, \dot{h} \rangle_{\mathfrak{H}}|^{pr} \right]^{2/r} \right)^{1/2}$$

which goes to 0 as  $\varepsilon$  goes to 0, since we recall that we have chosen  $p, \bar{r} > 1$  so that  $\kappa p \bar{r} < 2$ , which implies by (6.2), Lemma 6.1 and the Cameron-Martin formula that

$$\int_0^T \mathbb{E} \left[ (1 + |Y_t|^{\kappa p} + |Z_t|^{\kappa p} + |X_t|^{pq} + |X_t \circ \tau_{\varepsilon h}|^{pq \bar{r}})^{2/\bar{r}} \right] dt < \infty.$$

Concerning the term  $\int_0^T A_t^{2, \varepsilon} dt$ , choosing  $\tilde{p} > 1$  so that  $p\tilde{p} < 2$ , it holds by Hölder and by Jensen inequalities that

$$\int_0^T A_t^{2, \varepsilon} dt \leq C \left( \int_0^T \mathbb{E} \left[ |f_x(t, \bar{X}_t, Y_t, Z_t) - f_x(t, X_t, Y_t, Z_t)|^{p\tilde{p}} \right] dt \right)^{1/\tilde{p}},$$

since

$$\mathbb{E} \left[ \sup_{t \in [0, T]} |\langle DX_t, \dot{h} \rangle_{\mathfrak{H}}|^q \right] < \infty, \quad \forall q > 1.$$

As

$$\lim_{\varepsilon \rightarrow 0} \mathbb{E} \left[ \sup_{t \in [0, T]} |X_t \circ \tau_{\varepsilon h} - X_t|^q \right] = 0, \quad \forall q \geq 1$$

it holds that

$$\lim_{\varepsilon \rightarrow 0} \int_0^T |f_x(t, \bar{X}_t, Y_t, Z_t) - f_x(t, X_t, Y_t, Z_t)|^{p\tilde{p}} dt = 0, \quad \mathbb{P} \otimes dt - a.e.$$

Furthermore, for any  $2 > \rho > 1$ ,

$$\sup_{\varepsilon \in (0, 1)} \int_0^T \mathbb{E} \left[ |f_x(t, \bar{X}_t, Y_t, Z_t) - f_x(t, X_t, Y_t, Z_t)|^{\rho p \tilde{p}} \right] dt \\ \leq C \sup_{\varepsilon \in (0, 1)} \int_0^T \mathbb{E} \left[ (1 + |X_t|^q + |X_t \circ \tau_{\varepsilon h}|^q + |Y_t|^\kappa + |Z_t|^\kappa)^{\rho p \tilde{p}} \right] dt < \infty,$$

by choosing  $p$  small enough so that  $\kappa\rho p\tilde{p} \leq 2$ . So by Lebesgue's dominated convergence theorem,

$$\lim_{\varepsilon \rightarrow 0} \int_0^T A_t^{2,\varepsilon} dt = 0,$$

which proves  $(H_1)$ . Concerning,  $(H_2)$  we just mention that  $f_y$  (respectively  $f_z$ ) is bounded, jointly continuous in  $(x, y, z)$  and we make use of Lemma 3.2.  $\square$

## 6.2 Affine BSDEs

The aim of this section is to prove that with our condition, we can provide weaker conditions compared to [5] for affine BSDEs. We take a driver of the form

$$f(t, \omega, y, z) := \alpha_t(\omega) + \beta_t(\omega)y + \gamma_t(\omega)z$$

with bounded  $\mathbb{F}$ -progressively measurable processes, and  $\xi$  in  $\mathbb{D}^{1,2}$ . The conditions given in [5, Proposition 5.3] for proving that the associated solution  $(Y, Z)$  is Malliavin differentiable read as follows (together with some measurability conditions):

$$\exists \eta > 0 \text{ such that } \mathbb{E}[|\xi|^{2+\eta}] < \infty \quad \underline{\text{and}} \quad \int_0^T \mathbb{E} \left[ \left( \int_\theta^T |K_\theta(s)|^2 ds \right)^{2+\eta} \right]^{1/(2+\eta)} d\theta < +\infty, \quad (6.4)$$

with  $K_\theta(s) := |D_\theta \beta(s)| + |D_\theta \gamma(s)|$ .

In our setting, one needs to check assumptions (L), (D),  $(H_1)$  and  $(H_2)$ . As mentioned below by Lemma 3.2 Condition  $(H_2)$  comes for free, and Assumptions (D) and (L) are also trivially satisfied. The interesting point is that  $(H_1)$  is true as soon as (6.4) is replaced with:

$$\exists \eta > 0 \text{ such that } \lim_{\varepsilon \rightarrow 0} \int_0^T \mathbb{E} \left[ \left| \varepsilon^{-1}(\mu_t \circ \tau_{\varepsilon h} - \mu_t) - \langle D\mu_t, \dot{h} \rangle_{\mathfrak{H}} \right|^{2+\eta} \right] dt = 0, \text{ for } \mu \in \{\beta, \gamma\}. \quad (6.5)$$

Hence our condition only involve a condition on  $\gamma$  and  $\beta$  and not on  $\xi$ . For instance if  $\beta$  and  $\gamma$  are given as:

$$\beta_t = \varphi_1(X_t), \quad \gamma_t := \varphi_2(X_t), \quad t \in [0, T],$$

with  $\varphi_1, \varphi_2$  two smooth functions with polynomial growth and  $X$  is the solution to an SDE of the form of the one considered in Section 6.1, then the requirements of Conditions (6.4) and (6.5) are satisfied for  $\beta$  and  $\gamma$ , however in contradistinction to Condition (6.4), Assumption (6.5) does not put extra regularity on the terminal condition  $\xi$ .

We make precise our result.

**Proposition 6.1.** *Let  $\xi$  in  $\mathbb{D}^{1,2}$ , and  $\alpha, \beta, \gamma$  bounded  $\mathbb{F}$ -progressively processes in  $L^2([0, T]; \mathbb{D}^{1,2})$  such that  $D\alpha, D\beta$  and  $D\gamma$  are  $\mathbb{F}$ -progressively measurable. Assume that Assumption (6.5) is in force. Then for any  $t$  in  $[0, T]$ ,  $Y_t$  belongs to  $\mathbb{D}^{1,2}$ ,  $Z \in L^2([t, T]; \mathbb{D}^{1,2})$  where  $(Y, Z)$  is the unique solution in  $\mathbb{S}^2 \times \mathbb{H}^2$  to the affine BSDE:*

$$Y_t = \xi + \int_t^T \alpha_s + \beta_s Y_s + \gamma_s Z_s ds - \int_t^T Z_s dW_s, \quad t \in [0, T].$$

**Proof.** Once again we check that assumptions of Theorem 4.1 are in force. Properties (D) and (L) are immediately satisfied. Let  $f(t, \omega, y, z) := \alpha_t(\omega) + \beta_t(\omega)y + \gamma_t(\omega)z$ . Since  $f_y(t, \omega, y, z) = \beta_t(\omega)$ , and  $f_z(t, \omega, y, z) = \gamma_t(\omega)$  we immediately get by Lemma 3.2 and since  $\beta, \gamma$  are bounded that  $(H_2)$  is in force. Concerning  $(H_1)$ , we have for any  $1 < p < 2$  and  $h$  in  $H$ , that

$$\begin{aligned}
& \mathbb{E} \left[ \left( \int_0^T |\varepsilon^{-1}(f(t, \cdot + \varepsilon h, Y_t, Z_t) - f(t, \cdot, Y_t, Z_t)) - \langle Df(t, \cdot, Y_t, Z_t), \dot{h} \rangle_{\mathfrak{H}}| dt \right)^p \right] \\
& \leq C \mathbb{E} \left[ \left( \int_0^T |\varepsilon^{-1}(\alpha_t \circ \tau_{\varepsilon h} - \alpha_t) - \langle D\alpha_t, \dot{h} \rangle_{\mathfrak{H}}| dt \right)^p \right] \\
& \quad + C \mathbb{E} \left[ \left( \int_0^T |Y_t(\varepsilon^{-1}(\beta_t \circ \tau_{\varepsilon h} - \beta_t) - \langle D\beta_t, \dot{h} \rangle_{\mathfrak{H}})| dt \right)^p \right] \\
& \quad + C \mathbb{E} \left[ \left( \int_0^T |Z_t(\varepsilon^{-1}(\gamma_t \circ \tau_{\varepsilon h} - \gamma_t) - \langle D\gamma_t, \dot{h} \rangle_{\mathfrak{H}})| dt \right)^p \right] \\
& =: A_1^\varepsilon + A_2^\varepsilon + A_3^\varepsilon,
\end{aligned} \tag{6.6}$$

where  $C$  is a constant. By Lemma A.2 we have that  $\lim_{\varepsilon \rightarrow 0} A_1^\varepsilon = 0$ . We consider the term  $A_3^\varepsilon$ . We have that:

$$\begin{aligned}
A_3^\varepsilon & \leq \mathbb{E} \left[ \int_0^T \left| Z_t(\varepsilon^{-1}(\gamma_t \circ \tau_{\varepsilon h} - \gamma_t) - \langle D\gamma_t, \dot{h} \rangle_{\mathfrak{H}}) \right|^p dt \right] \\
& \leq C \int_0^T \mathbb{E} [|Z_t|^2]^{p/2} \mathbb{E} \left[ \left| \varepsilon^{-1}(\gamma_t \circ \tau_{\varepsilon h} - \gamma_t) - \langle D\gamma_t, \dot{h} \rangle_{\mathfrak{H}} \right|^{\frac{2p}{2-p}} \right]^{\frac{2-p}{2}} dt \\
& \leq C \left( \int_0^T \mathbb{E} [|Z_t|^2] dt \right)^{p/2} \left( \int_0^T \mathbb{E} \left[ \left| \varepsilon^{-1}(\gamma_t \circ \tau_{\varepsilon h} - \gamma_t) - \langle D\gamma_t, \dot{h} \rangle_{\mathfrak{H}} \right|^{\frac{2p}{2-p}} \right] dt \right)^{\frac{2-p}{2}}.
\end{aligned}$$

Choosing  $p$  such that  $\frac{2p}{2-p} = 2 + \eta$  we get that  $A_3^\varepsilon$  converges to 0 as  $\varepsilon$  tends to 0 by (6.5). Similarly,  $\lim_{\varepsilon \rightarrow 0} A_2^\varepsilon = 0$  for this choice of  $p$ .  $\square$

**Remark 6.1.** Note that, since the BSDE is affine,  $Y_t$  can be expressed explicitly as:

$$Y_t = \mathbb{E} \left[ M_{t,T} \xi - \int_t^T M_{t,s} \alpha_s ds | \mathcal{F}_t \right],$$

where

$$M_{t,s} := \exp \left( \int_t^s \gamma_u dW_u - \frac{1}{2} \int_t^s |\gamma_u|^2 du + \int_t^s \beta_u du \right), \quad s \in [t, T].$$

Hence, on the one hand,  $Y_t$  belongs to  $\mathbb{D}^{1,2}$  if and only if the coefficients  $\alpha, \beta, \gamma$  belong to  $L^2([0, T]; \mathbb{D}^{1,2})$  and  $\xi$  is in  $\mathbb{D}^{1,2}$ . The same conclusion follows for the  $Z$  component. Hence, neither our condition (6.5) nor the one of [5] namely (6.4) are sharp. However, both are sharp in the case where  $\beta = \gamma = 0$ . On the other hand, Conditions (6.4) or (6.5) give more information that the simple fact that  $Y, Z$  are Malliavin differentiable, since they imply that the BSDE solved by  $(DY, DZ)$  is limit in  $\mathbb{S}^2 \times \mathbb{H}^2$  of respectively  $(DY^n, DZ^n)$  (where  $(Y^n, Z^n)$  is the solution to the Picard iteration equation at order  $n$  approximating  $(Y, Z)$ ) for (6.4), and of the difference quotient  $(\varepsilon^{-1}(Y \circ \tau_{\varepsilon h} - Y), \varepsilon^{-1}(Z \circ \tau_{\varepsilon h} - Z))$  in our case (6.5).

### 6.3 Discussion and comparison of the results

We would like before going to the quadratic BSDE case to make a comment about the difference between our approach and the one of [12, 5] and our approach. In these references, the authors consider the sequence of BSDEs:

$$Y_t^n = \xi + \int_t^T f(s, Y_s^{n-1}, Z_s^{n-1})ds - \int_t^T Z_s^n dW_s, \quad t \in [0, T],$$

which approximate in  $\mathbb{S}^2 \times \mathbb{H}^2$  the solution to the original BSDE:

$$Y_t = \xi + \int_t^T f(s, Y_s, Z_s)ds - \int_t^T Z_s dW_s, \quad t \in [0, T].$$

Now, under mild assumptions on  $f$ , the processes  $(Y^n, Z^n)$  are Malliavin differentiable and it holds that a version of  $(D_r Y_t^n, D_r Z_t^n)$  satisfies for  $t \in [0, T]$ ,  $r \leq t$ :

$$D_r Y_t^n = \xi + \int_t^T [D_r f(s, \Theta_s^{n-1}) + \partial_y f(s, \Theta_s^{n-1})D_r Y_s^{n-1} + \partial_z f(s, \Theta_s^{n-1})D_r Z_s^{n-1}]ds - \int_t^T Z_s^n dW_s,$$

with  $\Theta_s^{n-1} := (Y_s^{n-1}, Z_s^{n-1})$ ,  $y, z$ . On the other if  $(Y, Z)$  where Malliavin differentiable we would have that a version of  $(D_r Y_t, D_r Z_t)$  would satisfy for  $t \in [0, T]$ ,  $r \leq t$ :

$$D_r Y_t = \xi + \int_t^T [D_r f(s, Y_s, Z_s) + \partial_y f(s, Y_s, Z_s)D_r Y_s + \partial_z f(s, Y_s, Z_s)D_r Z_s]ds - \int_t^T Z_s dW_s.$$

In other words, assuming that  $\partial_y f$  and  $\partial_z f$  to be continuous, we would get formally that  $(DY^n, DZ^n)$  converges to  $(DY, DZ)$  (in  $\mathbb{S}^2 \times \mathbb{H}^2$ ) as  $n$  goes to infinity provided that at the limit one can replace  $D_r f(s, Y_s^{n-1}, Z_s^{n-1})$  by  $D_r f(s, Y_s, Z_s)$  which is exactly where comes the main assumption in [5, 12] which impose  $D_r f$  to be (stochastic) Lipschitz continuous in  $(y, z)$  with integrability conditions on the Lipschitz constant to make the aforementioned argument rigorous. However, it is not a necessary condition for  $(Y, Z)$  to be Malliavin differentiable that  $(DY^n, DZ^n)$  to converge to  $(DY, DZ)$ , this is why this assumption is somehow arbitrary. However, for  $Y_t$  to be in  $\mathbb{D}^{1,2}$ , it is necessary (and sufficient) to have that  $\varepsilon^{-1}(Y_t \circ \tau_{\varepsilon h} - Y_t)$  to converge in  $L^p$  for some  $p < 2$  to  $\langle DY_t, \dot{h} \rangle_{\mathfrak{H}}$  for any  $h$  in  $H$  (according to Theorem 4.1). Hence, we feel that our conditions are more precise.

## 7 Extension to quadratic growth BSDEs

The aim of this section is to extend our previous results to so-called quadratic growth BSDEs. Such results already exist in the literature, see in particular [1, 7] or the thesis [4], however they are generally limited to specific forms of the generators or to a Markovian setting. We will show that our approach to the Malliavin differentiability is flexible enough to be able to treat this problem without major modifications to our proofs.

We will now list our assumptions in this quadratic setting

$(D_\infty)$   $\xi$  is bounded, belongs to  $\mathbb{D}^{1,\infty}$  and its Malliavin derivative  $D\xi$  is bounded, for any  $(y, z) \in \mathbb{R}^2$ ,  $(t, \omega) \longmapsto f(t, \omega, y, z)$  is in  $L^2([0, T]; \mathbb{D}^{1,\infty})$ ,  $f(\cdot, y, z)$  and  $Df(\cdot, y, z)$  are  $\mathbb{F}$ -progressively measurable,  $Df(\cdot, y, z)$  is uniformly bounded in  $y, z$ .

(Q) The map  $(y, z) \mapsto f(\cdot, y, z)$  is continuously differentiable and there exists some constant  $C > 0$  such that for any  $(s, \omega, y, z, z') \in [0, T] \times \Omega \times \mathbb{R}^3$

$$\begin{aligned} |f(s, \omega, y, z) - f(s, \omega, y, z')| &\leq C(1 + |z| + |z'|)|z - z'|, \quad |f(s, \omega, 0, 0)| \leq C, \\ |f_y(s, \omega, y, z)| &\leq C, \quad |f_z(s, \omega, y, z)| \leq C(1 + |z|). \end{aligned}$$

( $H_{1,\infty}$ ) For any  $p > 1$  and for any  $h \in H$

$$\lim_{\varepsilon \rightarrow 0} \mathbb{E} \left[ \left( \int_0^T \left| \frac{f(s, \cdot + \varepsilon h, Y_s, Z_s) - f(s, \cdot, Y_s, Z_s)}{\varepsilon} - \langle Df(s, \cdot, Y_s, Z_s), \dot{h} \rangle_{\mathfrak{H}} \right| ds \right)^p \right] = 0.$$

( $H_{2,\infty}$ ) Let  $(\varepsilon_n)_{n \in \mathbb{N}}$  be a sequence in  $(0, 1]$  such that  $\lim_{n \rightarrow +\infty} \varepsilon_n = 0$ , and let  $(Y^n, Z^n)_n$  be a sequence of random variables which converges in  $\mathbb{S}^p \times \mathbb{H}^p$  for any  $p > 1$  to some  $(Y, Z)$ . Then for all  $h \in H$ , the following convergences hold in probability

$$\begin{aligned} \|f_y(\cdot, \omega + \varepsilon_n h, Y^n, Z^n) - f_y(\cdot, \omega, Y, Z)\|_{\mathfrak{H}} &\xrightarrow{n \rightarrow +\infty} 0 \\ \|f_z(\cdot, \omega + \varepsilon_n h, Y^n, Z^n) - f_z(\cdot, \omega, Y, Z)\|_{\mathfrak{H}} &\xrightarrow{n \rightarrow +\infty} 0, \end{aligned} \quad (7.1)$$

or

$$\begin{aligned} \|f_y(\cdot, \omega + \varepsilon_n h, Y^n, Z^n) - f_y(\cdot, \omega, Y, Z)\|_{\mathfrak{H}} &\xrightarrow{n \rightarrow +\infty} 0 \\ \|f_z(\cdot, \omega + \varepsilon_n h, Y^n, Z^n) - f_z(\cdot, \omega, Y, Z)\|_{\mathfrak{H}} &\xrightarrow{n \rightarrow +\infty} 0. \end{aligned} \quad (7.2)$$

Let  $\mathbb{S}^\infty$  be the set of  $\mathbb{F}$ -progressively measurable processes  $Y$  such that  $\sup_{t \in [0, T]} |Y_t|$  is bounded and  $\mathbb{H}_{\text{BMO}}^2$  the set of predictable processes  $Z$  such that:

$$\text{esssup}_{\tau \in \mathcal{T}} \mathbb{E} \left[ \int_\tau^T |Z_s|^2 ds \middle| \mathcal{F}_\tau \right] < +\infty, \quad \mathbb{P}_0 - a.s.,$$

where  $\mathcal{T}$  denotes the set of  $\mathbb{F}$ -stopping times with values in  $[0, T]$ . We start by recalling the following by now classical results on quadratic growth BSDEs and stochastic Lipschitz BSDEs, which can be found among others in [7].

**Proposition 7.1.** *Under Assumptions ( $D_\infty$ ) and (Q), the BSDEs (5.1) and (5.5) both admit a unique solution in  $\mathbb{S}^\infty \times \mathbb{H}_{\text{BMO}}^2$ .*

We have the following extension of Theorem 5.1.

**Theorem 7.1.** *Let  $t$  be in  $[0, T]$ . Under Assumptions ( $D_\infty$ ), (Q), ( $H_{1,\infty}$ ) and ( $H_{2,\infty}$ ),  $Y_t$  belongs to  $\mathbb{D}^{1,\infty}$  and  $Z \in L^\infty([t, T]; \mathbb{D}^{1,2})$ .*

**Proof.** We follow the proof of Theorem (5.1), using the same notations. Since the BSDEs are now quadratic, we can use the *a priori* estimates of Lemma A.1 in [7] to obtain that for any

$p > 1$ , there exists some  $q > 1$  such that

$$\begin{aligned}
& \mathbb{E} \left[ \sup_{s \in [t, T]} |Y_s^\varepsilon - \tilde{Y}_s^h|^{2p} \right] + \mathbb{E} \left[ \left( \int_t^T |Z_s^\varepsilon - \tilde{Z}_s^h|^2 ds \right)^p \right] \\
& \leq C_p \left( \mathbb{E} \left[ |\xi^\varepsilon - \langle D\xi, \dot{h} \rangle_{\mathfrak{H}}|^{pq} \right]^{1/q} + \mathbb{E} \left[ \left( \int_0^T |\tilde{A}_s^\varepsilon - \langle Df(s, \cdot, Y_s, Z_s), \dot{h} \rangle_{\mathfrak{H}}| ds \right)^{pq} \right]^{1/q} \right) \\
& \quad + C_p \mathbb{E} \left[ \left( \int_0^T |\tilde{A}_s^{y, \varepsilon} - f_y(s, \cdot, Y_s, Z_s)| |\tilde{Y}_s^h| ds \right)^{pq} \right]^{1/q} \\
& \quad + C_p \mathbb{E} \left[ \left( \int_0^T |\tilde{A}_s^{z, \varepsilon} - f_z(s, \cdot, Y_s, Z_s)| |\tilde{Z}_s^h| ds \right)^{pq} \right]^{1/q}. \tag{7.3}
\end{aligned}$$

Since  $\xi \in \mathbb{D}^{1, \infty}$ , the first term on the right-hand side above goes to 0 thanks to Theorem 4.1. Moreover, the second term also goes to 0 thanks to Assumption  $(H_\infty)$ . Then, since  $f_y$  is bounded by Assumption  $(Q)$  and since  $\tilde{Y}^h \in \mathbb{S}^\infty$  by Proposition 7.1, we can easily conclude with Assumption  $(H_2)$  and the dominated convergence theorem that the third term on the right-hand side also goes to 0. Let us now concentrate on the fourth term involving the control variable. By Cauchy-Schwarz inequality we have that

$$\begin{aligned}
& \mathbb{E} \left[ \left( \int_0^T |\tilde{A}_s^{z, \varepsilon} - f_z(s, \cdot, Y_s, Z_s)| |\tilde{Z}_s^h| ds \right)^{pq} \right] \\
& \leq \mathbb{E} \left[ \left( \int_0^T |\tilde{A}_s^{z, \varepsilon} - f_z(s, \cdot, Y_s, Z_s)|^2 ds \right)^{pq} \right]^{1/2} \mathbb{E} \left[ \left( \int_0^T |\tilde{Z}_s^h|^2 ds \right)^{pq} \right]^{1/2}. \tag{7.4}
\end{aligned}$$

Since  $(\tilde{Y}^h, \tilde{Z}^h)$  is the solution to the stochastic linear BSDE (5.5) with bounded coefficients  $Df$  and  $f_y$  (by  $(D_\infty)$ ) and  $f_z(s, Y_s, Z_s)$  is in  $\mathbb{H}_{\text{BMO}}^2$  since  $|f_z(s, Y_s, Z_s)| \leq C(1 + |Z_s|)$  (by Assumption  $(Q)$ ), we deduce that  $\tilde{Z}^h \in \mathbb{H}_{\text{BMO}}^2$  which implies that  $\tilde{Z}^h \in \mathbb{H}^m$  for any  $m > 1$  by the energy Inequalities. Furthermore, for any  $\eta > 0$  it holds that

$$\begin{aligned}
& \mathbb{E} \left[ \left( \int_0^T |\tilde{A}_s^{z, \varepsilon} - f_z(s, \cdot, Y_s, Z_s)| |\tilde{Z}_s^h| ds \right)^{pq+\eta} \right] \\
& \leq C \mathbb{E} \left[ \left( \int_0^T (1 + |Z_s| + |Z_s \circ \tau_{\varepsilon h}|) |\tilde{Z}_s^h| ds \right)^{pq+\eta} \right] \\
& \leq C \mathbb{E} \left[ \left( \int_0^T (1 + |Z_s| + |Z_s \circ \tau_{\varepsilon h}|)^2 ds \right)^{\frac{pq+\eta}{2}} \left( \int_0^T |\tilde{Z}_s^h|^2 ds \right)^{\frac{pq+\eta}{2}} \right] \\
& \leq C \mathbb{E} \left[ \left( \int_0^T (1 + |Z_s| + |Z_s \circ \tau_{\varepsilon h}|)^2 ds \right)^{pq+\eta} \right]^{1/2} \mathbb{E} \left[ \left( \int_0^T |\tilde{Z}_s^h|^2 ds \right)^{pq+\eta} \right]^{1/2} \\
& \leq C \left( 1 + \mathbb{E} \left[ \left( \int_0^T |Z_s|^2 ds \right)^{p'} \right]^{1/q'} \right) \mathbb{E} \left[ \left( \int_0^T |\tilde{Z}_s^h|^2 ds \right)^{pq+\eta} \right]^{1/2} < \infty,
\end{aligned}$$

where  $p', q' > 1$  using Hölder Inequality and Proposition 3.1. Hence, taking limit as  $\varepsilon$  goes to 0 in (7.4) we get that  $\lim_{\varepsilon \rightarrow 0} \mathbb{E} \left[ \sup_{s \in [t, T]} |Y_s^\varepsilon - \tilde{Y}_s^h|^{2p} \right] + \mathbb{E} \left[ \left( \int_t^T |Z_s^\varepsilon - \tilde{Z}_s^h|^2 ds \right)^p \right] = 0$ . This proves that  $Y_t$  and  $\int_t^T Z_s dW_s$  belongs to  $\mathbb{D}^{1, \infty}$  by Theorem 4.1. In particular,  $Z\mathbf{1}_{[t, T]}$  belongs to  $L^2([t, T]; \mathbb{D}^{1, 2})$  (see [12]). Moreover, since  $(D_t Y, D_t Z)$  is the solution of the stochastic linear BSDE (5.7) for any  $t \in [0, T]$  and Assumptions  $(D_\infty)$  and  $(Q)$  hold, from the relation  $D_t Y_t = Z_t$  for all  $t \in [0, T]$  we obtain  $Z\mathbf{1}_{[t, T]} \in L^\infty([t, T]; \mathbb{D}^{1, 2})$ .  $\square$

**Remark 7.1.** *We would like to point out that our conditions cover the case of Markovian quadratic BSDEs presented in [6, Theorem 2.9]. Indeed, assume that we consider a forward-backward system of the form (6.1) under assumptions,  $(D_\infty)$ ,  $(Q)$ ,  $(A_1)$ ,  $(A_2)(i)$  and where  $(A_2)(ii)$  is replaced by the following assumption:*

$(A_2)(ii')$   $f : [0, T] \times \mathbb{R}^3 \rightarrow \mathbb{R}$  is continuously differentiable in  $(x, y, z)$  and satisfying for some  $C > 0$

$$\exists q \in \mathbb{R}_+, |f_x(t, x, y, z)| \leq C(1 + |y| + |z|^2 + |x|^q), \forall (t, x, y, z) \in [0, T] \times \mathbb{R}^3.$$

Under these assumptions, we can check that  $(H_{1, \infty})$  and  $(H_{2, \infty})$  are in force. To see this we just make a comment about how the proof of Theorem 6.1 has to be modified to obtain  $(H_{1, \infty})$ , whereas  $(H_{2, \infty})$  is met trivially. Using the notations of this proof one can manage a term of the form:

$$\mathbb{E} \left[ \left( \int_0^T \left| \varepsilon^{-1}(X_t \circ \tau_{\varepsilon h} - X_t) - \langle DX_t, \dot{h} \rangle_{\mathfrak{H}} \right| |f_x(t, \bar{X}_t, Y_t, Z_t)| dt \right)^p \right]$$

as follows:

$$\begin{aligned} & \mathbb{E} \left[ \left( \int_0^T \left| \varepsilon^{-1}(X_t \circ \tau_{\varepsilon h} - X_t) - \langle DX_t, \dot{h} \rangle_{\mathfrak{H}} \right| |f_x(t, \bar{X}_t, Y_t, Z_t)| dt \right)^p \right] \\ & \leq C \mathbb{E} \left[ \sup_{t \in [0, T]} \left| \frac{X_t \circ \tau_{\varepsilon h} - X_t}{\varepsilon} - \langle DX_t, \dot{h} \rangle_{\mathfrak{H}} \right|^p \left( \int_0^T (1 + |X_t|^q + |X_t \circ \tau_{\varepsilon h}|^q + |Y_t| + |Z_t|^2) dt \right)^p \right] \\ & \leq C \mathbb{E} \left[ \sup_{t \in [0, T]} \left| \varepsilon^{-1}(X_t \circ \tau_{\varepsilon h} - X_t) - \langle DX_t, \dot{h} \rangle_{\mathfrak{H}} \right|^{2p} \right]^{1/2} \\ & \quad \times \mathbb{E} \left[ \left( \int_0^T (1 + |X_t|^q + |X_t \circ \tau_{\varepsilon h}|^q + |Y_t| + |Z_t|^2) dt \right)^{2p} \right]^{1/2}, \end{aligned}$$

which goes to 0 as  $\varepsilon$  goes to 0 since  $Z$  belongs to  $\mathbb{H}_{\text{BMO}}^2$  and since  $Y$  is bounded. The term involving  $A^{2, \varepsilon}$  can be treated similarly.

## A Appendix

The following lemma was remarked in [16, Remark 2], we provide a proof for the comfort of the reader.

**Lemma A.1.** *Let  $p > 1$  and  $F$  be in  $L^p(\mathbb{R})$ ,  $G \in \mathcal{S}$  and  $h \in H$ . The mapping  $t \mapsto \mathbb{E}[F \circ \tau_{th} G]$  is differentiable in  $t$  and*

$$\frac{d}{dt} \mathbb{E}[F \circ \tau_{th} G] = \mathbb{E}[F \circ \tau_{th} \delta(Gh)]. \quad (\text{A.1})$$



**Proof.** Let  $s > 0$ , by the Cameron-Martin formula, we have that

$$\begin{aligned} & s^{-1} \left( \mathbb{E}[F \circ \tau_{(s+t)h} G] - \mathbb{E}[F \circ \tau_{th} G] \right) \\ &= \mathbb{E} \left[ F \circ \tau_{th} \frac{G \circ \tau_{-sh} \exp \left( s \int_0^T \dot{h}(u) dW_u - \frac{|s|^2}{2} \int_0^T |\dot{h}(u)|^2 du \right) - G}{s} \right]. \end{aligned}$$

Hence

$$\begin{aligned} & \lim_{s \rightarrow 0} s^{-1} \left( \mathbb{E}[F \circ \tau_{(s+t)h} G] - \mathbb{E}[F \circ \tau_{th} G] \right) \\ &= \mathbb{E} \left[ F \circ \tau_{th} \lim_{s \rightarrow 0} \frac{G \circ \tau_{-sh} \exp \left( s \int_0^T \dot{h}(u) dW_u - \frac{|s|^2}{2} \int_0^T |\dot{h}(u)|^2 du \right) - G}{s} \right] \\ &= \mathbb{E} \left[ F \circ \tau_{th} \lim_{s \rightarrow 0} \left( \frac{G \circ \tau_{-sh} - G}{s} + G \circ \tau_{-sh} \frac{\exp \left( s \int_0^T \dot{h}(u) dW_u - \frac{|s|^2}{2} \int_0^T |\dot{h}(u)|^2 du \right) - 1}{s} \right) \right], \end{aligned}$$

where the exchange between the limit and the expectation is justified by the fact that

$$\sup_{s \in (0,1]} s^{-q} \mathbb{E} \left[ \left| G \circ \tau_{-sh} \exp \left( s \int_0^T \dot{h}(u) dW_u - \frac{|s|^2}{2} \int_0^T |\dot{h}(u)|^2 du \right) - G \right|^q \right] < \infty \quad (\text{A.2})$$

for any  $q > 1$  and by the fact  $F \circ \tau_{th}$  is in  $L^p$  by the Cameron-Martin formula. Indeed for any  $r$  in  $(1, p)$  we have by Hölder Inequality:

$$\begin{aligned} & \mathbb{E} \left[ \left| F \circ \tau_{th} \frac{G \circ \tau_{-sh} \exp \left( s \int_0^T \dot{h}(u) dW_u - \frac{|s|^2}{2} \int_0^T |\dot{h}(u)|^2 du \right) - G}{s} \right|^r \right] \\ & \leq \underbrace{\mathbb{E} [|F \circ \tau_{th}|^{p_1}]^{r/p_1}}_{=: E_1} \underbrace{\mathbb{E} \left[ \left| \frac{G \circ \tau_{-sh} \exp \left( s \int_0^T \dot{h}(u) dW_u - \frac{|s|^2}{2} \int_0^T |\dot{h}(u)|^2 du \right) - G}{s} \right|^{rp_2} \right]^{1/p_2}}_{=: E_2}, \end{aligned}$$

where  $r < p_1 < p$  and  $p_2$  is the Hölder conjugate of  $p_1/r$ . Using Cameron-Martin Formula for  $E_1$ , Relation (3.1) and Hölder Inequality with  $r_1 = \frac{p}{p_1}$  and  $r_2$  such that  $\frac{1}{r_1} + \frac{1}{r_2} = 1$ , we deduce that:

$$E_1 \leq \mathbb{E}[|F|^p]^{r/p} \mathbb{E} \left[ \left| \mathcal{E} \left( \int_0^T \dot{h}_s dW_s \right) \right|^{r_2} \right]^{1/r_2} < +\infty.$$

We now turn to  $E_2$ , for any  $q > 1$

$$\begin{aligned} & \sup_{s \in (0,1]} \mathbb{E} \left[ \left| \frac{G \circ \tau_{-sh} \mathcal{E} \left( s \int_0^T \dot{h}(u) dW_u \right) - G}{s} \right|^q \right] \\ & \leq \underbrace{\sup_{s \in (0,1]} \mathbb{E} \left[ \left| \frac{G \circ \tau_{-sh} - G}{s} \mathcal{E} \left( s \int_0^T \dot{h}(u) dW_u \right) \right|^q \right]}_{=: A_1} + \underbrace{\sup_{s \in (0,1]} \mathbb{E} \left[ \left| \frac{\mathcal{E} \left( s \int_0^T \dot{h}(u) dW_u \right) - 1}{s} G \right|^q \right]}_{=: A_2}, \end{aligned}$$

hence, on the one hand there exists  $\alpha_1, \alpha_2 > 1$  such that:

$$A_1 \leq \sup_{s \in (0,1]} s^{-q} \mathbb{E} [|G \circ \tau_{-sh} - G|^{q\alpha_1}]^{\frac{1}{\alpha_1}} \mathbb{E} \left[ \left| \mathcal{E} \left( s \int_0^T \dot{h}(u) dW_u \right) \right|^{q\alpha_2} \right]^{\frac{1}{\alpha_2}} < +\infty,$$

using the fact that  $G$  is polynomial, so  $G$  is locally Lipschitz and we conclude by Relation (3.1). On the other hand, using the mean value Theorem and Relation 3.1, we obtain also  $A_2 < +\infty$ . We deduce that Relation (A.2) holds. Moreover, given that  $G \in \mathcal{S}$  is polynomial, we deduce that  $\frac{G \circ \tau_{-sh} - G}{s} \xrightarrow{s \rightarrow 0} -\langle \nabla G, h \rangle_H$  a.s.. Hence,

$$\begin{aligned} & \lim_{s \rightarrow 0} s^{-1} (\mathbb{E}[F \circ \tau_{(s+t)h} G] - \mathbb{E}[F \circ \tau_{th} G]) \\ &= \mathbb{E} \left[ F \circ \tau_{th} \lim_{s \rightarrow 0} \left( \frac{G \circ \tau_{-sh} - G}{s} + G \circ \tau_{-sh} \frac{\exp \left( s \int_0^T \dot{h}(u) dW_u - \frac{|s|^2}{2} \int_0^T |\dot{h}(u)|^2 du \right) - 1}{s} \right) \right] \\ &= \mathbb{E} [F \circ \tau_{th} (-\langle \nabla G, h \rangle_H + G\delta(h))] = \mathbb{E} [F \circ \tau_{th} \delta(Gh)], \end{aligned}$$

by (2.3), so (A.1) holds.  $\square$

**Lemma A.2.** *Let  $\alpha$  in  $L^2([0, T]; \mathbb{D}^{1,2})$ . Then for any  $p$  in  $(1, 2)$ ,*

$$\lim_{\varepsilon \rightarrow 0} \mathbb{E} \left[ \int_0^T \left| \frac{\alpha_s \circ \tau_{\varepsilon h} - \alpha_s}{\varepsilon} - \langle \nabla \alpha_s, h \rangle_H \right|^p ds \right] = 0.$$

**Proof.** Note first that the space  $L^2([0, T]; \mathbb{D}^{1,2})$  can be identified with the space  $\mathbb{D}^{1,2}(\mathfrak{H})$  which is the completion of the set of  $\mathfrak{H}$ -valued r.v. of the form:

$$\sum_{i=1}^n F_i u_i, \quad F_i \in \mathcal{S}, \quad u_i \in L^2([0, T]), \quad n \geq 1,$$

with respect to the norm  $\|\cdot\|_{1,2,2}$  defined as:

$$\|u\|_{1,2,2}^2 := \mathbb{E}[\|u\|_{L^2([0,T])}^2] + \mathbb{E}[\|\nabla u\|_{H \otimes L^2([0,T])}^2].$$

Alternatively, an element  $u$  in  $\mathbb{D}^{1,2}(L^2([0, T]))$  is identified with a stochastic process such that for almost every  $t$  in  $[0, T]$ ,  $u_t$  belongs to  $\mathbb{D}^{1,2}$  and such that

$$\mathbb{E}[\|\nabla u\|_{H \otimes L^2([0,T])}^2] = \mathbb{E} \left[ \int_0^T \int_0^T |D_s u_t|^2 ds dt \right] < +\infty.$$

Hence we can assume that  $\alpha$  belongs to  $\mathbb{D}^{1,2}(L^2([0, T]))$ . Thus by [16, Theorem 3.1],  $\alpha$  satisfies (RAC) and (SGD), which entails in this setting that for any  $h$  in  $H$ , there exists a  $L^2([0, T])$ -valued r.v.  $\tilde{\alpha}_h$  such that  $\tilde{\alpha}_h = \alpha$  in  $L^2([0, T])$ ,  $\mathbb{P}_0$ -a.s., and for any  $\varepsilon > 0$

$$\frac{\tilde{\alpha}_h \circ \tau_{\varepsilon h} - \tilde{\alpha}_h}{\varepsilon} = \varepsilon^{-1} \int_0^\varepsilon \langle \nabla \alpha \circ \tau_{sh}, h \rangle_H ds, \quad \text{in } L^2([0, T]), \quad \mathbb{P}_0 - a.s..$$

Using Lemma 3.1 we thus get that for any  $r \in (p, 2)$ , it holds that:

$$\begin{aligned}
\mathbb{E} \left[ \int_0^T |\varepsilon^{-1}(\alpha_s \circ \tau_{\varepsilon h} - \alpha_s)|^r ds \right] &= \mathbb{E} \left[ \int_0^T |\varepsilon^{-1}((\tilde{\alpha}_h)(s) \circ \tau_{\varepsilon h} - (\tilde{\alpha}_h)(s))|^r ds \right] \\
&\leq \mathbb{E} \left[ \int_0^T \varepsilon^{-1} \int_0^\varepsilon |\langle \nabla \alpha_s \circ \tau_{uh}, h \rangle_H|^r dud s \right] \\
&\leq C \int_0^T \mathbb{E} [|\langle \nabla \alpha_s, h \rangle_H|^p]^{r/p} ds \\
&\leq C \mathbb{E} \left[ \int_0^T |\langle \nabla \alpha_s, h \rangle_H|^p ds \right]^{r/p} \\
&\leq C \|h\|_H^r \mathbb{E} [\|\nabla \alpha\|_{H \otimes \mathfrak{H}}^p]^{r/p} < +\infty,
\end{aligned}$$

where we have used Cameron-Martin formula and similar computations to those of the proof of Lemma 4.2, and  $C$  denotes a positive constant which can differ from line to line. Hence, the family  $\left( \int_0^T |\varepsilon^{-1}(\alpha_s \circ \tau_{\varepsilon h} - \alpha_s) - \langle \nabla \alpha_s, h \rangle_H|^p ds \right)_{\varepsilon \in (0,1)}$  is uniformly integrable. In addition, by Property (SGD),  $\varepsilon^{-1}(\alpha \circ \tau_{\varepsilon h} - \alpha)$  converges in probability to  $\langle \nabla \alpha, h \rangle_H$  (with respect to the norm  $L^2([0, T])$ ) which implies that  $\int_0^T |\varepsilon^{-1}(\alpha_s \circ \tau_{\varepsilon h} - \alpha_s) - \langle \nabla \alpha_s, h \rangle_H|^p ds$  converges in probability to 0 as  $\varepsilon$  goes to 0, which provides the result.  $\square$

## Acknowledgments

Thibaut Mastrolia is grateful to Région Ile-De-France for financial support.

## References

- [1] S. Ankirchner, P. Imkeller, and G. Dos Reis. Classical and variational differentiability of bsdes with quadratic growth. *Electron. J. Probab.*, 12(53):1418–1453, 2007.
- [2] J.-M. Bismut. Contrôle des systèmes linéaires quadratiques: applications de l’intégrale stochastique. In *Séminaire de Probabilités, XII (Univ. Strasbourg, Strasbourg, 1976/1977)*, volume 649 of *Lecture Notes in Math.*, pages 180–264. Springer, Berlin, 1978.
- [3] P. Briand, B. Delyon, Y. Hu, E. Pardoux, and L. Stoica.  $L^p$  solutions of backward stochastic differential equations. *Stochastic Process. Appl.*, 108(1):109–129, 2003.
- [4] G. dos Reis. *On some properties of solutions of quadratic growth BSDE and applications in finance and insurance*. PhD thesis, Humboldt University, 2010.
- [5] N. El Karoui, S. Peng, and M.C. Quenez. Backward stochastic differential equations in finance. *Mathematical finance*, 7(1):1–71, 1997.
- [6] P. Imkeller and G. Dos Reis. Path regularity and explicit convergence rate for BSDE with truncated quadratic growth. *Stochastic Process. Appl.*, 120(3):348–379, 2010.
- [7] P. Imkeller, A. Réveillac, and A. Richter. Differentiability of quadratic bsdes generated by continuous martingales. *Ann. Appl. Probab.*, 22(1):285–336, 2012.

- [8] S. Kusuoka. Dirichlet forms and diffusion processes on Banach spaces. *J. Fac. Sci. Univ. Tokyo Sect. IA Math.*, 29(1):79–95, 1982.
- [9] P. Malliavin. Stochastic calculus of variation and hypoelliptic operators. In *Proceedings of the International Symposium on Stochastic Differential Equations (Res. Inst. Math. Sci., Kyoto Univ., Kyoto, 1976)*, pages 195–263. Wiley, New York-Chichester-Brisbane, 1978.
- [10] D. Nualart. *The Malliavin calculus and related topics*. Probability and its Applications (New York). Springer-Verlag, Berlin, second edition, 2006.
- [11] E. Pardoux and S. Peng. Adapted solution of a backward stochastic differential equation. *Systems Control Lett.*, 14(1):55–61, 1990.
- [12] E. Pardoux and S. Peng. Backward stochastic differential equations and quasilinear parabolic partial differential equations. In *Stochastic partial differential equations and their applications*, pages 200–217. Springer, 1992.
- [13] P. E. Protter. *Stochastic integration and differential equations*, volume 21 of *Stochastic Modelling and Applied Probability*. Springer-Verlag, Berlin, 2005. Second edition. Version 2.1, Corrected third printing.
- [14] I. Shigekawa. Derivatives of Wiener functionals and absolute continuity of induced measures. *J. Math. Kyoto Univ.*, 20(2):263–289, 1980.
- [15] D.W. Stroock. The Malliavin calculus and its application to second order parabolic differential equations. I. *Math. Systems Theory*, 14(1):25–65, 1981.
- [16] H. Sugita et al. On a characterization of the sobolev spaces over an abstract wiener space. *Journal of Mathematics of Kyoto University*, 25(4):717–725, 1985.
- [17] A.S. Üstünel and M. Zakai. *Transformation of measure on Wiener space*. Springer, 2000.