



**HAL**  
open science

## Genome Sequence of a Hyperthermophilic Archaeon, *Thermococcus nautili* 30-1, That Produces Viral Vesicles

Jacques Oberto, Marie Gaudin, Matteo Cossu, Aurore Gorlas, Alexei Slesarev,  
Evelyne Marguet, Patrick Forterre

### ► To cite this version:

Jacques Oberto, Marie Gaudin, Matteo Cossu, Aurore Gorlas, Alexei Slesarev, et al.. Genome Sequence of a Hyperthermophilic Archaeon, *Thermococcus nautili* 30-1, That Produces Viral Vesicles. *Genome Announcements*, 2014, 2 (2), epub ahead of print. 10.1128/genomeA.00243-14. hal-00975406

**HAL Id: hal-00975406**

**<https://hal.science/hal-00975406>**

Submitted on 11 Apr 2014

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# Genome Sequence of a Hyperthermophilic Archaeon, *Thermococcus nautili* 30-1, That Produces Viral Vesicles

Jacques Oberto,<sup>a</sup> Marie Gaudin,<sup>a</sup> Matteo Cossu,<sup>a</sup> Aurore Gorlas,<sup>a</sup> Alexei Slesarev,<sup>b</sup> Evelyne Marguet,<sup>a</sup> Patrick Forterre<sup>a,c</sup>

Université Paris-Sud, CNRS, UMR8621, Institut de Génétique et Microbiologie, Orsay, France<sup>a</sup>; Fidelity Systems, Inc., Gaithersburg, Maryland, USA<sup>b</sup>; Institut Pasteur, Paris, France<sup>c</sup>

***Thermococcus nautili* 30-1 (formerly *Thermococcus nautilus*), an anaerobic hyperthermophilic marine archaeon, was isolated in 1999 from a deep-sea hydrothermal vent during the Amistad campaign. Here, we present the complete sequence of *T. nautili*, which is able to produce membrane vesicles containing plasmid DNA. This property makes *T. nautili* a model organism to study horizontal gene transfer.**

Received 28 February 2014 Accepted 13 March 2014 Published 27 March 2014

**Citation** Oberto J, Gaudin M, Cossu M, Gorlas A, Slesarev A, Marguet E, Forterre P. 2014. Genome sequence of a hyperthermophilic archaeon, *Thermococcus nautili* 30-1, that produces viral vesicles. *Genome Announc.* 2(2):e00243-14. doi:10.1128/genomeA.00243-14.

**Copyright** © 2014 Oberto et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 3.0 Unported license](http://creativecommons.org/licenses/by/3.0/).

Address correspondence to Jacques Oberto, jacques.oberto@igmors.u-psud.fr.

*Thermococcus nautili* 30-1 (formerly *Thermococcus nautilus*) was isolated on the East Pacific ridge near a black smoker, 2,600 m below the ocean surface (1). *T. nautili* qualifies as a new species and the characterization of its physiology and metabolism has been reported (2).

*T. nautili* was grown anaerobically in ZB medium at 85°C (1), and genomic DNA was purified and sequenced using paired-end Illumina libraries to generate  $16 \times 10^6$  reads assembled with Velvet (3). Genome assembly was performed with Consed (4). The remaining gaps were filled using bacterial artificial chromosome (BAC) sequencing with Thermofidélase and Fimers (5). Genes were annotated using FGENESB (Softberry). Additional functional annotations were obtained with Integrated Microbial Genomes (6). The chromosome of *T. nautili* is 1,976,356 bp long and harbors 2,288 genes. The noncoding genes comprise 4 rRNA genes (23S, BD01\_1261, and 16S, BD01\_1259), with a duplication of the 5S rRNA gene (BD01\_1590 and BD01\_2049), and 46 tRNA genes. The chromosomal origin of replication was identified with FIT-BAR (7) using published origin recognition complex (ORC) sequences (8). It is positioned upstream of the CDC6 gene (BD01\_1824). A similar approach located the replication termination region at coordinate 772,784 using archaeal *dif* sites (9).

Several strains of *Thermococcus* contain either one or two plasmids (10–13). Our group previously sequenced and characterized two plasmids of 3.6 and 13 kb from *T. nautili* that were named pTN1 and pTN2, respectively (14–16). The rolling-circle plasmid pTN1 was used to construct the first shuttle vector able to replicate in both *Escherichia coli* and the model organism *Thermococcus kodakaraensis* (17) and contributed to the development of *T. kodakaraensis* as a host for genetics studies (18). Plasmid pTN2 encodes a new family of DNA primase formed by the fusion of protein domains homologous to the archaeo-eukaryotic primases PriS and PriL (19). DNA sequencing revealed the presence of an additional plasmid of 18 kb, pTN3 (16). This extrachromosomal element, related to the virus-like element TKV4 from *T. kodakaraensis* (20), is found as a freely replicating plasmid and also

integrated at a specific position on the chromosome into a tRNA<sup>Leu</sup> gene (BD01\_0018). The plasmid attachment site (*attP*) is located within the integrase-encoding gene TnaP3-13 (16). The *attP* and *attB* sites share an identical stretch of 41 nucleotides (nt), as observed previously in the case of the TKV4 integron of *T. kodakaraensis* (20). In the chromosome of *T. nautili*, 6 clustered regularly interspaced short palindromic repeat (CRISPR) loci were found using CRISPRFinder (21). Interestingly, two CRISPR spacers are directed against resident plasmid pTN3.

A major interest in the study of *T. nautili* is its ability to generate membrane vesicles containing plasmid DNA (22), including the virus-like genome of pTN3 (viral vesicles) (16). The discovery that these vesicles can transfer genetic information between related *Thermococcales* (23) provides an important tool for the study of horizontal gene transfer across species or even domains, from which genomic evolution models can be inferred. The genetic determinism of the biosynthesis of these viral vesicles is presently being investigated and relies primarily on the *T. nautili* genome sequence presented here.

**Nucleotide sequence accession number.** The *T. nautili* genome sequence has been deposited in the NCBI repository under the accession number CP007264.

## ACKNOWLEDGMENTS

This work was supported by the Centre National de la Recherche Scientifique (Program Marine Genome Biology) and the Agence Nationale de la Recherche (ANR 12-BSV3-0023-01).

## REFERENCES

1. Lepage E, Marguet E, Geslin C, Matte-Tailliez O, Zillig W, Forterre P, Tailliez P. 2004. Molecular diversity of new *Thermococcales* isolates from a single area of hydrothermal deep-sea vents as revealed by randomly amplified polymorphic DNA fingerprinting and 16S rRNA gene sequence analysis. *Appl. Environ. Microbiol.* 70:1277–1286. <http://dx.doi.org/10.1128/AEM.70.3.1277-1286.2004>.
2. Gorlas A, Croce O, Oberto J, Gaudiard E, Forterre P, Marguet E. 20 February 2014. *Thermococcus nautili* sp. nov., a hyperthermophilic ar-

- chaeon isolated from a hydrothermal deep sea vent (East Pacific Ridge). *Int. J. Syst. Evol. Microbiol.* <http://dx.doi.org/10.1099/ijs.0.060376-0>.
3. Zerbino DR, Birney E. 2008. Velvet: algorithms for de novo short read assembly using de Bruijn graphs. *Genome Res.* 18:821–829. <http://dx.doi.org/10.1101/gr.074492.107>.
  4. Gordon D, Green P. 2013. ConSeq: a graphical editor for next-generation sequencing. *Bioinformatics* 29:2936–2937. <http://dx.doi.org/10.1093/bioinformatics/btt515>.
  5. Malykh A, Malykh O, Polushin N, Kozyavkin S, Slesarev A. 2004. Finishing “working draft” BAC projects by directed sequencing with ThermoFidelase and Fimers. *Methods Mol. Biol.* 255:295–308. <http://dx.doi.org/10.1385/1-59259-752-1:295>.
  6. Markowitz VM, Mavromatis K, Ivanova NN, Chen IM, Chu K, Kyrpides NC. 2009. IMG ER: a system for microbial genome annotation expert review and curation. *Bioinformatics* 25:2271–2278. <http://dx.doi.org/10.1093/bioinformatics/btp393>.
  7. Oberto J. 2010. FITBAR: a web tool for the robust prediction of prokaryotic regulons. *BMC Bioinformatics* 11:554. <http://dx.doi.org/10.1186/1471-2105-11-554>.
  8. Matsunaga F, Glatigny A, Mucchielli-Giorgi MH, Agier N, Delacroix H, Marisa L, Durosay P, Ishino Y, Aggerbeck L, Forterre P. 2007. Genome-wide and biochemical analyses of DNA-binding activity of Cdc6/Orc1 and Mcm proteins in *Pyrococcus* sp. *Nucleic Acids Res.* 35:3214–3222. <http://dx.doi.org/10.1093/nar/gkm212>.
  9. Cortez D, Quevillon-Cheruel S, Gribaldo S, Desnoues N, Sezonov G, Forterre P, Serre MC. 2010. Evidence for a Xer/dif system for chromosome resolution in archaea. *PLoS Genet.* 6:e1001166. <http://dx.doi.org/10.1371/journal.pgen.1001166>.
  10. Gonnet M, Erauso G, Prieur D, Le Romancer M. 2011. pAMT11, a novel plasmid isolated from a *Thermococcus* sp. strain closely related to the virus-like integrated element TKV1 of the *Thermococcus kodakaraensis* genome. *Res. Microbiol.* 162:132–143. <http://dx.doi.org/10.1016/j.resmic.2010.11.003>.
  11. Krupovic M, Gonnet M, Hania WB, Forterre P, Erauso G. 2013. Insights into dynamics of mobile genetic elements in hyperthermophilic environments from five new *Thermococcus* plasmids. *PLoS One* 8:e49044. <http://dx.doi.org/10.1371/journal.pone.0049044>.
  12. Gorlas A, Krupovic M, Forterre P, Geslin C. 2013. Living side by side with a virus: characterization of two novel plasmids from *Thermococcus prieurii*, a host for the spindle-shaped virus TPV1. *Appl. Environ. Microbiol.* 79:3822–3828. <http://dx.doi.org/10.1128/AEM.00525-13>.
  13. Vannier P, Marteinsson VT, Fridjonsson OH, Oger P, Jebbar M. 2011. Complete genome sequence of the hyperthermophilic, piezophilic, heterotrophic, and carboxydophilic archaeon *Thermococcus barophilus* MP. *J. Bacteriol.* 193:1481–1482. <http://dx.doi.org/10.1128/JB.01490-10>.
  14. Soler N, Justome A, Quevillon-Cheruel S, Lorieux F, Le Cam E, Marguet E, Forterre P. 2007. The rolling-circle plasmid pTN1 from the hyperthermophilic archaeon *Thermococcus nautilus*. *Mol. Microbiol.* 66:357–370. <http://dx.doi.org/10.1111/j.1365-2958.2007.05912.x>.
  15. Soler N, Marguet E, Cortez D, Desnoues N, Keller J, van Tilbeurgh H, Sezonov G, Forterre P. 2010. Two novel families of plasmids from hyperthermophilic archaea encoding new families of replication proteins. *Nucleic Acids Res.* 38:5088–5104. <http://dx.doi.org/10.1093/nar/gkq236>.
  16. Gaudin M, Krupovic M, Marguet E, Gaudiard E, Cvirkaite-Krupovic V, Le Cam E, Oberto J, Forterre P. 1 August 2013. Extracellular membrane vesicles harbouring viral genomes. *Environ. Microbiol.* <http://dx.doi.org/10.1111/1462-2920.12235>.
  17. Santangelo TJ, Cubonová L, Reeve JN. 2008. Shuttle vector expression in *Thermococcus kodakaraensis*: contributions of cis elements to protein synthesis in a hyperthermophilic archaeon. *Appl. Environ. Microbiol.* 74:3099–3104. <http://dx.doi.org/10.1128/AEM.00305-08>.
  18. Farkas JA, Picking JW, Santangelo TJ. 2013. Genetic techniques for the archaea. *Annu. Rev. Genet.* 47:539–561. <http://dx.doi.org/10.1146/annurev-genet-111212-133225>.
  19. Gill S, Krupovic M, Desnoues N, Beguin P, Sezonov G, Forterre P. 20 January 2014. A highly divergent archaeo-eukaryotic primase from the *Thermococcus nautilus* plasmid, pTN2. *Nucleic Acids Res.* <http://dx.doi.org/10.1093/nar/gkt1385>.
  20. Krupovic M, Bamford DH. 2008. Archaeal proviruses TKV4 and MVV extend the PRD1-adenovirus lineage to the phylum *Euryarchaeota*. *Virology* 375:292–300. <http://dx.doi.org/10.1016/j.virol.2008.01.043>.
  21. Grissa I, Vergnaud G, Pourcel C. 2007. CRISPRFinder: a web tool to identify clustered regularly interspaced short palindromic repeats. *Nucleic Acids Res.* 35:W52–W57. <http://dx.doi.org/10.1093/nar/gkm360>.
  22. Soler N, Gaudin M, Marguet E, Forterre P. 2011. Plasmids, viruses and virus-like membrane vesicles from *Thermococcales*. *Biochem. Soc. Trans.* 39:36–44. <http://dx.doi.org/10.1042/BST0390036>.
  23. Gaudin M, Gaudiard E, Schouten S, Houel-Renault L, Lenormand P, Marguet E, Forterre P. 2013. Hyperthermophilic archaea produce membrane vesicles that can transfer DNA. *Environ. Microbiol. Rep.* 5:109–116. <http://dx.doi.org/10.1111/j.1758-2229.2012.00348.x>.