



HAL
open science

Optimal Values of Multidimensional Mean-Payoff Games

Romain Brenguier, Jean-François Raskin

► **To cite this version:**

Romain Brenguier, Jean-François Raskin. Optimal Values of Multidimensional Mean-Payoff Games. 2014. hal-00977352v1

HAL Id: hal-00977352

<https://hal.science/hal-00977352v1>

Submitted on 11 Apr 2014 (v1), last revised 22 May 2015 (v6)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Optimal Values of Multidimensional Mean-Payoff Games*

Romain Brenguier, Jean-François Raskin

Université Libre de Bruxelles (U.L.B.), Belgium

Abstract We study multidimensional mean-payoff games. More precisely, we want to compute all the values that can be ensured by the first player. Because weights are given in multiple dimensions, there can be incomparable such values, and even an infinite number of incomparable ones. We show that the set the values can be represented by a finite union of convex sets. From the computational point of view, we show that deciding if the intersection of the set of values with points satisfying a set of linear inequations can be done in Σ_2 -P. The problem is both NP-hard and coNP-hard.

1 Definitions

1.1 Arena

We define weighted arenas, played by two players **Eve** and **Adam**.

Definition 1 (Arena). An arena \mathcal{A} is a tuple $\langle \text{Stat}_{\exists}, \text{Stat}_{\forall}, s_0, \text{Edg} \rangle$, where:

- $\text{Stat} = \text{Stat}_{\exists} \cup \text{Stat}_{\forall}$ is a finite set of states;
- $s_0 \in \text{Stat}$ is the initial state;
- $\text{Edg} \subseteq \text{Stat} \times \text{Stat}$ is a finite set of edges;

In an arena \mathcal{A} , whenever we arrive at a state s :

- if $s \in \text{Stat}_{\exists}$, then **Eve** selects a state s' such that $(s, s') \in \text{Edg}$.
- if $s \in \text{Stat}_{\forall}$, then **Adam** selects a state s' such that $(s, s') \in \text{Edg}$.

The game then continues from s' . This process starts from s_0 and is repeated *ad infinitum* to form an infinite sequence of states.

Definition 2 (History and plays). A history of the game \mathcal{G} is a finite non-empty sequence of states i.e. an element of $\text{Stat}^* \cdot \text{Stat}$. We write h_i the i -th state of h , starting from 0, i.e. $h = h_0 \cdot h_1 \cdots h_n$. The length $|h|$ of such an history is $n + 1$. We write $\text{last}(h)$ the last state of h , i.e. $h_{|h|-1}$. A play ρ is an infinite sequence of states, i.e. an element of Stat^ω . We write $\rho_{\leq n}$ for the prefix of ρ of length $n + 1$, i.e. $\rho_0 \cdots \rho_n$. We also write $\rho_{\geq n}$ for the suffix of ρ after $\rho_{\leq n-1}$, i.e. $\rho_n \cdot \rho_{n+1} \cdots$ and $\rho_{\llbracket m, n \rrbracket}$ for $(\rho_{\leq n})_{\geq m}$.

* Work supported by ERC Starting Grant inVEST (279499).

Definition 3 (Strategies). Let \mathcal{A} be an arena, a strategy for *Eve* maps histories ending in a state of Stat_\exists to a successor of that state. Formally it is a function $\sigma_\exists: \text{Stat}^* \cdot \text{Stat}_\exists \rightarrow \text{Stat}$, such that for all history h and state s , $(s, \sigma_\exists(h \cdot s)) \in \text{Edg}$. Similarly, a strategy for *Adam* is a function $\sigma_\forall: \text{Stat}^* \cdot \text{Stat}_\forall \rightarrow \text{Act}$, such that for all history h and state s , $(s, \sigma_\forall(h \cdot s)) \in \text{Edg}$. A strategy profile is a pair of a strategy for *Eve* and a strategy for *Adam*.

Definition 4 (Outcomes). Let \mathcal{A} be an arena, and σ_\exists a strategy for *Eve*. A play ρ is compatible with the strategy σ_\exists if, for all $k \in \mathbb{N}$, if $\rho_k \in \text{Stat}_\exists$ then $\rho_{k+1} = \sigma_\exists(\rho_{\leq k})$. We write $\text{Out}_{\mathcal{G}}(s, \sigma_\exists)$ for the set of paths in \mathcal{G} that are compatible with strategy σ_\exists and have initial state s (i.e. ρ such that $\rho_0 = s$), these plays are called outcomes of σ_\exists from s . We simply write $\text{Out}_{\mathcal{G}}(\sigma_\exists)$ when $s = s_0$.

1.2 Multidimensional Mean Payoff

Definition 5 (Multidimensional weighted games). A weighted game is an arena equipped with a weight function. Formally it is a tuple $\langle \mathcal{A}, w \rangle$, where \mathcal{A} is an arena and $w: \text{Edg} \mapsto \mathbb{Z}^d$.

Given a weight function w , we write w_i the function that associate to an edge e the projection to the i -th dimension of $w(e)$. We now present mean payoff preferences which are given by the long term average of the weights.

Definition 6 (Mean payoff). Given a weight function w , the mean payoff inferior over dimension i of a play ρ is:

$$\underline{\text{MP}}_i(\rho) = \liminf_{n \rightarrow \infty} \frac{1}{n} \sum_{0 \leq k < n} w_i(\rho_k, \rho_{k+1}).$$

Similarly the mean payoff superior over dimension i of a play ρ is:

$$\overline{\text{MP}}_i(\rho) = \limsup_{n \rightarrow \infty} \frac{1}{n} \sum_{0 \leq k < n} w_i(\rho_k, \rho_{k+1}).$$

Definition 7. Let $v \in \mathbb{R}^{2 \cdot d}$, we say that a strategy σ_\exists ensures value v from state s if all outcomes ρ of σ_\exists from state s are such that on each dimension i , $\underline{\text{MP}}_i(\rho) \geq v_i$ and $\overline{\text{MP}}_i(\rho) \geq v_{d+i}$. We write $\text{value}(s)$ for the set of vectors v such that there is a strategy of *Eve* that ensures value v from state s .

1.3 The Value Problem

Our goal is to find the strategy of *Eve* that ensures a value as height as possible. However, since the weight are multidimensional it is not clear how to compare different values. A natural problem, is to try to optimize the value we can ensure with respect to a linear function.

Definition 8. A linear function is given by a vector $a \in \mathbb{R}^d$. The associated function $\alpha_a: \mathbb{R}^d \mapsto \mathbb{R}$ is the function such that $\alpha_a(x) = \sum_{i \in \llbracket 1, d \rrbracket} a_i \cdot x_i$. A linear equation is a pair (a, b) where $a \in \mathbb{R}^d \setminus \{0\}$ and $b \in \mathbb{R}$. The half-space satisfying (a, b) is the set $\mathbf{hsp}(a, b) = \{x \in \mathbb{R}^d \mid \alpha_a(x) \geq b\}$. The hyperplane defined by (a, b) is the set $\mathbf{hp}(a, b) = \{x \in \mathbb{R}^d \mid \alpha_a(x) = b\}$.

Given a , we are looking for a strategy σ_{\exists} that ensures a value $v \in \mathbb{R}^d$, and such that there is no σ'_{\exists} that ensures a value v' , with $\alpha(v') > \alpha(v)$. To make this into a decision problem, we fix a real b , and ask if it is possible to ensure a value v such that $\alpha(v) \geq b$.

Definition 9 (Value problem). Given a mean payoff game \mathcal{G} and a linear equation (a, b) over elements of $\mathbb{R}^{2 \cdot d}$, is there a strategy σ_{\exists} and a value $v \in \mathbf{hsp}(a, b)$ such that σ_{\exists} ensures v ?

Remark 1. The answer to this problem is also useful when computing Nash equilibria in (single dimension) mean payoff games. Indeed, thanks to the suspect transformation [2], a multiplayer concurrent game with mean payoff objectives can be transformed into a multidimensional mean payoff turn-based game. The existence of a Nash equilibria in the original game is then equivalent to the existence of a run which has a value which can be ensured by the first player of the game in the suspect game.

We also define a generalization of this problem that considers a system of equation instead of a single one.

Definition 10 (Threshold problem). Given a mean payoff game \mathcal{G} , a system of linear equation $\lambda = ((a_1, b_1), \dots, (a_l, b_l))$ over elements of $\mathbb{R}^{2 \cdot d}$, is there a strategy σ_{\exists} and a value $v \in \bigcap_{k=1}^l \mathbf{hsp}(a_k, b_k)$ such that σ_{\exists} ensures v ?

Remark 2. Other works ([10] for instance) usually focus on the 0-threshold problem, which can be specified here by $a_1 = (1, 0, \dots, 0)$, $a_2 = (0, 1, 0, \dots, 0)$, \dots and all b_k are equal to 0. In such a case, it is only needed to check that the value $\mathbf{0}$ can be ensured. This is indeed simpler than having to find the values that can be ensured.

1.4 Decomposition in Simple Cycles

Our analysis of mean payoff game will rely on finding good cycles for **Eve**. Our proof will use as a tool the decomposition of a play into *simple cycles*.

Definition 11 (Simple cycles). A simple cycle is sequence $s_0 \cdot s_1 \cdots s_n$, such that $s_0 = s_n$ and for all i and j , $0 \leq i < j < n$, $s_i \neq s_j$. We write $\mathbb{C}(\mathcal{G})$ (\mathbb{C} when \mathcal{G} is clear from the context) for the set of simple cycles in the concurrent game structure \mathcal{G} .

In this paragraph, we recall the notion of decomposition of a play into *simple cycles* [6]. A history $h = s_0 \cdot s_1 \cdots s_n$ is a *cycle* if $s_0 = s_n, n \geq 1$. A *simple cycle* is a cycle such that for all i and $j, 0 \leq i < j < n, s_i \neq s_j$. We write $\mathbb{C}(\mathcal{A})$ (\mathbb{C} when \mathcal{A} is clear from the context) for the set of simple cycles in the concurrent game \mathcal{A} .

Every history h of a finite game can be uniquely decomposed into a sequence of simple cycles, except for a finite part. The decomposition process maintains a *stack*, $\mathbf{st}(h)$, of distinct states and moves. We write the stack content $s_1 \cdot s_2 \cdots s_n$ where s_1 is at the bottom of the stack and s_n the top. We use the notation $s \in \mathbf{st}(h)$ for $s \in \{s_1, s_2, \dots, s_n\}$. The *decomposition*, $\mathbf{dec}(h)$, is a set of simple cycles. We define $\mathbf{dec}(h)$ and $\mathbf{st}(h)$ inductively as follows:

- for the single state history s , $\mathbf{dec}(s) = \emptyset$ and $\mathbf{st}(s) = s$.
- let $h' = h \cdot s$ be a history.
 - If $s \in \mathbf{st}(h)$, and $\mathbf{st}(h) = \alpha \cdot s \cdot \beta$, then $\mathbf{st}(h') = \alpha \cdot s$ and $\mathbf{dec}(h') = \mathbf{dec}(h) \cup \{s \cdot \beta \cdot s\}$.
 - else $\mathbf{dec}(h') = \mathbf{dec}(h)$, $\mathbf{st}(h') = \mathbf{st}(h) \cdot s$.

Note that the stack always contains distinct elements, therefore only simple cycles are added to the decomposition. The elements in the stack from the bottom to the top, form a history $s_0 \cdot s_1 \cdots s_n$, where $n+1$ is the height of the stack. The decomposition of a play is the union of the decompositions of the finite prefixes of the play.

2 Case of limit inferior

We first analyze the case where all the mean-payoff values would be taken as limit inferior of the average cost. We define value by:

$$\mathbf{value}(s) = \{v \in \mathbb{R}^d \mid \exists \sigma_{\exists}. \forall \rho \in \text{Out}(s, \sigma_{\exists}). \forall i \in \llbracket 1, d \rrbracket. \underline{\text{MP}}_i(\rho) \geq v_i\}$$

2.1 Downward closure

Definition 12. Let $\downarrow S$ be the downward closure of S , i.e. $\{x \mid \exists x' \in S. \forall i \in \llbracket 1, n \rrbracket. x_i \leq x'_i\}$.

Lemma 1. – Operator \downarrow is monotonic, i.e. for any set X and Y such that $X \subseteq Y$, we have that $\downarrow X \subseteq \downarrow Y$.

- Operator \downarrow is compatible with union, i.e. for any set X and Y : $(\downarrow X) \cup (\downarrow Y) = \downarrow (X \cup Y)$.

Proof. Assume $X \subseteq Y$. Let $x \in \downarrow X$, there is $x' \in X$ such that for all $i, x'_i \geq x_i$. As $X \subseteq Y, x' \in Y$, and therefore $x \in \downarrow Y$.

To show $(\downarrow X) \cup (\downarrow Y) \subseteq \downarrow (X \cup Y)$ We use the monotonicity property (Lem. 1): since $X \subseteq X \cup Y, \downarrow X \subseteq \downarrow (X \cup Y)$.

In the other direction, let $x \in \downarrow (X \cup Y)$, there exists $x' \in X \cup Y$ such that $x \leq x'$. If $x' \in X$ then $x \in \downarrow X$ and otherwise $x \in \downarrow Y$.

2.2 Characterizing values

We define memoryless strategies. We will use them to characterize values.

Definition 13 (Memoryless strategies). *A strategy σ_{\forall} is said memoryless if for every history h and h' , and all state s , $\sigma_{\forall}(h \cdot s) = \sigma_{\forall}(h' \cdot s)$. We write $\mathbb{M} = \{\sigma_{\forall}^1, \dots, \sigma_{\forall}^{|\mathbb{M}|}\}$ for the set of the memoryless strategies of Adam in \mathcal{G} .*

Let σ_{\forall} be a memoryless strategy. Strategy σ_{\forall} determines a subgraph of \mathcal{G} , let $\text{SSC}(\sigma_{\forall})$ be the set of *strongly connected component* in this subgraph. We write:

$$\mathbb{C}(s, \sigma_{\forall}) = \{C \mid \exists S \in \text{SSC}(\sigma_{\forall}). C = \{c \mid c \text{ simple cycle of } S\} \wedge S \text{ reachable from } s\}.$$

Definition 14 (Convex hull). *The convex hull of a set of points V is the set*

$$\text{conv}(V) = \left\{ \sum_{v \in V} t_v \cdot v \mid \forall v \in V. t_v \in [0, 1] \wedge \sum_{v \in V} t_v = 1 \right\}.$$

We now give a characterization of the set of values that Eve is able to ensure.

Theorem 1.

$$\underline{\text{value}}(s) = \bigcap_{\sigma_{\forall} \in \mathbb{M}} \bigcup_{C \in \mathbb{C}(s, \sigma_{\forall})} \downarrow \text{conv} \left(\left\{ \frac{1}{|c|} \cdot w(c) \mid c \in C \right\} \right)$$

We start by recalling two lemmas of [10] which we will use in our proof.

Lemma 2 ([10, Lem. 10(2)]). *Let S be a strongly connected graph. If S does not have a non-negative multi-cycle (i.e. $C \in \mathbb{C}(S)$ and $\lambda \in \mathbb{N}^C$ such that $\sum_{c \in C} \lambda(c) \cdot w(c) \geq 0$), then there exists a constant $m_G \in \mathbb{N}$ and a real $c_G > 0$ such that for all history h in the graph S we have $\min\{w_i(h) \mid i \in I\} \leq m_G - c_G \cdot |h|$.*

Lemma 3 ([10, Lem. 11]). *If there is a strongly connected component S reachable from s_0 that has a non-negative multi-cycle, then Eve has a strategy to satisfy the mean-payoff-inf (i.e. $\forall i \in I \cup J. \forall \rho \in \text{Out}_{s_0}(\sigma_{\exists}). \underline{\text{MP}}_i(\rho) \geq 0$).*

Lemma 4. *Let ρ be a path. If $\text{Inf}(\rho) \subseteq C$ for some $C \in \mathbb{C}(\sigma_{\forall})$, then $\underline{\text{MP}}(\rho) \in \text{conv} \left(\left\{ \frac{1}{|c|} \cdot w(c) \mid c \in C \right\} \right)$.*

Proof. We consider the lexicographic order \leq_{lex} defined by $v \leq_{\text{lex}} v' \Leftrightarrow v = v' \vee \exists i \leq d. v_i < v'_i \wedge \forall j < i. v_j = v'_j$. We also consider for each $i \in \llbracket 1, d \rrbracket$, the preorder \leq_i defined by $v \leq_i v' \Leftrightarrow v_i \leq v'_i$. We define $L(n) = \min_{\leq_{\text{lex}}} \left\{ \frac{1}{m} \cdot w(\rho_{\leq m}) \mid m \geq n \right\}$ and $K_i(n) = \min_{\leq_i} \left\{ \frac{1}{m} \cdot w(\rho_{\leq m}) \mid m \geq n \right\}$. We write $\text{MPL}(\rho) = \lim_{n \rightarrow \infty} L(n)$, this limit exists because $(\mathbb{R}^d, \leq_{\text{lex}})$ forms a complete lattice.

We show that for all coordinate i , $\text{MPL}(\rho)_i \geq \underline{\text{MP}}_i(\rho)$. We write 1_i for the vector in which all coordinates are equal to 0 except the i -th coordinate which is equal to 1. Let $\varepsilon > 0$ and consider the vector $L(n) + \varepsilon \cdot 1_i$. Since $L(n) <_{\text{lex}} L(n) + \varepsilon \cdot 1_i$, there exists $m \geq n$ such that $\frac{1}{m} \cdot w(\rho_{\leq m}) \leq_{\text{lex}} L(n) + \varepsilon \cdot 1_i$. We have that

$\frac{w_i(\rho_{\leq m})}{m} \leq L(n)_i + \varepsilon$ because for all coordinate $j < i$, $\frac{w_j(\rho_{\leq m})}{m}$ has to be equal to $L(n)_j$. We also have that $\frac{1}{m} \cdot w(\rho_{\leq m}) \geq_i K_i(n)$. Hence for all $\varepsilon > 0$, $K_i(n) < L(n)_i + \varepsilon$, which shows that $\text{MPL}(\rho)_i \geq \text{MP}_i(\rho)$.

We now show that $\text{MPL}(\rho) \in \text{conv} \left(\left\{ \frac{1}{|c|} \cdot w(c) \mid c \in C \right\} \right)$. After a finite prefix $\rho_{\leq m}$, the play stays in the component C . We consider the decomposition in simple cycle of $\rho_{\geq m}$, for $n > m$:

$$w(\rho_{\leq n}) = w(\rho_{\leq m}) + w(\text{st}(\rho_{[[m, n]]})) + \sum_{c \in \text{dec}(\rho_{[[m, n]])} w(c)$$

and for all $c \in \text{dec}(\rho_{[[m, n]])}$, c belongs to C . Since the length of the stack is bounded by the number of states, we have that:

$$w(\rho_{\leq m}) - W \cdot |V| + \sum_{c \in \text{dec}(\rho_{[[m, n]])} w(c) \leq w(\rho_{\leq n}) \leq w(\rho_{\leq m}) + W \cdot |V| + \sum_{c \in \text{dec}(\rho_{[[m, n]])} w(c)$$

where W contains on each coordinate the maximum of absolute value of weights in this coordinate.

$$\frac{w(\rho_{\leq m})}{n} - \frac{W \cdot |V|}{n} + \sum_{c \in \text{dec}(\rho_{[[m, n]])} \frac{1}{n} \cdot w(c) \leq \frac{w(\rho_{\leq n})}{n} \leq \frac{w(\rho_{\leq m})}{n} + \frac{W \cdot |V|}{n} + \sum_{c \in \text{dec}(\rho_{[[m, n]])} \frac{1}{n} \cdot w(c)$$

$$\begin{aligned} \sum_{c \in \text{dec}(\rho_{[[m, n]])} \frac{1}{n} \cdot w(c) &= \sum_{c \in \text{dec}(\rho_{[[m, n]])} \frac{|c|}{n} \cdot \frac{1}{|c|} \cdot w(c) \\ &= \frac{\sum_{c \in \text{dec}(\rho_{[[m, n]])} |c|}{n} \cdot \sum_{c \in \text{dec}(\rho_{[[m, n]])} \frac{|c|}{\sum_{c \in \text{dec}(\rho_{[[m, n]])} |c|} \cdot \frac{1}{|c|} \cdot w(c) \end{aligned}$$

We write v_n for $\sum_{c \in \text{dec}(\rho_{[[m, n]])} \frac{|c|}{\sum_{c \in \text{dec}(\rho_{[[m, n]])} |c|} \cdot \frac{1}{|c|} \cdot w(c)$, we have that $v_n \in \text{conv} \left(\left\{ \frac{1}{|c|} \cdot w(c) \mid c \in C \right\} \right)$. We will show that $\frac{1}{n} \cdot w(\rho_n)$ comes closer and closer to v_n .

$$\frac{w(\rho_{\leq m})}{n} - \frac{W \cdot |V|}{n} + \frac{\sum_{c \in \text{dec}(\rho_{[[m, n]])} |c|}{n} \cdot v_n \leq \frac{w(\rho_{\leq n})}{n} \leq \frac{1}{n} \cdot w(\rho_{\leq m}) + \frac{W \cdot |V|}{n} + \frac{\sum_{c \in \text{dec}(\rho_{[[m, n]])} |c|}{n} \cdot v_n$$

Let i be a coordinate:

$$\begin{aligned} \left| \frac{1}{n} \cdot w_i(\rho_n) - v_n \right| &\leq \frac{w_i(\rho_{\leq m})}{n} + \frac{W_i \cdot |V|}{n} + \frac{n - \sum_{c \in \text{dec}(\rho_{[[m, n]])} |c|}{n} \cdot |(v_n)_i| \\ &\leq \frac{m \cdot W_i}{n} + \frac{W_i \cdot |V|}{n} + \frac{m + |V|}{n} \cdot |W_i| \\ &\leq \frac{2 \cdot (m + |V|) \cdot W_i}{n} \end{aligned}$$

We show that there is a strictly increasing sequence u_n such that $\text{MP}(\rho_{u_n})$ converges to $\text{MPL}(\rho)$. We define u_0 by 0 then give u_n we set u_{n+1} to be the smallest $m \geq u_n + 1$ such that $L(u_n + 1) = \frac{1}{m} \cdot w(\rho_{\leq m})$. The sequence u_n is strictly increasing and $L(n) = \frac{w(\rho_{\leq u_n})}{n} = \text{MP}(\rho_{\leq u_n})$ hence $\text{MP}(\rho_{u_n})$ converges to $\text{MPL}(\rho)$.

Since $\text{MP}(\rho_{u_n})$ converges and is $\frac{2 \cdot (m+|V|) \cdot W_i}{n}$ close to v_n , and v_n belongs to $\text{conv} \left(\left\{ \frac{1}{|c|} \cdot w(c) \mid c \in C \right\} \right)$ which is a closed set, the limit of $\text{MP}(\rho_{u_n})$ is within $\text{conv} \left(\left\{ \frac{1}{|c|} \cdot w(c) \mid c \in C \right\} \right)$ and so is $\text{MPL}(\rho)$. \square

Proof (of Thm. 1). \square Let $v \in \text{value}(s)$, then there is a strategy σ_{\exists} of Eve, such that for all memoryless strategy $\sigma_{\forall} \in \mathbb{M}$ of Adam, $\forall i \in I \cup J$. $\text{MP}_i(\text{Out}(\sigma_{\exists}, \sigma_{\forall})) \geq v_i$. Let $\rho = \text{Out}(\sigma_{\exists}, \sigma_{\forall})$. We have that $\text{Inf}(\rho) \subseteq C$ for some $C \in \mathbb{C}(\sigma_{\forall})$. Lem. 4 shows that $\text{MP}(\rho) \in \downarrow \text{conv} \left(\left\{ \frac{1}{|c|} \cdot w(c) \mid c \in C \right\} \right)$ which implies that $v \in \downarrow \text{conv} \left(\left\{ \frac{1}{|c|} \cdot w(c) \mid c \in C \right\} \right)$.

As $\text{MPL}(\rho) \geq \text{MP}(\rho)$, we have that $\text{MP}(\rho) \in \downarrow \text{conv} \left(\left\{ \frac{1}{|c|} \cdot w(c) \mid c \in C \right\} \right)$. Moreover, since $\text{MP}(\text{Out}(\sigma_{\exists}, \sigma_{\forall})) \geq v$, it follows that $v \in \downarrow \text{conv} \left(\left\{ \frac{1}{|c|} \cdot w(c) \mid c \in C \right\} \right)$.

\square Let $v \in \bigcap_{\sigma_{\forall} \in \mathbb{M}} \bigcup_{C \in \mathbb{C}(\sigma_{\forall})} \downarrow \text{conv} \left(\left\{ \frac{1}{|c|} \cdot w(c) \mid c \in C \right\} \right)$. We fix a memoryless strategy σ_{\forall} of Adam. We have that v is in $\downarrow \text{conv} \left(\left\{ \frac{1}{|c|} \cdot w(c) \mid c \in C \right\} \right)$ for some $C \in \mathbb{C}(\sigma_{\forall})$.

We replace the weights of the game by w' such that $w'_i(s) = w_i(s) - v_i$. We have for all path ρ that if the weight $w(\rho)$ is greater than v then $w'(\rho) \geq 0$. We know there exists some $v' \geq v$ of the form $\sum_{k=1}^{|C|} \frac{\lambda_k}{|c_k|} \cdot w(c_k)$. Hence there exists $\lambda_1, \dots, \lambda_{|C|}$ such that $\sum_{k=1}^{|C|} \frac{\lambda_k}{|c_k|} \cdot w'(c_k) \geq 0$.

The idea is to construct for all n a finite path of length greater than n that has a total weight by w' close to 0. Let n be an integer, there exist $p_k \in \mathbb{N}$ such that $\left| \frac{p_k}{n} - \lambda_k \right| \leq \frac{1}{n}$. Let $l_k = \prod_{k' \neq k} |c_{k'}|$, we construct the history h that follows $p_1 \cdot l_1$ times the cycle c_1 , then goes to the beginning of cycle c_2 with a path of length at most $|\text{Stat}|$ then follows $p_2 \cdot l_2$ times the cycle c_2 and so on until the $|C|$ -th cycle.

First, lets look at the length of this path.

$$\begin{aligned} \sum_{k=1}^{|C|} \frac{p_k}{n} &\geq \sum_{k=1}^{|C|} \lambda_k - \frac{|C|}{n} \\ \sum_{k=1}^{|C|} p_k &\geq n \sum_{k=1}^{|C|} \lambda_k - |C| = n - |C| \end{aligned}$$

Hence $|h| \geq n - |C|$, hence it is not bounded when n grows toward infinity.

Let W' be the maximum weight that appears in w' . On dimension i , the total weight of h is such that:

$$\begin{aligned} & \sum_{k=1}^{|C|} p_k \cdot l_k \cdot w'_i(c_k) - |\text{Stat}| \cdot |C| \cdot |W'| \leq w'_i(h) \\ & \prod_{k=1}^{|C|} |c_k| \cdot \sum_{k=1}^{|C|} p_k \cdot \frac{w'_i(c_k)}{|c_k|} - |\text{Stat}| \cdot |C| \cdot |W'| \leq w'_i(h) \\ & \prod_{k=1}^{|C|} |c_k| \cdot \sum_{k=1}^{|C|} \left(\lambda_k - \frac{1}{n} \right) \cdot \frac{w'_i(c_k)}{|c_k|} - |\text{Stat}| \cdot |C| \cdot |W'| \leq w'_i(h) \\ & - \frac{1}{n} \cdot \prod_{k=1}^{|C|} |c_k| \cdot \sum_{k=1}^{|C|} \frac{w'_i(c_k)}{|c_k|} - |\text{Stat}| \cdot |C| \cdot |W'| \leq w'_i(h) \end{aligned}$$

Hence $w'_i(h)$ is bounded on all dimension i when we make n grow.

By Lem. 10(2) of [10]¹ this implies there is a non negative cycle in C for the weight w' . By Lem. 11 of [10], σ_\forall is not a winning strategy. We know thanks to Theorem 8 of [10] that multi-mean payoff games are determined under memoryless strategies of **Adam**. Therefore **Adam** has no winning strategy. Since these games are also determined [7], this means that **Eve** has a winning strategy. Therefore $v \in \underline{\text{value}}(s)$.

2.3 Half-spaces representation

The goal of this section is to show that the equation: $\underline{\text{value}}(s) \cap \text{hsp}(\lambda) \neq \emptyset$ can be solved using a relatively simple half-space representation (it might still be exponential compared with the size of the arena).

The set $\text{conv} \left(\left\{ \frac{1}{|c|} \cdot w(c) \mid c \in C \right\} \right)$ is a convex hull and it is well known these can be written as finite intersection of half-spaces [8].

Definition 15 (Polyhedron). *The boundary of $\text{hsp}(a, b)$ is the hyperplane: $\text{bndr}(a, b) = \{x \in \mathbb{R}^d \mid \sum_{i \in [1, d]} a_i \cdot x_i = b\}$. If $P = \bigcap_{i=1}^k H_i$ where for all i , H_i is a half-space, P is called a polyhedron. A bounded polyhedron is called a polytope. A face F of P is a subset of P of the form $F = P \cap \text{bndr}(H_F)$, where H_F is a half-space such that $P \subseteq H_F$. In that case, say that H_F defines face F of P . A face of dimension 1 is called a vertex. If P has dimension d' , then a face of dimension $d' - 1$ is called a facet. A complete set of facet defining half-spaces \mathcal{F} contains for each facet F a half-space H_F such that $P \cap \text{bndr}(H) = F$ and $P \subseteq H$.*

¹ Note that in [10] the weights are integer, which may not be the case for us. However the proof of Lem. 10(2) still works in the case of arbitrary reals.

Definition 16 (Affine Hull). A affine subspace is a set of the form $x + L$ with $x \in \mathbb{R}^d$ and L is a linear subspace of \mathbb{R}^d (i.e. a subset closed under addition of vectors and under multiplication by real numbers). The affine hull of a set $X \subseteq \mathbb{R}^d$ is the intersection of all affine subspaces of \mathbb{R}^d containing X , it is written $\mathbf{aff}(X)$. It can be described as the set of linear combinations of points of X [8].

We prove a proposition, similar in spirit to Helly's theorem (see [8]), that we will use to restrict the search to certain points.

Lemma 5. Let H_1, \dots, H_n be n half-spaces of \mathbb{R}^d with $n \geq d \geq 1$, and S be an affine subspace of \mathbb{R}^d of dimension k , with $1 \leq k \leq d$. If $S \cap \bigcap_{i=1}^n H_i \neq \emptyset$ then there is i_1, \dots, i_k such that:

$$S \cap \bigcap_{j=1}^k \mathbf{bndr}(H_{i_j}) \subseteq S \cap \bigcap_{i=1}^n H_i.$$

Notice in particular that this lemma applies to the case where $S = \mathbb{R}^d$.

Proof. We show the proposition by induction over k . If $k = 0$, then if $S \cap \bigcap_{i=1}^n H_i \neq \emptyset$ then every H_i contain S since it is reduced to one point. Therefore $S \cap \bigcap_{i=1}^{k+1} H_i = S = S \cap \bigcap_{i=1}^n H_i$.

Assume now $k > 0$ and that the property holds for all affine subspace of dimension $k' < k$. We proceed by induction over n . If $n = k$, then since $\mathbf{bndr}(H_i) \subseteq H_i$ for all i , we have that $\bigcap_{i=1}^k \mathbf{bndr}(H_i) \subseteq \bigcap_{i=1}^k H_i$ which shows the property.

Assume now that the property holds for n , we consider a half-space H_{n+1} .

– If $S \subseteq \mathbf{bndr}(H_{n+1})$ then we use the induction hypothesis for n , we have

$$\begin{aligned} \bigcap_{j=1}^k (\mathbf{bndr}(H_{i_j}) \cap S) &\subseteq S \cap \bigcap_{i=1}^n H_i \\ &\subseteq S \cap \mathbf{bndr}(H_{n+1}) \cap \bigcap_{i=1}^n H_i \\ &\subseteq S \cap H_{n+1} \cap \bigcap_{i=1}^n H_i \\ &\subseteq S \cap \bigcap_{i=1}^{n+1} H_i \end{aligned}$$

– Otherwise $\mathbf{bndr}(H_{n+1}) \cap S$ forms a affine subspace of dimension $k - 1$, to which we can apply the induction hypothesis.

- If $S \cap \text{bndr}(H_{n+1}) \cap \bigcap_{i=1}^n H_i \neq \emptyset$ then there is i_1, \dots, i_{k-1} such that:

$$\begin{aligned} (\text{bndr}(H_{n+1}) \cap S) \cap \bigcap_{j=1}^{k-1} \text{bndr}(H_{i_j}) &\subseteq S \cap \text{bndr}(H_{n+1}) \cap \bigcap_{i=1}^n H_i \\ S \cap \bigcap_{j=1}^{k-1} \text{bndr}(H_{i_j}) \cap \text{bndr}(H_{n+1}) &\subseteq S \cap \bigcap_{i=1}^{n+1} H_i \end{aligned}$$

Which shows the property.

- Otherwise $(\bigcap_{i=1}^n H_i \cap S) \cap \text{bndr}(H_{n+1}) = \emptyset$, then since $\bigcap_{i=1}^n H_i \cap S$ is convex it is either totally inside H_{n+1} or totally outside of it. If it is outside then the intersection is empty. Otherwise, $\bigcap_{i=1}^n H_i \cap S \subseteq H_{n+1}$. We apply the induction hypothesis over n : there is i_1, \dots, i_k such that:

$$\bigcap_{j=1}^k \text{bndr}(H_{i_j}) \cap S \subseteq \bigcap_{i=1}^n H_i \cap S \subseteq S \cap \bigcap_{i=1}^{n+1} H_i$$

Which proves the result.

Theorem 2 ([5, Thm. 3.(b)]). *Let P be a polytope. Let \mathcal{F} be complete set of facet defining half-spaces, then P is the intersection of $\text{aff}(P)$ with all half-spaces in \mathcal{F} .*

The set $\text{conv}\left(\left\{\frac{1}{|c|} \cdot w(c) \mid c \in C\right\}\right)$ is a convex hull and can be written as finite intersection of half-spaces [8] but the number of half-spaces that is required can be exponential with respect to the dimension d : it is of the order of n^d where n is the number of points given as input. In the following, we will describe how these half-spaces are obtained.

Definition 17. *We write $\pi_j(x)$ for the projection of x over the plane $\{x \mid x_j = 0\}$, that is $\pi_j(x) = x'$ such that for all $i \neq j$, $x'_i = x_i$ and $x'_j = 0$. We also write $\pi_J(x)$ for the projection of x over the plane $\{x \mid \forall j \in J. x_j = 0\}$. We write $\downarrow_i X$ for $\{v \mid \exists x \in X, \lambda \geq 0. v = x - \lambda \cdot \mathbf{1}_i\}$. Let also $\uparrow_K S$ be the set $\{x \mid \exists x' \in S. \forall j \in [1, n] \setminus K. x'_j = x_j\}$.*

Lemma 6. *Let P be a polyhedron, and $\mathcal{F}(\pi_j(P))$ be a set of half-spaces of $\text{aff}(\pi_j(P))$ such that $\pi_j(P) = \bigcap_{H \in \mathcal{F}(\pi_j(P))} H$.*

$$\uparrow_j P = \bigcap_{H \in \mathcal{F}(\pi_j(P))} H + \mathbb{R} \cdot \mathbf{1}_j$$

Proof.

$$\begin{aligned}
x \in \downarrow_j P &\Leftrightarrow \exists x' \in P. \pi_j(x) = \pi_j(x') \\
&\Leftrightarrow \pi_j(x) \in \bigcap_{H \in \mathcal{F}(\pi_j(P))} H \\
&\Leftrightarrow \pi_j(x) + \mathbb{R} \cdot \mathbf{1}_j \subseteq \bigcap_{H \in \mathcal{F}(\pi_j(P))} H + \mathbb{R} \cdot \mathbf{1}_j \\
&\Leftrightarrow x \in \bigcap_{H \in \mathcal{F}(\pi_j(P))} H + \mathbb{R} \cdot \mathbf{1}_j
\end{aligned}$$

□

Lemma 7. *Let P be a polyhedron and $\mathcal{F}(P)$ be set of half-spaces such that $P = \bigcap_{H \in \mathcal{F}(P)} H$. Let $\mathcal{F}_-^j(P) \subseteq \mathcal{F}(P)$ be the set of half-spaces of $\mathcal{F}(P)$ that contain the direction $-\mathbf{1}_j$, i.e. $H - \mathbb{R}^+ \cdot \mathbf{1}_j = H$.*

$$\downarrow_j P = \downarrow_j P \cap \bigcap_{H \in \mathcal{F}_-^j(P)} H$$

Proof. $\boxed{\subseteq}$ Let $x \in \downarrow_j P$. We have that $x \in \downarrow_j P$. There is $x' \in P$ such that $x = x' - \lambda \cdot \mathbf{1}_j$ with $\lambda \geq 0$. Let $f \in \mathcal{F}_-^j$, since $x' \in H_f$ we have that $x \in H_f - \mathbb{R}^+ \cdot \mathbf{1}_j = H_f$. Therefore $x \in \downarrow_j P \cap \bigcap_{f \in \mathcal{F}_-^j} H_f$.

$\boxed{\supseteq}$ Let $x \in \downarrow_j P \cap \bigcap_{f \in \mathcal{F}_-^j} H_f$. There exists $x' \in P$ such that $\pi_j(x) = \pi_j(x')$. Consider the line segment $[x, x']$, it does not intersect any $\text{bndr}(H_f)$ with $f \in \mathcal{F}_-^j$ since both x and x' are inside these half-spaces.

If $[x, x']$ does not intersect any $\text{bndr}(H_f)$ where f is a facet of P then x is in the intersection of these half-spaces, and using Thm. 2, this means that $x \in P$.

Otherwise $[x, x']$ intersects some $\text{bndr}(H_f)$ with f facet of P which is not in \mathcal{F}_-^j . Since H_f contains x' but not x , and these two only differs by their j coordinates, this means that $x_j \leq x'_j$ and therefore $x \in \downarrow_j P$. □

We write B_J for the subspace $\sum_{j \in J} \mathbb{R} \cdot \mathbf{1}_j$ and C_J for the cone $\sum_{j \in J} \mathbb{R}_+ \cdot \mathbf{1}_j$. If \mathcal{F} is a set of half-spaces, we write $\mathcal{F}_-^J \subseteq \mathcal{F}$ the half-spaces of \mathcal{F} that contain the direction $-C_J$, i.e. $H - C_J = H$.

Lemma 8.

$$\downarrow_K P = \bigcap_{J \subseteq K} \bigcap_{H \in \mathcal{F}_-^{K \setminus J}(\pi_J(P))} H + B_J$$

with the convention that $\pi_\emptyset(P) = P$, $\mathcal{F}_-^\emptyset(P) = \mathcal{F}(P)$ and $B_\emptyset = \mathbf{0}$.

Proof. We prove the result by induction over the size of K . The case where $|K| = 1$ is a direct consequence of Lem. 6 and 7.

Assume the equality holds for some set K . Consider a dimension $j \notin K$.

$$\pi_j(\downarrow_K P) = \downarrow_K \pi_j(P) = \bigcap_{J \subseteq K} \bigcap_{H \in \mathcal{F}_-^{K \setminus J}(\pi_{J \cup \{j\}}(P))} H + B_J$$

So we can chose:

$$\mathcal{F}(\pi_j(\downarrow_K P)) = \bigcup_{J \subseteq K} \{H + B_J \mid H \in \mathcal{F}_-^{K \setminus J}(\pi_{J \cup \{j\}}(P))\} \quad (1)$$

We also have by induction hypothesis:

$$\begin{aligned} \mathcal{F}(\downarrow_K P) &= \bigcup_{J \subseteq K} \{H + B_J \mid H \in \mathcal{F}_-^{K \setminus J}(\pi_J(P))\} \\ \mathcal{F}_-^j(\downarrow_K P) &= \bigcup_{J \subseteq K} \{H + B_J \mid H \in \mathcal{F}_-^{K \cup \{j\} \setminus J}(\pi_J(P))\} \end{aligned} \quad (2)$$

Using Lem. 6 and 7:

$$\begin{aligned} \downarrow_{K \cup \{j\}} P &= \bigcap_{H \in \mathcal{F}(\pi_j(\downarrow_K P))} H + B_j \cap \bigcap_{H \in \mathcal{F}_-^j(\downarrow_K P)} H \\ &= \bigcap_{J \subseteq K} \bigcap_{H \in \mathcal{F}_-^{K \setminus J}(\pi_{J \cup \{j\}}(P))} H + B_J + B_j \cap \bigcap_{H \in \mathcal{F}_-^j(\downarrow_K P)} H \quad \text{By (1)} \\ &= \bigcap_{J \subseteq K} \bigcap_{H \in \mathcal{F}_-^{K \setminus J}(\pi_{J \cup \{j\}}(P))} H + B_{J \cup \{j\}} \cap \bigcap_{J \subseteq K} \bigcap_{H \in \mathcal{F}_-^{K \cup \{j\} \setminus J}(\pi_J(P))} H + B_J \quad \text{By (2)} \\ &= \bigcap_{J \subseteq K \cup \{j\}} \bigcap_{H \in \mathcal{F}_-^{K \cup \{j\} \setminus J}(\pi_J(P))} H + B_J \end{aligned}$$

Which proves the property for $K \cup \{j\}$ and concludes the induction. \square

Corollary 1.

$$\downarrow P = \bigcap_{J \subseteq \llbracket 1, d \rrbracket} \bigcap_{H \in \mathcal{F}_-^{\llbracket 1, d \rrbracket \setminus J}(\pi_J(P))} H + B_J$$

2.4 Algorithm

In order to solve the value problem, our algorithm is going to guess a vector of a certain form which satisfies the linear equation and check that it belong to value. For this last step, we reuse an algorithm from [10], which is **coNP** as we recall here.

Theorem 3. [10, Thm. 7.2] *For multi-weighted games with objective $\{\rho \mid \text{MP}(\rho) \geq (0, \dots, 0)\}$, the problem of deciding whether a given state is winning for *Eve* is **coNP**-complete.*

Corollary 2. *Given a vector $x \in \mathbb{R}^d$ and a state s , we can decide with a coNP algorithm whether $x \in \underline{\text{value}}(s)$.*

Proof. This is done by subtracting x from all the weights and using the preceding algorithm to decide if s is winning for the objective $\{\rho \mid \underline{\text{MP}}(\rho) \geq (0, \dots, 0)\}$.

$\underline{\text{value}}(s) \cap \text{hsp}(\lambda) \neq \emptyset$ if, and only if, there is a function $f: \mathbb{M} \mapsto \mathbb{C}$, such that $f(\sigma_\forall) \in \mathbb{C}(s, \sigma_\forall)$ for all strategy σ_\forall and

$$\text{hsp}(\lambda) \cap \bigcap_{\sigma_\forall \in \mathbb{M}} \downarrow \text{conv} \left(\left\{ \frac{1}{|c|} \cdot w(c) \mid c \in f(\sigma_\forall) \right\} \right) \neq \emptyset \quad (3)$$

By Corollary 1, $\downarrow \text{conv} \left(\left\{ \frac{1}{|c|} \cdot w(c) \mid c \in f(\sigma_\forall) \right\} \right)$ is the intersection of the half-spaces in $\mathcal{H}^{f(\sigma_\forall)}$, where for a set of simple cycles C :

$$\mathcal{H}^C = \left\{ H + B_J \mid J \subseteq [1, d] \wedge H \in \mathcal{F}_-^{[1, d] \setminus J}(\pi_J(P)) \wedge P = \text{conv} \left(\left\{ \frac{1}{|c|} \cdot w(c) \mid c \in C \right\} \right) \right\}$$

Lemma 9. *Let $\mathcal{H} = \{\text{hsp}(\lambda)\} \cup \bigcup_{\sigma_\forall \in \mathbb{M}} \mathcal{H}^{f(\sigma_\forall)}$. The three following propositions are equivalent:*

1. $\underline{\text{value}}(s) \cap \text{hsp}(\lambda) \neq \emptyset$;
2. there exists d half-spaces H_1, \dots, H_d in \mathcal{H} such that $\bigcap_{1 \leq i \leq d} \text{bndr}(H_i) \neq \emptyset$ and $\exists x \in \bigcap_{1 \leq i \leq d} \text{bndr}(H_i)$, $x \in \underline{\text{value}}(s) \cap \text{hsp}(\lambda)$.
3. there exists \bar{d} half-spaces $H_1, \dots, H_{\bar{d}}$ in \mathcal{H} such that $\bigcap_{1 \leq i \leq \bar{d}} \text{bndr}(H_i) \neq \emptyset$ and $\forall x \in \bigcap_{1 \leq i \leq \bar{d}} \text{bndr}(H_i)$, $x \in \underline{\text{value}}(s) \cap \text{hsp}(\lambda)$.

Proof. The implications (3) \implies (2) \implies (1) are obvious.

We now prove the implication (1) \implies (3). By Thm. 1

$$\begin{aligned} \underline{\text{value}}(s) &= \bigcap_{\sigma_\forall \in \mathbb{M}} \bigcup_{C \in \mathbb{C}(s, \sigma_\forall)} \downarrow \text{conv} \left(\left\{ \frac{1}{|c|} \cdot w(c) \mid c \in C \right\} \right) \\ &= \bigcup_{f: \mathbb{M} \mapsto \mathbb{C}} \bigcap_{\sigma_\forall \in \mathbb{M}} \downarrow \text{conv} \left(\left\{ \frac{1}{|c|} \cdot w(c) \mid c \in C \right\} \right) \end{aligned}$$

The intersection of $\text{hsp}(\lambda)$ with $\underline{\text{value}}(s)$ is non-empty if, and only if, there exists $f: \mathbb{M} \mapsto \mathbb{C}$, such that $\text{hsp}(\lambda) \cap \bigcap_{\sigma_\forall \in \mathbb{M}} \downarrow \text{conv} \left(\left\{ \frac{1}{|c|} \cdot w(c) \mid c \in f(\sigma_\forall) \right\} \right)$ is non empty. By Corollary 1, $\downarrow \text{conv} \left(\left\{ \frac{1}{|c|} \cdot w(c) \mid c \in f(\sigma_\forall) \right\} \right)$ is the intersection of the half-spaces in $\mathcal{H}^{f(\sigma_\forall)}$. So $\text{hsp}(\lambda) \cap \bigcap_{\sigma_\forall \in \mathbb{M}} \downarrow \text{conv} \left(\left\{ \frac{1}{|c|} \cdot w(c) \mid c \in f(\sigma_\forall) \right\} \right)$ is non empty if and only if $\bigcap_{H \in \mathcal{H}^{f(\sigma_\forall)}} H$ is non empty. By Lem. 5 this is the case if and only if there is d half-spaces H_1, \dots, H_d in $\mathcal{H}^{f(\sigma_\forall)}$ such that $\bigcap_{1 \leq i \leq d} \text{bndr}(H_i) \subseteq \text{hsp}(\lambda) \cap \bigcap_{\sigma_\forall \in \mathbb{M}} \downarrow \text{conv} \left(\left\{ \frac{1}{|c|} \cdot w(c) \mid c \in f(\sigma_\forall) \right\} \right)$. This implies $\bigcap_{1 \leq i \leq d} \text{bndr}(H_i) \subseteq \text{hsp}(\lambda) \cap \underline{\text{value}}(s)$ which shows that (3) holds. \square

Algorithm 1: Algorithm for the value problem in multi-mean-payoff

Existentially guess d half-space $H_1, \dots, H_d \in \mathcal{H}$
 if $\bigcap_{j=1}^d \text{bndr}(H_j) = \emptyset$ **then return false**;
 else take x one arbitrary point in $\bigcap_{j=1}^d \text{bndr}(H_j)$
 if $x \notin \text{hsp}(\lambda)$ **then return false**;
 if $x \in \text{value}(s)$ **then return true**

Based on this property we devise Algorithm 1 to decide the value problem. Its correctness holds from Lem. 9. Algorithm 1 is in $\Sigma_2\text{P} = \text{NP}^{\text{NP}} = \text{NP}^{\text{coNP}}$, since checking whether $x \in \text{value}(s)$ can be done in coNP (by Corollary 2) and computing the intersection of hyperplanes can be done in polynomial time [9, Thm. 3.3]. We obtain from this, the following theorem.

Theorem 4. *The threshold problem is in $\Sigma_2\text{P}$ for limit inferior objectives.*

2.5 Hardness

Proposition 1. *The threshold problem is NP-hard for limit inferior objectives.*

Proof. Let $\phi = \exists x_1. \exists x_2. \dots \exists x_n. C_1 \wedge \dots \wedge C_n$ be a formula, where $C_i = \ell_{i,1} \vee \ell_{i,2} \vee \ell_{i,3}$ with $\ell_{i,j} \in \{x_k, \neg x_k \mid k \in \llbracket 1, n \rrbracket\}$. We have two dimensions for each variable x_i , so $d = 2 \cdot n$. We write 0_i for the vector v such that $v_i = 0$ and $v_j = 1$ for $j \neq i$. We define a family of vectors $v_{i,j}$ where if $\ell_{i,j} = x_k$ then $v_{i,j} = 0_{2 \cdot k+1}$ and if $\ell_{i,j} = \neg x_k$ then $v_{i,j} = 0_{2 \cdot k}$. We have one initial state s_0 controlled by **Adam**, one state for each clause C_i controlled by **Eve**, and one state for each literal of the formula. The construction is illustrated in Figure 1.

We consider the constraints $\lambda_i = (v_{2 \cdot i} + v_{2 \cdot i+1} \geq 1)$.

In the example, valuation $x_1 \rightarrow \text{true}, x_2 \rightarrow \text{false}, x_3 \rightarrow \text{true}$ makes the formula valid, the corresponding value is $v = (1, 0, 0, 1, 1, 0)$. Consider the strategy of **Eve** that in C_1 chooses x_1 , in C_2 chooses x_1 and in C_3 chooses $\neg x_2$. In all the cases this strategy ensures to **Eve** a payoff better than v . In contrast, valuation $x_1 \rightarrow \text{false}, x_2 \rightarrow \text{false}, x_3 \rightarrow \text{true}$ does not make the formula valid, the corresponding value is $v' = (0, 1, 0, 1, 1, 0)$, and if **Adam** chooses C_1 then **Eve** cannot ensure v' .

Let ξ be a valuation, we associate to it a vector v_ξ such that if $\xi(x_i) = \text{true}$ then $(v_\xi)_{2 \cdot i} = 1$ and $(v_\xi)_{2 \cdot i+1} = 0$; if $\xi(x_i) = \text{false}$ then $(v_\xi)_{2 \cdot i} = 0$ and $(v_\xi)_{2 \cdot i+1} = 1$. We prove that if ξ makes ϕ valid then **Eve** can ensure v_ξ .

In the other direction, assume v_ξ makes ϕ valid. We consider a strategy for **Eve** that in each C_i chooses a state $\ell_{i,j}$ such that ξ makes $\ell_{i,j}$ true, this is possible because v_ξ makes ϕ valid. If $\ell_{i,j} = x_k$, then $\xi(x_k) = \text{true}$ and $(v_\xi)_{2 \cdot k+1} = 0$. The payoff is then $0_{2 \cdot k+1}$ which is greater than v_ξ . Similarly, if $\ell_{i,j} = \neg x_k$, the payoff $0_{2 \cdot k}$ is greater than v_ξ . This shows that **Eve** has a strategy to ensure v_ξ .

We can already conclude that if ϕ is satisfiable then **Eve** can ensure a value that satisfies the equations in λ .

Now, reciprocally, assume there is some v that satisfies λ and that can be ensured by **Eve**. For each $k \in \llbracket 1, n \rrbracket$, $v_{2 \cdot k} > 0$ or $v_{2 \cdot k+1} > 0$. We select a valuation




Figure 1. Example of the encoding of 3SAT into the threshold problem, for formula $\phi = (x_1 \vee x_2 \vee \neg x_3) \wedge (x_1 \vee \neg x_2 \vee \neg x_3) \wedge (\neg x_1 \vee \neg x_2 \vee x_3)$.

ξ_v such that $\xi_v(x_k) = \text{true} \Leftrightarrow v_{2,k} > 0$. We prove that if Eve ensures v then ξ_v makes ϕ valid.

Assume Eve ensures v . This means that from every C_i , she can find a $\ell_{i,j}$ such that $v_{i,j} \geq v$. If $\ell_{i,j} = x_k$, then $v_{i,j} = 0_{2,k+1}$, hence $(v_\xi)_{2,k+1} = 0$. Therefore $\xi_v(x_k) = \text{true}$. Similarly if $\ell_{i,j} = \neg x_k$, then $\xi_v(x_k) = \text{false}$. This means that in each C_i , we can find a literal made true by ξ_v and therefore ξ_v makes ϕ valid.

This proves that ϕ is satisfiable if, and only if, Eve can ensure a value that satisfies the equations in λ . \square

3 General case

3.1 Characterizing values

Proposition 2. *If $J \neq \emptyset$, then:*

$$\text{value}(s) = \left(\bigcap_{j \in J} \downarrow_{J \setminus \{j\}} \bigcap_{\sigma_\forall \in \mathbb{M}} \bigcup_{C \in \mathcal{C}(\sigma_\forall)} \downarrow \text{conv} \left(\left\{ \frac{1}{|c|} \cdot w(c) \mid c \in C \right\} \right) \right)$$

In order to prove this result, we will reuse some properties of multi-mean-payoff games that were proved in [10].

Lemma 10 ([10] Lem. 17).

$$\begin{aligned} & \exists \sigma_\exists. \forall \sigma_\forall. \forall i \in I. \text{MP}_i(\rho) \geq v_i \text{ and } \forall j \in J. \overline{\text{MP}}_j(\rho) \geq v_j \\ & \Leftrightarrow \forall j \in J. \exists \sigma_\exists. \forall \sigma_\forall. \forall i \in I. \text{MP}_i(\rho) \geq v_i \text{ and } \overline{\text{MP}}_j(\rho) \geq v_j. \end{aligned}$$

Lemma 11 ([10] **Lem. 14**). *Let $j \in J$.*

$$\begin{aligned} & \exists \sigma_{\exists}. \forall \sigma_{\forall}. \forall i \in I. \text{MP}_i(\rho) \geq v_i \text{ and } \overline{\text{MP}}_j(\rho) \geq v_j \\ \Leftrightarrow & \exists \sigma_{\exists}. \forall \sigma_{\forall}. \forall i \in I. \text{MP}_i(\rho) \geq v_i \text{ and } \text{MP}_j(\rho) \geq v_j. \end{aligned}$$

We now proceed to the proof of the proposition.

Proof (of Prop. 2). Let v be a vector of $\mathbb{R}^{2 \cdot d}$.

$$\begin{aligned} v \in \text{value}(s) & \Leftrightarrow \exists \sigma_{\exists}. \forall \sigma_{\forall}. \forall i \in I. \text{MP}_i(\rho) \geq v_i \text{ and } \forall j \in J. \overline{\text{MP}}_j(\rho) \geq v_j \\ & \Leftrightarrow \forall j \in J. \exists \sigma_{\exists}. \forall \sigma_{\forall}. \forall i \in I. \text{MP}_i(\rho) \geq v_i \text{ and } \overline{\text{MP}}_j(\rho) \geq v_j \text{ (by Lem. 10)} \\ & \Leftrightarrow \forall j \in J. \exists \sigma_{\exists}. \forall \sigma_{\forall}. \forall i \in I. \text{MP}_i(\rho) \geq v_i \text{ and } \text{MP}_j(\rho) \geq v_j \text{ (by Lem. 11)} \\ & \Leftrightarrow \forall j \in J. v \in \downarrow_{J \setminus \{j\}} \text{value}(s) \\ & \Leftrightarrow v \in \left(\bigcap_{j \in J} \downarrow_{J \setminus \{j\}} \text{value}(s) \right) \\ & \Leftrightarrow v \in \downarrow_{J \setminus \{j\}} \bigcap_{\sigma_{\forall} \in \mathbb{M}} \bigcup_{C \in \mathcal{C}(\sigma_{\forall})} \downarrow \text{conv} \left(\left\{ \frac{1}{|c|} \cdot w(c) \mid c \in C \right\} \right) \text{ (by Thm. 1)} \end{aligned}$$

3.2 Algorithm

In the following proposition, the goal is to express the value problem in terms of intersection of convex sets in a space of dimension $d = |I| + |J|$. The number of such convex sets is exponential but if the intersection is not empty, there is a witness whose coordinates are expressible as a small rational. This leads the way toward a non-deterministic polynomial algorithm.

Definition 18 (Sizes). *The size of a rational number $r = \frac{p}{q}$ where $p \in \mathbb{Z}, q \in \mathbb{Z}$, p and q are relatively prime, is: $\|r\| = 1 + \lceil \log_2(|p| + 1) \rceil + \lceil \log_2(q + 1) \rceil$. The size of a vector $v = (r_1, \dots, r_d)$ is $\|v\| = d + \sum_{i \in [1, d]} \|r_i\|$. The size of a matrix $A = (a_{ij})_{i=1, j=1}^d$ is $\|A\| = m \cdot n + \sum_{i, j \in [1, d] \times [1, n]} \|a_{ij}\|$.*

Theorem 5. *$\text{value}(s) \cap [-W, W]^d$ is of the form $\bigcup_{f \in F} \text{conv}(E_f)$ where F is finite and vertices of E_f have polynomially bounded size.*

Proposition 3 ([9, Corollary 3.2b]). *If the system $Ax = b$ of rational linear equations has a solution, it has one of size polynomially bounded by the sizes of A and b .*

Proposition 4. *Given n hyperplanes $H_1 = \text{hp}(a_1, b_1)$, dots, $H_n = \text{hp}(a_n, b_n)$, if there intersection is non empty we can find a point in the intersection expressed by $\frac{\alpha}{\beta}$ with α, β polynomially bounded with respect to the size of the equations $\sum_{j \in [1, n]} |b_j| + \sum_{i \in [1, d]} |a_{i, j}|$.*

Proof. Points at the intersection are the solutions of the system

$$\begin{pmatrix} a_1 \\ \vdots \\ a_n \end{pmatrix} X = \begin{pmatrix} b_1 \\ \vdots \\ b_n \end{pmatrix}.$$

Then Prop. 3 guarantees that we can find a solution with polynomially bounded size.

Proposition 5. *Let P be a polyhedron and \mathcal{F} be a complete set of facet defining half-spaces. If v is a vertex of P , then v is the intersection of at most $d + 1$ boundary of half-spaces in \mathcal{F} , i.e. there is $H_1, \dots, H_m \in \mathcal{F}$ with $m \leq d + 1$ such that $\{v\} = \cap_{j \in \llbracket 1, m \rrbracket} \mathbf{bndr}(H_j)$.*

Proof. A vertex is the intersection of the facets containing it [8, Chapter 5]. Let v be a vertex of P and let $F_v \subset \mathcal{F}$ be the set of facet defining half-spaces, corresponding to facets containing v . We have that $\cap_{H \in F_v} \mathbf{bndr}(H) = \{v\}$. We then use Prop. 5, to conclude that d half-spaces are enough.

Proposition 6. *Let V_1, \dots, V_n be set of points, and let $P = \cap_{i=1}^n \mathbf{conv}(V_i)$. Then $P = \mathbf{conv}(X)$ where X is a finite set of points whose size of coordinates are polynomial with respect to d and size coordinates of points of $\cup_i V_i$.*

Proof. By Prop. 5, vertices of P are at the intersection of at most $d+1$ boundaries of facet defining half-spaces of P . These half-spaces can be taken as a subset of facet defining half-spaces of $\{V_i \mid i \in \llbracket 1, n \rrbracket\}$.

We first show that equation describing these half-spaces are not too complicated. Facets of $\mathbf{conv}(V_i)$ are given by taking at most $d + 1$, vertices of $\mathbf{conv}(V_i)$. If H is given by vertices v_1, \dots, v_d of V_i , then $H = \{v_1 + \lambda_2 \cdot (v_2 - v_1) + \dots + \lambda_{d-J} \cdot (v_{d-J} - v_1) \mid \lambda_2, \dots, \lambda_d \in \mathbb{R}\}$. This is equivalently described by an equation

(l_1, l_2, \dots, l_d) where l_j is a solution of the equation $\begin{pmatrix} v_1 \\ \vdots \\ v_d \end{pmatrix} l_j^T = \mathbf{1}_j^T$. Note

that this is only valid when v_1, \dots, v_d are linearly independent. Using Prop. 3, coordinates in this equation are polynomial with respect to the coordinates of vertices v_1, \dots, v_d .

Now, vertices of P are at the intersection of at most d of these hyperplanes. Therefore, using Prop.3, there coordinates are polynomially bounded. \square

Proposition 7. $[-W, W]^d \cap \cap_{\sigma_{\forall} \in \mathbb{M}} \downarrow \mathbf{conv} \left(\left\{ \frac{1}{|c|} \cdot w(c) \mid c \in f(j, \sigma_{\forall}) \right\} \right)$ is the convex hull of a finite set $V(j, \sigma_{\forall})$ where all points of $V(j, \sigma_{\forall})$ have coordinate polynomial with respect to W , d and $|\mathbf{Stat}|$.

Proof. By Corollary 1:

$$\bigcap_{\sigma_{\forall} \in \mathbb{M}} \downarrow \mathbf{conv} \left(\left\{ \frac{1}{|c|} \cdot w(c) \mid c \in f(j, \sigma_{\forall}) \right\} \right) = \bigcap_{\sigma_{\forall} \in \mathbb{M}, H \in \mathcal{H}^f(\sigma_{\forall})} H$$

We also have $[-W, W]^d$ as the intersection of half-spaces in $\{F_i \mid i \in \llbracket 1, d \rrbracket\} \cup \{G_i \mid i \in \llbracket 1, d \rrbracket\}$ where $F_i = \{x \in \mathbb{R}^d \mid x_i \geq -W\}$ and $G_i = \{x \in \mathbb{R}^d \mid x_i \leq W\}$. As a consequence, the set $[-W, W]^d \cap \bigcap_{\sigma_V \in \mathbb{M}} \downarrow \mathbf{conv} \left(\left\{ \frac{1}{|c|} \cdot w(c) \mid c \in f(j, \sigma_V) \right\} \right)$ the intersection of half-spaces and it is bounded, thus it is a convex polytope. Therefore, it is the convex hull of its set of vertices V [8, Prop.5.3.2]. Moreover $\mathcal{F} = \mathcal{H} \cup \bigcup_{i \in \llbracket 1, d \rrbracket} F_i \cup \bigcup_{i \in \llbracket 1, d \rrbracket} G_i$ is a complete set of facet defining half-spaces where $\mathcal{H} = \bigcup_{\sigma_V \in \mathbb{M}} \mathcal{H}^{J(\sigma_V)}$. Vertices in V are at the intersection of at most $d + 1$ hyperplanes in \mathcal{F} by Prop. 5. Therefore by Prop. 3 their coordinates have size polynomially bounded by the size of W and coefficients appearing in the equations of half-spaces in \mathcal{H} .

The half-spaces of \mathcal{H} are of the form $H + B_J$ where $J \subseteq \llbracket 1, d \rrbracket$ and H belong to $\mathcal{F}(\pi_J(\mathbf{conv}(\frac{1}{|c|} \cdot w(c) \mid c \in f(j, \sigma_V))))$. Vertices of the projection of the convex hull are projection of vertices of the convex hull, so equation for these hyperplanes are given by $d - |J|$ vertices of the projection. If H is given by vertices v_1, \dots, v_{d-J} of the projection, then $H = \{v_1 + \lambda_2 \cdot (v_2 - v_1) + \dots + \lambda_{d-J} \cdot (v_{d-J} - v_1) \mid \lambda_2, \dots, \lambda_{d-J} \in \mathbb{R}\}$. This is equivalently described by an equation

(l_1, l_2, \dots, l_d) where l_i is a solution of the equation $\begin{pmatrix} v_1 \\ \vdots \\ v_{d-J} \end{pmatrix} l_i^T = \mathbf{1}_i^T$. Note that

this is only valid when v_1, \dots, v_{d-J} are linearly independent. Using Prop. 3, coordinates in this equation are polynomial with respect to the coordinates of vertices v_1, \dots, v_{d-J} which are themselves of the form $\frac{1}{|c|} \cdot w(c)$ for some $c \in \mathbb{C}$ or equal to 0.

We can therefore conclude that coordinates of vectors in V are polynomial with respect to W and $|\mathbf{Stat}|$. \square

Proposition 8. $[-W, W]^d \cap \uparrow_{J \setminus \{j\}} \bigcap_{\sigma_V \in \mathbb{M}} \downarrow \mathbf{conv} \left(\left\{ \frac{1}{|c|} \cdot w(c) \mid c \in f(j, \sigma_V) \right\} \right) = [-W, W]^d \cap \uparrow_{J \setminus \{j\}} [-W, W]^d \cap \bigcap_{\sigma_V \in \mathbb{M}} \downarrow \mathbf{conv} \left(\left\{ \frac{1}{|c|} \cdot w(c) \mid c \in f(j, \sigma_V) \right\} \right)$

Proof. Let $v \in [-W, W]^d \cap \uparrow_{J \setminus \{j\}} \bigcap_{\sigma_V \in \mathbb{M}} \downarrow \mathbf{conv} \left(\left\{ \frac{1}{|c|} \cdot w(c) \mid c \in f(j, \sigma_V) \right\} \right)$ then for all $j \in J$ there is some v^j such that $v^j \in \bigcap_{\sigma_V \in \mathbb{M}} \downarrow \mathbf{conv} \left(\left\{ \frac{1}{|c|} \cdot w(c) \mid c \in f(j, \sigma_V) \right\} \right)$ and for all coordinates $k \in J \setminus \{j\}$, $v_k^j = v_k$. We take a vector $v'_k = -W$ if $k \in J \setminus \{j\}$ and v_k otherwise. Obviously v' belongs to $[-W, W]^d$.

We show that v' also belongs to $\downarrow \mathbf{conv} \left(\left\{ \frac{1}{|c|} \cdot w(c) \mid c \in f(j, \sigma_V) \right\} \right)$. There is $x \geq v^j$ such that $x \in \mathbf{conv} \left(\left\{ \frac{1}{|c|} \cdot w(c) \mid c \in f(j, \sigma_V) \right\} \right)$. We have that for all coordinate k , $x_k \geq -W$. Hence, we have that $v' \leq x$ and therefore it belongs to $\downarrow \mathbf{conv} \left(\left\{ \frac{1}{|c|} \cdot w(c) \mid c \in f(j, \sigma_V) \right\} \right)$.

Hence $[-W, W]^d \cap \uparrow_{J \setminus \{j\}} [-W, W]^d \cap \bigcap_{\sigma_V \in \mathbb{M}} \downarrow \mathbf{conv} \left(\left\{ \frac{1}{|c|} \cdot w(c) \mid c \in f(j, \sigma_V) \right\} \right)$ contains v . The inclusion in the other direction being trivial, this shows the equality. \square

Proposition 9. *If $Y \subseteq [-W, W]^d$ then $[-W, W]^d \cap \downarrow_J \text{conv}(Y) = \text{conv}(Z)$ where $Z = \{z \mid \exists y \in Y. \forall i \notin J. z_i = y_i \wedge \forall i \in J. z_i \in \{-W, W\}\}$.*

Proof. \square Let $x \in [-W, W]^d \cap \downarrow_J \text{conv}(Y)$, there is $y \in \text{conv}(Y)$ such that for all $i \notin J$, $x_i = y_i$.

For $j \in J$, we define $\mu_j = \frac{x_j + W}{2 \cdot W}$, notice that $\mu_j \in [0, 1]$ since $x_j \in [-W, W]$. We then define for $v \in \{0, 1\}^{|J|}$ the coefficient $\lambda_v = \prod_{v_j=1} \mu_j \prod_{v_j=0} (1 - \mu_j)$. We have that for all v , $\lambda_v \in [0, 1]$ since this is the cause of all μ_j . Moreover $\sum_{v \in \{0, 1\}^{|J|}} \lambda_v = 1$, this is proved by a simple induction:

- if $|J| = 1$, then let $\{j\} = J$, $\sum_{v \in \{0, 1\}^{|J|}} \lambda_v = \mu_j + (1 - \mu_j) = 1$;
- if $|J| > 1$, then let $k \in J$:

$$\begin{aligned} \sum_{v \in \{0, 1\}^{|J|}} \lambda_v &= \mu_k \cdot \sum_{v|v_k=1} \frac{\lambda_v}{\mu_k} + (1 - \mu_k) \cdot \sum_{v|v_k=0} \frac{\lambda_v}{1 - \mu_k} \\ &= \mu_k \cdot \sum_{v|v_k=1} \prod_{v_j=1|j \neq k} \mu_j \prod_{v_j=0|j \neq k} (1 - \mu_j) \\ &\quad + (1 - \mu_k) \cdot \sum_{v|v_k=0} \prod_{v_j=1|j \neq k} \mu_j \prod_{v_j=0|j \neq k} (1 - \mu_j) \\ &= \mu_k + (1 - \mu_k) \quad (\text{By induction hypothesis}) \\ &= 1 \end{aligned}$$

We can also prove that $\sum_{v \in \{0, 1\}^{|J|}} \lambda_v \cdot v = (\mu_1, \dots, \mu_j)$: let $k \in J$:

$$\begin{aligned} \left(\sum_{v \in \{0, 1\}^{|J|}} \lambda_v \cdot v \right)_k &= \sum_{v \in \{0, 1\}^{|J|} | v_k=1} \lambda_v \\ &= \mu_k \cdot \sum_{v \in \{0, 1\}^{|J|} | v_k=1} \frac{\lambda_v}{\mu_k} \\ &= \mu_k \end{aligned}$$

Lets write $y = \sum_{j=1}^n \nu_j \cdot y^j$ with $\nu_j \in [0, 1]$, $\sum_{j=1}^n \nu_j = 1$, and $y^j \in Y$. Then given $v \in \{0, 1\}^{|J|}$, we write $z(j, v)$ for z such that $\forall i \notin J. z_i = y_i^j$ and $\forall i \in J. z_i \in -W$ if $v_i = 0$ and $z_i \in W$ if $v_i = 1$. We have that $\pi_J(z(j, v)) = \pi_J(y^j)$. Now, let $x' = \sum_{j \in J, v \in \{0, 1\}^{|J|}} \nu_j \cdot \lambda_v \cdot z(j, v)$.

- If $i \notin J$, then:

$$\begin{aligned} x'_i &= \sum_{j \in J, v \in \{0, 1\}^{|J|}} \nu_j \cdot \lambda_v \cdot y_i^j \\ &= \sum_{j \in J} \nu_j \cdot y_i^j = y_i = x_i. \end{aligned}$$

– If $i \in J$, then:

$$\begin{aligned}
x'_i &= \sum_{j \in J, v \in \{0,1\}^{|J|}} \nu_j \cdot \lambda_v \cdot z(j, v) \\
&= \sum_{j \in J} \nu_j \cdot \left(\sum_{v \in \{0,1\}^{|J|} | v_i=1} \lambda_v \cdot W - \sum_{v \in \{0,1\}^{|J|} | v_i=0} \lambda_v \cdot W \right) \\
&= \sum_{j \in J} \nu_j \cdot \left(-W + 2 \cdot W \cdot \left(\sum_{v \in \{0,1\}^{|J|} | v_i=1} \lambda_v \right) \right) \\
&= \sum_{j \in J} \nu_j \cdot (-W + 2 \cdot W \cdot \mu_i) \\
&= \sum_{j \in J} \nu_j \cdot (-W + x_i + W) \\
&= \sum_{j \in J} \nu_j \cdot x_i \\
&= x_i
\end{aligned}$$

Hence $x = x'$ which shows that x is in $\text{conv}(Z)$.

\square All points $z \in Z$ are in $[-W, W] \cap \downarrow J \setminus \{j\} \cap_{\sigma_v \in \mathbb{M}} \downarrow \text{conv} \left(\left\{ \frac{1}{|c|} \cdot w(c) \mid c \in f(j, \sigma_v) \right\} \right)$ therefore their convex hull also is. \square

Proof (of Thm. 5). By Prop. 2:

$$\text{value}(s) = \bigcap_{j \in J} \downarrow J \setminus \{j\} \bigcap_{\sigma_v \in \mathbb{M}} \bigcup_{C \in \mathbb{C}(\sigma_v)} \downarrow \text{conv} \left(\left\{ \frac{1}{|c|} \cdot w(c) \mid c \in C \right\} \right).$$

We can then reorder intersection and union operators by considering functions that selects for pairs (j, σ_v) a simple cycle among $\mathbb{C}(\sigma_v)$, i.e. functions of F , where $F = \{f: J \times \mathbb{M} \mapsto \mathbb{C} \mid \forall j \in J. \forall \sigma_v \in \mathbb{M}. f(j, \sigma_v) \in \mathbb{C}(\sigma_v)\}$.

We have:

$$\text{value}(s) = \bigcup_{f \in F} \bigcap_{j \in J} \downarrow J \setminus \{j\} \bigcap_{\sigma_v \in \mathbb{M}} \downarrow \text{conv} \left(\left\{ \frac{1}{|c|} \cdot w(c) \mid c \in f(j, \sigma_v) \right\} \right)$$

$$\begin{aligned}
\text{value}(s) \cap [-W, W]^d &= [-W, W]^d \cap \bigcup_{f \in F} \bigcap_{j \in J} \downarrow J \setminus \{j\} \bigcap_{\sigma_v \in \mathbb{M}} \downarrow \text{conv} \left(\left\{ \frac{1}{|c|} \cdot w(c) \mid c \in f(j, \sigma_v) \right\} \right) \\
&= \bigcup_{f \in F} \bigcap_{j \in J} [-W, W]^d \cap \downarrow J \setminus \{j\} \bigcap_{\sigma_v \in \mathbb{M}} \downarrow \text{conv} \left(\left\{ \frac{1}{|c|} \cdot w(c) \mid c \in f(j, \sigma_v) \right\} \right)
\end{aligned}$$

Using Prop. 8, this equals to:

$$\bigcup_{f \in F} \bigcap_{j \in J} [-W, W]^d \cap \downarrow_{J \setminus \{j\}} [-W, W]^d \cap \bigcap_{\sigma_v \in \mathbb{M}} \downarrow \text{conv} \left(\left\{ \frac{1}{|c|} \cdot w(c) \mid c \in f(j, \sigma_v) \right\} \right)$$

Using Prop. 7, this equals to

$$\bigcup_{f \in F} \bigcap_{j \in J} [-W, W]^d \cap \downarrow_{J \setminus \{j\}} \text{conv}(V(f, j))$$

Where all points of $\bigcup_{f \in F, j \in J} V(f, j)$ have their coordinates polynomially bounded.

Using Prop. 9, $[-W, W]^d \cap \downarrow_{J \setminus \{j\}} \text{conv}(Y(f, j)) = \text{conv}(Z(j, f))$ and points of $Z(j, f)$ also have their coordinates polynomially bounded.

Then using Prop. 6, the vertices of $\bigcap_{j \in J} [-W, W]^d \cap \downarrow_{J \setminus \{j\}} \text{conv}(V(f, j))$ have polynomially bounded coordinates. This concludes the proof of Thm. 5. \square

We write X for the set of points that have coordinates polynomially bounded by the same polynomial as for Thm. 5 We give a $\Sigma_2\text{P}$ algorithm (Algorithm 2) that given a constraint λ on the payoffs as a system of linear equation, solves the value problem. Its correctness is a direct consequence of Thm. 5. It therefore implies the following theorem.

Theorem 6. *The threshold problem is in $\Sigma_2\text{P}$.*

Algorithm 2: Algorithm for the value problem in multi-mean-payoff

Existentially guess v in X
 | if $x \in \text{value}(s) \cap \text{hsp}(\lambda)$ then return true;
 | else return false;

References

1. E. H. Bareiss. Sylvester's identity and multistep integer-preserving Gaussian elimination. *Mathematics of computation*, 22(103):565–578, 1968.
2. P. Bouyer, R. Brenguier, N. Markey, and M. Ummels. Concurrent games with ordered objectives. In L. Birkedal, editor, *FoSSaCS'12*, volume 7213 of *Lecture Notes in Computer Science*, pages 301–315. Springer-Verlag, Mar. 2012.
3. T. Brázdil, P. Jancar, and A. Kucera. Reachability games on extended vector addition systems with states. In *ICALP (2)*, pages 478–489, 2010.
4. L. Brim, J. Chaloupka, L. Doyen, R. Gentilini, and J.-F. Raskin. Faster algorithms for mean-payoff games. *Formal methods in system design*, 38(2):97–118, 2011.
5. M. Joswig. Beneath-and-beyond revisited. In *Algebra, Geometry and Software Systems*, pages 1–21. Springer, 2003.

6. M. Jurdziński. Deciding the winner in parity games is in UP and in co-UP. *Information Processing Letters*, 68(3):119–124, 1998.
7. D. A. Martin. Borel determinacy. *The annals of Mathematics*, 102(2):363–371, 1975.
8. J. Matoušek. *Lectures on discrete geometry*, volume 212. Springer, 2002.
9. A. Schrijver. *Theory of linear and integer programming*. John Wiley & Sons, 1998.
10. Y. Velner, K. Chatterjee, L. Doyen, T. A. Henzinger, A. Rabinovich, and J.-F. Raskin. The complexity of multi-mean-payoff and multi-energy games. *CoRR*, abs/1209.3234, 2012.