



HAL
open science

Optimality issues for a class of controlled singularly perturbed stochastic systems

Dan Goreac, Oana Silvia Serea

► **To cite this version:**

Dan Goreac, Oana Silvia Serea. Optimality issues for a class of controlled singularly perturbed stochastic systems. 2014. hal-00987521v1

HAL Id: hal-00987521

<https://hal.science/hal-00987521v1>

Preprint submitted on 6 May 2014 (v1), last revised 27 Jul 2014 (v2)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Optimality issues for a class of controlled singularly perturbed stochastic systems

Dan Goreac* Oana-Silvia Serea†

April 11, 2014

Abstract

The present paper aims at studying stochastic singularly perturbed control systems. We begin by recalling the linear (primal and dual) formulations for classical control problems. In this framework, we give necessary and sufficient support criteria for optimality of the measures intervening in these formulations. Motivated by these remarks, in a first step, we provide linearized formulations associated to the value function in the averaged dynamics setting. Second, these formulations are used to infer criteria allowing to identify the optimal trajectory of the averaged stochastic system.

Key words: Optimal stochastic control, singularly perturbed Brownian diffusions, occupation measures, linear programming.

AMS Classification: 93E20, 49J45, 49L25.

1 Preliminaries

1.1 Introduction

The present paper aims at studying stochastic singularly perturbed control systems. We begin by recalling the linear (primal and dual) formulations for classical control problems. In this framework, we give necessary and sufficient support criteria for optimality of the measures intervening in these formulations. Motivated by these remarks, in a first step, we provide linearized formulations associated to the value

*Université Paris-Est, LAMA (UMR 8050), UPEMLV, UPEC, CNRS, F-77454, Marne-la-Valle, France, Email : dan.goreac@univ-mlv.fr.

†Université de Perpignan Via Domitia, Laboratoire de Mathématiques et de Physique, EA 4217, 52 avenue Paul Alduy, 66860 Perpignan Cedex, France. Email: oana-silvia.serea@univ-perp.fr

function in the averaged dynamics setting. Second, these formulations are used to infer criteria allowing to identify the optimal trajectory of the averaged stochastic system.

Linear programming techniques have proved to be very useful in dealing with deterministic and stochastic control problems. A wide literature is available on the subject both in the deterministic and in the stochastic setting ([10, 3, 14, 15, 13, 17, 4, 9]).

One of the advantages of transforming a nonlinear control problem into a linear optimization problem consists in the possibility of obtaining approximation results for the value function. Following the methods presented in [9] and [16] for the deterministic controlled dynamics, one can approximate the occupational measures by Dirac measures and construct an optimal feedback control. Moreover, when considering the ergodic control problem (see, e.g. [1]), the study of the behavior of the value function is simplified whenever this value is expressed by a linear problem. Recently, linearized versions of the standard continuous infinite horizon discounted control problems have been provided in [16, 6].

When dealing with controlled perturbed dynamics, if the associated system is fully nonlinear, then it is very difficult to characterize the optimal trajectories using the classical methods. Indeed, these criteria involve Pontryagin's maximum principle which is difficult to study if one does not fully understand the averaged dynamics. We recall [11, 22, 12] and references therein dealing with this kind of problems.

We propose an alternative to these classical methods. Our approach consists in embedding the controlled trajectories into a space of probability measures satisfying a convenient constraint. This condition is given in terms of the coefficient functions (and involves the infinitesimal generator of the underlying process). The results allow to characterize the set of constraints as the closed convex hull of occupational measures associated to controls. We first consider general control problems with Lipschitz continuous running and final costs allowing to explain the approach. Using linearization techniques and the dual formulations, we characterize the optimal occupational measures by describing their support set. Next, we extend the linear formulations to singularly perturbed Brownian systems. Finally, we propose support criteria for the optimality of measures in this setting. To our best knowledge, this work is the first to propose a linearization approach to the existence of the optimal policy in the singularly perturbed setting. We emphasize that it does not require to effectively compute the averaged dynamics.

This paper is organized as follows. We briefly state our problem in subsection 1.2. In section 2, we present the main ingredients allowing to deal with classical control problems. We begin with recalling the linear formulations in this setting (cf. [18], see also [19]). In subsection 2.2, we provide a support condition for the optimality of measures appearing in the primal linear formulation. We distinguish between the regular and the general case. The final section aims at presenting singularly perturbed control systems and the averaging method and some important results concerning the singularly perturbed systems and the

value functions associated to this problem. We begin by recalling the basic assumptions and ingredients in subsection 3.1. These results are mainly taken from [4] (see also [5]). Combined with the results in the classical framework, these ingredients allow one to infer linear formulations for the control problems with stochastic singularly perturbed systems (in subsection 3.2). Finally, in subsection 3.3, we give some criteria for optimality in the singularly perturbed setting.

1.2 Singularly perturbed control systems

In the following we shortly present our problem. We consider the following dynamics:

$$\begin{cases} dX_s^{x,y,u;\varepsilon} = f(X_s^{x,y,u;\varepsilon}, Y_s^{x,y,u;\varepsilon}, u_s) ds + \sigma(X_s^{x,y,u;\varepsilon}, Y_s^{x,y,u;\varepsilon}, u_s) dW_s, \\ dY_s^{x,y,u;\varepsilon} = \frac{1}{\varepsilon} g(X_s^{x,y,u;\varepsilon}, Y_s^{x,y,u;\varepsilon}, u_s) ds + \frac{1}{\sqrt{\varepsilon}} \beta(X_s^{x,y,u;\varepsilon}, Y_s^{x,y,u;\varepsilon}, u_s) dB_s, \end{cases} \quad (1)$$

for all $s \geq 0$, $(x, y) \in \mathbb{R}^M \times \mathbb{R}^N$ for some positive integers $M, N > 0$. Here, $\varepsilon > 0$ is a small real parameter. The evolutions of the two state variables X and Y of the system are of different scale. We call x the "slow" variable and y the "fast" variable.

The control space U is assumed to be a compact metric space. The functions $f : \mathbb{R}^M \times \mathbb{R}^N \times U \rightarrow \mathbb{R}^M$, $\sigma : \mathbb{R}^M \times \mathbb{R}^N \times U \rightarrow \mathbb{R}^{M \times d}$ and $g : \mathbb{R}^M \times \mathbb{R}^N \times U \rightarrow \mathbb{R}^N$, $\beta : \mathbb{R}^M \times \mathbb{R}^N \times U \rightarrow \mathbb{R}^{N \times d'}$ are assumed to be uniformly continuous on their domains and Lipschitz-continuous in (x, y) , uniformly with respect to the control parameter $u \in U$. We consider the family of weak control processes :

$$\pi = \left(\Omega, \mathcal{F}, (\mathcal{F}_t)_{t \geq 0}, \mathbb{P}, (W, B), u \right) = \left(\Omega^\pi, \mathcal{F}^\pi, (\mathcal{F}_t^\pi)_{t \geq 0}, \mathbb{P}^\pi, (W^\pi, B^\pi), u^\pi \right)$$

is called a weakly-admissible control and, for every $(x, y) \in \mathbb{R}^{M+N}$, $(X^{x,y,u^\pi;\varepsilon}, Y^{x,y,u^\pi;\varepsilon}, u^\pi)$ is called a weakly-admissible pair, if

- (i) The quadruple $(\Omega, \mathcal{F}, (\mathcal{F}_t)_{t \geq 0}, \mathbb{P})$ is a filtered probability space satisfying the usual assumptions;
- (ii) The process W is a d -dimensional standard Brownian motion defined on $(\Omega, \mathcal{F}, (\mathcal{F}_t)_{t \geq 0}, \mathbb{P})$; the process B is a d' -dimensional standard Brownian motion defined on $(\Omega, \mathcal{F}, (\mathcal{F}_t)_{t \geq 0}, \mathbb{P})$ and independent of W ;
- (iii) The process u is an $(\mathcal{F}_t)_{t \geq 0}$ -progressively measurable process on $(\Omega, \mathcal{F}, \mathbb{P})$ taking its values in U ;
- (iv) The process $(X^{x,y,u^\pi;\varepsilon}, Y^{x,y,u^\pi;\varepsilon}, u^\pi)$ is the unique solution of (1) on $(\Omega, \mathcal{F}, (\mathcal{F}_t)_{t \geq 0}, \mathbb{P})$ satisfying $X_0^{x,y,u^\pi;\varepsilon} = x$ and $Y_0^{x,y,u^\pi;\varepsilon} = y$.

The set of weakly-admissible controls is denoted by \mathcal{U}^w . We denote by $(X_{(\cdot)}^{x,y,u;\varepsilon}, Y_{(\cdot)}^{x,y,u;\varepsilon})$ the solution of (1) starting from $(x, y) \in \mathbb{R}^M \times \mathbb{R}^N$ for some $\pi \in \mathcal{U}^w$.

We let $h : \mathbb{R}^M \rightarrow \mathbb{R}$ be a given bounded function and $T > 0$ a finite time horizon and define the

following payoff

$$C_{x,y;\varepsilon}(\pi) = \mathbb{E}^\pi \left[h \left(X_T^{x,y,u^\pi;\varepsilon} \right) \right], \quad (2)$$

for all $(x, y) \in \mathbb{R}^M \times \mathbb{R}^N$ and all $\pi \in \mathcal{U}^w$. The value function associated with (1) and (2) is

$$W_{\varepsilon,h}(x, y) = \inf_{\pi \in \mathcal{U}^w} C_{x,y;\varepsilon}(\pi), \quad (3)$$

for all $(x, y) \in \mathbb{R}^M \times \mathbb{R}^N$.

The asymptotic behavior of the value function (3) when $\varepsilon \rightarrow 0$ is a very interesting problem. Whenever the control system (1) has some stability property, it is possible to prove that the trajectories $\left(X_{(\cdot)}^{x,y,u^\pi;\varepsilon}, Y_{(\cdot)}^{x,y,u^\pi;\varepsilon} \right)$ of (1) converge towards some solution of some system obtained by formally replacing ε by 0 in (1). This is the so called Tikhonov approach which has been successfully developed in [23, 24], for instance.

When (1) is not stable, another approach consists in investigating relationships between the system (1) and a new differential equation

$$\begin{cases} dX_s^{x,y,u} = \bar{f}(X_s^{x,y,u}, \mu_s) ds + \bar{\sigma}(X_s^{x,y,u}, \mu_s) dW_s, \\ \mu_s \in D_{X_s^{x,y,u}} \text{ for (almost) all } s \in [0, T]. \end{cases} \quad (4)$$

obtained by an averaging method, that will be described later on. We emphasize that, in general, the averaged system is set-valued. We refer the reader to [12, 20] for averaging methods. It is important to notice that only the behavior of the "slow" variable $X_{(\cdot)}^{x,y,u^\pi;\varepsilon}$ is concerned by this approach.

2 Classical control problems

We let $\pi = \left(\Omega, \mathcal{F}, (\mathcal{F}_t)_{t \geq 0}, \mathbb{P}, W, u \right)$ be a weak control consisting of a complete probability space $(\Omega, \mathcal{F}, \mathbb{P})$ endowed with a filtration $\mathbb{F} = (\mathcal{F}_t)_{t \geq 0}$ satisfying the usual assumptions, a standard p -dimensional Brownian motion with respect to this filtration denoted W . We recall that an admissible control process u is any \mathbb{F} -progressively measurable process with values in the compact metric space \mathbb{A} . We denote by $T > 0$ a finite time horizon and let \mathcal{U}^w denote the class of all admissible (weak) controls on $[0, T]$. We consider the stochastic control system

$$\begin{cases} dX_s^{t,x,u} = b(X_s^{t,x,u}, u_s) ds + \rho(X_s^{t,x,u}, u_s) dW_s, \text{ for all } s \in [t, T], \\ X_t^{t,x,u} = x \in \mathbb{R}^m, \end{cases} \quad (5)$$

where $t \in [0, T]$. Throughout the section, we use the following standard assumption on the coefficient functions $b : \mathbb{R}^m \times \mathbb{A} \rightarrow \mathbb{R}^m$ and $\rho : \mathbb{R}^m \times \mathbb{A} \rightarrow \mathbb{R}^{m \times p}$:

$$\left\{ \begin{array}{l} \text{(i) the functions } b \text{ and } \rho \text{ are bounded and uniformly continuous on } \mathbb{R}^m \times \mathbb{A}, \\ \text{(ii) there exists a real constant } c > 0 \text{ such that} \\ |b(x, u) - b(y, u)| + |\rho(x, u) - \rho(y, u)| \leq c|x - y|, \end{array} \right. \quad (6)$$

for all $(x, y, u) \in \mathbb{R}^{2m} \times \mathbb{A}$. Under the Assumption (6), for every $(t, x) \in [0, T] \times \mathbb{R}^m$ and every admissible control $\pi \in \mathcal{U}^w$, there exists a unique solution to (5) starting from (t, x) denoted by X^{t,x,u^π} .

2.1 Lipschitz continuous cost functionals

In this subsection, we recall the basic tools that allow to identify the primal and dual linear formulations associated to (finite horizon) stochastic control problems. The results can be found in [18] (see also [6] for the infinite time horizon).

To any $(t, x) \in [0, T] \times \mathbb{R}^m$ and any $\pi \in \mathcal{U}^w$, we associate the (expectation of the) occupation measures

$$\gamma_{t,T,x,\pi}^1(A \times B \times C) = \frac{1}{T-t} \mathbb{E}^\pi \left[\int_t^T 1_{A \times B \times C}(s, X_s^{t,x,u^\pi}, u_s^\pi) ds \right], \quad \gamma_{t,T,x,\pi}^2(D) = \mathbb{E}^\pi \left[1_D(X_T^{t,x,u^\pi}) \right],$$

for all Borel subsets $A \times B \times C \times D \subset [t, T] \times \mathbb{R}^m \times \mathbb{A} \times \mathbb{R}^m$. Also, we can define

$$\gamma_{t,T,x,\pi}^1(\cdot \times C) = \delta_{(t,x)}(\cdot) \times \mathbb{P}^\pi(u_T^\pi \in C), \quad \gamma_{t,T,x,\pi}^2 = \delta_x,$$

where δ denotes the Dirac measure. We denote by

$$\Gamma_{(b,\rho)}(t, T, x) = \left\{ \gamma_{t,T,x,\pi} = (\gamma_{t,T,x,\pi}^1, \gamma_{t,T,x,\pi}^2) \in \mathcal{P}([t, T] \times \mathbb{R}^m \times \mathbb{A}) \times \mathcal{P}(\mathbb{R}^m) : \pi \in \mathcal{U}^w \right\}.$$

Here, $\mathcal{P}(\mathcal{X})$ stands for the set of probability measures on the metric space \mathcal{X} . Due to the Assumption (6), there exists a positive constant C_0 (depending on $T > 0$) such that, for every $(t, x) \in [0, T] \times \mathbb{R}^m$ and every $\pi \in \mathcal{U}^w$, one has

$$\sup_{s \in [t, T]} \mathbb{E}^\pi \left[\left| X_s^{t,x,u^\pi} \right|^4 \right] \leq C_0 (|x|^4 + 1). \quad (7)$$

Therefore,

$$\left\{ \begin{array}{l} \int_{\mathbb{R}^m} |y|^4 \gamma_{t,T,x,\pi}^1([t, T], dy, \mathbb{A}) \leq C_0 (|x|^4 + 1), \\ \int_{\mathbb{R}^m} |y|^4 \gamma_{t,T,x,\pi}^2(dy) \leq C_0 (|x|^4 + 1). \end{array} \right. \quad (8)$$

We define

$$\Theta_{(b,\rho)}(t, T, x) = \left\{ \begin{array}{l} \gamma \in \mathcal{P}([t, T] \times \mathbb{R}^m \times \mathbb{A}) \times \mathcal{P}(\mathbb{R}^m) : \forall \phi \in C_b^{1,2}([0, T] \times \mathbb{R}^m), \\ \int_{[t, T] \times \mathbb{R}^m \times \mathbb{A} \times \mathbb{R}^m} \begin{bmatrix} (T-t) \mathcal{L}_{(b,\rho)}^v \phi(s, y) \\ + \phi(t, x) - \phi(T, z) \end{bmatrix} \gamma^1(ds, dy, dv) \gamma^2(dz) = 0. \end{array} \right\}, \quad (9)$$

where

$$\mathcal{L}_{(b,\rho)}^v \phi(s, y) = \frac{1}{2} \text{Tr} [(\rho \rho^*)(y, v) D^2 \phi(s, y)] + \langle b(y, v), D\phi(s, y) \rangle + \partial_t \phi(s, y),$$

for all $(s, y) \in [0, T] \times \mathbb{R}^m$, $v \in \mathbb{A}$ and all $\phi \in C^{1,2}([0, T] \times \mathbb{R}^m)$. Itô's formula applied to test functions $\phi \in C_b^{1,2}([0, T] \times \mathbb{R}^m)$ yields

$$\Gamma_{(b,\rho)}(t, T, x) \subset \Theta_{(b,\rho)}(t, T, x).$$

Moreover, the set $\Theta_{(b,\rho)}(t, T, x)$ is convex and a closed subset of $\mathcal{P}([t, T] \times \mathbb{R}^m \times \mathbb{A}) \times \mathcal{P}(\mathbb{R}^m)$. For further details, the reader is referred to [18].

Let us suppose that $l_1 : \mathbb{R} \times \mathbb{R}^m \times \mathbb{A} \rightarrow \mathbb{R}$, $l_2 : \mathbb{R}^m \rightarrow \mathbb{R}$ are bounded and uniformly continuous such that

$$|l_1(t, x, u) - l_1(s, y, u)| + |l_2(x) - l_2(y)| \leq c(|x - y| + |t - s|), \quad (10)$$

for all $(s, t, x, y, u) \in \mathbb{R}^2 \times \mathbb{R}^{2m} \times \mathbb{A}$, and for some positive $c > 0$. We introduce the usual value function

$$\begin{aligned} V_{l_1, l_2, (b, \rho)}(t, x) &= \inf_{\pi \in \mathcal{U}^w} \mathbb{E}^\pi \left[\int_t^T l_1(s, X_s^{t,x,u^\pi}, u_s^\pi) ds + l_2(X_T^{t,x,u^\pi}) \right] \\ &= \inf_{\gamma \in \Gamma_{(b,\rho)}(t, T, x)} \left((T-t) \int_{[t, T] \times \mathbb{R}^m \times \mathbb{A}} l_1(s, y, u) \gamma^1(ds, dy, du) + \int_{\mathbb{R}^m} l_2(y) \gamma^2(dy) \right), \end{aligned} \quad (11)$$

and the primal linearized value function

$$\Lambda_{l_1, l_2, (b, \rho)}(t, x) = \inf_{\gamma \in \Theta_{(b,\rho)}(t, T, x)} \left((T-t) \int_{[t, T] \times \mathbb{R}^m \times \mathbb{A}} l_1(s, y, u) \gamma^1(ds, dy, du) + \int_{\mathbb{R}^m} l_2(y) \gamma^2(dy) \right), \quad (12)$$

for all $(t, x) \in [0, T] \times \mathbb{R}^m$. We also consider the dual value function

$$\mu_{l_1, l_2, (b, \rho)}(t, x) = \sup \left\{ \begin{array}{l} \mu \in \mathbb{R} : \exists \phi \in C_b^{1,2}([0, T] \times \mathbb{R}^m) \text{ s.t. } \forall (s, y, v, z) \in [t, T] \times \mathbb{R}^m \times \mathbb{A} \times \mathbb{R}^m, \\ \mu \leq (T-t) \left(\mathcal{L}_{(b,\rho)}^v \phi(s, y) + l_1(s, y, v) \right) + l_2(z) - \phi(T, z) + \phi(t, x), \end{array} \right\} \quad (13)$$

for all $(t, x) \in [0, T] \times \mathbb{R}^m$. The following result is a slight generalization of Theorem 4 in [18]. The proof

is very similar and will be omitted.

Theorem 2.1 (see [18], Theorem 4). *Under the Assumptions (6) and (10),*

$$V_{l_1, l_2, (b, \rho)} = \Lambda_{l_1, l_2, (b, \rho)} = \mu_{l_1, l_2, (b, \rho)}.$$

Since this result holds true for arbitrary (regular) functions l_1 and l_2 , a standard separation argument yields:

Corollary 2.1 *The set of constraints $\Theta_{(b, \rho)}(t, T, x)$ is the closed, convex hull of $\Gamma_{(b, \rho)}(t, T, x)$:*

$$\Theta_{(b, \rho)}(t, T, x) = \overline{\text{co}}\Gamma_{(b, \rho)}(t, T, x). \quad (14)$$

The closure is taken with respect to the usual (narrow) convergence of probability measures.

Remark 2.1 *Due to the inequality (8), Prohorov's theorem yields that $\text{co}\Gamma(t, T, x)$ is relatively compact and, thus, $\Theta_{(b, \rho)}(t, T, x)$ is compact. Moreover,*

$$\begin{cases} \int_{\mathbb{R}^m} |y|^4 \gamma^1([t, T], dy, \mathbb{A}) \leq C_0 (|x|^4 + 1), \\ \int_{\mathbb{R}^m} |y|^4 \gamma^2(dy) \leq C_0 (|x|^4 + 1), \end{cases} \quad (15)$$

for all $\gamma = (\gamma^1, \gamma^2) \in \Theta_{(b, \rho)}(t, T, x)$.

We equally mention the following result due to N. V. Krylov (cf. [21], Theorem 2.1). It is both an essential ingredient in proving Theorem 2.1 and a tool for further developments.

Proposition 2.1 *There exists a constant $C > 0$ such that, for every $\delta \in (0, 1]$, there exists a function $V^\delta \in C_b^{1,2}([0, T + \delta^2] \times \mathbb{R}^m)$ such that*

$$\mathcal{L}_{(b, \rho)}^v \phi(s, y) + l_1(s, y, v) \geq 0,$$

for all $(s, y, v) \in [0, T + \delta^2] \times \mathbb{R}^m \times \mathbb{A}$ and

- (i) $|V^\delta(t, \cdot) - l_2(\cdot)| \leq C\delta$, for $t \in [T, T + \delta^2]$, and
- (ii) $|V^\delta(\cdot) - V_{l_1, l_2, (b, \rho)}(\cdot)| \leq C\delta$, on $[0, T] \times \mathbb{R}^N$.

Remark 2.2 (i) The constant C only depends on the Lipschitz constants and the bounds of (b, ρ) :

$$C \leq c_0 (1 + |b|_\infty + \text{Lip}(b) + |\rho|_\infty + \text{Lip}(\rho)),$$

where c_0 is a constant (depending, eventually on T).

(ii) We assume that $l_1 = 0$. Then, the functions V^δ are obtained by the "shaking of coefficients" method as $V_\delta * \psi_\delta$, where $V_\delta = V_{0, l_2, (b^\delta, \rho^\delta)}$ with

$$b^\delta(x, u, v) = b(x + \delta v, u), \quad \rho^\delta(x, u, v) = \rho(x + \delta v, u), \quad u \in \mathbb{A}, \quad v \in \mathbb{R}^m, \quad |v| \leq 1$$

and $(\psi_\delta)_\delta$ a sequence of standard mollifiers $\psi_\delta(y) = \frac{1}{\delta^m} \psi\left(\frac{y}{\delta}\right)$, $y \in \mathbb{R}^m$, $\delta > 0$, where $\psi \in C^\infty(\mathbb{R}^m)$ is a positive function such that

$$\text{Supp}(\psi) \subset \overline{B}(0, 1) \quad \text{and} \quad \int_{\mathbb{R}^m} \psi(x) dx = 1.$$

2.2 Characterization of optimal measures

In this subsection we present necessary and sufficient conditions for characterizing optimal occupational measures. We consider that $l_1 \equiv 0$, $T > 0$ is fixed and we set

$$\Theta(x) := \Theta_{(b, \rho)}(0, T, x), \quad V_{l_2}(x) := V_{0, l_2, (b, \rho)}(0, x), \quad \Lambda_{l_2}(x) := \Lambda_{0, l_2, (b, \rho)}(0, x), \quad \eta_{l_2}(x) := \eta_{0, l_2, (b, \rho)}(0, x),$$

for simplicity. Recall that, with the above notations,

$$V_{l_2}(x) = \Lambda_{l_2}(x) = \eta_{l_2}(x),$$

for all initial data $x \in \mathbb{R}^m$ and

$$\eta_{l_2}(x) = \sup \left\{ \begin{array}{l} \eta \in \mathbb{R} : \exists \phi \in C_b^{1,2}(\mathbb{R}_+ \times \mathbb{R}^m) \text{ s.t. } \forall (s, y, v, z) \in [0, T] \times \mathbb{R}^m \times \mathbb{A} \times \mathbb{R}^m, \\ \eta \leq T \mathcal{L}^v \phi(s, y) + l_2(z) - \phi(T, z) + \phi(0, x) \end{array} \right\}, \quad (16)$$

for all $x \in \mathbb{R}^m$. We denote by

$$D_{l_2}(x) = \left\{ \begin{array}{l} (\eta, \phi) \in \mathbb{R} \times C_b^{1,2}(\mathbb{R}_+ \times \mathbb{R}^m) \text{ s.t.} \\ \eta = \inf_{(s, y, v, z) \in [0, T] \times \mathbb{R}^m \times \mathbb{A} \times \mathbb{R}^m} \{T \mathcal{L}^v \phi(s, y) + l_2(z) - \phi(T, z) + \phi(0, x)\} \end{array} \right\}, \quad (17)$$

for all $x \in \mathbb{R}^m$. By our assumptions, the coefficient functions are bounded and, thus, the set $D_{l_2}(x)$ is well defined.

The dual formulation yields

$$V_{l_2}(x) = \sup\{\eta, (\eta, \bar{\phi}) \in D_{l_2}(x)\}. \quad (18)$$

2.2.1 The regular case

We introduce the following.

Definition 2.1 *Whenever $x \in \mathbb{R}^m$, we say that $(\bar{\eta}, \bar{\phi}) \in D_{l_2}(x)$ is an optimal pair whenever we have $V_{l_2}(x) = \bar{\eta}$.*

We denote by

$$\Omega_{l_2, (\bar{\eta}, \bar{\phi})}(x) = \left\{ \begin{array}{l} (s, y, v, z) \in [0, T] \times \mathbb{R}^m \times \mathbb{A} \times \mathbb{R}^m, s.t. \\ \bar{\eta} = T\mathcal{L}^v \bar{\phi}(s, y) + l_2(z) - \bar{\phi}(T, z) + \bar{\phi}(0, x). \end{array} \right\} \quad (19)$$

Of course, neither the set of optimal pairs, nor $\Omega_{l_2, (\bar{\eta}, \bar{\phi})}$ are a priori non empty. It is the case if $V_{0, l_2}(\cdot, \cdot)$ belongs to $C_b^{1,2}(\mathbb{R}_+ \times \mathbb{R}^m)$ and we consider the setting of the problem to be some invariant compact set $K \subset \mathbb{R}^m$. In this framework, one can guarantee that optimal pairs exists for every $x \in K$. Indeed, it suffices to consider $\bar{\phi} = V_{0, l_2}$ and get, using the fact that it is a (regular) subsolution of the associated HJB equation,

$$T\mathcal{L}^v \bar{\phi}(s, y) \geq 0, \quad l_2(z) \geq \bar{\phi}(T, z),$$

for all $(s, y, v, z) \in [0, T] \times K \times \mathbb{A} \times K$. Hence,

$$V_{l_2}(x) \leq T\mathcal{L}^v \bar{\phi}(s, y) + l_2(z) - \bar{\phi}(T, z) + \bar{\phi}(0, x),$$

for all $(s, y, v, z) \in [0, T] \times K \times \mathbb{A} \times K$. The fact that $\Omega_{l_2, (\bar{\eta}, \bar{\phi})}(x)$ is nonempty follows from the compactness of K .

Proposition 2.2 *Let $x \in \mathbb{R}^m$ be fixed and assume that $(\bar{\eta}, \bar{\phi}) \in D_{l_2}(x)$ is an optimal pair. Then, $\gamma \in \Theta(x)$ is optimal for $\Lambda_{l_2}(x)$ if and only if $\Omega_{l_2, (\bar{\eta}, \bar{\phi})}(x)$ is nonempty and $\gamma\left(\Omega_{l_2, (\bar{\eta}, \bar{\phi})}(x)\right) = 1$.*

Proof. We begin with assuming that $\gamma \in \Theta(x)$ is such that $\gamma\left(\Omega_{l_2, (\bar{\eta}, \bar{\phi})}(x)\right) = 1$, i.e. the support of γ is included in $\Omega_{l_2, (\bar{\eta}, \bar{\phi})}(x)$. Then, by definition, we have the following equality

$$\bar{\eta} = T\mathcal{L}^v \bar{\phi}(s, y) + l_2(z) - \bar{\phi}(T, z) + \bar{\phi}(t, x),$$

on $\Omega_{l_2,(\bar{\eta},\bar{\phi})}(x)$. Consequently, recalling the definition of $\Theta(x)$, one gets

$$V_{l_2}(x) = \int_{[0,T] \times \mathbb{R}^m \times \mathbb{A} \times \mathbb{R}^m} \bar{\eta} \gamma(ds dx dudz) = \int_{[0,T] \times \mathbb{R}^m \times \mathbb{A} \times \mathbb{R}^m} l_2(z) \gamma(ds dx dudz),$$

i.e. $\gamma \in \Theta(x)$ is optimal. Conversely, let us consider some optimal $\gamma \in \Theta(x)$. One writes

$$\begin{aligned} V_{l_2}(x) &= \int_{[t,T] \times \mathbb{R}^m \times \mathbb{A} \times \mathbb{R}^m} l_2(z) \gamma(ds dx dudz) = \\ &= \int_{[0,T] \times \mathbb{R}^N \times \mathbb{A} \times \mathbb{R}^N} [T \mathcal{L}^v \bar{\phi}(s, y) + l_2(z) - \bar{\phi}(T, z) + \bar{\phi}(t, x)] \gamma(ds dx dudz). \end{aligned}$$

for all optimal pairs $(\bar{\eta} = V_{l_2}(x), \bar{\phi}) \in D_{l_2}(x)$. By the definition of $D_{l_2}(x)$, it follows that

$$[T \mathcal{L}^v \bar{\phi}(s, y) + l_2(z) - \bar{\phi}(T, z) + \bar{\phi}(t, x)] \geq V_{l_2}(x).$$

Hence, in order for the previous equality to hold, one has

$$[T \mathcal{L}^v \bar{\phi}(s, y) + l_2(z) - \bar{\phi}(T, z) + \bar{\phi}(t, x)] = V_{l_2}(x),$$

γ -almost everywhere. Hence, the support of γ is included in $\Omega_{l_2,(\bar{\eta},\bar{\phi})}(x)$ and the proof is now complete.

□

Remark 2.3 *One can construct a set which is independent of the choice of optimal pairs $(\bar{\eta}, \bar{\phi}) \in D_{l_2}(x)$. Indeed, in the case where the state space K is compact, the set $C_b^{1,2}([0, T] \times K)$ is compact. The family of optimal test functions is denoted by $Opt_{l_2}(x)$ and is totally bounded with respect to the usual topology of $C_b^{1,2}$. For every $n \geq 1$, we select a finite family $(\bar{\phi}_j^n)_{1 \leq j \leq k^n} \subset Opt_{l_2}(x)$ such that, for every $\bar{\phi} \in Opt_{l_2}(x)$*

$$d(\bar{\phi}, \bar{\phi}_j^n) \leq \frac{1}{n^2},$$

for some $1 \leq j \leq k^n$. The distance is given in the sense of $C_b^{1,2}([0, T] \times K)$ functions. We define

$$\Omega_{l_2}(x) := \bigcap_{n \geq 1, 1 \leq j \leq k^n} \Omega_{l_2,(\bar{\eta}, \bar{\phi}_j^n)}(x).$$

Due to the previous proposition, whenever γ is optimal, $\gamma(\Omega_{l_2}(x)) = 1$. The converse also holds true. If no invariant compact can be found for the system, a localization procedure can be developed starting from Remark 2.1.

2.2.2 The general framework

If the value function is not smooth, optimal pairs may not exist. However, if optimal pairs do not exist, one finds some sequence $(\eta_n, \phi_n) \in D_{l_2}(x)$ such that $(\eta_n)_n$ is strictly increasing and converging to $V_{l_2}(x)$. The functions ϕ_n can be chosen to be uniformly bounded (e.g. Theorem 2.1 in [21], see also Proposition 3 in [18]). We define the nonempty, closed sets

$$\Omega_{l_2}^n(x) = \left\{ \begin{array}{l} (s, y, v, z) \in [0, T] \times \mathbb{R}^m \times \mathbb{A} \times \mathbb{R}^m, \text{ s.t.} \\ V_{l_2}(x) + \sqrt{V_{l_2}(x) - \eta_n} \geq T\mathcal{L}^v \phi_n(s, y) + l_2(z) - \phi_n(T, z) + \phi_n(0, x). \end{array} \right\} \quad (20)$$

and

$$\begin{aligned} \Omega_{l_2}^{in}(x) &:= \liminf_{n \rightarrow \infty} \Omega_{l_2}^n(x) = \bigcup_{n \geq 1} \bigcap_{k \geq n} \Omega_{l_2}^k(x), \quad \Omega_{l_2}^{out}(x) := \limsup_{n \rightarrow \infty} \Omega_{l_2}^n(x) = \bigcap_{n \geq 1} \bigcup_{k \geq n} \Omega_{l_2}^k(x), \\ \Omega_{l_2}^{out, cl}(x) &:= \bigcap_{n \geq 1} cl \left(\bigcup_{k \geq n} \Omega_{l_2}^k(x) \right), \end{aligned}$$

where cl is the usual Kuratowski closure operator.

Remark 2.4 *If an optimal pair $(V_{l_2}(x), \bar{\phi})$ exists, we pick $\phi_n = \bar{\phi}$. In this case, $\eta_n = V_{l_2}(x)$. The sets $\Omega_{l_2}^n(x)$ coincide. Hence, $\Omega_{l_2}^{out}(x) = \Omega_{l_2}^{in}(x) = \Omega_{l_2, (V_{l_2}(x), \bar{\phi})}(x)$ as in the previous case.*

We get the following characterization of the support of optimal measures.

Proposition 2.3 *Let us consider $x \in \mathbb{R}^m$.*

(i) *If $\gamma \in \Theta(x)$ is optimal, then*

$$\gamma \left(\Omega_{l_2}^{out, cl}(x) \right) = \gamma \left(\Omega_{l_2}^{out}(x) \right) = 1,$$

(i.e. the support of γ is included in $\Omega_{l_2}^{out}(x)$). In particular, when the limit of the sets exists (i.e. $\Omega_{l_2}^{in}(x) = \Omega_{l_2}^{out}(x)$), one gets

$$\sup_{n \geq 1} \gamma \left(\bigcap_{k \geq n} \Omega_{l_2}^k(x) \right) = 1.$$

(ii) *Conversely, if $\gamma \in \Theta(x)$ is such that the supremum can be replaced with maximum (i.e. if there exists some n_0 such that $\gamma \left(\bigcap_{k \geq n_0} \Omega_{l_2}^k(x) \right) = 1$), then γ is optimal.*

Proof. We begin with assuming that $\gamma \in \Theta(x)$ is optimal (should it exist). Then

$$\begin{aligned} V_{l_2}(x) &= \int_{[t, T] \times \mathbb{R}^m \times \mathbb{A} \times \mathbb{R}^m} l_2(z) \gamma(ds dx dudz) = \\ &= \int_{[0, T] \times \mathbb{R}^N \times \mathbb{A} \times \mathbb{R}^N} [T\mathcal{L}^v \phi_k(s, y) + l_2(z) - \phi_k(T, z) + \phi_k(0, x)] \gamma(ds dx dudz), \end{aligned}$$

for all $k \geq 1$. The definition of $D_{l_2}(x)$ yields

$$[T\mathcal{L}^v \phi_k(s, y) + l_2(z) - \phi_k(T, z) + \phi_k(0, x)] \geq \eta_k,$$

for all $(s, y, v, z) \in [0, T] \times \mathbb{R}^N \times \mathbb{A} \times \mathbb{R}^N$ and all $k \geq 1$. Thus,

$$V_{l_2}(x) \geq \eta_k \gamma(\Omega_{l_2}^k(x)) + \left(V_{l_2}(x) + \sqrt{V_{l_2}(x) - \eta_k} \right) \gamma\left((\Omega_{l_2}^k(x))^c \right).$$

It follows that

$$\left(1 + \sqrt{V_{l_2}(x) - \eta_k} \right) \gamma(\Omega_{l_2}^k(x)) \geq 1.$$

Passing to the lim sup, one gets $\gamma(\Omega_{l_2}^{out}(x)) = 1$ and the proof is complete.

For the converse, let us assume that $\gamma\left(\bigcap_{k \geq n_0} \Omega_{l_2}^k(x)\right) = 1$. Then $\gamma(\Omega_{l_2}^k(x)) = 1$, for all $k \geq n_0$. It follows that

$$\begin{aligned} V_{l_2}(x) &\leq \int_{[0, T] \times \mathbb{R}^m \times \mathbb{A} \times \mathbb{R}^m} l_2(z) \gamma(dsdxdudz) \\ &= \int_{[0, T] \times \mathbb{R}^m \times \mathbb{A} \times \mathbb{R}^m} (T\mathcal{L}^v \phi_k(s, y) + l_2(z) - \phi_k(T, z) + \phi_k(0, x)) \gamma(dsdxdudz) \\ &\leq V_{l_2}(x) + \sqrt{V_{l_2}(x) - \eta_k}, \end{aligned}$$

for all $k \geq n_0$. Passing to the limit as $k \rightarrow \infty$, one gets that $\int_{[0, T] \times \mathbb{R}^m \times \mathbb{A} \times \mathbb{R}^m} l_2(z) \gamma(dsdxdudz) = V_{l_2}(x)$, i.e. $\gamma \in \Theta(x)$ is optimal. \square

Remark 2.5 *If an optimal pair exists for our control problem then, due to the Remark 2.4, a measure $\gamma \in \Theta(x)$ is optimal if and only if $\gamma(\Omega_{l_2}^{in/out}(x)) = 1$.*

3 The averaging method

3.1 General considerations

All the assumptions and ideas of this preliminary part can be found in [4]. Let us shortly explain the behavior of the perturbed system (1) as $\varepsilon \rightarrow 0$. To this purpose, let us fix, for the time being, $\varepsilon > 0$ and the weak control $\pi = \left(\Omega, \mathcal{F}, (\mathcal{F}_t)_{t \geq 0}, \mathbb{P}, (W, B), u \right)$. If one makes the change of variables $\tau = \frac{s}{\varepsilon}$ in the system (1) and sets $\left(\tilde{X}_\tau, \tilde{Y}_\tau, \tilde{u}_\tau \right) = (X_{\varepsilon\tau}, Y_{\varepsilon\tau}, u_{\varepsilon\tau})$, $B'_\tau = \frac{1}{\sqrt{\varepsilon}} B_{\varepsilon\tau}$, $W'_\tau = \frac{1}{\sqrt{\varepsilon}} W_{\varepsilon\tau}$ for $\tau \in [0, \frac{T}{\varepsilon}]$, one gets

$$\begin{cases} d\tilde{X}_\tau^{x,y,u} = \varepsilon f\left(\tilde{X}_\tau^{x,y,u}, \tilde{Y}_\tau^{x,y,u}, \tilde{u}_\tau\right) d\tau + \sqrt{\varepsilon}\sigma\left(\tilde{X}_\tau^{x,y,u}, \tilde{Y}_\tau^{x,y,u}, \tilde{u}_\tau\right) dW'_\tau, \\ d\tilde{Y}_\tau^{x,y,u} = g\left(\tilde{X}_\tau^{x,y,u}, \tilde{Y}_\tau^{x,y,u}, \tilde{u}_\tau\right) d\tau + \beta\left(\tilde{X}_\tau^{x,y,u}, \tilde{Y}_\tau^{x,y,u}, \tilde{u}_\tau\right) dB'_\tau, \end{cases} \quad (21)$$

When ε tends to 0, we are led to consider the following associated system:

$$dY_\tau^{x,y,u} = g(x, Y_\tau^{x,y,u}, u_\tau) d\tau + \beta(x, Y_\tau^{x,y,u}, u_\tau) dB'_\tau \quad (22)$$

for $\tau \in [0, +\infty)$, where x (resp. y) is a fixed \mathbb{R}^M (resp. \mathbb{R}^N)-valued random variable independent of B' . We denote by $y_{(\cdot)}^{y,u;x}$ the unique solution of (22) corresponding to the control u and to the initial value y . The framework will still be that of weak controls.

Assumption 1 *Following the approach in [4] (see also [5]), throughout the paper, unless stated otherwise, we will assume that*

$$\begin{aligned} \sup_{\varepsilon>0, t \in [0, T], \pi \in \mathcal{U}^w} \mathbb{E}^\pi \left[\left| Y_t^{x,y,u^\pi; \varepsilon} \right|^4 \right] &< c \left(|x|^4 + |y|^4 + 1 \right), \\ \sup_{t \in [0, T], \pi \in \mathcal{U}^w} \mathbb{E}^\pi \left[\left| y_t^{x,y,u^\pi} \right|^4 \right] &< c \left(|x|^4 + |y|^4 + 1 \right), \end{aligned} \quad (A1)$$

for all initial data $(x, y) \in \mathbb{R}^M \times \mathbb{R}^N$.

For explicit conditions (e.g. asymptotic exponential stability for the fourth order moment) implying the above inequalities, the reader is referred to [4], page 172.

Whenever $x \in \mathbb{R}^M$, we let

$$D_x := \left\{ \begin{array}{l} \mu \in \mathcal{P}(\mathbb{R}^N \times U) : \\ \int \langle g(x, y, u), D\phi(y) \rangle + \frac{1}{2} \text{Tr}(\beta\beta^*(x, y, u) D^2\phi(x)) \mu(dydu) = 0. \end{array} \right\}$$

It turns out that $x \rightsquigarrow D_x$ is an upper semicontinuous set-valued function with nonempty, closed, convex values (cf. [4], Lemma 2.1).

The averaged system is given by

$$\begin{cases} d\bar{X}_s^{x,u} = \bar{f}\left(\bar{X}_s^{x,\mu}, \mu_s\right) ds + \bar{\sigma}\left(\bar{X}_s^{x,\mu}, \mu_s\right) dW_s, \\ \mu_s \in D_{\bar{X}_s^{x,\mu}} \text{ for (almost) all } s \in [0, T], \end{cases} \quad (23)$$

where $\bar{f}(x, \mu) := \int f(x, y, u) \mu(dydu)$, $\bar{\sigma}(x, \mu) := \int \sigma(x, y, u) \mu(dydu)$ and the control processes are $\mathcal{P}(\bar{\mathbb{R}}^N \times U)$ -valued. For further considerations on the compactness issues on $\mathcal{P}(\bar{\mathbb{R}}^N \times U)$, the reader is referred to [4], section 2. In particular, one can introduce a metric (denoted by d) on $\mathcal{P}(\bar{\mathbb{R}}^N \times U)$

which is consistent with the weak convergence of probability measures. The set of $\mathcal{P}(\overline{\mathbb{R}}^N \times U)$ -valued weakly-admissible controls will be denoted by $\overline{\mathcal{U}}_N^w$.

Following Assumption 2 in [4], we ask that

Assumption 2 *There exists some $\omega_c \in C(\mathbb{R}_+; \mathbb{R}_+)$ satisfying $\lim_{S \rightarrow \infty} \omega_c(S) = 0$ such that, whenever $x \in \mathbb{R}^M, y \in \mathbb{R}^N$ satisfy $|x| \leq c$ and $\mu \in D_x$, there exists an admissible weak control π such that*

$$\mathbb{E}^\pi [d(\mu, \mu_{0,S,x,\pi}^1)] \leq \omega_c(S).$$

The measure $\mu_{0,S,x,\pi}^1$ is similar to the occupation measures $\gamma_{0,S,x,\pi}^1$ but it does not involve the expectation i.e.

$$\mu_{0,S,x,\pi}^1(B \times C) = \frac{1}{S} \int_0^S 1_{B \times C}(s, y_s^{y, u^\pi; x}, u_s^\pi) ds,$$

for all Borel subsets $B \times C \subset \mathbb{R}^N \times U$. The previous assumption is implied by classical mixing conditions (cf. [4], Proposition 4.1) if one further assumes that the noise coefficient is control independent.

We recall that $h : \mathbb{R}^M \rightarrow \mathbb{R}$ is a given bounded function and $T > 0$ a finite time horizon and define the following payoff

$$C_{x,y;\varepsilon}(\pi) = \mathbb{E}^\pi \left[h \left(X_T^{x,y,u^\pi;\varepsilon} \right) \right],$$

for all $(x, y) \in \mathbb{R}^M \times \mathbb{R}^N$ and all $\pi \in \mathcal{U}^w$. The value function associated with (1) and (2) is

$$W_{\varepsilon,h}(x, y) = \inf_{\pi \in \mathcal{U}^w} C_{x,y;\varepsilon}(\pi),$$

for all $(x, y) \in \mathbb{R}^M \times \mathbb{R}^N$. We also consider the optimal control problem

$$W_h(x) = \inf_{\overline{\pi} \in \overline{\mathcal{U}}_N^w} \mathbb{E}^{\overline{\pi}} \left[h \left(\overline{X}_T^{x, \mu^{\overline{\pi}}} \right) \right], \quad (24)$$

for all initial data $x \in \mathbb{R}^M$.

We endow the space $\mathbb{R}^M \times \mathcal{P}(\overline{\mathbb{R}}^N \times U)$ with the metric \tilde{d} given by

$$\tilde{d}((x, \mu), (x', \mu')) = |x - x'| + d(\mu, \mu'),$$

for all $(x, \mu), (x', \mu') \in \mathbb{R}^M \times \mathcal{P}(\overline{\mathbb{R}}^N \times U)$. We introduce the set valued function with nonempty, convex, compact values

$$\mathbb{R}^M \ni x \rightsquigarrow Q_x := \{(\overline{b}(x, \mu), \mu) : \mu \in D_x\}$$

and make the following (see Assumption 3 in [4])

Assumption 3 *The set valued function Q is Lipschitz continuous on \mathbb{R}^M (i.e. there exists $c_0 \in \mathbb{R}$ such that*

$$\tilde{d}_{\text{Hausdorff}}(Q_x, Q_{x'}) \leq c_0 |x - x'|, \text{ for all } x, x' \in \mathbb{R}^M.$$

Here, $\tilde{d}_{\text{Hausdorff}}$ denotes the Hausdorff distance constructed from \tilde{d}).

Remark 3.1 *Both the Assumption 2 and Assumption 3 hold true if the system (22) satisfies an exponential ergodicity condition, uniformly with respect to the control process (cf. [4], Assumption 4 and Proposition 5.2). This condition can be obtained if dissipativity is assumed for the stochastic system (22). Alternatively, it is possible to adapt the arguments in [7] to deal with nonexpansive (yet nondissipative) systems. However, this generalization is not within the scopus of the present paper.*

Under the above conditions (cf. [4], Theorem 3.3, Theorem 4.2 and [5], Theorem 5.1), every partial limit of solutions $\left(X_{(\cdot)}^{x,y,u^{\pi^\varepsilon};\varepsilon}\right)_{\varepsilon>0}$ satisfies (23) and, conversely, for every solution $\bar{X}^{x,u^{\bar{\pi}}}$ of (23), one finds a suitable sequence $\left(X_{(\cdot)}^{x,y,u^{\pi^\varepsilon};\varepsilon}\right)_{\varepsilon>0}$ converging to $\bar{X}^{x,u^{\bar{\pi}}}$. Due to Assumption 2, the distance is given uniformly with respect to x within a compact set. To simplify our presentation, let us assume that

Assumption 4 *There exists some compact set $K \subset \mathbb{R}^M$ such that $K \times \mathbb{R}^N$ is invariant with respect to (5).*

For explicit criteria of invariance, the reader is referred to [2] (also see [8]). We note that these criteria only involve the coefficients f and σ .

If the cost functional h is bounded and uniformly continuous, the convergence of the value functions is a direct consequence of the convergence of trajectories. More precisely, we have $W_{\varepsilon,h} \rightarrow W_h$ with respect with the uniform convergence :

There exists $\omega \in C(\mathbb{R}_+; \mathbb{R}_+)$ satisfying $\lim_{\varepsilon \rightarrow 0} \omega(\varepsilon) = 0$ such that

$$|W_{\varepsilon,h}(x,y) - W_h(x)| \leq \omega(\varepsilon), \tag{25}$$

for all $x \in K$ and all $y \in \mathbb{R}^N$ (see [4] Corollaries 3.4 and 4.3).

Remark 3.2 *The estimates in [4] show that ω depends on the bounds of the coefficient and cost functions and their continuity moduli, but not on the functions themselves. Thus, if $\delta > 0$ and $W_{\varepsilon,h,\delta}$ is the value function associated with the "shaked" problem (i.e. in which $\varphi \in \{f, \sigma, g, \beta\}$ are replaced with*

$\varphi^\delta(x, y, u, v) = \varphi(x + \delta v, y + \delta v', u)$, $(v, v') \in \mathbb{R}^M \times \mathbb{R}^N$, $|(v, v')| \leq 1$) under analogous assumptions, the inequality (25) holds true for some $W_{h,\delta}$ constructed as before replacing W_h . In particular,

$$|W_{\varepsilon,h,\delta}(x, y) - W_{\varepsilon,h,\delta}(x, y')| \leq 2\omega(\varepsilon),$$

for all $x \in K$ and all $y, y' \in \mathbb{R}^N$. Now, let us consider $(\psi_\delta)_\delta$ to be a sequence of standard mollifiers $\psi_\delta(x, y) = \frac{1}{\delta^{M+N}} \psi\left(\frac{x}{\delta}, \frac{y}{\delta}\right)$, $(x, y) \in \mathbb{R}^{M+N}$, $\delta > 0$, where $\psi \in C^\infty(\mathbb{R}^{M+N})$ is a positive function such that

$$\text{Supp}(\psi) \subset \bar{B}(0, 1) \quad \text{and} \quad \int_{\mathbb{R}^{M+N}} \psi(x) dx = 1.$$

Then, (cf. Remark 2.2 (i) and (25)), the convoluted function $W_{\varepsilon,h}^\delta := W_{\varepsilon,h,\delta} * \psi_\delta$ satisfy :

$$\left\{ \begin{array}{l} |W_{\varepsilon,h}^\delta(x, y) - W_{\varepsilon,h}(x, y)| \leq c_0 \left(1 + \frac{1}{\varepsilon}\right) \delta, \\ |W_{\varepsilon,h}^\delta(x, y) - W_{\varepsilon,h}^\delta(x, y')| \leq 2c_0 \left(1 + \frac{1}{\varepsilon}\right) \delta + 2|W_{\varepsilon,h}(x, \cdot) - W_h(x)| \\ \leq 2c_0 \left(1 + \frac{1}{\varepsilon}\right) \delta + 2\omega(\varepsilon) \end{array} \right. \quad (26)$$

where c_0 is independent of δ and ε . Moreover, since $D_x W_{\varepsilon,h}^\delta = \frac{1}{\delta} W_{\varepsilon,h,\delta} * D_x \psi_\delta$, one gets

$$|D_x W_{\varepsilon,h}^\delta(x, y) - D_x W_{\varepsilon,h}^\delta(x, y')| \leq \frac{1}{\delta} 2\omega(\varepsilon).$$

Similar assertions are valid for $|D_x^2 W_{\varepsilon,h}^\delta(x, y) - D_x^2 W_{\varepsilon,h}^\delta(x, y')|$. The approach equally works for the time dependent problem $W_{\varepsilon,h}^\delta(t, x, y), W_h(t, x)$ (see Remark 2.2). Also (see [21], Theorem 2.1, estimate 2.3) one can prove that

$$\|W_{\varepsilon,h}^\delta\|_\infty + \|\partial_t W_{\varepsilon,h}^\delta\|_\infty + \|DW_{\varepsilon,h}^\delta\|_\infty + \|D^2 W_{\varepsilon,h}^\delta\|_\infty \leq c_0 \frac{1}{\delta^2}, \quad (27)$$

where c_0 depends only on T (but not on δ).

3.2 Linear formulations for the averaged system

As previously, let us consider that $T > 0$ is a fixed time horizon. We fix $\varepsilon > 0$ and $(x_0, y_0) \in \mathbb{R}^M \times \mathbb{R}^N$. To every $\pi \in \mathcal{U}^w$, one can associate a couple of occupation measures $\gamma_{x_0, y_0, \pi; \varepsilon} = (\gamma_{x_0, y_0, \pi; \varepsilon}^1, \gamma_{x_0, y_0, \pi; \varepsilon}^2) \in \mathcal{P}([0, T] \times \mathbb{R}^M \times \mathbb{R}^N \times U) \times \mathcal{P}(\mathbb{R}^M \times \mathbb{R}^N)$ defined by

$$\left\{ \begin{array}{l} \gamma_{x_0, y_0, \pi; \varepsilon}^1(A \times B \times C \times D) = \frac{1}{T} \mathbb{E}^\pi \left[\int_0^T 1_{A \times B \times C \times D}(s, X_s^{x_0, y_0, u^\pi; \varepsilon}, Y_s^{x_0, y_0, u^\pi; \varepsilon}, u_s^\pi) ds \right], \\ \gamma_{x_0, y_0, \pi; \varepsilon}^2(E \times F) = \mathbb{E}^\pi \left[1_{E \times F} \left(X_T^{x_0, y_0, u^\pi; \varepsilon}, Y_T^{x_0, y_0, u^\pi; \varepsilon} \right) \right], \end{array} \right.$$

for all Borel sets $A \subset [0, T]$, $B \subset \mathbb{R}^M$, $C \subset \mathbb{R}^N$ and $D \subset U$. The family of occupation measures associated to weak controls

$$\Gamma(x_0, y_0; \varepsilon) := \left\{ (\gamma_{x_0, y_0, \pi; \varepsilon}^1, \gamma_{x_0, y_0, \pi; \varepsilon}^2), \text{ for all } \pi \in \mathcal{U}^w \right\} \quad (28)$$

can be embedded into a larger set

$$\Theta(x_0, y_0; \varepsilon) = \left\{ \begin{array}{l} (\gamma^1, \gamma^2) \in \mathcal{P}([0, T] \times \mathbb{R}^M \times \mathbb{R}^N \times U) \times \mathcal{P}(\mathbb{R}^M \times \mathbb{R}^N) \\ \forall \phi \in C_b^{1,2}(\mathbb{R}_+ \times \mathbb{R}^M \times \mathbb{R}^N), \\ \int_{[0, T] \times \mathbb{R}^M \times \mathbb{R}^N \times U \times \mathbb{R}^M \times \mathbb{R}^N} \left(\begin{array}{c} \phi(0, x_0, y_0) + T \mathcal{L}^{u; \varepsilon} \phi(s, x, y) \\ -\phi(T, z, w) \end{array} \right) \gamma^1(ds dx dy du) \gamma^2(dz dw) = 0. \end{array} \right\}, \quad (29)$$

where

$$\begin{aligned} \mathcal{L}^{u; \varepsilon} \phi(s, x, y) &= \frac{1}{2} \text{Tr} [(\sigma \sigma^*)(x, y, u) D_x^2 \phi(s, x, y)] + \frac{1}{2\varepsilon} \text{Tr} [(\beta \beta^*)(x, y, u) D_y^2 \phi(s, x, y)] \\ &+ \langle f(x, y, u), D_x \phi(s, x, y) \rangle + \frac{1}{\varepsilon} \langle g(x, y, u), D_y \phi(s, x, y) \rangle + \partial_t \phi(s, x, y), \end{aligned}$$

for all $\phi \in C^{1,2}(\mathbb{R}_+ \times \mathbb{R}^M \times \mathbb{R}^N)$ and all $s \geq 0$, $(x, y) \in \mathbb{R}^M \times \mathbb{R}^N$, $u \in U$.

Remark 3.3 *Using similar arguments as in the previous sections, the set $\Theta(x_0, y_0; \varepsilon)$ contains all occupation measures issued from (x_0, y_0) at time t . Moreover, it is also convex and relatively compact with respect to the weak convergence of probability measures (due to Prohorov's Theorem).*

Throughout the remaining of the paper, h is assumed to be bounded and Lipschitz-continuous. The linearized value function is given by

$$\Lambda_{\varepsilon, h}(x_0, y_0) = \inf_{\gamma = (\gamma^1, \gamma^2) \in \Theta(x_0, y_0; \varepsilon)} \int_{\mathbb{R}^M \times \mathbb{R}^N} h(z) \gamma^2(dz dw),$$

and its dual by

$$\eta_{\varepsilon,h}(x_0, y_0) = \sup \left\{ \begin{array}{l} \eta \in \mathbb{R} : \exists \phi \in C_b^{1,2}(\mathbb{R}_+ \times \mathbb{R}^M \times \mathbb{R}^N) \text{ s.t.} \\ \forall (s, x, y, v, z, w) \in [0, T] \times \mathbb{R}^M \times \mathbb{R}^N \times U \times \mathbb{R}^M \times \mathbb{R}^N, \\ \eta \leq T \mathcal{L}^{v;\varepsilon} \phi(s, x, y) + h(z) - \phi(T, z, w) + \phi(0, x_0, y_0). \end{array} \right\}, \quad (30)$$

for all $(x_0, y_0) \in \mathbb{R}^M \times \mathbb{R}^N$. This is a particular case of systems considered in subsection 2.2. Hence, for every $\varepsilon > 0$, one gets (applying Theorem 2.1),

$$W_{\varepsilon,h}(x_0, y_0) = \Lambda_{\varepsilon,h}(x_0, y_0) = \eta_{\varepsilon,h}(x_0, y_0),$$

for all initial data $(x_0, y_0) \in \mathbb{R}^M \times \mathbb{R}^N$.

At this point, we wish to give the intuition leading to the linear formulation for the averaged problem : if one thinks of the y component as being some penalization term, as $\varepsilon \rightarrow 0$, the corresponding part in $\mathcal{L}^{u;\varepsilon}$ should be 0 on the support of admissible measures. For the remaining component, y would be indifferent. We denote by

$$\Theta(x_0, y_0) = \left\{ \begin{array}{l} \gamma = (\gamma^1, \gamma^2) \in \mathcal{P}([0, T] \times \mathbb{R}^M \times \mathbb{R}^N \times U) \times \mathcal{P}(\mathbb{R}^M \times \mathbb{R}^N) : \\ \exists \gamma_\varepsilon \in \Theta(x_0, y_0; \varepsilon), \gamma_\varepsilon \rightarrow \gamma \text{ along some subsequence } \varepsilon_n \xrightarrow{n \rightarrow \infty} 0 \end{array} \right\},$$

for all $(x_0, y_0) \in \mathbb{R}^M \times \mathbb{R}^N$. Whenever $\gamma_\varepsilon = (\gamma_\varepsilon^1, \gamma_\varepsilon^2) \in \Theta(x_0, y_0; \varepsilon)$ for all $\varepsilon > 0$, one can find a subsequence (still indexed by $\varepsilon > 0$, for notation purposes) and a probability measure γ such that $\gamma_\varepsilon \rightarrow \gamma$ (cf. (A1) and Prohorov's theorem). Hence, the set $\Theta(x_0, y_0)$ is nonempty. One can also prove that it is closed (see Corollary 14).

Proposition 3.1 *The following inclusion holds true*

$$\Theta(x_0, y_0) \subset \left\{ \begin{array}{l} (\gamma^1, \gamma^2) \in \mathcal{P}([0, T] \times \mathbb{R}^M \times \mathbb{R}^N \times U) \times \mathcal{P}(\mathbb{R}^M \times \mathbb{R}^N) \text{ s.t.} \\ \forall \psi \in C_b^{1,2}(\mathbb{R}_+ \times \mathbb{R}^M) \text{ and } \forall \phi \in C_b^{1,2}(\mathbb{R}_+ \times \mathbb{R}^M \times \mathbb{R}^N), \\ \int_{[0, T] \times \mathbb{R}^M \times \mathbb{R}^N \times U \times \mathbb{R}^M \times \mathbb{R}^N} \begin{pmatrix} \psi(0, x_0) + T \mathcal{L}^{u,f} \psi(s, x, y) \\ -\psi(T, z) \end{pmatrix} \gamma^1(ds dx dy du) \gamma^2(dz dw) = 0 \text{ and} \\ \int_{[0, T] \times \mathbb{R}^M \times \mathbb{R}^N \times U \times \mathbb{R}^M \times \mathbb{R}^N} \mathcal{L}^{u,g} \phi(s, x, y) \gamma^1(ds dx dy du) \gamma^2(dz dw) = 0 \end{array} \right\},$$

where

$$\mathcal{L}^{u,f}\psi(s,x,y) = \frac{1}{2}Tr[(\sigma\sigma^*)(x,y,u)D^2\psi(s,x)] + \langle f(x,y,u), D_x\psi(s,x) \rangle + \partial_t\psi(s,x)$$

and

$$\mathcal{L}^{u,g}\phi(s,x,y) = \frac{1}{2}Tr[(\beta\beta^*)(x,y,u)D^2\phi(s,x,y)] + \langle g(x,y,u), D_y\phi(s,x,y) \rangle,$$

for all $\phi \in C^{1,2}(\mathbb{R}_+ \times \mathbb{R}^M \times \mathbb{R}^N)$, $\psi \in C^{1,2}(\mathbb{R}_+ \times \mathbb{R}^M)$ and all $s \geq 0$, $(x,y) \in \mathbb{R}^M \times \mathbb{R}^N$, $u \in U$.

Proof. Let us fix $\gamma \in \Theta(x_0, y_0)$ and $\gamma_\varepsilon = (\gamma_\varepsilon^1, \gamma_\varepsilon^2) \in \Theta(x_0, y_0; \varepsilon)$ such that $\gamma_\varepsilon \rightarrow \gamma$. Whenever $\psi \in C_b^{1,2}(\mathbb{R}_+ \times \mathbb{R}^M)$, the definition of $\Theta(x_0, y_0; \varepsilon)$ yields

$$\int_{[0,T] \times \mathbb{R}^M \times \mathbb{R}^N \times U \times \mathbb{R}^M \times \mathbb{R}^N} [\psi(0, x_0) + T\mathcal{L}^{u,f}\psi(s, x, y) - \psi(T, z)] \gamma_\varepsilon^1(dsdx dy du) \gamma_\varepsilon^2(dzdw) = 0.$$

Moreover, if one considers any fixed (though arbitrary) $\phi \in C_b^{1,2}(\mathbb{R}_+ \times \mathbb{R}^M \times \mathbb{R}^N)$, then

$$\begin{aligned} & \int_{[0,T] \times \mathbb{R}^M \times \mathbb{R}^N \times U \times \mathbb{R}^M \times \mathbb{R}^N} \mathcal{L}^{u,g}\phi(s, x, y) \gamma_\varepsilon^1(dsdx dy du) \gamma_\varepsilon^2(dzdw) \\ &= -\varepsilon \int_{[0,T] \times \mathbb{R}^M \times \mathbb{R}^N \times U \times \mathbb{R}^M \times \mathbb{R}^N} [\phi(0, x_0, y_0) + T\mathcal{L}^{u,f}\phi(s, x, y) - \phi(T, z, w)] \gamma_\varepsilon^1(dsdx dy du) \gamma_\varepsilon^2(dzdw) \end{aligned}$$

and the conclusion follows by letting $\varepsilon \rightarrow 0$ and recalling that $\phi \in C_b^{1,2}(\mathbb{R}_+ \times \mathbb{R}^M \times \mathbb{R}^N)$, resp. $\psi \in C_b^{1,2}(\mathbb{R}_+ \times \mathbb{R}^M)$. \square

We define the following linearized problem

$$\Lambda_h(x_0, y_0) = \inf_{\gamma=(\gamma^1, \gamma^2) \in \Theta(x_0, y_0)} \int_{\mathbb{R}^M \times \mathbb{R}^N} h(z) \gamma^2(dzdw),$$

and denote by

$$\eta_h(x_0) = \sup \left\{ \begin{array}{l} \eta \in \mathbb{R} : \exists \alpha \in C(\mathbb{R}_+; \mathbb{R}_+), \lim_{\varepsilon \rightarrow 0} \alpha(\varepsilon) = 0 \text{ s.t. } \forall \varepsilon > 0, \\ \exists \phi \in C_b^{1,2}(\mathbb{R}_+ \times \mathbb{R}^M \times \mathbb{R}^N) \text{ s.t.} \\ \sup_{y, y' \in \mathbb{R}^N} \|\phi(\cdot, \cdot, y) - \phi(\cdot, \cdot, y')\|_\infty \leq \alpha(\varepsilon) \text{ and s.t.} \\ \forall (s, x, y, v, z) \in [0, T] \times \mathbb{R}^M \times \mathbb{R}^N \times U \times \mathbb{R}^M, \\ \eta \leq T\mathcal{L}^{v;\varepsilon}\phi(s, x, y) + h(z) + \|\phi(T, z, \cdot)\|_\infty + \|\phi(0, x_0, \cdot)\|_\infty \end{array} \right\}, \quad (31)$$

for all $(x_0, y_0) \in \mathbb{R}^M \times \mathbb{R}^N$.

Remark 3.4 In the previous definition one can, equivalently, ask that $\|\phi(\cdot, \cdot, \cdot) - \phi(\cdot, \cdot, y_0)\|_\infty \leq \alpha(\varepsilon)$ for some fixed $y_0 \in \mathbb{R}^M$.

Consequently, we can formulate the main result of this section:

Theorem 3.1 We assume (A1) and (25) to hold true. Moreover, we assume the invariance condition (4) to be satisfied. Then the following equalities hold true

$$W_h(x_0) = \Lambda_h(x_0, y_0) = \eta_h(x_0),$$

for all $(x_0, y_0) \in K \times \mathbb{R}^N$.

Remark 3.5 As we have hinted in the previous subsection, whenever the Assumptions 1 - 3 hold true, then (25) holds true (cf. [4] (see also [5])).

Proof. Let us fix $(x_0, y_0) \in K \times \mathbb{R}^N$. In a first step, we recall that there exists an optimal measure $\bar{\gamma}_{(x_0, y_0); \varepsilon} = (\bar{\gamma}_\varepsilon^1, \bar{\gamma}_\varepsilon^2) \in \Theta(x_0, y_0; \varepsilon)$ such that

$$\Lambda_{\varepsilon, h}(x_0, y_0) = \int_{\mathbb{R}^M \times \mathbb{R}^N} h(z) \bar{\gamma}_\varepsilon^2(dzdw),$$

for all $\varepsilon > 0$. One can find a subsequence (still indexed by $\varepsilon > 0$, for notation purposes) and a probability measure γ such that $\bar{\gamma}_\varepsilon \rightarrow \gamma$ (cf. (A1) and Prohorov's theorem). Consequently,

$$\begin{aligned} \Lambda_h(x_0, y_0) &\leq \int_{\mathbb{R}^M \times \mathbb{R}^N} h(z) \gamma^2(dzdw) = \lim_{\varepsilon \rightarrow 0} \int_{\mathbb{R}^M \times \mathbb{R}^N} h(z) \bar{\gamma}_\varepsilon^2(dzdw) \\ &= \lim_{\varepsilon \rightarrow 0} \Lambda_{\varepsilon, h}(x_0, y_0) = \lim_{\varepsilon \rightarrow 0} W_{\varepsilon, h}(x_0, y_0) = W_h(x_0). \end{aligned} \quad (32)$$

for all $(x_0, y_0) \in \mathbb{R}^M \times \mathbb{R}^N$. The converse inequality is similar.

We continue by considering $\gamma \in \Theta(x_0, y_0)$ and $\eta \in \mathbb{R}$ such that

$$\begin{aligned} \exists \alpha \in C(\mathbb{R}_+; \mathbb{R}_+) \text{ with } \lim_{\varepsilon \rightarrow 0} \alpha(\varepsilon) = 0, \text{ s.t. } \forall \varepsilon > 0, \exists \phi \in C_b^{1,2}(\mathbb{R}_+ \times \mathbb{R}^M \times \mathbb{R}^N) \text{ s.t.} \\ \sup_{y, y' \in \mathbb{R}^N} \|\phi(\cdot, \cdot, y) - \phi(\cdot, \cdot, y')\|_\infty \leq \alpha(\varepsilon) \text{ and } \forall (s, x, y, v, z) \in [0, T] \times \mathbb{R}^M \times \mathbb{R}^N \times U \times \mathbb{R}^M, \\ \eta \leq T\mathcal{L}^{v; \varepsilon} \phi(s, x, y) + h(z) - \inf_{y' \in \mathbb{R}^N} \phi(T, z, y') + \sup_{y' \in \mathbb{R}^N} \phi(0, x_0, y') \end{aligned}$$

Then,

$$\eta \leq T\mathcal{L}^{v; \varepsilon} \phi(s, x, y) + h(z) - \phi(T, z, w) + \phi(0, x_0, y_0) + 2\alpha(\varepsilon), \quad (33)$$

for all $\forall (s, x, y, v, z, w) \in [0, T] \times \mathbb{R}^M \times \mathbb{R}^N \times U \times \mathbb{R}^M \times \mathbb{R}^N$. By the definition of $\Theta(x_0, y_0)$, there exists some sequence $\gamma_\varepsilon \in \Theta(x_0, y_0; \varepsilon)$ converging to γ . By integrating with respect to γ_ε the inequality (33), we obtain that

$$\eta \leq \int_{\mathbb{R}^M \times \mathbb{R}^N} h(z) \gamma_\varepsilon^2(dzdw) + 2\alpha(\varepsilon),$$

and, consequently, recalling that $\gamma \in \Theta(x_0, y_0)$, $\varepsilon > 0$ are arbitrary and $\lim_{\varepsilon \rightarrow 0} \alpha(\varepsilon) = 0$, it follows that

$$\eta_h(x_0) \leq \Lambda_h(x_0, y_0). \quad (34)$$

Let $\varepsilon > 0$ be fixed. Using Proposition 2.1 (see Remark 3.2 for the specific details; in particular the inequality (26)), there exists a family of functions $W_{\varepsilon, h}^\delta \in C_b^{1,2}([0, T + \delta^2] \times \mathbb{R}^{M+N})$ such that, for every $(s, x, y, v, z, w) \in [t, T] \times \mathbb{R}^M \times \mathbb{R}^N \times U \times \mathbb{R}^M \times \mathbb{R}^N$,

$$\begin{aligned} \mathcal{L}^{v; \varepsilon} W_{\varepsilon, h}^\delta(s, x, y) &\geq 0 \text{ and} \\ h(z) - W_{\varepsilon, h}^\delta(T, z, w) &\geq h(z) - W_{\varepsilon, h}(T, z, w) - c_0 \left(1 + \frac{1}{\varepsilon}\right) \delta \geq -c_0 \left(1 + \frac{1}{\varepsilon}\right) \delta. \end{aligned}$$

Hence,

$$\begin{aligned} W_{\varepsilon, h}^\delta(0, x_0, y_0) - c_0 \left(1 + \frac{1}{\varepsilon}\right) \delta &\leq \mathcal{L}^{v; \varepsilon} W_{\varepsilon, h}^\delta(s, x, y) + h(z) - W_{\varepsilon, h}^\delta(T, z, w) + W_{\varepsilon, h}^\delta(0, x_0, y_0) \\ &\leq \mathcal{L}^{v; \varepsilon} W_{\varepsilon, h}^\delta(s, x, y) + h(z) - \inf_w W_{\varepsilon, h}^\delta(T, z, w) + \sup_w W_{\varepsilon, h}^\delta(0, x_0, w) \end{aligned} \quad (35)$$

Thus, $W_{\varepsilon, h}^{\varepsilon^2}(0, x_0, y_0) - c_0 \left(1 + \frac{1}{\varepsilon}\right) \varepsilon^2 \leq \eta_h(x_0)$. The first inequality in (26) and (25) yield that

$$\begin{aligned} \left| W_{\varepsilon, h}^{\varepsilon^2}(0, x_0, y_0) - W_h(x_0) \right| &\leq \left| W_{\varepsilon, h}^{\varepsilon^2}(0, x_0, y_0) - W_{\varepsilon, h}(x_0) \right| + |W_{\varepsilon, h}(x_0, y_0) - W_h(x_0)| \\ &\leq c_0 \left(1 + \frac{1}{\varepsilon}\right) \varepsilon^2 + \omega(\varepsilon). \end{aligned} \quad (36)$$

Consequently, passing to the limit as $\varepsilon \rightarrow 0$, we get

$$W_h(x_0) \leq \eta_h(x_0). \quad (37)$$

By combining the inequalities (34) and (37) and recalling we have already proven that $W_h(x_0) = \Lambda_h(x_0, y_0)$, we complete the proof. \square

Remark 3.6 *If the estimates in (26) are independent of ε (e.g. by imposing a dissipativity condition on (g, β)), then one can prove that Λ_h can be defined with respect to the (explicit) set appearing in Proposition*

3.1.

A careful look at the proof (cf. (35) and (36)) tells us that

$$\begin{aligned}
W_h(x_0) &= \lim_{n \rightarrow \infty} W_{\frac{1}{n}, h}^{\frac{1}{n^2}}(0, x_0, y_0) \\
&\leq \liminf_{n \rightarrow \infty} \inf_{(s, x, y, v, z, w) \in [t, T] \times K \times \mathbb{R}^N \times U \times K \times \mathbb{R}^N} \left(\begin{array}{c} \mathcal{L}^{v; \frac{1}{n}} W_{\frac{1}{n}, h}^{\frac{1}{n^2}}(s, x, y) + h(z) \\ -W_{\frac{1}{n}, h}^{\frac{1}{n^2}}(T, z, w) + W_{\frac{1}{n}, h}^{\frac{1}{n^2}}(0, x_0, y_0) \end{array} \right) \quad (38)
\end{aligned}$$

In particular, we deduce that $\Theta(x_0, y_0)$ can be replaced with

$$\tilde{\Theta}(x_0, y_0) = \left\{ \begin{array}{l} \gamma = (\gamma^1, \gamma^2) \in \mathcal{P}([0, T] \times \mathbb{R}^M \times \mathbb{R}^N \times U) \times \mathcal{P}(\mathbb{R}^M \times \mathbb{R}^N) : \\ \exists \gamma_n \in \Theta(x_0, y_0; \frac{1}{n}), \gamma_n \rightarrow \gamma \text{ along some subsequence} \end{array} \right\}. \quad (39)$$

Moreover, if γ_n is an optimal measure for $W_{\frac{1}{n}, h}$, one can find a subsequence converging to an optimal measure in $\Theta(x_0, y_0)$. Hence, one can also replace $\Theta(x_0, y_0)$ with

$$\Theta^{opt}(x_0, y_0) = \left\{ \begin{array}{l} \gamma = (\gamma^1, \gamma^2) \in \mathcal{P}([0, T] \times \mathbb{R}^M \times \mathbb{R}^N \times U) \times \mathcal{P}(\mathbb{R}^M \times \mathbb{R}^N) : \\ \exists \gamma_n \in \Theta(x_0, y_0; \frac{1}{n}), \gamma_n \text{ is optimal for } W_{\frac{1}{n}, h} \\ \left(\text{i.e.} \int_{[0, T] \times \mathbb{R}^M \times \mathbb{R}^N \times U \times \mathbb{R}^M \times \mathbb{R}^N} h(z) \gamma_n(ds dx dy dz dw) = W_{\frac{1}{n}, h}(x_0, y_0) \right), \\ \gamma_n \rightarrow \gamma \text{ along some subsequence.} \end{array} \right\}. \quad (40)$$

3.3 Characterization of optimal trajectories for the averaged system

As already mentioned in the introduction, when the perturbed system is fully nonlinear it is very difficult to characterize the optimal trajectories using the Pontryagin maximum principle because we do not know exactly the form of the averaged dynamics. An alternative to this method is to look at the support of the occupational measures contained in the set $\Theta(x_0, y_0)$ in order to obtain optimal trajectories from every $x_0 \in K$. Following the approach already introduced in Subsection 2.2, we denote by

$$D_{\varepsilon, h}(x_0, y_0) = \left\{ \begin{array}{l} (\eta, \phi) \in \mathbb{R} \times C_b^{1,2}(\mathbb{R}_+ \times \mathbb{R}^M \times \mathbb{R}^N) \text{ s.t.} \\ \eta = \inf_{(s, x, y, v, z, w) \in [t, T] \times \mathbb{R}^M \times \mathbb{R}^N \times U \times \mathbb{R}^M \times \mathbb{R}^N} \left(\begin{array}{c} T\mathcal{L}^{v; \varepsilon} \phi(s, x, y) + h(z) \\ -\phi(T, z, w) + \phi(0, x_0, y_0) \end{array} \right) \end{array} \right\}, \quad (41)$$

for all $(x_0, y_0) \in K \times \mathbb{R}^N$. We can write

$$W_{\varepsilon, h}(x_0, y_0) = \sup \{ \eta, (\eta, \phi) \in D_{\varepsilon, h}(x_0, y_0) \} \text{ and}$$

$$W_h(x_0) = \sup \left\{ \begin{array}{l} \limsup_{\varepsilon \rightarrow 0} \eta_\varepsilon : (\eta_\varepsilon, \phi_\varepsilon) \in D_{\varepsilon, h}(x_0, y_0), \\ \lim_{\varepsilon \rightarrow 0} \|\phi_\varepsilon(\cdot, \cdot, \cdot) - \phi_\varepsilon(\cdot, \cdot, y_0)\|_\infty = 0 \end{array} \right\}.$$

At this point, we pick $\left(\eta_n, W_{\frac{1}{n}, h}^{\frac{1}{n^2}} \right) \in D_{\frac{1}{n}, h}(x_0, y_0)$ and recall that

$$\left| W_{\varepsilon, h}^{\varepsilon^2}(\cdot, \cdot, \cdot) - W_{\varepsilon, h}^{\varepsilon^2}(\cdot, \cdot, y_0) \right|_\infty \leq 2c_0 \left(1 + \frac{1}{\varepsilon} \right) \varepsilon^2 + 2\omega(\varepsilon),$$

(cf. the second inequality in (26)). Then $W_h(x_0) \geq \limsup_{n \rightarrow \infty} \eta_n$. Combining this inequality with (38) yields

$$W_h(x_0) = \lim_{n \rightarrow \infty} \inf_{(s, x, y, v, z, w) \in [t, T] \times \mathbb{R}^M \times \mathbb{R}^N \times U \times \mathbb{R}^M \times \mathbb{R}^N} \left(\begin{array}{l} \mathcal{L}^{v; \frac{1}{n}} W_{\frac{1}{n}, h}^{\frac{1}{n^2}}(s, x, y) + h(z) \\ -W_{\frac{1}{n}, h}^{\frac{1}{n^2}}(T, z, w) + W_{\frac{1}{n}, h}^{\frac{1}{n^2}}(0, x_0, y_0) \end{array} \right).$$

According to the approach of Subsection 2.2, we introduce the following.

Whenever $(x_0, y_0) \in K \times \mathbb{R}^N$, we denote by

$$\Omega_{\frac{1}{n}, h}^{simple}(x_0, y_0) = \left\{ \begin{array}{l} (s, x, y, v, z, w) \in [t, T] \times \mathbb{R}^M \times \mathbb{R}^N \times U \times \mathbb{R}^M \times \mathbb{R}^N \text{ s.t.} \\ W_h(x_0) + \sqrt{|W_h(x_0) - \eta_n|} \\ \geq \mathcal{L}^{v; \frac{1}{n}} W_{\frac{1}{n}, h}^{\frac{1}{n^2}}(s, x, y) + h(z) - W_{\frac{1}{n}, h}^{\frac{1}{n^2}}(T, z, w) + W_{\frac{1}{n}, h}^{\frac{1}{n^2}}(0, x_0, y_0) \end{array} \right\} \quad (42)$$

$$\Omega_{\frac{1}{n}, h}^{double}(x_0, y_0) = \left\{ \begin{array}{l} (s, x, y, v, z, w) \in [t, T] \times \mathbb{R}^M \times \mathbb{R}^N \times U \times \mathbb{R}^M \times \mathbb{R}^N \text{ s.t.} \\ W_{\frac{1}{n}, h}(x_0) + \sqrt{W_{\frac{1}{n}, h}(x_0) - \eta_n} \\ \geq \mathcal{L}^{v; \frac{1}{n}} W_{\frac{1}{n}, h}^{\frac{1}{n^2}}(s, x, y) + h(z) - W_{\frac{1}{n}, h}^{\frac{1}{n^2}}(T, z, w) + W_{\frac{1}{n}, h}^{\frac{1}{n^2}}(0, x_0, y_0) \end{array} \right\} \quad (43)$$

and by

$$\Omega_h^{in}(x_0, y_0) := \bigcup_{n \geq 1} \bigcap_{k \geq n} \Omega_{\frac{1}{k}, h}^{double}(x_0, y_0), \quad \Omega_h^{out, cl}(x_0, y_0) := \bigcap_{n \geq 1} cl \left(\bigcup_{k \geq n} \Omega_{\frac{1}{k}, h}^{double}(x_0, y_0) \right).$$

We get the following criteria of optimality.

Proposition 3.2 *Let $(x_0, y_0) \in K \times \mathbb{R}^N$ be fixed.*

(i) If $\gamma_n \in \Theta(x_0, y_0; \frac{1}{n})$ is a (sub)sequence such that

$$\lim_{n \rightarrow \infty} n^3 \gamma_n \left(\left(\Omega_{\frac{1}{n}, h}^{simple}(x_0, y_0) \right)^c \right) = 0,$$

then any limit of γ_n is optimal.

(ii) Every $\gamma \in \Theta^{opt}(x_0, y_0)$ is optimal for W_h and

$$\gamma \left(\Omega_h^{out, cl}(x_0, y_0) \right) = 1 \quad (44)$$

(i.e. the support of γ is included in $\Omega_h^{out, cl}(x_0, y_0)$). Moreover, if $\Omega_h^{in}(x_0, y_0) = \Omega_h^{out, cl}(x_0, y_0)$ (i.e. the limit of $\Omega_{\frac{1}{n}, h}(x_0, y_0)$ exists), then

$$\sup_{n \geq 1} \gamma \left(\bigcap_{k \geq n} \Omega_{\frac{1}{k}, h}^{double}(x_0, y_0) \right) = 1.$$

Remark 3.7 (i) The sufficient condition (ii) relies on finding (near) optimal measures $\gamma_n \in \Theta(x_0, y_0; \frac{1}{n})$.

(ii) The reader is invited to note that the condition (44) is the same as in the classical framework (cf. Proposition 2.3).

Proof. (of Proposition 3.2). (i) Let us fix $\Theta(x_0, y_0; \frac{1}{n}) \ni \gamma_n$ as in our assertion and converging (along some subsequence) to some γ . The inequality (27) yields

$$\mathcal{L}^{v; \frac{1}{n}} W_{\frac{1}{n}, h}^{\frac{1}{n^2}}(s, x, y) \leq c(1+n)n^2,$$

for some constant c independent of n . Then, recalling the definition of $\Omega_{\frac{1}{n}, h}^{simple}(x_0, y_0)$, we get

$$\begin{aligned} W_h(x_0) &\leq \int_{[0, T] \times \mathbb{R}^M \times \mathbb{R}^N \times U \times \mathbb{R}^M \times \mathbb{R}^N} h(z) \gamma(ds dx dy dz dw) \\ &= \lim_{n \rightarrow \infty} \int_{[0, T] \times \mathbb{R}^M \times \mathbb{R}^N \times U \times \mathbb{R}^M \times \mathbb{R}^N} h(z) \gamma_n(ds dx dy dz dw) \\ &\leq \lim_{n \rightarrow \infty} \int_{[0, T] \times \mathbb{R}^M \times \mathbb{R}^N \times U \times \mathbb{R}^M \times \mathbb{R}^N} \left[\begin{array}{c} T \mathcal{L}^{v; \frac{1}{n}} W_{\frac{1}{n}, h}^{\frac{1}{n^2}}(s, x, y) + h(z) \\ -W_{\frac{1}{n}, h}^{\frac{1}{n^2}}(T, z, w) + W_{\frac{1}{n}, h}^{\frac{1}{n^2}}(0, x_0, y_0) \end{array} \right] \gamma_n(ds dx dy dz dw) \\ &\leq \limsup_{n \rightarrow \infty} \left[W_h(x_0) + \sqrt{|W_h(x_0) - \eta_n|} + (3\|h\|_\infty + c(1+n)n^2) \gamma_n \left(\left(\Omega_{\frac{1}{n}, h}^{simple}(x_0, y_0) \right)^c \right) \right] \\ &= W_h(x_0). \end{aligned}$$

It follows that γ is optimal.

(ii) If $\gamma \in \Theta^{opt}(x_0, y_0)$, then $\gamma \in \tilde{\Theta}(x_0, y_0)$ is the limit of some (sub)sequence $\gamma_n \in \Theta(x_0, y_0; \frac{1}{n})$ of optimal measures for $W_{\frac{1}{n}, h}(x_0, y_0)$ (cf. (40)). It is obvious that

$$\begin{aligned} & \int_{[0, T] \times \mathbb{R}^M \times \mathbb{R}^N \times U \times \mathbb{R}^M \times \mathbb{R}^N} h(z) \gamma(ds dx dy dz dw) \\ &= \lim_{n \rightarrow \infty} \int_{[0, T] \times \mathbb{R}^M \times \mathbb{R}^N \times U \times \mathbb{R}^M \times \mathbb{R}^N} h(z) \gamma_n(ds dx dy dz dw) = \lim_{n \rightarrow \infty} W_{\frac{1}{n}, h}(x_0, y_0), \end{aligned}$$

hence γ is optimal. Since $\gamma_n \in \Theta(x_0, y_0; \frac{1}{n})$ is optimal for $W_{\frac{1}{n}, h}(x_0, y_0)$, it follows that

$$\begin{aligned} W_{\frac{1}{n}, h}(x_0) &= \int_{[0, T] \times \mathbb{R}^M \times \mathbb{R}^N \times U \times \mathbb{R}^M \times \mathbb{R}^N} h(z) \gamma_n(ds dx dy dz dw) \\ &= \int_{[0, T] \times \mathbb{R}^M \times \mathbb{R}^N \times U \times \mathbb{R}^M \times \mathbb{R}^N} \begin{bmatrix} T \mathcal{L}^{v; \frac{1}{n}} W_{\frac{1}{n}, h}^{\frac{1}{2}}(s, x, y) + h(z) \\ -W_{\frac{1}{n}, h}^{\frac{1}{2}}(T, z, w) + W_{\frac{1}{n}, h}^{\frac{1}{2}}(0, x_0, y_0) \end{bmatrix} \gamma_n(ds dx dy dz dw) \\ &\geq \eta_n \gamma_n \left(\Omega_{\frac{1}{n}, h}^{double}(x_0, y_0) \right) + \left(W_{\frac{1}{n}, h}(x_0) + \sqrt{W_{\frac{1}{n}, h}(x_0) - \eta_n} \right) \gamma_n \left(\Omega_{\frac{1}{n}, h}^{double}(x_0, y_0) \right). \end{aligned}$$

Then

$$\gamma_n \left(\Omega_{\frac{1}{n}, h}^{double}(x_0, y_0) \right) \geq \frac{1}{1 + \sqrt{W_{\frac{1}{n}, h}(x_0) - \eta_n}},$$

for all $n \geq 1$. Hence,

$$\gamma_n \left(cl \left(\bigcup_{k \geq n_0} \Omega_{\frac{1}{k}, h}^{double}(x_0, y_0) \right) \right) \geq \frac{1}{\left(1 + \sqrt{W_{\frac{1}{n}, h}(x_0) - \eta_n} \right)},$$

for all $n \geq n_0$. Passing to the $\lim_{n \rightarrow \infty}$, one gets $\gamma \left(cl \left(\bigcup_{k \geq n_0} \Omega_{\frac{1}{k}, h}^{double}(x_0, y_0) \right) \right) = 1$, for all $n_0 \geq 1$ and the proof is complete. \square

Remark 3.8 (i) If a suitable monotonicity can be established for the approximating problems $W_{\frac{1}{n}, h}(x_0, y_0)$, then, one can envisage the use of the dual formulation in Theorem 3.1 to infer necessary conditions similar to those in Proposition 2.3.

(ii) When the inclusion in Proposition (3.1) is an equality (see Remark 3.6 (i)), one can employ convex duality arguments to get another dual formulation for the limit value. One can, for instance, adapt the same methods as [6], Theorem 1. This dual formulation would be very similar to the classical case and the ingredients of Proposition 2.3 apply. The main drawback in this approach is that, unlike the classical case, we have no information on the structure of the test functions ϕ_n in the (almost-) optimal pairs.

References

- [1] M. Bardi and I. Capuzzo Dolcetta. *Optimal control and viscosity solutions of Hamilton-Jacobi-Bellman equations*. Systems and Control: Foundations and Applications, Birkhäuser, Boston, 1997.
- [2] M. Bardi and P. Goatin. Invariant sets for controlled degenerate diffusions: A viscosity solutions approach. In WilliamM. McEneaney, G.George Yin, and Qing Zhang, editors, *Stochastic Analysis, Control, Optimization and Applications*, Systems Control: Foundations and Applications, pages 191–208. Birkhäuser Boston, 1999.
- [3] A.G. Bhatt and V.S. Borkar. Occupation measures for controlled Markov processes: Characterization and optimality. *Ann. of Probability*, 24:1531–1562, 1996.
- [4] V. Borkar and V. Gaitsgory. Averaging of singularly perturbed controlled stochastic differential equations. *Appl. Math. Optimization*, 56(2):169–209, 2007.
- [5] Vivek Borkar and Vladimir Gaitsgory. On existence of limit occupational measures set of a controlled stochastic differential equation. *SIAM J. Control Optim.*, 44(4):1436–1473 (electronic), 2005.
- [6] R. Buckdahn, D. Goreac, and M. Quincampoix. Stochastic optimal control and linear programming approach. *Appl. Math. Optimization*, 63(2):257–276, 2011.
- [7] R. Buckdahn, D. Goreac, and M. Quincampoix. Existence of Asymptotic Values for Nonexpansive Stochastic Control Systems. *Applied Mathematics and Optimization*, pages 1–28, 2014.
- [8] R. Buckdahn, S. Peng, M. Quincampoix, and C. Rainer. Existence of stochastic control under state constraints. *C. R. Acad. Sci. Paris, Sér. I Math*, 327:17–22, 1998.
- [9] L. Finlay, V. Gaitsgory, and I. Lebedev. Linear programming solutions of periodic optimization problems: approximation of the optimal control. *J. Ind. Manag. Optim.*, 3(2):399–413, 2007.
- [10] W. Fleming and D. Vermes. Convex duality approach to the optimal control of diffusions. *SIAM J. Control Optimization*, 36(2):1136–1155, 1989.
- [11] E. Fridman. Exact slow-fast decomposition of a class of non-linear singularly perturbed optimal control problems via invariant manifolds. *International Journal of Control*, 72(17):1609–1618, 1999.
- [12] V. Gaitsgory. Suboptimization of singularly perturbed control systems. *SIAM J. Control and Optim.*, 30(5):1228–1249., 1992.

- [13] V. Gaitsgory. On a representation of the limit occupational measures set of a control system with applications to singularly perturbed control systems. *SIAM J. Control Optim.*, 43(1):325–340, 2004.
- [14] V. Gaitsgory and A. Leizarowitz. Limit occupational measures set for a control system and averaging of singularity perturbed control systems. *J. Math. Anal. Appl.*, 233(2):461–475, 1999.
- [15] V. Gaitsgory and M.-T. Nguyen. Multiscale singularly perturbed control systems: Limit occupational measures sets and averaging. *SIAM J. Control Optimization*, 41(3):954–974, 2002.
- [16] V. Gaitsgory and M. Quincampoix. Linear programming approach to deterministic infinite horizon optimal control problems with discounting. *SIAM J. Control Optimization*, 48(4):2480–2512, 2009.
- [17] V. Gaitsgory and S. Rossomakhine. Linear programming approach to deterministic long run average problems of optimal control. *SIAM J. Control Optim.*, 44(6):2006–2037, 2006.
- [18] D. Goreac and O.S. Serea. Mayer and optimal stopping stochastic control problems with discontinuous cost. *Journal of Mathematical Analysis and Applications*, 380(1):327–342, 2011.
- [19] D. Goreac and O.S. Serea. A note on linearization methods and dynamic programming principles for stochastic discontinuous control problems. *Electron. Commun. Probab.*, 17:no. 12, 1–12, 2012.
- [20] G. Grammel. Singularly perturbed differential inclusions. an averaging approach. *Set-valued Analysis*, 4(4):361–374, 1996.
- [21] N. V. Krylov. On the rate of convergence of finite-difference approximations for Bellman’s equations with variable coefficients. *Probab. Theory Related Fields*, 117(1):1–16, 2000.
- [22] D. Naidu. Singular perturbations and time scales in control theory and applications: an overview. *Dynamics of Continuous, Discrete and Impulsive Systems, Series B*, (24):233–278, 2002.
- [23] A.N. Tichonov. Systems of differential equations containing small parameter near derivatives. *Mat. Sbornik.*, 31:575–586, 1952.
- [24] V. Veliiov. A generalization of the tikhonov theorem for singularly perturbed differential inclusions. *J. Dynam. Control Systems*, 3(3):291–319, 1997.