



HAL
open science

Rate of convergence to equilibrium of fractional driven stochastic differential equations with some multiplicative noise

Joaquin Fontbona, Fabien Panloup

► To cite this version:

Joaquin Fontbona, Fabien Panloup. Rate of convergence to equilibrium of fractional driven stochastic differential equations with some multiplicative noise. 2014. hal-00989414v1

HAL Id: hal-00989414

<https://hal.science/hal-00989414v1>

Preprint submitted on 11 May 2014 (v1), last revised 14 Jan 2016 (v2)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Rate of convergence to equilibrium of fractional driven stochastic differential equations with some multiplicative noise

Joaquin Fontbona* and Fabien Panloup†

May 11, 2014

Abstract

We investigate the problem of the rate of convergence to equilibrium for ergodic stochastic differential equations driven by fractional Brownian motion with Hurst parameter $H > 1/2$ and multiplicative noise component σ . When σ is constant and for every $H \in (0, 1)$, it was proved in [9] that, under some mean-reverting assumptions, such a process converges to its equilibrium at a rate of order $t^{-\alpha}$ where $\alpha \in (0, 1)$ (depending on H). The aim of this paper is to extend such types of results to some multiplicative noise setting. More precisely, we show that we can recover such convergence rates when $H > 1/2$ and the inverse of the diffusion coefficient σ is a Jacobian matrix. The main novelty of this work is a type of extension of Foster-Lyapunov like techniques to this non-Markovian setting, which allows us to put in place an asymptotic coupling scheme as that of [9] without resorting to deterministic contracting properties.

Keywords: Stochastic Differential Equations; Fractional Brownian Motion; Multiplicative noise; Ergodicity; Rate of convergence to equilibrium; Lyapunov function; Total variation distance.

AMS classification (2010): 60G22, 37A25.

1 Introduction

Stochastic Differential Equations (SDEs) driven by a fractional Brownian motion (fBm) have been introduced to model random evolution phenomena whose noise has long range dependence properties. Indeed, beyond the historical motivations in Hydrology and Telecommunication for the use of fBm (highlighted e.g in [14]), recent applications of dynamical systems driven by this process include challenging issues in Finance [8], Biotechnology [17] or Biophysics [12, 13].

The study of the long-time behavior (under some stability properties) for fractional SDEs has been developed by Hairer [9], Hairer and Ohashi [10], and by Hairer and Pillai [11] (see also [1, 5, 7] for another setting called random dynamical systems and [2, 3] for some results of approximations of stationary solutions) who built a way to define stationary solutions of these a priori non-Markov processes and to extend some of the tools of the Markovian theory to this setting. In particular, criterions for uniqueness of the invariant distribution are proved in the three above papers in some different settings, respectively: additive noise, multiplicative noise with $H > 1/2$ and multiplicative noise with $H \in (1/3, 1/2)$ in the last one (in an hypoelliptic context).

When uniqueness holds for the invariant distribution, a challenging question is that of the rate of convergence to this equilibrium. In [9], the author proved that in the additive noise setting,

*Department of Mathematical Engineering and Center for Mathematical Modeling, UMI(2807) UCHILE-CNRS, Universidad de Chile, Casilla 170-3, Correo 3, Santiago-Chile. E-mail:fontbona@dim-uchile.cl.

†Institut de Mathématiques de Toulouse, Université Paul Sabatier & INSA Toulouse, 135, av. de Rangueil, F-31077 Toulouse Cedex 4, France. E-mail: fabien.panloup@math.univ-toulouse.fr

the process converges in total variation to the stationary regime with a rate upper-bounded by $C_\varepsilon t^{-(\alpha-\varepsilon)}$ for any $\varepsilon > 0$, with

$$\alpha = \begin{cases} \frac{1}{8} & \text{if } H \in (\frac{1}{4}, 1) \setminus \{\frac{1}{2}\} \\ H(1-2H) & \text{if } H \in (0, \frac{1}{4}]. \end{cases} \quad (1.1)$$

But, to the best of our knowledge, no result of rate of convergence exists in the multiplicative setting. The aim of the current paper is to extend the results of [9] in the multiplicative setting when $H > 1/2$.

More precisely, we deal with an \mathbb{R}^d -valued process $(X_t)_{t \geq 0}$ which is a solution to the following SDE

$$dX_t = b(X_t)dt + \sigma(X_t)dB_t \quad (1.2)$$

where $b : \mathbb{R}^d \rightarrow \mathbb{R}^d$ and $\sigma : \mathbb{R}^d \rightarrow \mathbb{M}_{d,d}$ are (at least) continuous functions, and where $\mathbb{M}_{d,d}$ is the set of $d \times d$ real matrices. In (1.2), $(B_t)_{t \geq 0}$ is a d -dimensional H -fBm with Hurst parameter $H \in (\frac{1}{2}, 1)$. Note that under some Hölder regularity assumptions on the coefficients (see *e.g.* [16, 4] for background), (strong) existence and uniqueness hold for the solution to (1.2) starting from $x_0 \in \mathbb{R}^d$.

Introducing the Mandelbrot-Van Ness representation of the fractional Brownian motion,

$$B_t = \alpha_H \int_{-\infty}^0 (-r)^{H-\frac{1}{2}} (dW_{r+t} - dW_r), \quad t \geq 0, \quad (1.3)$$

where $(W_t)_{t \in \mathbb{R}}$ is a two-sided \mathbb{R}^d -valued Brownian Motion and α_H is a normalization coefficient depending on H , $(X_t, (B_{s+t})_{s \leq 0})_{t \geq 0}$ can be realized through a Feller transformation $(\mathcal{Q}_t)_{t \geq 0}$ on the product space $\mathbb{R}^d \times \mathcal{W}_{\theta, \delta}$ ($\theta \in (1/2, H)$ and $\theta + \delta \in (H, 1)$) whose definition is recalled in (2.4) (we refer to [10] for more rigorous background on this topic). In particular, an initial distribution of this dynamical system is a distribution μ_0 on $\mathbb{R}^d \times \mathcal{W}_{\theta, \delta}$. With probabilistic words, an initial distribution is the distribution of a couple $(X_0, (B_s)_{s \leq 0})$ where $(B_s)_{s \leq 0}$ is an \mathbb{R}^d -valued fBm on $(-\infty, 0]$.

Then, such an initial distribution is classically called an invariant distribution if it is invariant by the transformation \mathcal{Q}_t for every $t \geq 0$. However, the concept of uniqueness of invariant distribution is slightly different of the classical setting. Actually, one says that uniqueness of the invariant distribution holds if the stationary regime, we mean, the distribution $\bar{Q}\mu$ of the whole process $(X_t^\mu)_{t \geq 0}$ with initial distribution μ , is unique (in other words, this concept of uniqueness corresponds to the classical one conditioned by the equivalence relation: $\mu \sim \nu \iff \bar{Q}\mu \sim \bar{Q}\nu$, see [10] for background). In harmony with the previous concept, coupling two paths issued of μ_0 and μ where the second one denotes an invariant distribution of $(\mathcal{Q}_t)_{t \geq 0}$ consists (classically) in finding a stopping time τ_∞ such that $(X_{t+\tau_\infty}^{\mu_0})_{t \geq 0} = (X_{t+\tau_\infty}^\mu)_{t \geq 0}$. Thus, the rate of convergence in total variation will be obtained by some bounds on $\mathbb{P}(\tau_\infty > t)$, $t \geq 0$.

Now, let us briefly recall the coupling strategy of [9]. First, one classically waits that the paths get close. Then, at each trial, the coupling attempt is divided in two steps. First, one tries in Step 1 to cluster the positions on an interval of length 1. Then, in Step 2, one tries to ensure that the paths stay clustered until $+\infty$. Actually, oppositely to the Markovian case where the paths stay naturally together after a clustering (by putting the same noise on each coordinate), the main difficulty here is that, due to the memory, staying together is costly. In other words, this property can be ensured only with the help of a non trivial coupling of the noises. One thus talks of *asymptotic coupling*. If one of the two previous steps fails, we will begin a new attempt but only after a (long) waiting time which is called Step 3. During this step, one again waits that the paths get close but one also expects the memory of the coupling cost to vanish sufficiently in order to begin the new trial with a weak weight of the memory.

In the previous construction, the fact that σ is constant is fundamental for ensuring the two following properties:

- If two paths B^1 and B^2 of the fBm differ from a drift term, then two paths X^1 and X^2 of (1.2) respectively directed by B^1 and B^2 also differ from a drift term, which allows in particular to use Girsanov Theorem to build the coupling in Step 1.
- Under some “convexity” assumptions on the drift apart from a compact set, two paths X^1 and X^2 directed by the same fBm (or more precisely, by two slightly different paths) get closer and the distance between the two paths can be controlled deterministically.

In the current paper, σ is not constant and the two above properties are no longer valid. The challenge then is to extend the applicability of the previous scheme of coupling to such a situation. To cope with the lack of the first property, we will need to introduce an assumption on the function $x \mapsto \sigma(x)$ which ensures the existence of an injective function of the path which does have it. The most natural assumption is that $x \mapsto \sigma^{-1}(x)$ is (well-defined and is) a Jacobian matrix. We note that without such an assumption, the lack of smoothness of the difference between two paths seems to yield a cost of coupling which could be problematical for Step 2.

As concerns a suitable substitution of the second lacking property, a natural (but to our knowledge so far not explored) idea would be to try to extend Meyn-Tweedie techniques (see *e.g.* [6] for background) to the fractional setting. More precisely, even if the paths do not get closer to each other deterministically, one could expect that some Lyapunov assumption could eventually make the two paths return in some compact set simultaneously. The main contribution of the present paper is to incorporate such a Lyapunov-type approach into the study of long-time convergence in the fractional diffusion setting. As one could expect, compared to the Markovian case, the problem is much more involved. Actually, the return time to a compact set after a (failed) coupling attempt does not only depend on the positions of the processes after it, but also on all the past of the fBm. Therefore, in order that the coupling attempt succeeds with lower-bounded probability, one needs to establish some controls on the past behavior of the fBms that drive the two copies of the process, conditionally to the failure of the previous attempts. This point is one of the main difficulties of the paper, since, in the corresponding estimates, we carefully have to take into account all the deformations of the distribution that previously failed attempts induce. Then, we show that after a sufficiently long waiting time, conditionally on previous fails the probability that the two paths be in a compact set and that the influence of past noise on the future be controlled, is lower-bounded. Bringing all the estimates together yields a global control of the coupling time and a rate of convergence which is similar to the one in [9] in the additive noise case.

In Section 2 we detail our assumptions and state our main result, namely Theorem 2.1. The scheme of its proof, based on the previous described coupling strategy, is then given. The proof of Theorem 2.1 is achieved in Sections 3, 4 and 5, which are outlined at the end of Section 2.

2 Assumptions and main result

We begin by listing a series of notations and definitions.

- The scalar product and the Euclidean norm on \mathbb{R}^d are respectively denoted by (\cdot) and $|\cdot|$.
- The non explicit constants will be usually denoted by C and may change from line to line.
- The space $\mathcal{C}([0, +\infty), \mathbb{R}^d)$ denotes the space of continuous functions on $[0, +\infty)$ endowed with the topology of uniform convergence on compact spaces.
- For some given $a, b \in \mathbb{R}$, with $a, b, L^2([a, b], \mathbb{R}^d)$ denotes the space of Lebesgue-measurable functions such that $\|g\|_{[a,b],2} = \sqrt{\int_a^b |g(s)|^2 ds} < +\infty$.
- For some positive θ and δ such that $\theta \in (1/2, H)$ and $\theta + \delta \in (H, 1)$, $\mathcal{W}_{\theta,\delta}$ denotes the Polish space $\mathcal{W}_{\theta,\delta}$ which is the completion of $\mathcal{C}_0^\infty((-\infty, 0], \mathbb{R}^d)$ (the space of \mathcal{C}^∞ -functions $f : (-\infty, 0] \rightarrow \mathbb{R}^d$ with compact support and $f(0) = 0$) for the norm

$$\|f\|_{\mathcal{W}_{\theta,\delta}} = \sup_{-\infty < s < t \leq 0} \frac{|f(t) - f(s)|}{|t - s|^{\theta}(1 + |t|^{\delta} + |s|^{\delta})}. \quad (2.4)$$

- For some real $a < b$ and for $\theta \in (0, H)$, we denote by $\mathcal{C}^\theta([a, b], \mathbb{R}^d)$ the set of functions $f : \mathbb{R}_+ \rightarrow \mathbb{R}^d$ such that

$$\|f\|_\theta^{a,b} = \sup_{0 \leq s < t \leq T} \frac{|f(t) - f(s)|}{(t - s)^\theta} < +\infty,$$

- Let $\sigma : \mathbb{R}^d \rightarrow \mathbb{M}_{d,d}$ be a \mathcal{C}^1 -function and $0 < \gamma < 1$. We say that σ is $(1 + \gamma)$ -Lipschitz if for every $i, j \in \{1, \dots, d\}$, the following norm is finite:

$$\|\sigma_{i,j}\|_{1+\alpha} = \sup_{x \in \mathbb{R}^d} |\nabla \sigma_{i,j}(x)| + \sup_{x, y \in \mathbb{R}^d} \frac{|\nabla \sigma_{i,j}(x) - \nabla \sigma_{i,j}(y)|}{|x - y|^\gamma}, \quad (2.5)$$

where for a given \mathcal{C}^1 -function $f : \mathbb{R}^d \rightarrow \mathbb{R}$, $\nabla f = (\partial_{x_1} f, \dots, \partial_{x_d} f)'$.

- We also denote by $\mathcal{EQ}(\mathbb{R}^d)$ the set of *Essentially Quadratic* functions, that is \mathcal{C}^1 -functions $V : \mathbb{R}^d \rightarrow (0, \infty)$ such that ∇V is Lipschitz continuous,

$$\liminf \frac{V(x)}{|x|^2} > 0, \quad \text{and} \quad |\nabla V| \leq C\sqrt{V}.$$

where C is a positive constant. Note that these assumptions ensure that $\inf V = \min V$ is positive and that \sqrt{V} is Lipschitz continuous (since it has a bounded gradient) which in turns implies that V is subquadratic.

Now, let us introduce the assumptions:

(H₀): b is a locally Lipschitz and sublinear function and σ is a bounded $(1 + \gamma)$ -Lipschitz continuous function ($\gamma > 0$).

This condition ensures existence and uniqueness of solutions for (1.2). Note that the condition on the derivative of σ only plays a role for uniqueness and in particular, is not fundamental for what follows. However, for the sake of simplicity, we choose to assume this assumption throughout the paper.

Now, we turn to some more specific assumptions **(H₁)** and **(H₂)**. The first one is a Lyapunov-stability assumption:

(H₁): There exists a function $V : \mathbb{R}^d \rightarrow \mathbb{R}$ of $\mathcal{EQ}(\mathbb{R}^d)$, there exists some positive β_0 and κ_0 such that

$$\forall x \in \mathbb{R}^d, \quad (\nabla V(x)|b(x)) \leq \beta_0 - \kappa_0 V(x).$$

REMARK 2.1. The above assumption will be used to ensure that the paths live in a compact set with a high probability. Note that in the classical diffusion setting, such a property holds with some less restrictive Lyapunov assumptions. Here, the assumptions essentially allow us to consider only (attractive) drift terms whose growth is linear at infinity. On the one hand, due to **(H₀)**, one can not consider drift terms with (strictly) superlinear growth at infinity and on the other hand, Assumption **(H₁)** combined with the fact that V is subquadratic implies more or less that b can not have (strictly) sublinear growth at infinity (this would be possible if V had an exponential growth). These restrictions are mainly due to the lack of martingale property for the integrals driven by fBms.

Then, when the paths are in this compact set, one tries classically to couple them with positive probability. But, as mentioned before, the specificity of the non-Markovian setting is that the coupling attempt generates a cost for the future (in a sense made precise later). In order to control this cost or more precisely in order to couple the paths with the help of a controlled drift term, we need to ensure the next assumption:

(H₂) $\forall x \in \mathbb{R}^d$, $\sigma(x)$ is invertible and there exists a \mathcal{C}^1 -function $h = (h_1, \dots, h_d) : \mathbb{R}^d \rightarrow \mathbb{R}^d$ such that the Jacobian matrix $\nabla h = (\partial_{x_j} h_i)_{i,j \in \{1, \dots, d\}}$ satisfies $\nabla h(x) = \sigma^{-1}(x)$ and such that ∇h is a locally Lipschitz function on \mathbb{R}^d .

REMARK 2.2. \triangleright Under (\mathbf{H}_0) and (\mathbf{H}_2) , h is a global \mathcal{C}^1 -diffeomorphism from \mathbb{R}^d to \mathbb{R}^d . Indeed, under these assumptions, ∇h is invertible everywhere and $x \mapsto [(\nabla h)(x)]^{-1} = \sigma(x)$ is bounded on \mathbb{R}^d . Then, the property (which will be important in the sequel), follows from the Hadamard-Lévy theorem (see e.g. [18]).

\triangleright As mentioned before, the main restriction here is to assume that $x \mapsto \sigma^{-1}(x)$ is a Jacobian matrix. However, note that there is no assumption on h (excepted smoothness). In particular, σ^{-1} does not need to be bounded. This allows us to consider for instance some cases where σ vanishes at infinity.

Let us exhibit some simple classes of SDEs for which (\mathbf{H}_2) is fulfilled. First, it contains the class of non-degenerated SDEs for which each coordinate is directed by one real-valued fBm. More precisely, if for every $i \in \{1, \dots, d\}$,

$$dX_t^i = b_i(X_t^1, \dots, X_t^d)dt + \sigma_i(X_t^1, \dots, X_t^d)dB_t^i$$

where $\sigma_i : \mathbb{R}^d \rightarrow \mathbb{R}$ is a \mathcal{C}^1 positive function, Assumption (\mathbf{H}_2) holds. Now, let us also remark that since, for a given constant matrix, $\nabla(P_h) = P\nabla h$, we have the following equivalence:

$$\exists h \text{ such that } \nabla h = \sigma^{-1} \iff \exists \tilde{h}, \exists \text{ an invertible matrix } P \text{ such that } \sigma^{-1} = P\nabla \tilde{h}.$$

One deduces from this property that (\mathbf{H}_2) also holds true if:

$$\sigma(x) = P \text{Diag}(\sigma_1(x_1, \dots, x_d), \dots, \sigma_d(x_1, \dots, x_d))$$

where P is a given invertible $d \times d$ -matrix and for every $i \in \{1, \dots, d\}$ σ_i has the same properties as before.

We are now able to state our main result. One denotes by $\mathcal{L}((X_t^{\mu_0})_{t \geq 0})$ the distribution of the process on $\mathcal{C}([0, +\infty), \mathbb{R}^d)$ starting from an initial distribution μ_0 and by $\bar{\mathcal{Q}}\mu$ the distribution of the stationary solution (starting from an invariant distribution μ). The distribution $\bar{\mu}_0(dx)$ denotes the first marginal of $\mu_0(dx, dw)$.

THEOREM 2.1. *Let $H \in (1/2, 1)$. Assume (\mathbf{H}_0) , (\mathbf{H}_1) and (\mathbf{H}_2) . Then, existence and uniqueness hold for the invariant distribution μ (up to equivalence). Furthermore, for every initial distribution μ_0 for which there exists $r > 0$ such that $\int |x|^r \bar{\mu}_0(dx) < \infty$, for each $\varepsilon > 0$ there exists $C_\varepsilon > 0$ such that*

$$\|\mathcal{L}((X_{t+s}^{\mu_0})_{s \geq 0}) - \bar{\mathcal{Q}}\mu\|_{TV} \leq C_\varepsilon t^{-(\frac{1}{8} - \varepsilon)}.$$

REMARK 2.3. In the previous result, the main contribution is the fact that one is able to recover the rates of the additive case. Existence and uniqueness results are not really new. However, compared with the assumptions of [10], one observes that when $x \mapsto \sigma^{-1}(x)$ is a Jacobian matrix (assumption which does not appear in [10]), our other assumptions are slightly less constraining. In particular, b is assumed to be locally Lipschitz and sublinear (instead of Lipschitz continuous) and, as mentioned before, $x \mapsto \sigma^{-1}(x)$ does not need to be bounded.

2.1 Scheme of coupling

As explained before, the proof of Theorem 2.1 is based on a coupling strategy similar to that of [9]. Let $(B_t^1)_{t \in \mathbb{R}}$ and $(B_t^2)_{t \in \mathbb{R}}$ denote two fractional Brownian motions with Hurst parameter $H > 1/2$. Then, denote by (X_t^1, X_t^2) , a couple of solutions of (1.2):

$$\begin{cases} dX_t^1 = b(X_t^1)dt + \sigma(X_t^1)dB_t^1 \\ dX_t^2 = b(X_t^2)dt + \sigma(X_t^2)dB_t^2 \end{cases} \quad (2.6)$$

with initial conditions $(X_0^1, (B_t^1)_{t \leq 0})$ $(X_0^2, (B_t^2)_{t \leq 0})$. We denote by $(\mathcal{F}_t)_{t \geq 0}$ the usual augmentation of the filtration $(\sigma(B_s^1, B_s^2, (X_0^1, X_0^2)_{s \leq t}))_{t \geq 0}$. To begin the coupling procedure without “weight of the past”, we will certainly assume that

$$(B_t^1)_{t \leq 0} = (B_t^2)_{t \leq 0}$$

and that the initial distribution $\tilde{\mu}$ of the initial distribution of (X^1, X^2) is of the form

$$\tilde{\mu}(dx, dw) = \mu_1(w, dx_1)\mu_2(w, dx_2)\mathbb{P}_H(dw) \quad (2.7)$$

where \mathbb{P}_H denotes the distribution of a fBm $(B_t)_{t \leq 0}$ on $\mathcal{W}_{\gamma, \delta}$ and the transitions probabilities $\mu_1(\cdot, dx_1)$ and $\mu_2(\cdot, dx_2)$ correspond respectively to the conditional distributions of X_0^1 and X_0^2 given $(B_t^1)_{t \leq 0}$. The processes $(B_t^1)_{t \in \mathbb{R}}$ and $(B_t^2)_{t \in \mathbb{R}}$ can be realized through the Mandelbrot-Van Ness representation (see (1.3)) with the help of some two-sided Brownian motions respectively denoted by W^1 and W^2 . In particular, the filtration $(\mathcal{F}_t)_{t \geq 0}$ is also generated by $(\sigma(W_s^1, W_s^2, (X_0^1, X_0^2))_{s \leq t})_{t \geq 0}$.

Furthermore, we will assume in all the proof that on $[0, \infty)$, W^1 and W^2 (resp. B^1 and B^2) differ by a (random) drift term denoted by g_w (resp. g_B):

$$dW_t^2 = dW_t^1 + g_w(t)dt \quad \text{and} \quad dB_t^2 = dB_t^1 + g_B(t)dt. \quad (2.8)$$

Note that the functions g_w and g_B are linked by some inversion formulas (see [9], Lemma 4.2 for details).

The idea is to build g_w (resp. g_B) in order to stick X^1 and X^2 . We set

$$\tau_\infty := \inf\{t \geq 0, X_s^1 = X_s^2 \forall s \geq t\}.$$

As usual, this coupling will be achieved after a series of trials. As mentioned in the introduction, each trial is decomposed in three steps:

- Step 1: Try to couple the positions with a *controlled cost* (in a sense made precise below).
- Step 2 (specific to non-Markov processes): Try to keep the paths fastened together.
- Step 3: If Step 2 fails, wait a sufficiently long time in order that in the next trial, Step 1 be achieved with a *controlled cost* and with (uniformly lower-bounded away from 0) probability. During this step, we suppose that $g_w(t) = 0$.

Let us make a few precisions:

▷ We denote by $\tau_0 \geq 0$ the beginning of the first trial and by τ_k , $k \geq 1$, the end of each trial. If $\tau_k = +\infty$, the coupling tentative has been successful. Otherwise, τ_k is the end of Step 3 of trial k . We will assume that

$$\forall t \in (-\infty, \tau_0], \quad W_t^1 = W_t^2 \quad a.s. \quad \text{or equivalently that} \quad g_w(t) = g_B(t) = 0 \quad \text{on} \quad [-\infty, \tau_0].$$

▷ About Step 1 and the “controlled cost”: Step 1 is carried out on each interval $[\tau_{k-1}, \tau_{k-1} + 1]$. The “cost” of coupling is represented by the function g_w that one needs to build on $[\tau_{k-1}, \tau_{k-1} + 1]$ in order to get X^{x_1} and X^{x_2} stuck together at time $\tau_{k-1} + 1$. Oppositely to the Markovian case, this cost does not only depend on the positions of $X_{\tau_{k-1}}^1$ and $X_{\tau_{k-1}}^2$ but also on the past of the Brownian motions, which have a (strong) influence on the dynamics of B^1 and B^2 . This is the reason why one needs in Step 3 to wait enough before beginning a new attempt of coupling.

In [9], the “controlled cost” concept is called “admissibility” (see Definition 5.6). Here, we slightly modify it and we will say that one is in position to attempt a coupling if the system is (K, α) -admissible. We define this concept below but need before to introduce notations. For $T \geq 0$ and a measurable function $g : \mathbb{R} \rightarrow \mathbb{R}$, we denote by $\mathcal{R}_T g$ the function defined (when it makes sense) by

$$(\mathcal{R}_T g)(t) = \int_{-\infty}^0 \frac{t^{\frac{1}{2}-H}(T-s)^{H-\frac{1}{2}}}{t+T-s} g(s) ds, \quad t \in (0, +\infty).$$

Let g_w be the (random) function defined by (2.8). For a positive time τ , we denote by g_w^τ the function defined by $g_w^\tau(t) = g_w(t + \tau)$, $t \in \mathbb{R}$.

The following definition is relative to a fixed $\theta \in (1/2, H)$.

DEFINITION 2.1. Let K and α be some positive constants and τ denote stopping time with respect to $(\mathcal{F}_t)_{t \in \mathbb{R}}$. We say that the system is (K, α) -admissible at time τ if $\tau(\omega) < +\infty$ and if $(X_\tau^1(\omega), X_\tau^2(\omega), (W^1(\omega), W^2(\omega))_{t \leq \tau})$ satisfies:

$$\sup_{T \geq 0} \int_0^{+\infty} (1+t)^{2\alpha} |(\mathcal{R}_T g_w^\tau)(t)|^2 dt \leq 1. \quad (2.9)$$

and

$$|X_\tau^1(\omega)| \leq K, \quad |X_\tau^2(\omega)| \leq K, \quad \varphi_{\tau, \varepsilon_\theta}(W^1(\omega)) \leq K \quad \text{and} \quad \varphi_{\tau, \varepsilon_\theta}(W^2(\omega)) \leq K, \quad (2.10)$$

where $\varepsilon_\theta = \frac{H-\theta}{2}$ and for a given positive ε ,

$$\varphi_{\tau, \varepsilon}(w) = \sup_{\tau \leq s \leq t \leq \tau+1} \left| \frac{1}{t-s} \int_{-s}^{\tau-1} (t-r)^{H-\frac{1}{2}} - (s-r)^{H-\frac{1}{2}} dw_r \right| + \|w\|_{\frac{1}{2}-\varepsilon}^{\tau-1, \tau}$$

If these two conditions hold, we will show that the coupling attempt is successful with lower-bounded probability. Thus, we will need to ensure that at each time τ_k , the (K, α) -admissibility also holds with lower-bounded probability. We set $\Omega_{K, \alpha, \tau} = \Omega_{\alpha, \tau}^1 \cup \Omega_{K, \tau}^2$ where

$$\Omega_{\alpha, \tau}^1 := \left\{ \omega, \tau(\omega) < +\infty, \sup_{T \geq 0} \int_0^{+\infty} (1+t)^{2\alpha} |(\mathcal{R}_T g_w^\tau)(t)|^2 dt \leq 1 \right\}, \quad (2.11)$$

and

$$\Omega_{K, \tau}^2 := \left\{ \omega, \tau(\omega) < +\infty, |X_\tau^1| \leq K, |X_\tau^2| \leq K, \varphi_{\tau, \varepsilon_\theta}(W^1) \leq K, \varphi_{\tau, \varepsilon_\theta}(W^2) \leq K \right\}, \quad (2.12)$$

The novelty here is the event defined in (2.12). Since, contrarily to the additive noise case, we are not able to reduce here the distance between the positions deterministically, we ask X^{x_1} and X^{x_2} to be in the compact set $\bar{B}(0, K) = \{y, |y| \leq K\}$ with positive probability. The same type of assumption is needed on the past of the fractional Brownian motion (which is represented by the functionals $\varphi_{\tau, \varepsilon_\theta}(W^j)$, $j = 1, 2$). Note that, oppositely to the event $\Omega_{\alpha, \tau}^1$, which comes from [9], $\Omega_{K, \tau}^2$ can certainly not have a probability equal to 1. We will attempt the coupling on $[\tau_{k-1}, \tau_{k-1} + 1]$ only if $\omega \in \Omega_{K, \alpha, \tau_{k-1}}$. Otherwise, we set $g_w(t) = 0$ on $[\tau_{k-1}, \tau_{k-1} + 1]$ (and, in this case, we certainly say that Step 1 fails).

▷ If Step 1 fails (which includes the case where one does not attempt the coupling), one begins Step 3 (see below). Otherwise, one begins Step 2. Step 2 is in fact a series of trials on some intervals I_ℓ with length

$$|I_\ell| = c_2 2^\ell \quad (2.13)$$

where c_2 is a constant greater than one which will be calibrated in the sequel. More precisely, one successively tries to keep X^1 and X^2 as being equal on intervals $[\tau_{k-1} + 1 + c_2 \sum_{u=1}^{\ell-1} 2^k, \tau_{k-1} + c_2 \sum_{u=1}^{\ell} 2^k]$ (with the convention $\sum_{\emptyset} = 0$). The exponential increase of the length of the intervals will be of first importance to ensure the success of Step 2.

▷ If Step 2 fails at trial ℓ with $\ell \geq 0$ ($\ell = 0$ corresponds to the case where Step 1 fails), one begins Step 3. We denote by τ_k^3 , the beginning of Step 3. As mentioned before, the aim on this interval is to wait a sufficiently long time in order to be in an (K, α) -admissible state with positive probability (Step 3 ends at time τ_k , *i.e.* at the beginning of the next attempt). This has two natural consequences. On the one hand, one assumes that

$$g_w(t) = 0 \quad \text{on } [\tau_k^3, \tau_k], \quad \text{so that} \quad W_t^1 - W_{\tau_k^3}^1 = W_t^2 - W_{\tau_k^3}^2 \quad \text{on } [\tau_k^3, \tau_k]. \quad (2.14)$$

On the other hand, the waiting time will strongly depend on the length of Step 2. Longer is Step 2, longer is the waiting time. We set

$$\mathcal{A}_{k, \ell} = \{\text{At trial } k, \text{ Step 2 fails after } \ell \text{ attempts}\}. \quad (2.15)$$

We assume in the sequel that

$$\forall \omega \in \mathcal{A}_{k,\ell}, \quad \tau_k - \tau_k^3 = \Delta_3(\ell, k) \quad \text{with} \quad \Delta_3(\ell, k) := c_3 a_k 2^{\beta \ell} \quad (2.16)$$

where $c_3 \geq 2c_2$, $\beta \in [1, +\infty)$ and $(a_k)_{k \geq 1}$ is an increasing deterministic sequence. We will calibrate these quantities later (see Proposition 4.6). At this stage, we can however remark a useful property for the sequel: conditionally to $\mathcal{A}_{k,\ell}$, the length of each step is deterministic. We are now ready to begin the proof. In Section 3, we focus on Steps 1 and 2 and prove that we can achieve the coupling scheme in such a way that for every positive K and α , the probability of coupling can be lower-bounded by a constant which does not depend on k . Then, in Section 4, we focus on the (K, α) -admissibility condition. In particular, we show that for K large enough, (2.12) holds with high probability (which does not depend on k). Finally, in Section 5, we prove Theorem 2.1.

3 Lower-bound for the successful-coupling probability

In this section, we detail the construction of Steps 1 and 2 with the aim of proving that if the system is (K, α) -admissible at time τ_{k-1} , then the probability that $\Delta\tau_k$ be infinite (*i.e.* that the coupling be successful) can be lower-bounded. The main result of this section is the next proposition.

PROPOSITION 3.1. *Let $H > 1/2$. Assume that (\mathbf{H}_0) and (\mathbf{H}_2) hold. Then, for every $K > 0$ and $\alpha \in (0, H)$, Steps 1 and 2 can be achieved in such a way that there exists δ_0 and δ_1 in $(0, 1)$ such that for every $k \geq 1$, $\delta_0 \leq \mathbb{P}(\Delta\tau_k = +\infty | \Omega_{K,\alpha,\tau_{k-1}}) \leq 1 - \delta_1$. Furthermore, δ_1 can be chosen independently of K .*

The (uniform) upper-bound is almost obvious. Actually, at the beginning of Step 1, it is always possible (if necessary) to attempt the coupling with probability $1 - \delta_1$ only (and to put $W^1 = W^2$ otherwise). This upper-bound may appear of weak interest but in fact, it will play an important role in Section 4.

The lower-bound is a consequence of the combination of Equation (3.30) with Lemmas 3.1 and 3.2 below.

3.1 Step 1

LEMMA 3.1. *Assume that (\mathbf{H}_0) and (\mathbf{H}_2) hold. Let K and α denote two positive constants and $\theta \in (1/2, H)$ be fixed. Then, for each $k \geq 1$, (W^1, W^2) can be built on $[\tau_{k-1}, \tau_{k-1} + 1]$ in such a way that the following properties hold:*

(a) *There exists $\tilde{\delta}_0 > 0$ depending only on K , α and $\theta \in (1/2, H)$ such that for all $k \geq 0$,*

$$\mathbb{P}(X_{\tau_{k-1}+1}^1 = X_{\tau_{k-1}+1}^2 | \Omega_{K,\alpha,\tau_{k-1}}) \geq \tilde{\delta}_0.$$

(b) *There exists $C_K > 0$ such that $\int_{\tau_{k-1}}^{\tau_{k-1}+1} |g_w(s)|^2 ds \leq C_K$ a.s.*

(c) *If Step 1 is successful, $t \mapsto g_B(t)$ is a \mathcal{C}^1 -function on $[\tau_{k-1}, \tau_{k-1} + 1]$ which is such that*

$$\sup_{t \in [0,1]} |g_B(t)| \leq C \quad \text{and} \quad \forall t \in [1/2, 1], \quad g_B(t + \tau_{k-1}) = 0.$$

where C is a deterministic constant which does not depend on k .

Proof. (a) The proof of this statement is divided in five parts:

(i) Let $\theta \in (1/2, H)$ and set $\varepsilon_\theta = \frac{H-\theta}{2}$. Let \tilde{K} be a positive constant. Then, there exists a deterministic constant depending only on θ , K and \tilde{K} denoted $C(K, \tilde{K})$ such that

$$\forall \omega \in \Omega_{K,\tau_{k-1}}^2 \cap \{ \|W^1\|_{\frac{1}{2}-\varepsilon_\theta}^{\tau_{k-1}, \tau_{k-1}+1} \leq \tilde{K} \}, \quad \sup_{t \in [0,1]} |X_{\tau_{k-1}+t}^1(\omega)| \leq C(K, \tilde{K}).$$

The proof of this property (whose arguments are close to some of the next sections) is given in Appendix A.

(ii) Building a function g_B to couple (X_t^1) and (X_t^2) at time $\tau_{k-1} + 1$. First, note that this step is strongly based on Assumption **(H₂)** and that the construction is a modified version of Lemma 5.8 of [9]. For a given past on $(-\infty, \tau_{k-1}]$ and a given innovation path $(W_{t+\tau_{k-1}}^1, W_{t+\tau_{k-1}}^2) = (w_1(t) + W_{\tau_{k-1}}^1, w_2(t) + W_{\tau_{k-1}}^2)_{t \in [0,1]}$ of $\mathcal{C}^{\frac{1}{2}-\varepsilon_\theta}([0,1], \mathbb{R}^d)^2$ (w_1 and w_2 satisfy $w_1(0) = w_2(0) = 0$), set $(x^{w_1}(t), x^{w_2}(t)) = (X_{t+\tau_{k-1}}^1, X_{t+\tau_{k-1}}^2)$, $t \in [0,1]$. Also, with the same conditions as before, set $(B_t^{w_1}, B_t^{w_2}) := (B_{t+\tau_{k-1}}^1, B_{t+\tau_{k-1}}^2)$.

The aim is now to build, conditionally to $\Omega_{K,\alpha,\tau_{k-1}}$ and $(X_{\tau_{k-1}}^1, X_{\tau_{k-1}}^2, (W^1, W^2)_{t \leq \tau_{k-1}})$, a function denoted by g_h on $[0,1]$ such that if for every $t \in [0,1]$, $w_2(t) = w_1(t) + \int_0^t g_h(s) ds$, then $x^{w_1}(1) = x^{w_2}(1)$. In fact, it is more convenient to build the associated function f_h which is defined by $dB_t^{w_2} = dB_t^{w_1} + f_h(t)dt$ (see (2.8) for background).

With the previous notations, we deduce from Assumption **(H₂)** and from a change of variable formula for Hölder functions with exponent greater than 1/2 (see *e.g.* [20], Theorem 4.3.1) that, for every $t \in [0,1]$,

$$h(x^{w_2}(t)) - h(x^{w_1}(t)) = h(x^{w_1}(0)) - h(x^{w_1}(0)) + \int_0^t \nabla h b(x^{w_2}(u)) - \nabla h b(x^{w_1}(u)) du + \int_0^t f_h(u) du.$$

Set

$$\rho_h(t) = h(x^{w_2}(t)) - h(x^{w_1}(t)) \quad \text{and} \quad f_h(t) = -\kappa_1 \rho_h(t) - \kappa_2 \frac{\rho_h(t)}{\sqrt{|\rho_h(t)|}},$$

where κ_1 and κ_2 are some positive constants, and with $x \mapsto x/\sqrt{|x|}$ extended by 0 at $x = 0$. Notice that if $\rho_h(0) = 0$, then $\rho_h(t) = 0$ on $[0,1]$. We thus assume without loss of generality that $\rho_h(0) \neq 0$. By the global inversion theorem, $h : \mathbb{R}^d \rightarrow \mathbb{R}^d$ is a \mathcal{C}^1 -diffeomorphism under **(H₂)**. Denote by h^{-1} its inverse function. The function $(\rho_h(t))_{t \geq 0}$ then satisfies (for a given realization of $(x^{w_1}(t))_{t \in [0,1]}$)

$$\frac{d\rho_h}{dt} = F(t, \rho_h(t)) \quad \text{with} \quad F(t, \rho) = \nabla h b(h^{-1}(h(x^{w_1}(t)) + \rho)) - \nabla h b(x^{w_1}(t)) - \kappa_1 \rho - \kappa_2 \frac{\rho}{\sqrt{|\rho|}}. \quad (3.17)$$

The function $\rho \mapsto F(t, \rho)$ being locally Lipschitz continuous on $\mathbb{R}^d \setminus \{0\}$, one deduces from Cauchy-Lipschitz Theorem that $(\rho_h(t))$ is well-defined on an interval $[0, t_0]$ with $t_0 > 0$. To extend it on $[0,1]$, we are going to define κ_1 in order to ensure that $(\rho_h(t))$ does not explode. For $M \geq 0$, set

$$A_M := \{w_1 \in \mathcal{C}^{\frac{1}{2}-\varepsilon_\theta}([0,1], \mathbb{R}^d), w_1(0) = 0 \text{ and, } \forall \omega \in \Omega_{K,\tau_{k-1}}^2, \sup_{t \in [0,1]} |h(x^{w_1}(t))| \leq M\}.$$

Remark that with the notations of (i), $\{w_1, \|w_1\| \leq \tilde{K}\} \subset A_{C(K,\tilde{K})}$ (one will come back on this property in (v)). Checking that $w_1 \mapsto B^{w_1}$ is *a.s.* continuous from $\mathcal{C}^{\frac{1}{2}-\varepsilon_\theta}([0,1], \mathbb{R}^d)$ to $\mathcal{C}^\theta([0,1], \mathbb{R}^d)$, one deduces from classical arguments of Young integration that $w_1 \mapsto h(x^{w_1})$ is also *a.s.* continuous from $\mathcal{C}^{\frac{1}{2}-\varepsilon_\theta}([0,1], \mathbb{R}^d)$ to $\mathcal{C}^\theta([0,1], \mathbb{R}^d)$. As a consequence, for almost every $\omega \in \Omega_{K,\tau_{k-1}}^2$, $w_1 \mapsto \sup_{t \in [0,1]} |h(x^{w_1}(t))|$ is also continuous (since for every $w, \tilde{w} \in \mathcal{C}^{\frac{1}{2}-\varepsilon_\theta}([0,1], \mathbb{R}^d)$, $|\|w\|_\infty - \|\tilde{w}\|_\infty| \leq \|w - \tilde{w}\|_{\frac{1}{2}-\varepsilon_\theta}^{0,1}$). At the price of a potential removal of a negligible set in its definition, one can assume without loss of generality that A_M is an intersection of closed subsets of $\mathcal{C}^{\frac{1}{2}-\varepsilon_\theta}([0,1], \mathbb{R}^d)$ and thus is closed.

Set $\bar{K} = \sup_{x,y \in \bar{B}(0,K)} |h(x) - h(y)|$ where $\bar{B}(0,K) := \{y \in \mathbb{R}^d, |y| \leq K\}$. For a given M , we define κ_1 as follows:

$$\kappa_1 = \sup_{y_1, y_2 \in \bar{B}(0, M + \bar{K} + 1)} \frac{|\nabla h b(h^{-1}(y_2)) - \nabla h b(h^{-1}(y_1))|}{|y_2 - y_1|} \quad (3.18)$$

The constant κ_1 is finite since $b, \nabla h$ and h^{-1} are locally Lipschitz continuous functions. From the definition, if $w_1 \in A_M$, for every $\omega \in \Omega_{K,\tau_{k-1}}^2$ and $t \in [0,1]$, $|h(x^{w_1}(t))| \leq M$ and $|h(x^{w_2}(0))| \leq$

$M + \bar{K}$ (since $|\rho_h(0)| \leq \bar{K}$). Owing to the continuity of ρ_h on $[0, t_0]$, there exists $t_1 \in (0, t_0]$ such that for every $t \in [0, t_1]$, $|h(x^{w_2}(t))| \leq M + \bar{K} + 1$. Then, it follows from the definition of κ_1 that

$$\forall t \in [0, t_1], \quad \frac{d|\rho_h(t)|^2}{dt} \leq -2\kappa_2|\rho_h(t)|^{\frac{3}{2}}.$$

In particular, the function $t \mapsto |\rho_h(t)|^2$ is non-increasing on $[0, t_1]$ and thus, $\forall t \in [0, t_1]$, $|h(x^{w_2}(t))| \leq M + \bar{K}$. Then, a classical argument about maximal solutions of ordinary differential equations shows that existence and uniqueness of solutions of (3.17) holds on whole the interval $[0, 1]$ and that the previous inequality is true with $t_0 = 1$. A standard computation yields

$$|\rho_h(t)| \leq \begin{cases} (\frac{\kappa_2}{2}t - \sqrt{|\rho_h(0)|})^2 & \text{if } t \leq 2\sqrt{|\rho_h(0)|}/\kappa_2 \\ 0 & \text{if } t \geq 2\sqrt{|\rho_h(0)|}/\kappa_2. \end{cases} \quad (3.19)$$

Then, we set $\kappa_2 = 4\sqrt{\bar{K}}$ in order that $\rho_h(t) = 0$ on $[1/2, 1]$ (which implies that $f_h(t) = 0$ on $[1/2, 1]$; this fact will be used in Step 2). This concludes the construction of the function f_h .

(iii) About f_h and g_h : let $\omega \in \Omega_{K, \alpha, \tau_{k-1}}$ and $w_1 \in A_M$ and consider the \mathcal{C}^1 -function $(f_h(t))_{t \in [0, 1]}$ built in the previous subsection. Given this function, let us recall how one defines a function $(g_h(t))_{t \in [0, 1]}$ which is such that

$$dw_t^2 = dw_t^1 + g_h(t)dt \text{ on } [0, 1] \implies dB_t^{w_2} = dB_t^{w_1} + f_h(t)dt \text{ on } [0, 1].$$

The function g_B of (2.8) being known on $(-\infty, \tau_{k-1}]$, one can define it on $(-\infty, \tau_{k-1} + 1]$ by setting $g_B(t) = f_h(t - \tau_{k-1})$ on $[\tau_{k-1}, \tau_{k-1} + 1]$. Then, by an inversion formula (see (4.11a) of [9]), one obtains a unique g_w on $(-\infty, \tau_{k-1} + 1]$ (where g_w is defined in (2.8)). Then, g_h can be defined by $g_h(t) = g_w(t + \tau_{k-1}) = g_w^{\tau_{k-1}}(t)$, $t \in [0, 1]$. In fact, it can be shown (see proof of Lemma 5.9 of [9] for details) that the function g_h is given by

$$g_h(t) = C\mathcal{R}_0 g_w^{\tau_{k-1}}(t) + \alpha_H \frac{d}{dt} \left(\int_0^t (t-s)^{H-\frac{1}{2}} f_h(s) ds \right), \quad t \in (0, 1]. \quad (3.20)$$

Note that owing to the (K, α) -admissibility condition and to the differentiability of f_h , the function g_h is measurable and integrable on $[0, 1]$. We can thus define $\phi_1 : A_M \rightarrow \mathcal{C}^{\frac{1}{2}-\varepsilon_\theta}([0, 1], \mathbb{R}^d)$ by

$$\phi_1(w_1)_t = w_1(t) + \int_0^t g_h(s) ds, \quad t \in [0, 1]. \quad (3.21)$$

Using again that $w_1 \mapsto h(x^{w_1}(\cdot))$ is continuous from $\mathcal{C}^{\frac{1}{2}-\varepsilon_\theta}([0, 1], \mathbb{R}^d)$ to $\mathcal{C}^\theta([0, 1], \mathbb{R}^d)$, one can check that application ϕ_1 is measurable on A_M . The function ϕ_1 is also injective. Actually, let $(w_1, w_2) \in A_M \times \phi_1(A_M)$ such that $\phi_1(w_1) = w_2$. From the very definition, $x^{w_1}(t) = x^{w_2}(t) - \rho_h(t)$ where ρ_h is the unique solution of (3.17). But by construction, $(\rho_h(t))_{t \in [0, 1]}$ is also the unique solution of

$$\frac{d\rho_h}{dt} = \tilde{F}(t, \rho_h(t)) \quad \text{with} \quad \tilde{F}(t, \rho) = \nabla h b(x^{w_2}(t)) - \nabla h b(h^{-1}(h(x^{w_2}(t)) - \rho) - \kappa_1 \rho - \kappa_2 \frac{\rho}{\sqrt{|\rho|}}). \quad (3.22)$$

Thus, f_h and g_h defined above can be obtained as some measurable functions of w_2 . Denote them by \tilde{f}_h and \tilde{g}_h and set $\phi_2(w_2) = w_2(\cdot) - \int_0^\cdot \tilde{g}_h(s) ds$. By construction, we have $\phi_2 \circ \phi_1(w_1) = w_1$ so that ϕ_1 is a bijection from A_M to $\phi_1(A_M)$ with measurable inverse ϕ_2 .

Now, remark that, owing to the definition of \tilde{F} and to the arguments of (ii), existence and uniqueness for the solutions of (3.22) also hold on $[0, 1]$ (since the solutions do not explode) so that one can define \tilde{f}_h and \tilde{g}_h on A_M . In other words, one can extend ϕ_2 into a measurable function ψ on $A_M \cup \phi_1(A_M)$ defined by $\psi(w_2) = w_2(\cdot) + \int_0^\cdot \tilde{g}_h(s) ds$. Owing to the same arguments as above, ψ is injective. The function ϕ_1 can in turn be extended into a bijective measurable function $\varphi : A_M \cup \psi(A_M) \rightarrow A_M \cup \phi_1(A_M)$ defined by

$$\varphi(w_1)_t = w_1(t) + \int_0^t g_h(s) ds, \quad t \in [0, 1]. \quad (3.23)$$

Once again, $\varphi^{-1} = \psi$ is measurable and $\forall(w_1, w_2)$ such that $\varphi(w_1) = w_2$ with $w_1 \in A_M \cup \psi(A_M)$, $x_1^{w_1}(1) = x_1^{w_2}(1)$.

REMARK 3.4. Note that with this construction, we have built a function φ which is well-defined on $\varphi^{-1}(A_M)$ and which satisfies

$$\varphi(A_M) \subset A_{M+\bar{K}}. \quad (3.24)$$

The inclusion holds since $|h(x_t^{w_2})| \leq |h(x_t^{w_1})| + |\rho_h(0)| \leq M + \bar{K}$, when $w_2 = \varphi(w_1)$ with $w_1 \in A_M$. Note that these properties are important for the sequel. If $(\nabla hb) \circ h^{-1}$ was Lipschitz continuous, the function φ could be defined on the whole space and it would simplify the construction of the coupling (see (v)).

(iv) Control of the function g_h and Girsanov: For $\omega \in \Omega_{K,\alpha,\tau_{k-1}}$ and $w_1 \in A_M \cup \psi(A_M)$, consider the explicit expression of g_h given by (3.20).

By (2.9), the L^2 -norm of $((\mathcal{R}_0 g_w^{\tau_{k-1}})(t))_{t \in (0,1]}$ is bounded by 1. As concerns that of the second term in (3.20), it follows from Lemma 5.1 of [9] that it is enough to bound $f_h(0)$ and $|df_h/dt|$. But, by (3.17) and (3.19), one can check that there exists a deterministic constant \tilde{C} depending on K , M and θ such that $|f_h(0)| \leq \tilde{C}$ and

$$\forall t \in [0, 1], \quad \left| \frac{df_h(t)}{dt} \right| \leq C(|\rho_h(t)| + |\sqrt{\rho_h(t)}|) \leq \tilde{C}.$$

We deduce that for every positive M and K , there exists another finite constant $\bar{C}(M, K)$ such that

$$\forall \omega \in \Omega_{K,\alpha,\tau_k}, \quad \forall w_1 \in A_M \cup \psi(A_M), \quad \int_0^1 |g_h(s)|^2 ds < \bar{C}(M, K). \quad (3.25)$$

In particular, $w_1 \mapsto \int_0^1 |g_h(s)|^2 ds$ being finite on $A_M \cup \psi(A_M)$, this allows us to apply Girsanov Theorem. More precisely, extend φ into a measurable application $\hat{\varphi}$ on $\mathcal{C}^{\frac{1}{2}-\varepsilon\theta}([0, 1], \mathbb{R}^d)$ by setting $\hat{\varphi}(w_1) = w_1$ if $w_1 \in (A_M \cup \psi(A_M))^c$. Then, denoting by \mathbb{P}_W the Wiener measure and by $\hat{\varphi}^* \mathbb{P}_W$ the image measure of \mathbb{P}_W by the application $\hat{\varphi}$, one deduces from Girsanov Theorem that $\hat{\varphi}^* \mathbb{P}_W(dw) = D_{\hat{\varphi}}(w) \mathbb{P}_W(dw)$ where

$$D_{\hat{\varphi}}(w) = \exp \left(\int_0^1 (\bar{g}_h(s) |dw(s)) - \frac{1}{2} \int_0^1 |\bar{g}_h(s)|^2 ds \right) \quad (3.26)$$

with $\bar{g}_h = g_h$ if $w \in A_M \cup \psi(A_M)$ and $\bar{g}_h = 0$ otherwise. We can now explicit the coupling strategy.

(v) Construction of (W^1, W^2) on $[\tau_{k-1}, \tau_{k-1} + 1]$. First, we recall that we set $W_t^1 = W_t^2$ on $[\tau_{k-1}, \tau_{k-1} + 1]$ if $\omega \in \Omega_{K,\alpha,\tau_k}^c$ (in this case, attempting a coupling would generate a too important cost for the future). Now, if $\omega \in \Omega_{K,\alpha,\tau_{k-1}}$, the construction follows the lines of [9] but with the constraint that one can attempt the coupling only on a subset of $\mathcal{C}^{\frac{1}{2}-\varepsilon\theta}([0, 1], \mathbb{R}^d)$.

More precisely, for a Borel subset A of $\mathcal{C}^{\frac{1}{2}-\varepsilon\theta}([0, 1], \mathbb{R}^d)$, denote by \mathbb{P}_W^A the restriction of \mathbb{P}_W to A . For positive measures μ_1 and μ_2 with densities D_1 and D_2 with respect to another measure μ , denote by $\mu_1 \wedge \mu_2$ the measure defined by $(\mu_1 \wedge \mu_2)(dw) = D_1(w) \wedge D_2(w) \mu(dw)$. With the help of the function φ introduced in (iii) (see (3.23)), we define a non-negative measure \mathbf{P}_1^M on $\mathcal{C}^{\frac{1}{2}-\varepsilon\theta}([0, 1], \mathbb{R}^d)^2$ by

$$\mathbf{P}_1^M = \frac{1}{2} \varphi_1^* \mathbb{P}_W^{B_M} \wedge \varphi_2^* \mathbb{P}_W^{A_M}$$

where φ_1 and φ_2 are the functions from respectively $A_M \cup \psi(A_M)$ and $A_M \cup \varphi(A_M)$ to $\mathcal{C}^{\frac{1}{2}-\varepsilon\theta}([0, 1], \mathbb{R}^d)^2$ defined by

$$\varphi_1(w) = (w, \varphi(w)) \quad \text{and} \quad \varphi_2(w) = (\varphi^{-1}(w), w),$$

and $B_M = \varphi^{-1}(A_M)$ (note that A_M and B_M are Borel sets since A_M is closed and φ^{-1} is measurable).

Using that the restriction of $\hat{\varphi}$ to $\varphi^{-1}(A_M)$ is equal to φ and that $\varphi^{-1}(A_M) = \hat{\varphi}^{-1}(A_M)$, one can check that

$$\varphi_1^* \mathbb{P}_W^{B_M}(dw_1, dw_2) = \mathbf{1}_{\{(\varphi^{-1}(w), w)\}}(w_1, w_2) D_{\hat{\varphi}}(w) \mathbb{P}_W^{A_M}(dw),$$

where $D_{\hat{\varphi}}$ is defined by (3.26). This implies that \mathbf{P}_1^M satisfies

$$\mathbf{P}_1^M(dw_1, dw_2) = \frac{1}{2} \mathbf{1}_{\{(\varphi^{-1}(w), w)\}}(w_1, w_2) (D_{\hat{\varphi}}(w) \wedge 1) \mathbb{P}_W^{A_M}(dw). \quad (3.27)$$

Write $S(w_1, w_2) = (w_2, w_1)$ and denote by $\tilde{\mathbf{P}}_1^M$ the ‘‘symmetrized’’ non-negative measure induced by \mathbf{P}_1^M : $\tilde{\mathbf{P}}_1^M = \mathbf{P}_1^M + S^* \mathbf{P}_1^M$. We then define the coupling $(\tilde{W}_t^1, \tilde{W}_t^2) = (W_{t+\tau_{k-1}}^1 - W_{\tau_{k-1}}^1, W_{t+\tau_{k-1}}^2 - W_{\tau_{k-1}}^2)$ as follows:

$$\mathcal{L}((\tilde{W}_t^1, \tilde{W}_t^2)_{t \in [0,1]}) = \tilde{\mathbf{P}}_1^M + \Delta^*(\mathbb{P}_W - \Pi_1^* \tilde{\mathbf{P}}_1^M) = \mathbf{P}_1^M + \mathbf{P}_2^M$$

with $\Delta(w) = (w, w)$, $\Pi_1(w_1, w_2) = w_1$ and $\mathbf{P}_2^M = S^* \mathbf{P}_1^M + \Delta^*(\mathbb{P}_W - \Pi_1^* \tilde{\mathbf{P}}_1^M)$. Thanks to (3.27), one checks that

$$\begin{aligned} \Pi_1 \tilde{\mathbf{P}}_1^M(f) &\leq \frac{1}{2} \int_{A_M} (f(\varphi^{-1}(w)) D_{\hat{\varphi}}(w) + f(w)) \mathbb{P}_W(dw) \\ &\leq \frac{1}{2} \int f(w) (1_{\varphi^{-1}(A_M)}(w) + 1_{A_M}(w)) \mathbb{P}_W(dw) \end{aligned}$$

where we used that $\hat{\varphi}^{-1}(A_M) = \varphi^{-1}(A_M)$ and that $\varphi^{-1} \circ \hat{\varphi}(w) = w$ for every $w \in \varphi^{-1}(A_M)$. This implies that $\Pi_1 \tilde{\mathbf{P}}_1^M < \mathbb{P}_W$ and that \mathbf{P}_2^M is the sum of two positive measures. Owing to the symmetry property of \mathbf{P}_1^M and to the fact that $\Pi_1 \circ \Delta$ is the identity, one can also check that the marginals of $\mathbf{P}_1^M + \mathbf{P}_2^M$ are both equal to \mathbb{P}_W .

(v) Lower-bound for the probability of coupling: by construction, conditionally on $\Omega_{K, \alpha, \tau_{k-1}}$ and $(X_{\tau_{k-1}}^1, X_{\tau_{k-1}}^2, (W^1, W^2)_{t \leq \tau_{k-1}})$ the coupling is successful under the subprobability \mathbf{P}_1^M . In other words, if we assume that (W^1, W^2) is realized with the previous coupling construction, we have

$$\mathbb{P}(X_{\tau_{k-1}+1}^1 = X_{\tau_{k-1}+1}^2 | \Omega_{K, \alpha, \tau_{k-1}}) \geq \|\mathbf{P}_1^M\|_{TV}.$$

By (3.27) and Lemma C.1. of [15] (applied to $p = 2$, $\mu_1 = \hat{\varphi}^* \mathbb{P}_W$, $\mu_2 = \mathbb{P}_W$ and $X = A_M$)

$$\|\mathbf{P}_1^M\|_{TV} \geq \frac{\left[\int_{A_M} D_{\hat{\varphi}}(w) \mathbb{P}_W(dw) \right]^2}{4 \int_{A_M} D_{\hat{\varphi}}(w)^3 \mathbb{P}_W(dw)}.$$

We will now show that M can be chosen in such a way that the above quantity is bounded away from 0 independently of $k \in \mathbb{N}$. On the one hand, by exhibiting an exponential martingale and by using (3.25),

$$\begin{aligned} \int_{A_M} D_{\hat{\varphi}}(w)^3 \mathbb{P}_W(dw) &\leq \left(\sup_{w \in A_M} \exp\left(3 \int_0^1 |g_h(s)|^2 ds\right) \right) \times \\ &\int \exp\left(3 \int_0^1 (\bar{g}_h(s) |dw_s| - \frac{3^2}{2} \int_0^1 |\bar{g}_h(s)|^2 ds\right) \mathbb{P}_W(dw) \leq \exp(3\bar{C}(M, K)). \end{aligned}$$

On the other hand,

$$\int_{A_M} D_{\hat{\varphi}}(w) \mathbb{P}_W(dw) = \mathbb{P}_W(\varphi^{-1}(A_M)).$$

For every $M > \bar{K}$, we deduce from (3.24) that $\varphi(A_{M-\bar{K}}) \subset A_M$ so that $A_{M-\bar{K}} \subset \varphi^{-1}(A_M)$. Furthermore, we get from (i) for the choice

$$M := C(K, \tilde{K}) + \bar{K}$$

that

$$\{w, \|w\|_{\frac{1}{2}-\varepsilon_\theta}^{0,1} \leq \tilde{K}\} \subset A_{M-\bar{K}}.$$

As a consequence, $\int_{A_M} D_{\tilde{\varphi}}(w) \mathbb{P}_W(dw) \geq \mathbb{P}_W(\{w, \|w\|_{\frac{1}{2}-\varepsilon_\theta}^{0,1} \leq \tilde{K}\})$ and

$$\|\mathbf{P}_1^M\|_{TV} \geq \frac{\left[\mathbb{P}_W(\{w, \|w\|_{\frac{1}{2}-\varepsilon_\theta}^{0,1} \leq \tilde{K}\})\right]^2}{4 \exp(3\tilde{C}(C(K, \tilde{K}) + \tilde{K}, K))} > 0$$

which concludes the proof.

(b) When Step 1 is successful, this property follows from (3.25). If Step 1 is not attempted (and thus fails) since $\omega \in \Omega_{K, \alpha, \tau_{k-1}}^c$, $W^1 = W^2$ on $[\tau_{k-1}, \tau_{k-1} + 1]$ so that g_w is null on $[\tau_{k-1}, \tau_{k-1} + 1]$. If Step 1 is attempted and fails, it follows from the above construction of the coupling that $w_1 = w_2$ or $w_2 = \varphi^{-1}(w_1)$ with $w_1 \in A_M$. Then, since the control of the functions \tilde{f}_h (defined in (iii)) and its derivative are similar to that of f_h in (iv), we deduce that the L^2 -norm of $\tilde{g}_h(t) = g_w(t + \tau_{k-1})$ can be also bounded in a similar way.

(c) When Step 1 is successful, $g_B(t + \tau_{k-1}) = f_h(t)$ on $[0, 1]$ and the boundedness of f_h follows from that of ρ_h which is proved in (ii). \square

3.2 Step 2

As explained before, Step 2 is a series of trials on some intervals I_ℓ of length $c_2 2^\ell$ (the first one of length $2c_2$, the second one of length $4c_2, \dots$). We denote by $s_{k,\ell}$ the left extreme of each interval I_ℓ . More precisely, for every $k \geq 1$, we define $(s_{k,\ell})_{\ell \geq 0}$ by

$$s_{k,0} = s_{k,1} = \tau_{k-1} + 1 \quad \text{and for every } \ell \geq 1 \quad s_{k,\ell+1} = s_{k,\ell} + c_2 2^\ell. \quad (3.28)$$

Also denote by ℓ_k^* , the (first) trial after time τ_{k-1} where Step 2 fails. The case $\ell_k^* = 0$ and $\ell_k^* = +\infty$ correspond respectively to the failure of Step 1 and to the success of Step 2. For some given positive α and K , we set

$$\mathcal{B}_{k,\ell} := \Omega_{K,\alpha,\tau_{k-1}} \cap \{\ell_k^* > \ell\}, \quad k \geq 1, \ell \geq 0. \quad (3.29)$$

With this definition,

$$\mathbb{P}(\tau_k = +\infty | \Omega_{K,\alpha,\tau_{k-1}}) = \mathbb{P}(X_{\tau_{k-1}+1}^1 = X_{\tau_{k-1}+1}^2 | \Omega_{K,\alpha,\tau_{k-1}}) \prod_{\ell=1}^{+\infty} \mathbb{P}(\mathcal{B}_{k,\ell} | \mathcal{B}_{k,\ell-1}). \quad (3.30)$$

Consequently, the aim is now to lower-bound $\mathbb{P}(\mathcal{B}_{k,\ell} | \mathcal{B}_{k,\ell-1})$. This is the purpose of the next lemma. Before doing so, let us introduce some notations. Owing to the one-to-one correspondence between g_w and g_B , there is a unique choice for function g_w in $[\tau_{k-1} + 1, \infty)$ which ensures that $g_B(t) = 0$ after $\tau_{k-1} + 1$ (or equivalently that $B_t^1 = B_t^2$ after $\tau_{k-1} + 1$). We denote it by g_s in the next lemma (see the proof for an explicit expression of g_s).

LEMMA 3.2. *Let $K > 0$ and assume that $\alpha \in (0, H)$. There exists a constant $C_K \geq 1$ which does not depend on k such that,*

$$\int_0^{+\infty} (1+t)^{2\alpha} |g_s(\tau_{k-1} + 1 + t)|^2 dt \leq C_K.$$

Then, (W_1, W_2) can be constructed during Step 2 in such a way that for all k and ℓ ,

$$\rho_K^1 \leq \mathbb{P}(\mathcal{B}_{k,1} | \mathcal{B}_{k,0}) \leq \rho_K^2 \quad \text{and} \quad \forall \ell \geq 2, \quad (1 - \rho_K^3 2^{-\alpha\ell}) \leq \mathbb{P}(\mathcal{B}_{k,\ell} | \mathcal{B}_{k,\ell-1}) \leq (1 - \rho_K^3 2^{-\alpha\ell-1}) \quad (3.31)$$

where $\rho_K^1, \rho_K^2 \in (0, 1)$ do not depend on k and $\rho_K^3 = c_2^{-\alpha} \sqrt{C_K}$. In particular, if $c_2 = C_K^{\frac{1}{2\alpha}}$, $\rho_K^3 = 1$ and in this case, if $2 \leq \ell_k^* < +\infty$ one has

$$\int_{s_{k,\ell_k^*}}^{s_{k,\ell_k^*+1}} |g_w(t)|^2 dt \leq (2(\ell_k^* + 3))^2 \quad \text{and} \quad \forall \ell \in \{2, \dots, \ell_k^*\}, \quad \int_{s_{k,\ell-1}}^{s_{k,\ell}} |g_w(t)|^2 dt \leq 2^{-2\alpha\ell}.$$

and if $\ell_k^* = 1$, $\int_{s_{k,1}}^{s_{k,2}} |g_w(t)|^2 dt \leq C'_K$ where C'_K is a finite constant.

REMARK 3.5. The lower-bounds obtained in (3.31) ensure the strict positivity of $\mathbb{P}(\tau_k = +\infty | \Omega_{K,\alpha,\tau_{k-1}})$. The other properties will be needed for the sequel. Note that c_2 can be chosen in such a way that the involved quantities do not depend on K except if $\ell = 1$.

Proof. We first remark that if at a positive stopping time T_1 one has $X_{T_1}^1 = X_{T_1}^2$, then (X_t^1) and (X_t^2) remain equal on $[T_1, T_2]$ (where $T_2 > T_1$ is a second stopping time) if and only if $g_B(t) = 0$ on $(T_1, T_2]$. By Lemma 4.3 in [9], and its proof, this condition is satisfied if and only if

$$\forall t \in (0, T_2 - T_1], g_w(t + T_1) = g_s(t + T_1) := \mathcal{R}_0 g_w^{T_1}(t).$$

The interesting point is that the above function is \mathcal{F}_{T_1} -measurable (the context is thus different from Step 1, where the function denoted by g_h was defined in a dynamic way). In particular, by conditioning on $\mathcal{F}_{s_{k,\ell}}$ one can write:

$$\mathbb{P}(\mathcal{B}_{k,\ell} | \mathcal{B}_{k,\ell-1}) = \mathbb{E}(Q(\mathcal{R}_0 g_w^{\tau_{k-1}+1}(s_{k,\ell} + \cdot), c_2 2^\ell))$$

where for positive T and a (deterministic) measurable function g on $[0, +\infty)$ we denote

$$Q(g, T) = \mathbb{P}(\forall t \in [0, T], \tilde{W}_t^2 = \tilde{W}_t^1 + g(t)).$$

for $(\tilde{W}^1, \tilde{W}^2)$ a given couple of Brownian motions on $[0, T]$. By Lemma 3.4, if $\|g\|_{[0,T],2} \leq b \leq b_0$, we can build the couple $(\tilde{W}_1, \tilde{W}_2)$ in such a way that $Q(g, T) \geq (1 - b) \vee \rho$ where ρ depends only on b_0 .

Following carefully the proof of Lemma 5.12. of [9] (see in particular (5.36) therein), one deduces from Lemma 3.1(b) and Condition (2.9) that on $\mathcal{B}_{k,0}$,

$$\int_0^{+\infty} (1+t)^{2\alpha} \mathcal{R}_0 g_w^{\tau_{k-1}+1}(t) dt \leq C_K$$

for some positive constant C_K . Without loss of generality, we can assume that $C_K \geq 1$. This yields the first property of the lemma and this easily implies that for every $\ell \geq 1$,

$$\int_{s_{k,\ell}}^{s_{k,\ell+1}} |g_s(u)|^2 du \leq b_\ell^2$$

with $b_1 = \sqrt{C_K}$ and $b_\ell = c_2^{-\alpha} \sqrt{C_K} 2^{-\alpha\ell}$ if $\ell \geq 2$. It remains to apply Lemma 3.4 ((i) for $\ell = 1$ and (ii) for $\ell \geq 2$) to obtain (3.31). Finally, the bound for $\int_{s_{k,\ell_k}^*}^{s_{k,\ell_k+1}^*} |g_w(t)|^2 dt$ follows from the value of M_{b_ℓ} given by Lemma 3.4. □

LEMMA 3.3. *Let K and b be positive numbers.*

(i) *Then, there exist $M_b > 0$, ρ_b^1 and $\rho_b^2 \in (0, 1)$ such that for every $a \in [-b, b]$, we can build a random variable (U_1, U_2) with values in \mathbb{R}^2 such that*

$$\mathcal{L}(U_1) = \mathcal{L}(U_2) = \mathcal{N}(0, 1), \quad \rho_b^1 \leq \mathbb{P}(U_2 = U_1 + a) \leq \rho_b^2, \quad \mathbb{P}(|U_2 - U_1| \leq M_b) = 1$$

and on the event $\{U_2 = U_1 + a\}$, $|U_1| \leq \tilde{M}_b$ and $|U_2| \leq \tilde{M}_b$ hold a.s., where $\tilde{M}_b \leq \frac{M_b}{2} + b$.

(ii) *Furthermore, if $b \in (0, 1)$, the previous statement holds with $M_b \leq \max\{4b, -2 \log(b/8)\}$, $\rho_b^1 = 1 - b$ and $\rho_b^2 = 1 - \frac{b}{2}$.*

REMARK 3.6. In order to ensure the (K, α) -admissibility condition at the next trials, one needs to control the increments of W^1 and W^2 during Step 2. In particular, when Step 2 fails, we will need the probability of success to be not too large. This explains the property of domination of the probability of success $\mathbb{P}(U_2 = U_1 + a)$ (and $\mathbb{P}(W^2 = W^1 + g)$ in the next result) which may appear of poor interest. For the same reason, we give in the following result an explicit construction of W^1 and W^2 during Step 2.

Proof. (ii) is almost the statement of Lemma 5.13 of [9]. The only new points are the deterministic control of $|U_1|$ and $|U_2|$ on the event $\{U_2 = U_1 + a\}$ and the domination of the probability of success by $\rho_b^2 = 1 - \frac{b}{2}$. With the notations of [9], the first property follows from the construction of the measure \mathcal{N}_3 which is such that for every $a \in [-b, b]$, the support of \mathcal{N}_3 is included in $[-M_b/2, M_b/2 - a] \times [-M_b/2 - a, M_b/2]$. For the second one, it is enough to note that the probability of success introduced in Lemma 5.13 of [9] and denoted by $\mathcal{N}_3(L_3)$ is a non-decreasing continuous function of M and equal to 0 if $M_b = 0$. Thus, the domination of this probability can be obtained by reducing sufficiently the value of M .

On the other hand, (i) is in some sense a rough version of (ii). Its proof can also be done by following the lines of the lemma of [9] and by checking that for every $b > 0$, we can choose M_b large enough such that $\inf_{a \in [-b, b]} \mathcal{N}_3(L_3) > 0$. \square

The following lemma is a slightly modified version of Corollary 5.14 of [9].

LEMMA 3.4. *Let T and b be positive numbers and $g \in L^2([0, T], \mathbb{R})$ with $\|g\|_{2, [0, T]} \leq b$.*

(i) There exists $M_b > 0$, $\rho_b^1, \rho_b^2 \in (0, 1)$ and a couple of Wiener processes (W^1, W^2) defined in $[0, T]$ such that

$$\rho_b^1 \leq \mathbb{P}\left(W_t^2 = W_t^1 + \int_0^t g(s)ds, t \in [0, T]\right) \leq \rho_b^2 \quad \text{and} \quad \mathbb{P}(\|W^2 - W^1\|_{2, [0, T]} \leq M_b) = 1. \quad (3.32)$$

Furthermore, there exists a triple (U_1, U_2, V) of standard normally distributed random variables and a Brownian motion \tilde{W} such that (U_1, U_2) and (V, \tilde{W}) are independent,

$$W_t^i = (U^i + V) \frac{\int_0^t g(s)ds}{\|g\|_{2, [0, T]}} + \tilde{W}_t, \quad t \in [0, T] \quad \text{for } i = 1, 2, \quad (3.33)$$

and moreover $|U_i| \leq \tilde{M}_b := \frac{M_b}{2} + b$ on the event $\{W_t^2 = W_t^1 + \int_0^t g(s)ds, t \in [0, T]\}$.

(ii) Furthermore, if $b \in (0, 1)$, the previous statement holds with $M_b = \max\{4b, -2 \log(b/8)\}$, $\rho_b^1 = 1 - b$ and $\rho_b^2 = 1 - \frac{b}{2}$.

Proof. (i) Let $(f_k)_{k \geq 1}$ denote a complete orthonormal basis of $L^2([0, T], \mathbb{R})$ with $f_1 = g/\|g\|_{2, [0, T]}$. In some probability space $(\Omega', \mathcal{F}', \mathbb{P}')$ let (U_1, U_2) be a couple of random variables satisfying the properties of Lemma 3.3(i) and $(\xi_k)_{k \geq 2}$ be a sequence independent of (U_1, U_2) of i.i.d. random variables with $\mathcal{L}(\xi_2) = \mathcal{N}(0, 1)$. Defining for $i = 1, 2$ the process $W_t^i := \mathcal{W}^i(\mathbf{1}_{[0, t]}), t \geq 0$, where $\mathcal{W}^i : L^2([0, T], \mathbb{R}) \rightarrow \mathcal{W}^i(L^2([0, T], \mathbb{R})) \subset L^2(\Omega', \mathcal{F}', \mathbb{P}')$ is the isometry of Hilbert spaces such that $\mathcal{W}^i(f_1) = U^i$ and $\mathcal{W}^i(f_k) = \xi_k, k \geq 2$, one easily checks (by computing covariances) that

$$W_t^i = U_i \frac{\int_0^t g(s)ds}{\|g\|_{2, [0, T]}} + \sum_{k \geq 2} \xi_k \int_0^t f_k(s)ds$$

and that (W_t^i) is a standard Brownian Motion. It follows from Lemma 3.3(i) and from the previous construction that (3.32) holds. Furthermore, introducing artificially a last standard normally distributed random variable V independent of $\sigma(U_1, U_2, \xi_k, k \geq 2)$, we can write W^i as follows

$$W_t^i = (U_i + V) \frac{\int_0^t g(s)ds}{\|g\|_{2, [0, T]}} + \tilde{W}_t, \quad t \in [0, T]$$

where $\tilde{W}_t := -V \int_0^t g(s)ds/\|g\|_{2, [0, T]} + \sum_{k \geq 2} \xi_k \int_0^t f_k(s)ds$ is a standard Brownian motion independent of (U_1, U_2) . Finally, the boundedness property of U_i on $\{W_t^2 = W_t^1 + \int_0^t g(s)ds, t \in [0, T]\}$ again follows from that obtained in Lemma 3.3(i).

(ii) The proof is identical using the properties of Lemma 3.3(ii) instead of those of (i). \square

4 About the (K, α) -admissibility condition

In this section, we assume that Steps 1 and 2 are carried out as described previously, and the aim is to ensure that the system is (K, α) -admissible with positive probability at every times τ_k . This is the purpose of the next proposition:

PROPOSITION 4.2. *Let $(X_t^1, X_t^2)_{t \geq 0}$ denote a solution of (2.6) with initial condition $\tilde{\mu}$ satisfying $\tilde{\mu}(|x_1|^r + |x_2|^r) < +\infty$ for some $r > 0$. Assume (\mathbf{H}_0) , (\mathbf{H}_1) and (\mathbf{H}_2) . Let $\alpha \in (0, 1/2)$. Assume that for each $K > 0$, c_2 defined in (2.13) satisfies $c_2 = C_K^{\frac{1}{2\alpha}}$ (where C_K is a constant greater than 1 defined in Lemma 3.2) and that for every $k \geq 1$ and $\ell \geq 0$, $\Delta_3(\ell, k)$ introduced in (2.16) is defined by $\Delta_3(\ell, k) = c_3 a_k 2^{\beta \ell}$ with $\beta > (1 - 2\alpha)^{-1}$, $a_k = \zeta^k$ for some (arbitrary) fixed $\zeta > 1$, and c_3 an appropriate constant depending on the previous parameters (see Proposition 4.6 and Remark 4.8 for details). Then, for every $\varepsilon > 0$, there exists $K_\varepsilon > 0$ such that for every $k \geq 0$,*

$$\mathbb{P}(\Omega_{K_\varepsilon, \alpha, \tau_k} | \tau_k < +\infty) \geq 1 - \varepsilon.$$

The proof of this proposition is divided into two parts corresponding respectively to Conditions (2.9) and (2.10). The first concerns the coupling function g_w and the proof corresponding to this condition easily follows from [9] (see Subsection 4.4 for details).

REMARK 4.7. In the sequel of this section, we always assume that α is a fixed number in $(0, 1/2)$ and that $c_2 = C_K^{\frac{1}{2\alpha}}$. These properties are not recalled again in each statement.

The lower-bound for the second condition is obtained in the next subsections.

4.1 (K, α) -admissibility and Lyapunov

We denote in what follows

$$\mathcal{E}_k := \{\tau_k < \infty\} (= \{\tau_1 < \infty, \dots, \tau_k < \infty\}).$$

Let $\alpha \in (0, 1/2)$. We want to prove that for every $\varepsilon > 0$, there exists $K_\varepsilon > 0$ such that

$$\mathbb{P}(\Omega_{K_\varepsilon, \tau_k}^2 | \mathcal{E}_k) \geq 1 - \varepsilon.$$

But since for every events A_1, A_2, A_3 and A_4 , $\mathbb{P}(\cap_{i=1}^4 A_i) \geq \sum_{i=1}^4 \mathbb{P}(A_i) - 3$, it is enough to prove that for every $\varepsilon > 0$, there exists $K_\varepsilon > 0$ such that for $j = 1, 2$,

$$\mathbb{P}(\varphi_{\tau_k, \varepsilon_\theta}(W^j) \leq K_\varepsilon | \mathcal{E}_k) \geq 1 - \varepsilon, \quad j = 1, 2 \quad (4.34)$$

and

$$\mathbb{P}(|X_{\tau_k}^j| \leq K_\varepsilon | \mathcal{E}_k) \geq 1 - \varepsilon \quad j = 1, 2. \quad (4.35)$$

Since the arguments to prove (4.34) are contained in those needed for the (4.35), we defer the proof of the former to the appendix (see Appendix B) and we only focus on the second statement. The proof of this property is based on a Lyapunov-type argument: owing to the Markov inequality, it is obvious that (4.35) will be true if one exhibits a positive function $\Psi : \mathbb{R}^d \rightarrow \mathbb{R}$ such that $\lim_{|x| \rightarrow +\infty} \Psi(x) = +\infty$ and for which there exists a finite positive constant C such that for every $k \in \mathbb{N} \cup \{0\}$ and for every $K > 0$,

$$\mathbb{E}(\Psi(X_{\tau_k}^j) | \mathcal{E}_k) \leq C \quad j = 1, 2. \quad (4.36)$$

Note that since the construction of Step 1 depends on K , the independence of C with respect to K is primordial. To this end, we first introduce the following contraction assumption depending on $\theta \in (1/2, H)$.

H₁'(θ): There exists a subquadratic continuous function $\Psi : \mathbb{R}^d \mapsto \mathbb{R}_+^*$ satisfying

$$\lim_{|x| \rightarrow +\infty} \Psi(x) = +\infty \text{ and } \exists \rho \in (0, 1) \text{ and } C > 0 \text{ such that a.s., } \forall x \in \mathbb{R}^d,$$

$$\Psi(X_1) \leq \rho \Psi(x) + C(1 + \|B\|_\theta^{0,1}).$$

In the previous assumption, $(X_t)_{t \geq 0}$ denotes a solution of (1.2) and *subquadratic* means that there exists $C > 0$ such that for every $x \in \mathbb{R}^d$, $\Psi(x) \leq C(1 + |x|^2)$. In Subsection 4.2, we will prove that, under the Lyapunov assumption (\mathbf{H}_1) , $\mathbf{H}'_1(\theta)$ is true. As detailed in the next proposition, $\mathbf{H}'_1(\theta)$ leads to (4.36) if the following condition, which will be proved in subsection 4.3, is also true:

$\mathbf{H}'_2(\theta)$: For every $\rho \in (0, 1)$, there exists $C_\rho > 0$ such that for every $k \in \mathbb{N}$ and $K > 0$

$$\mathbb{E}\left[\sum_{u=1}^{\Delta\tau_k} \rho^{\Delta\tau_k - u} \|B\|_{\theta}^{\tau_{k-1} + u - 1, \tau_{k-1} + u} |\mathcal{E}_k|\right] \leq C_\rho.$$

PROPOSITION 4.3. *Let $\theta \in (1/2, H)$ and assume $\mathbf{H}'_1(\theta)$. Let $(X_t^1, X_t^2)_{t \geq 0}$ denote pair of solutions to (2.6) with initial condition $\tilde{\mu}$ satisfying $\tilde{\mu}(\Psi^2(x_1) + \Psi^2(x_2)) < +\infty$. For $x_1, x_2 \in \mathbb{R}^d$, set*

$$\tau_0(x_1, x_2) := \inf\{u \in \mathbb{N}, \rho^u(\Psi(x_1) + \Psi(x_2)) \leq 1\}. \quad (4.37)$$

Assume that $(\tau_k)_{k \geq 1}$ is built in such a way that, $\mathbf{H}'_2(\theta)$ holds, that for every $k \geq 1$, $\mathbb{P}(\mathcal{E}_k | \mathcal{E}_{k-1}) \geq \delta_1 > 0$ (where δ_1 is a positive number which does not depend on k) and that $\Delta\tau_k \geq \frac{\log(\delta_1/2)}{\log \rho}$. Then, there exists a positive constant C such that for every $k \in \mathbb{N} \cup \{0\}$ and $K > 0$,

$$\mathbb{E}(\Psi(X_{\tau_k}^j) | \mathcal{E}_k) \leq C, \quad j = 1, 2.$$

REMARK 4.8. \triangleright Under the assumptions of Proposition 4.2, $\Delta\tau_k \geq c_3$ (see (2.16)). To ensure that $\Delta\tau_k \geq \frac{\log(\delta_1/2)}{\log \rho}$, one can thus choose c_3 large enough in order that $c_3 \geq \frac{\log(\delta_1/2)}{\log \rho}$.

\triangleright By the elementary inequalities $|u + v|^p \leq |u|^p + |v|^p$ and $|u|^p \leq C(1 + |u|)$ for $p \in (0, 1)$, one remarks that if $\mathbf{H}'_1(\theta)$ holds for Ψ , it also holds for Ψ^p if $p < 1$. Since Ψ is subquadratic, it follows that one can assume without loss of generality that $\Psi^2(x) \leq C(1 + |x|^r)$ for some given $r > 0$. This explains the assumption $\tilde{\mu}(|x_1|^r + |x_2|^r) < +\infty$ in Proposition 4.2.

Proof. By $\mathbf{H}'_1(\theta)$ and an induction

$$\Psi(X_{\tau_k}^j) \leq \rho^{\Delta\tau_k} \Psi(X_{\tau_{k-1}}^j) + C \sum_{\ell=1}^{\Delta\tau_k} \rho^{\Delta\tau_k - \ell} (1 + \|B\|_{\theta}^{\tau_{k-1} + \ell - 1, \tau_{k-1} + \ell}).$$

First, since $\Delta\tau_k \geq \frac{\log(\delta_1/2)}{\log \rho}$, we deduce that $\rho^{\Delta\tau_k} \leq \frac{\delta_1}{2}$. Thus,

$$\mathbb{E}[\Psi(X_{\tau_k}^j) | \mathcal{E}_k] \leq \frac{\delta_1}{2} \mathbb{E}[\Psi(X_{\tau_{k-1}}^j) | \mathcal{E}_k] + C \sum_{u=0}^{+\infty} \rho^u + C \mathbb{E}\left[\sum_{\ell=1}^{\Delta\tau_k} \rho^{\Delta\tau_k - \ell} \|B\|_{\theta}^{\tau_{k-1} + \ell - 1, \tau_{k-1} + \ell} | \mathcal{E}_k\right].$$

Since $\mathcal{E}_k \subset \mathcal{E}_{k-1}$ and $\mathbb{P}(\mathcal{E}_k | \mathcal{E}_{k-1}) \geq \delta_1$, $\mathbb{E}[\Psi(X_{\tau_{k-1}}^j) | \mathcal{E}_k] \leq \delta_1^{-1} \mathbb{E}[\Psi(X_{\tau_{k-1}}^j) | \mathcal{E}_{k-1}]$. It follows that

$$\mathbb{E}[\Psi(X_{\tau_k}^j) | \mathcal{E}_k] \leq \frac{1}{2} \mathbb{E}[\Psi(X_{\tau_{k-1}}^j) | \mathcal{E}_{k-1}] + \frac{C}{1 - \rho} + C \mathbb{E}\left[\sum_{\ell=1}^{\Delta\tau_k} \rho^{\Delta\tau_k - \ell} \|B\|_{\theta}^{\tau_{k-1} + \ell - 1, \tau_{k-1} + \ell} | \mathcal{E}_k\right].$$

Assumption $\mathbf{H}'_2(\theta)$ combined with an induction then yields

$$\sup_{k \geq 0} \mathbb{E}[\Psi(X_{\tau_k}^j) | \mathcal{E}_k] \leq \mathbb{E}[\Psi(X_{\tau_0}^j) | \mathcal{E}_0] + \tilde{C}_\rho$$

where \tilde{C}_ρ neither depends on k and j nor on the starting condition $\tilde{\mu}$. Noticing that $\mathcal{E}_0 = \Omega$, it remains to bound $\mathbb{E}[\Psi(X_{\tau_0}^j)]$. By the definition of τ_0 (which is \mathcal{F}_0 -measurable) and the Cauchy-Schwarz inequality,

$$\mathbb{E}_{\tilde{\mu}}[\Psi(X_{\tau_0}^j)] \leq \sum_{u=0}^{+\infty} \mathbb{E}_{\tilde{\mu}}[\Psi^2(X_u^j)]^{\frac{1}{2}} (\tilde{\mu}(\tau_0 = u))^{\frac{1}{2}}. \quad (4.38)$$

On the one hand, checking that for $\varepsilon > 0$, there exists $C_\varepsilon > 0$ such that for all u, v of \mathbb{R}^d , $|u + v|^2 \leq (1 + \varepsilon)|u|^2 + C_\varepsilon|v|^2$, one deduces from $\mathbf{H}'_1(\theta)$ that there exists $0 < \tilde{\rho} < 1$ and $C_{\tilde{\rho}}$ such that for every starting point x ,

$$\Psi^2(X_1) \leq \tilde{\rho}\Psi^2(x) + C_{\tilde{\rho}}(1 + \|B\|_\theta^{0,1})^2.$$

Thus, it again follows from an induction and from the stationarity of the increments of the fBm that

$$\mathbb{E}_{\tilde{\mu}}[\Psi^2(X_u^j)] \leq \int \Psi^2(x_j)\tilde{\mu}_j(dx_j) + \frac{C_{\tilde{\rho}}}{1 - \tilde{\rho}}\mathbb{E}[(1 + \|B\|_\theta^{0,1})^2] < +\infty,$$

since $\int \Psi^2(x_j)\tilde{\mu}_j(dx_j) < +\infty$. It remains to control the queue of τ_0 . We have

$$\tilde{\mu}(\tau_0 \geq u) \leq \sum_{j=1}^2 \tilde{\mu}_j(\rho^u \Psi(x_j) > \frac{1}{2}) \leq 2 \sum_{j=1}^2 \rho^u \int \Psi(x_j)\tilde{\mu}_j(dx_j) \leq C\rho^u. \quad (4.39)$$

Plugging the previous inequality in yields the boundedness of $\mathbb{E}_{\tilde{\mu}}[\Psi(X_{\tau_0}^j)]$. \square

As a consequence, it remains now to prove $\mathbf{H}'_1(\theta)$ and $\mathbf{H}'_2(\theta)$. This is the purpose of the next subsections.

4.2 Proof of $\mathbf{H}'_1(\theta)$

PROPOSITION 4.4. *Assume (\mathbf{H}_1) . Then, $\mathbf{H}'_1(\theta)$ holds for every $\theta \in (\frac{1}{2}, H)$ with $\Psi = V^{\frac{2\theta-1}{4}}$.*

Proof. The proof is divided in four steps. In all of them, we assume that $0 \leq s < t \leq 1$.

Step 1. We prove the following statement: there exists $C > 0$ such that

$$|X_t| \leq C \left(|X_s| + C(t-s) + \left| \int_s^t \sigma(X_u)dB_u \right| \right) \quad a.s. \quad (4.40)$$

Actually, using that b is a sublinear function,

$$|X_t| \leq |X_s| + C(t-s) + \left| \int_s^t \sigma(X_u)dB_u \right| + \int_s^t |X_u|du.$$

The result then follows from Gronwall lemma (note that the time-dependence of the Gronwall constant does not appear since $s, t \in [0, 1]$).

Step 2. *Control of the Hölder norm of X in a small (random) interval:* Let $\theta \in (1/2, H)$. We show that there exist some positive constants c_0 and C such that for every $0 \leq s < t \leq 1$, satisfying $c_0(1 + \|B\|_\theta^{0,1})(t-s)^\theta \leq \frac{1}{2}$,

$$\|X\|_\theta^{s,t} \leq C \left(\|B\|_\theta^{0,1} + (1 + |X_s|)(t-s)^{1-\theta} \right). \quad (4.41)$$

Let us prove this property. Owing to the classical controls of Young integrals (see *e.g.* [19], Inequality (10.9)), for every $(p, q) \in (0, 1]^2$ with $p+q > 1$, there exists $C_{p,q} > 0$ such that for every p -Hölder and q -Hölder functions f and g (respectively), for every $0 \leq s < t \leq 1$,

$$\left| \int_s^t f(u)dg(u) - f(s)(g(t) - g(s)) \right| \leq C_{p,q} \|f\|_p^{0,1} \|g\|_q^{0,1} (t-s)^{p+q}. \quad (4.42)$$

Applying the previous inequality with $p = q = \theta$ and using that σ is Lipschitz continuous and bounded, we deduce that for every $0 \leq s \leq u < v \leq t \leq 1$,

$$\begin{aligned} \left| \int_u^v \sigma(X_w)dB_w \right| &\leq C \|B\|_\theta^{u,v} (v-u)^\theta (\|X\|_\theta^{u,v} (v-u)^\theta + \|\sigma\|_\infty) \\ &\leq C \|B\|_\theta^{0,1} (v-u)^\theta (\|X\|_\theta^{s,t} (t-s)^\theta + \|\sigma\|_\infty) \end{aligned} \quad (4.43)$$

By (4.40) and what precedes, we also have

$$\int_u^v |b(X_r)| dr \leq C(v-u) \left(1 + |X_s| + \|B\|_\theta^{0,1}(t-s)^\theta(1 + \|X\|_\theta^{s,t}(t-s)^\theta)\right).$$

Using the previous inequalities, we deduce that

$$\|X\|_\theta^{s,t} \leq C \left(\|B\|_\theta^{0,1} + (1 + |X_s|)(t-s)^{1-\theta}\right) + C\|X\|_\theta^{s,t}\|B\|_\theta^{0,1}(t-s)^\theta$$

and (4.41) follows.

Step 3. *Control of sup $|X_u|$ in a small (random) interval:* let $\theta \in (1/2, H)$. There exists $C > 0$ such that for every $0 \leq s \leq t \leq 1$ satisfying $c_0(1 + \|B\|_\theta)(t-s)^\theta \leq \frac{1}{2}$

$$\sup_{s \leq u \leq t} |X_u| \leq C(1 + |X_s|). \quad (4.44)$$

Actually, using that $\|B\|_\theta^{0,1}(t-s)^\theta \leq (2c_0)^{-1}$, we deduce from (4.43) that for every $0 \leq s \leq t \leq 1$ satisfying $c_0(1 + \|B\|_\theta^{0,1})(t-s)^\theta \leq \frac{1}{2}$

$$\left| \int_s^t \sigma(X_v) dB_v \right| \leq C (\|X\|_\theta^{s,t}(t-s)^\theta + 1).$$

Using again that $\|B\|_\theta^{0,1}(t-s)^\theta \leq (2c_0)^{-1}$, it follows from (4.41) that

$$\left| \int_s^t \sigma(X_v) B_v \right| \leq C(1 + (1 + |X_s|)(t-s)).$$

Then, it is enough to plug this control in (4.40) to obtain (4.44).

Step 4. *Use of the Lyapunov assumption.* Let V be such that Assumption (\mathbf{H}_1) holds. Let $\theta \in (1/2, H)$. Then, there exists $\bar{\rho} \in (0, 1)$ and $C > 0$ such that for every $x \in \mathbb{R}^d$,

$$V(X_1) \leq \bar{\rho}V(x) + C(1 + \|B\|_\theta)^{\frac{4}{2\theta-1}}. \quad (4.45)$$

Let us prove this statement. By e.g. [20] (see Theorem 4.3.1) and Assumption (\mathbf{H}_1) ,

$$\begin{aligned} e^{\kappa_0(t-s)}V(X_t) &= V(X_s) + \int_s^t e^{\kappa_0(u-s)} ((\nabla V|b)(X_u)\kappa_0V(X_u)) du + \int_s^t e^{\kappa_0(u-s)} (\nabla V(X_u)|\sigma(X_u)dB_u) \\ &\leq V(X_s) + \beta_0(t-s) + \int_s^t e^{\kappa_0(u-s)} (\nabla V(X_u)|\sigma(X_u)dB_u). \end{aligned} \quad (4.46)$$

Using that the functions ∇V and σ are Lipschitz continuous, that σ is bounded and that $u \mapsto e^{\kappa_0 u}$ is bounded and Lipschitz continuous on $[0, 1]$, we obtain that for every $0 \leq u < v \leq 1$,

$$|e^{\kappa_0(v-s)}\nabla V\sigma(X_v) - e^{\kappa_0(u-s)}\nabla V\sigma(X_u)| \leq C((1 + |\nabla V(X_v)|)(|X_v - X_u|) + |\nabla V(X_v)|(v-u)).$$

By (4.42), it follows that

$$\begin{aligned} \left| \int_s^t e^{\kappa_0(u-s)} (\nabla V(X_u)|\sigma(X_u)dB_u) \right| &\leq C \left((1 + \sup_{v \in [s,t]} |\nabla V(X_v)|) (\|X\|_\theta^{s,t} + (t-s)^{1-\theta})(t-s)^\theta \right. \\ &\quad \left. + |\nabla V(X_s)| \right) \|B\|_\theta^{s,t}(t-s)^\theta. \end{aligned}$$

From now on, assume that $(1 + \|B\|_\theta^{0,1}(t-s)^\theta) \leq (2c_0)^{-1}$. By (4.44) and the fact that $1 + |\nabla V(x)| \leq C_1(1 + |x|) \leq C_2\sqrt{V}(x)$, we have

$$1 + \sup_{v \in [s,t]} |\nabla V(X_v)| \leq C\sqrt{V(X_s)}.$$

Owing to (4.41) and to some reductions implied by the previous inequality, we obtain

$$\left| \int_s^t e^{\kappa_0(u-s)} (\nabla V(X_u) | \sigma(X_u) dB_u) \right| \leq C \left(V(X_s) \|B\|_\theta^{0,1} (t-s)^{1+\theta} + \sqrt{V}(X_s) \|B\|_\theta^{0,1} (t-s)^\theta \right).$$

Set $\tilde{\theta} := \frac{1}{2}(\theta - \frac{1}{2})$ (so that $2(\theta - \tilde{\theta}) = \frac{1}{2} + \theta$). By the inequality $|xy| \leq \frac{1}{2}(|x|^2 + |y|^2)$,

$$\sqrt{V}(X_s) \|B\|_\theta^{0,1} (t-s)^\theta \leq \frac{1}{2} \left((t-s)^{2(\theta-\tilde{\theta})} V(X_s) + (\|B\|_\theta^{0,1})^2 (t-s)^{2\tilde{\theta}} \right)$$

and on the other hand,

$$V(X_s) \|B\|_\theta^{0,1} (t-s)^{1+\theta} \leq (t-s)^{2(\theta-\tilde{\theta})} V(X_s) \|B\|_\theta^{0,1} (t-s)^{\frac{1}{2}}.$$

Now, we set

$$\eta = (2c_0(1 + \|B\|_\theta^{0,1}))^{-\frac{1}{\theta}} \wedge (1 + \|B\|_\theta^{0,1})^{-\frac{1}{\theta}} \quad (4.47)$$

in order that for every $0 \leq s < t \leq 1$ such that $t-s \leq \eta$,

$$c_0(1 + \|B\|_\theta^{0,1})\eta^\theta \leq \frac{1}{2} \quad \text{and} \quad (\|B\|_\theta^{0,1})^2 (t-s)^{2\tilde{\theta}} \leq 1.$$

For such s, t , we finally obtain (using that $1/2 \geq \tilde{\theta}$ and that $2(\theta - \tilde{\theta}) = \frac{1}{2} + \theta$),

$$\left| \int_s^t e^{\kappa_0(u-s)} (\nabla V(X_u) | \sigma(X_u) dB_u) \right| \leq C(t-s)^{\frac{1}{2}+\theta} V(X_s) + \tilde{\beta} \quad (4.48)$$

where $\tilde{\beta}$ is a positive constant. Plugging this control into (4.46), we deduce: for every $0 \leq s < t \leq 1$ such that $t-s \leq \eta$,

$$V(X_t) \leq e^{-\kappa_0(t-s)} V(X_s) (1 + C(t-s)^{\frac{1}{2}+\theta}) + \hat{\beta}$$

where $\hat{\beta} = \beta_0 \eta + \tilde{\beta}$. Using that $e^{-\kappa_0 u} \leq 1 - \kappa_0 u + \frac{(\kappa_0 u)^2}{2}$ in a right neighborhood of 0 and that $\frac{1}{2} + \theta > 1$, we can find $u_0 \in [0, 1]$ (depending on κ_0, θ and C) such that

$$\forall u \in [0, u_0], \quad e^{-\kappa_0 u} (1 + C u^{\frac{1}{2}+\theta}) \leq 1 - \frac{\kappa_0}{2} u.$$

Thus, for every $0 \leq s < t \leq 1$ such that $t-s \leq \tilde{\eta} := \eta \wedge u_0$,

$$V(X_t) \leq (1 - \frac{\kappa_0}{2}(t-s)) V(X_s) + \hat{\beta}.$$

In particular, applying this control on $[k\tilde{\eta}, ((k+1)\tilde{\eta}) \wedge 1]$ for $k \in \{0, \dots, \lfloor \frac{1}{\tilde{\eta}} \rfloor\}$ yields

$$V(X_1) \leq (1 - \frac{\kappa_0}{2}\tilde{\eta})^{\lfloor \frac{1}{\tilde{\eta}} \rfloor} V(x) + \sum_{k=1}^{\lfloor \frac{1}{\tilde{\eta}} \rfloor} (1 - \frac{\kappa_0}{2}\tilde{\eta})^{\lfloor \frac{1}{\tilde{\eta}} \rfloor - k} \hat{\beta}.$$

It follows from standard computations that

$$V(X_1) \leq \exp(-\frac{\kappa_0}{2} + \tilde{\eta}) V(x) + \frac{2\tilde{\beta}}{\kappa_0 \tilde{\eta}}.$$

We can assume without loss of generality that $u_0 \leq \kappa_0/4$ so that

$$\exp(-\frac{\kappa_0}{2} + \tilde{\eta}) \leq e^{-\frac{\kappa_0}{4}} =: \bar{\rho}.$$

Finally, since $2/\tilde{\theta} \geq 1/\theta$, one can check that there exists $C > 0$ such that

$$\tilde{\eta}^{-1} \leq C(1 + \|B\|_\theta^{0,1})^{\frac{1}{\theta}}.$$

Since $2/\tilde{\theta} = \frac{4}{2\theta-1}$, this concludes the proof of Step 4.

To prove the proposition, it remains now to set $\Psi = V^{\tilde{\theta}}$ and to apply the inequality $|u+v|^{\bar{p}} \leq |u|^{\bar{p}} + |v|^{\bar{p}}$ (which holds for every real numbers u, v and $\bar{p} \in (0, 1]$) with $\bar{p} = \tilde{\theta}$. \square

4.3 Proof of $\mathbf{H}'_2(\theta)$

The main result of this section is Proposition 4.5. Before, we need to establish several lemmas related to the control of the past of the fBm.

Let $j \in \{1, 2\}$. We recall that for every $0 \leq s < t$,

$$B_t - B_s = \alpha_H \left(\int_{-\infty}^s (t-r)^{H-\frac{1}{2}} - (s-r)^{H-\frac{1}{2}} dW_r^j + \int_s^t (t-r)^{H-\frac{1}{2}} dW_r^j \right).$$

This can be rewritten

$$B_t - B_s = \alpha_H \left(\int_{-\infty}^{\lfloor s \rfloor - 1} (t-r)^{H-\frac{1}{2}} - (s-r)^{H-\frac{1}{2}} dW_r^j + \Gamma_1(s, t, W^j) - \Gamma_2(s, t, W^j) + \Gamma_3(s, t, W^j) \right)$$

where, setting $h = t - s$,

$$\begin{aligned} \Gamma_1(s, t, W^j) &= \int_{\lfloor s \rfloor - 1}^{s-h} (t-r)^{H-\frac{1}{2}} - (s-r)^{H-\frac{1}{2}} dW_r^j, \\ \Gamma_2(s, t, W^j) &= \int_{s-h}^s (s-r)^{H-\frac{1}{2}} dW_r^j \quad \Gamma_3(s, t, W^j) = \int_{s-h}^t (t-r)^{H-\frac{1}{2}} dW_r^j. \end{aligned}$$

Let $k \geq 1$. Assume that $\tau_{k-1} < +\infty$ and that $\tau_{k-1} \leq s < t \leq \lfloor s \rfloor + 1$. Setting $\tau_{-1} = -\infty$, we choose to decompose the first right-hand member with respect to the sequence $(\tau_k)_{k \geq -1}$:

$$\int_{-\infty}^{\lfloor s \rfloor - 1} (t-r)^{H-\frac{1}{2}} - (s-r)^{H-\frac{1}{2}} dW_r^j = \sum_{m=0}^k \Lambda_{m,k}(s, t, W^j)$$

with

$$\Lambda_{m,k}(s, t, W^j) = \begin{cases} \int_{\tau_{m-1}}^{\tau_m \wedge \tau_{k-1} - 1} (t-r)^{H-\frac{1}{2}} - (s-r)^{H-\frac{1}{2}} dW_r^j & \text{if } m \in \{0, \dots, k-1\} \\ \int_{\tau_{k-1} - 1}^{\lfloor s \rfloor - 1} (t-r)^{H-\frac{1}{2}} - (s-r)^{H-\frac{1}{2}} dW_r^j & \text{if } m = k. \end{cases}$$

Note that for $i = 1, 2, 3$, Γ_i is related to the local behavior of the fBm whereas for $m = 0, \dots, k$, $\Lambda_{m,k}$ is a memory term. The idea of the sequel of the proof is to bound $\|B\|_{\theta}^{u, u+1}$ ($u \in \{\tau_{k-1}, \dots, \tau_k\}$) through the study of the Γ_i and the $\Lambda_{m,k}$. With a slight abuse of notation, we will sometimes write

$$\|\Gamma_i^j\|_{\theta}^{a,b} = \sup_{a \leq s < t \leq b} \frac{|\Gamma_i(s, t, W^j)|}{|t-s|^{\theta}} \quad \text{and} \quad \|\Lambda_{m,k}\|_{\theta}^{a,b} = \sup_{a \leq s < t \leq b} \frac{|\Lambda_{m,k}(s, t, W^j)|}{|t-s|^{\theta}}. \quad (4.49)$$

The starting point of the study of the $\Lambda_{m,k}$ is the following lemma:

LEMMA 4.5. *Let $a < b < s < t$. Let W be a two-sided Brownian motion. Then,*

$$\frac{1}{t-s} \left| \int_a^b (t-r)^{H-\frac{1}{2}} - (s-r)^{H-\frac{1}{2}} dW_r \right| \leq (t-a)^{H-\frac{3}{2}} |W_b - W_a| + \frac{1}{2} \int_a^b (s-r)^{H-\frac{5}{2}} |W_r - W_b| dr.$$

Proof. By an integration by parts,

$$\begin{aligned} \int_a^b (t-r)^{H-\frac{1}{2}} - (s-r)^{H-\frac{1}{2}} dW_r &= \left((t-a)^{H-\frac{1}{2}} - (s-a)^{H-\frac{1}{2}} \right) (W_a - W_b) \\ &\quad + (H - \frac{1}{2}) \int_a^b \left((t-r)^{H-\frac{3}{2}} - (s-r)^{H-\frac{3}{2}} \right) (W_r - W_b) dr. \end{aligned} \quad (4.50)$$

On the one hand, by the elementary inequality $(1+x)^\rho \geq 1+x$ for every $x \in (-1, 0]$ and $\rho \in (0, 1]$, we remark that

$$\begin{aligned} 0 \leq (t-a)^{H-\frac{1}{2}} - (s-a)^{H-\frac{1}{2}} &= (t-a)^{H-\frac{1}{2}} \left(1 - \left(1 + \frac{s-t}{t-a} \right)^{H-\frac{1}{2}} \right) \\ &\leq (t-s)(t-a)^{H-\frac{3}{2}} \end{aligned}$$

On the other hand, by the inequality $(1+x)^\rho \geq 1+\rho x$ for $x \geq 0$ and $\rho < 0$, we obtain similarly

$$(s-r)^{H-\frac{3}{2}} - (t-r)^{H-\frac{3}{2}} \leq \left(\frac{3}{2} - H \right) (t-s)(s-r)^{H-\frac{5}{2}}.$$

The result follows (using that $(3/2 - H)(H - 1/2) \leq 1/2$). \square

In the next lemma, we propose to bound some quantities which are related to those which appear in the previous lemma on some sub-intervals of $[\tau_{m-1}, \tau_m]$ where $m \in \mathbb{N}$. With the notations introduced in (2.16) and in (3.28), we set

$$\tau_m^0 = \tau_{m-1}, \quad \tau_m^1 = \tau_{m-1} + 1 + 2c_2, \quad \tau_m^2 = s_{m, \ell_m^*} \vee \tau_m^1, \quad \tau_m^3 = s_{m, \ell_m^*+1} \quad \text{and} \quad \tau_m^4 = \tau_m.$$

Since c_3 defined in (2.16) satisfies $c_3 \geq 2c_2$, Step 3 is longer than $2c_2$ and $i \mapsto \tau_m^i$ is non-decreasing. Furthermore, τ_m^0 is the beginning of Step 1, τ_m^1 denotes the end of the first trial of Step 2 (or some time during Step 3) if Step 1 is successful (resp. if Step 1 fails). If Step 1 and the first trial of Step 2 are successful, τ_m^2 and τ_m^3 correspond to the beginning and to the end of the failed trial of Step 2. If $\ell_m^* \in \{0, 1\}$, $\tau_m^1 = \tau_m^2 = \tau_m^3$.

Note that τ_m^1 is defined as the end of the first trial of Step 2, instead of the end of Step 1 as it could be expected. Without going into the technical details, let us remark that this particular cutting of the interval is due to the dependence in K (which appears in the (K, α) -admissibility condition) of the probability of success of the first trial of Step 2 (and that this dependence does not appear for the next trials, see Remark 3.5 for background) and that, in view of Assumption $\mathbf{H}'_2(\theta)$, it is of first importance that the next results be obtained independently of K .

LEMMA 4.6. *Assume that there exists $\delta_1 > 0$ such that for all $m \in \mathbb{N}$ and $K > 0$ $\mathbb{P}(\mathcal{E}_{m+1} | \mathcal{E}_m) \geq \delta_1 > 0$. Then, for every $p \geq 1$ and $\varepsilon \in (0, 1)$, there exists $C_{p, \varepsilon, \delta_1} \in \mathbb{R}_+^*$ such that for every $m \in \mathbb{N}$, $i \in \{0, \dots, 3\}$, $j \in \{1, 2\}$ and $K > 0$,*

(i)

$$\mathbb{E} \left[\left(\int_{\tau_m^i}^{\tau_m^{i+1}} \left| (1 + \tau_m^{i+1} - r)^{-\left(\frac{3}{2} + \varepsilon\right)} (W_{\tau_m^{i+1}}^j - W_r^j) \right| dr \right)^p \middle| \mathcal{E}_m \right] \leq C_{p, \varepsilon, \delta_1}. \quad (4.51)$$

(ii) If $\tau_m^i \neq \tau_m^{i+1}$,

$$\mathbb{E} \left[\left| (\tau_m^{i+1} - \tau_m^i)^{-\left(\frac{1}{2} + \varepsilon\right)} \left(W_{\tau_m^{i+1}}^j - W_{\tau_m^i}^j \right) \right|^p \middle| \mathcal{E}_m \right] \leq C_{p, \varepsilon, \delta_1}, \quad (4.52)$$

REMARK 4.9. The proof of this lemma could be shortened by using some rougher arguments similar to those of the proof of Proposition 4.5 below (see (4.69)). However, such arguments do provide an understanding of what implies the conditioning by $\{\tau_m < +\infty\}$, or in other words, to how the distribution of the Wiener process is deformed by the coupling attempt. To this end and when it is possible (especially in the case $i = 1$), we thus choose an approach by which we try to make explicit these distortions.

Proof. (i) By a change of variable, for every $i \in \{0, 1, 2, 3\}$,

$$\int_{\tau_m^i}^{\tau_m^{i+1}} (1 + \tau_m^{i+1} - r)^{-\left(\frac{3}{2} + \varepsilon\right)} |W_{\tau_m^{i+1}}^j - W_r^j| dr = H_i(\tau_m^{i+1} - \tau_m^i)$$

where for a given $c > 0$,

$$H_i(c) = \int_0^c (1+u)^{-\frac{1}{2}} |W_{\tau_m^j}^j - W_{\tau_m^j - u}^j| \nu(du) \quad \text{with} \quad \nu(dr) = (1+u)^{-1-\varepsilon} du.$$

Noticing that $\nu([0, c]) \leq \varepsilon^{-1}$, we deduce from Jensen inequality that for every $p \geq 1$,

$$(H_i(c))^p \leq \left(\frac{1}{\varepsilon}\right)^{p-1} \int_0^c (1+u)^{-\frac{p}{2}-1-\varepsilon} |W_{\tau_m^j}^j - W_{\tau_m^j - u}^j|^p du. \quad (4.53)$$

Now, we focus successively on cases $i = 0, 1, 2, 3$:

$i = 0$: In this case, $\tau_m^1 - \tau_m^0$ is deterministic and is equal to $\bar{c} := 1 + 2c_2$. Using that $\mathcal{E}_m \subset \mathcal{E}_{m-1}$ and the Cauchy-Schwarz inequality, we have for every $u \in [0, \bar{c}]$ and $m \geq 1$,

$$\mathbb{E}[|W_{\tau_m^1}^j - W_{\tau_m^1 - u}^j|^p | \mathcal{E}_m] \leq \frac{\mathbb{E}[|W_{\tau_m^1}^j - W_{\tau_m^1 - u}^j|^{2p} | \mathcal{E}_{m-1}]^{\frac{1}{2}}}{\mathbb{P}(\mathcal{E}_m | \mathcal{E}_{m-1})^{\frac{1}{2}}}.$$

But, conditionally on $\{\tau_{m-1} < +\infty\}$, $(W_{\tau_{m-1}+u}^j - W_{\tau_{m-1}}^j, u \geq 0)$ is a Brownian motion independent of τ_{m-1} so that

$$\mathbb{E}[|W_{\tau_m^1}^j - W_{\tau_m^1 - u}^j|^{2p} | \mathcal{E}_{m-1}] = u^p.$$

Then, since $\mathbb{P}(\mathcal{E}_m | \mathcal{E}_{m-1}) \geq \delta_1$, one deduces that

$$\sup_{u \in [0, \bar{c}]} u^{-\frac{p}{2}} \mathbb{E}[|W_{\tau_m^1}^j - W_{\tau_m^1 - u}^j|^p | \mathcal{E}_m] \leq \delta_1^{-\frac{1}{2}}. \quad (4.54)$$

Plugging this control into (4.53) yields the result when $i = 0$ with $C_{p,\varepsilon,\delta_1} = \delta_1^{-\frac{1}{2}} \varepsilon^{-p}$.

$i = 1$: If $\ell_k^* \in \{0, 1, 2\}$, $\tau_m^1 = \tau_m^2$. Otherwise, we first write $\mathcal{E}_m = \bigcup \mathcal{A}_{m,\ell}$ where $\mathcal{A}_{m,\ell} = \mathcal{B}_{m,\ell}^c \cap \mathcal{B}_{m,\ell-1}$. We recall that $\mathcal{A}_{m,0}$ corresponds to the failure of Step 1 and for every $\ell \geq 1$, $\mathcal{A}_{m,\ell}$ is the event that Step 2 failed after exactly ℓ trials.

With the notations introduced in (3.28), we recall that on $\mathcal{A}_{m,\ell}$, $\tau_m^1 = s_{m,2}$ and $\tau_m^2 = s_{m,\ell}$. By (4.53), it is enough to show that for every $\ell \geq 3$,

$$\int_0^{s_{m,\ell} - s_{m,2}} (1+u)^{-\frac{p}{2}-1-\varepsilon} \mathbb{E}[|(W_{s_{m,\ell}}^j - W_{s_{m,\ell}-u}^j)|^p | \mathcal{A}_{m,\ell}] du \leq C_{p,\varepsilon,\delta_1}. \quad (4.55)$$

where $C_{p,\varepsilon,\delta_1}$ does not depend on k, m, ℓ and K . With the notations of Lemma 3.2, we know that on the event $\mathcal{A}_{m,\ell}$ with $\ell \geq 2$, we have for all $v \in \{1, \dots, \ell-1\}$,

$$\forall t \in [s_{m,v}, s_{m,v+1}], \quad W_t^2 = W_t^1 + \int_{s_{m,v}}^t g_s(s) ds$$

where g_s is a $\mathcal{F}_{\tau_{m-1}+1}$ -measurable function (defined in Lemma 3.2). Moreover, by Lemma 3.4(ii), which can be applied with $b = 2^{-\alpha v}$ (owing to Lemma 3.2 and Remark 4.7), W^j , $j = 1, 2$ can be realized as follows on $[s_{m,v}, s_{m,v+1}]$:

$$\forall t \in [s_{m,v}, s_{m,v+1}], \quad W_t^j = (U_j^{m,v} + V_{m,v}) \frac{\int_{s_{m,v}}^t g_s(s) ds}{\|g_s\|_{[s_{m,v}, s_{m,v+1}], 2}} + \tilde{W}_{t-s_{m,2}}^m,$$

where $(\tilde{W}_t^m)_{t \geq 0}$ is a standard Brownian motion, $(V_{m,v})_{v \geq 1}$ is a sequence of *i.i.d.* normally distributed random variables, and

$$\forall v \in \{2, \dots, \ell-1\}, \forall \omega \in \mathcal{A}_{m,\ell}, |U_j^{m,v}(\omega)| \leq \frac{1}{2} \max\{2^{2-\alpha v}, -2 \log\left(\frac{2^{-\alpha v}}{8}\right)\} + 2^{-\alpha v} \leq C \log(2^{\alpha v})$$

where C does not only depend on α . Furthermore, $(\tilde{W}_t^m)_{t \geq 0}$ and $(V_{m,v})_{v \geq 1}$ are independent of $(U_1^{m,v}, U_2^{m,v})_v$ and g_S . In particular, $(\tilde{W}_t^m)_{t \geq 0}$ and $(V_{m,v})_{v \geq 1}$ are independent of $\mathcal{A}_{m,\ell}$. Set $s_{m,v}^u = s_{m,v} \vee (s_{m,\ell} - u)$. The above properties imply that for every $u \in [0, s_{m,\ell} - s_{m,2}]$,

$$\begin{aligned} \mathbb{E}[|W_{s_{m,\ell}}^j - W_{s_{m,\ell}-u}^j|^p | \mathcal{A}_{m,\ell}] &\leq C_p \mathbb{E} \left[\left(\sum_{v=2}^{\ell-1} \log(2^{\alpha v}) \frac{\int_{s_{m,v}^u}^{s_{m,v+1}^u} |g_S(s)| ds}{\|g_S\|_{[s_{m,v}, s_{m,v+1}], 2}} \right)^p \middle| \mathcal{A}_{m,\ell} \right] \\ &+ C_p \mathbb{E} \left[\left| \sum_{v=2}^{\ell} V_{m,v} \frac{\int_{s_{m,v}^u}^{s_{m,v+1}^u} g_S(s) ds}{\|g_S\|_{[s_{m,v}, s_{m,v+1}], 2}} \right|^p \middle| \mathcal{A}_{m,\ell} \right] \\ &+ C_p \mathbb{E}[|\tilde{W}_{s_{m,\ell}-s_{m,2}}^m - \tilde{W}_{s_{m,\ell}-s_{m,2}-u}^m|^p]. \end{aligned} \quad (4.56)$$

We focus successively on each term of the right-hand side of the above inequality. First,

$$\mathbb{E}[|\tilde{W}_{s_{m,\ell}-s_{m,2}}^m - \tilde{W}_{s_{m,\ell}-s_{m,2}-u}^m|^p] = u^{\frac{p}{2}} \mathbb{E}[|U|^p] \quad (4.57)$$

where U stands for a normally distributed random variable.

For the first right-hand member term of (4.56), we deduce from the Cauchy-Schwarz inequality that

$$\sum_{v=2}^{\ell-1} \log(2^{\alpha v}) \frac{\int_{s_{m,v}^u}^{s_{m,v+1}^u} |g_S(s)| ds}{\|g_S\|_{[s_{m,v}, s_{m,v+1}], 2}} \leq \left(\sum_{v=2}^{\ell-1} \log(2^{\alpha v})^2 \right)^{\frac{1}{2}} \left(\sum_{v=2}^{\ell-1} \left(\frac{\int_{s_{m,v}^u}^{s_{m,v+1}^u} |g_S(s)| ds}{\|g_S\|_{[s_{m,v}, s_{m,v+1}], 2}} \right)^2 \right)^{\frac{1}{2}}$$

Using that $v \mapsto \log(2^{\alpha v})$ is non-decreasing and that

$$\left(\int_{s_{m,v}^u}^{s_{m,v+1}^u} |g_S(s)| ds \right)^2 \leq (s_{m,v+1}^u - s_{m,v}^u) (\|g_S\|_{[s_{m,v}, s_{m,v+1}], 2})^2, \quad (4.58)$$

we deduce that

$$\sum_{v=2}^{\ell-1} (\log(2^{\alpha v})) \frac{\int_{s_{m,v}^u}^{s_{m,v+1}^u} |g_S(s)| ds}{\|g_S\|_{[s_{m,v}, s_{m,v+1}], 2}} \leq \sqrt{\ell} \log(2^{\alpha \ell}) u^{\frac{1}{2}}.$$

Using that for all $\ell \geq 3$, $s_{m,\ell} - s_{m,2} = c_2 2^{\ell-1}$ with $c_2 \geq 1$, one deduces that for every positive p and ε , there exists $C_{p,\varepsilon}$ such that for every $\ell \geq 3$, $\sqrt{\ell} \log(2^{\alpha \ell}) \leq C_{p,\varepsilon} (s_{m,\ell} - s_{m,2})^{\frac{\varepsilon}{p}}$ (we recall that α is a fixed number of $(0, 1/2)$). As a consequence, the first right-hand member term of (4.56) satisfies for every $u \in [0, s_{m,\ell} - s_{m,2}]$

$$\mathbb{E} \left[\left(\sum_{v=2}^{\ell-1} \log(2^{\alpha v}) \frac{\int_{s_{m,v}^u}^{s_{m,v+1}^u} |g_S(s)| ds}{\|g_S\|_{[s_{m,v}, s_{m,v+1}], 2}} \right)^p \middle| \mathcal{A}_{m,\ell} \right] \leq C_{p,\varepsilon} (s_{m,\ell} - s_{m,2})^\varepsilon u^{\frac{p}{2}} \quad (4.59)$$

where $C_{p,\varepsilon}$ is the constant defined above. Finally, for the second right-hand member term of (4.56), let us define $(X_{m,v}^u)_{v=2}^{\ell-1}$ by

$$\forall v \in \{2, \dots, \ell-1\}, \quad X_{m,v}^u = V_{m,v} \frac{\int_{s_{m,v}^u}^{s_{m,v+1}^u} g_S(s) ds}{\|g\|_{[s_{m,v}, s_{m,v+1}], 2}}.$$

Since $(V_{m,v})_{v \geq 1}$ is centered and independent of g_S and $\mathcal{A}_{m,\ell}$, it follows that $(X_{m,v}^u)_{v=2}^{\ell-1}$ is a sequence of martingale increments under $\mathbb{P}(\cdot | \mathcal{A}_{m,\ell})$. By the Doob inequality and (4.58), we deduce that,

$$\begin{aligned} \mathbb{E} \left[\left| \sum_{v=2}^{\ell-1} V_{m,v} \frac{\int_{s_{m,v}^u}^{s_{m,v+1}^u} g_S(s) ds}{\|g_S\|_{[s_{m,v}, s_{m,v+1}], 2}} \right|^p \middle| \mathcal{A}_{m,\ell} \right] &\leq C_p \mathbb{E} \left[\left(\sum_{v=2}^{\ell-1} \left(\frac{\int_{s_{m,v}^u}^{s_{m,v+1}^u} g_S(s) ds}{\|g_S\|_{[s_{m,v}, s_{m,v+1}], 2}} \right)^2 \right)^{\frac{p}{2}} \middle| \mathcal{A}_{m,\ell} \right] \\ &\leq C_p \left(\sum_{v=2}^{\ell-1} (s_{m,v+1}^u - s_{m,v}^u) \right)^{\frac{p}{2}} \leq C_p u^{\frac{p}{2}}. \end{aligned} \quad (4.60)$$

By (4.57), (4.59) and (4.60), we obtain that there exists $C_{p,\varepsilon} \in \mathbb{R}_+^*$ such that for all $m \in \mathbb{N}$, $\ell \geq 2$, $j \in \{1, 2\}$ and $u \in [0, s_{m,\ell} - s_{m,2}]$,

$$\mathbb{E}[|W_{s_{m,\ell}}^j - W_{s_{m,\ell}-u}^j|^p | \mathcal{A}_{m,\ell}] \leq C_{p,\varepsilon} (s_{m,\ell} - s_{m,2})^\varepsilon u^{\frac{p}{2}}. \quad (4.61)$$

The results follows by plugging this inequality into (4.55).

$\underline{i = 2}$: Here, we consider the interval where Step 2 fails. With the previous notations, $\tau_m^2 = s_{m,\ell}$ and $\tau_m^3 = s_{m,\ell+1}$ on $\mathcal{A}_{m,\ell}$ when $\ell \geq 2$. By a similar strategy as in the case $i = 1$ (see in particular (4.55) and (4.61)), it is enough to show that there exists $C_{p,\varepsilon}$ such that for all $m \in \mathbb{N}$, $\ell \geq 0$, $j \in \{1, 2\}$ and $u \in [0, s_{m,\ell+1} - s_{m,\ell}]$,

$$\mathbb{E}[|W_{s_{m,\ell+1}}^j - W_{s_{m,\ell+1}-u}^j|^p | \mathcal{A}_{m,\ell}] \leq C_{p,\varepsilon} (s_{m,\ell+1} - s_{m,\ell})^\varepsilon u^{\frac{p}{2}}. \quad (4.62)$$

When $\ell = 0, 1$, $\tau_m^3 - \tau_m^2 = 0$ so that the property is obvious. Let us consider the set $\mathcal{B}_{m,\ell-1}$ defined by (3.29). From the very definition, $\mathcal{A}_{m,\ell} \subset \mathcal{B}_{m,\ell-1}$.

By Hölder inequality applied with some $\tilde{p} > 1$ and $\tilde{q} > 1$ such that $1/\tilde{p} + 1/\tilde{q} = 1$, we have

$$\mathbb{E}[|W_{s_{m,\ell+1}}^j - W_{s_{m,\ell+1}-u}^j|^p | \mathcal{A}_{m,\ell}] \leq \mathbb{E}[|W_{s_{m,\ell+1}}^j - W_{s_{m,\ell+1}-u}^j|^{p\tilde{p}} | \mathcal{B}_{m,\ell-1}]^{\frac{1}{\tilde{p}}} \mathbb{P}(\mathcal{A}_{m,\ell} | \mathcal{B}_{m,\ell-1})^{\frac{1}{\tilde{q}}-1}.$$

On the one hand, we deduce from the independence of the increments of the Brownian motion that

$$\mathbb{E}[|W_{s_{m,\ell+1}}^j - W_{s_{m,\ell+1}-u}^j|^{p\tilde{p}} | \mathcal{B}_{m,\ell-1}]^{\frac{1}{\tilde{p}}} = u^{\frac{p}{2}} \mathbb{E}[|U|^{p\tilde{p}}]^{\frac{1}{\tilde{p}}}$$

where U stands for a normally distributed random variable. On the other hand, by (3.31),

$$\mathbb{P}(\mathcal{A}_{m,\ell} | \mathcal{B}_{m,\ell-1}) = \mathbb{P}(\mathcal{B}_{m,\ell}^c | \mathcal{B}_{m,\ell-1}) \geq 2^{-\alpha\ell-1}.$$

Then, for each $\varepsilon \in (0, 1)$, Inequality (4.62) follows by setting $\tilde{q} = (1 - \varepsilon)^{-1}$ (so that $1 - \frac{1}{\tilde{q}} = \varepsilon$).

$\underline{i = 3}$: This corresponds to Step 3. The key point here is that the increments of the Brownian motion after τ_m^3 are independent of the previous coupling attempt so that, denoting by $\Delta_3(m, \ell)$, the length of Step 3 under $\mathcal{A}_{m,\ell}$, we have

$$\mathbb{E}[|W_{\tau_m}^j - W_{\tau_m-u}^j|^p | \mathcal{A}_{m,\ell}] = \mathbb{E}[|W_{\tau_m^3 + \Delta_3(m,\ell)}^j - W_{\tau_m^3 + \Delta_3(m,\ell)-u}^j|^p] = C_p u^{\frac{p}{2}}. \quad (4.63)$$

The result then follows similarly to the case $i = 1$ (see (4.55)).

(ii) This result can be easily derived from the controls established previously. More precisely, cases $i = 0, 1, 2, 3$ can be viewed as particular cases of (4.54), (4.61), (4.62) and (4.63). \square

In the next lemma, we adopt the convention $\sum_{\emptyset} = 1$. Also, let us recall that by (2.16), $\Delta\tau_k \geq c_3 a_k \geq a_k$ since $c_3 \geq 2c_2 \geq 1$ (and that $c_2 \geq 1$ by Remark 4.7).

LEMMA 4.7. *Let Ψ satisfy $\int \Psi(x_1) \bar{\mu}_1(dx_1) + \int \Psi(x_2) \bar{\mu}_2(dx_2) < +\infty$ and assume that τ_0 is defined in terms of this function as in (4.37). Assume that there exists $\delta_1 > 0$ such that for every $m \geq 1$ and $K > 0$, $\mathbb{P}(\mathcal{E}_m | \mathcal{E}_{m-1}) \geq \delta_1 \in (0, 1)$. Then, for $j = 1, 2$ and for every $\rho \in (0, 1)$ and $\varepsilon \in (0, 1-H)$, there exists C such that for every $k \geq 1$, $m \in \{0, \dots, k-1\}$ and $K > 0$,*

$$\mathbb{E}\left[\sup_{(s,t), \tau_{k-1} \leq s < t \leq s+1} \frac{1}{t-s} |\Lambda_{m,k}|(s, t, W^j) | \mathcal{E}_k \right] \leq C \frac{\left(\sum_{\ell=m+1}^{k-1} a_\ell \right)^{H-1+\varepsilon}}{\delta_1^{\rho(k-m)}}. \quad (4.64)$$

As a consequence, if $a_k = \zeta^k$ where $\zeta \in (1, +\infty)$, there exists $\Delta \in (0, 1)$ and $C > 0$ such that for every integers m and k with $m \leq k$

$$\mathbb{E}\left[\sup_{(s,t), \tau_{k-1} \leq s < t \leq s+1} \frac{1}{t-s} |\Lambda_{m,k}|(s, t, W^j) | \mathcal{E}_k \right] \leq C \Delta^{k-m}. \quad (4.65)$$

REMARK 4.10. \triangleright The assumption on the moments of $\tilde{\mu}$ is only necessary for the case $m = 0$ which corresponds to the interval $[-\infty, \tau_0]$. Due to the memory, τ_0 is not independent of the past of the Brownian Motion before τ_0 . But the assumption on $\tilde{\mu}$ leads to a control of the queue of τ_0 which is sufficient to overcome the non-independence property.

\triangleright The fact that the quality of the estimate strongly decreases with $m - k$ may appear surprising. The main problem is that we do not have a sharp idea of the distribution of $\mathcal{L}(W_t - W_{\tau_{m-1}}, \tau_{m-1} \leq t \leq \tau_m)$ conditionally to the event $\{\Delta\tau_l < +\infty, m \leq l < k\}$ and thus, we compensate this failure by some Hölder-type inequalities.

\triangleright The second statement says that if one waits sufficiently between each trial, the influence of the past decreases geometrically with m . Note that this waiting time increases geometrically. This may be a problem for the sequel and the fact that ς can be chosen arbitrarily close to 1 will be of first importance.

Proof. First, note that if (4.64) is true, (4.65) easily follows: let $\varsigma > 1$ and let $\gamma_1 \in (0, +\infty)$ be such that $\varsigma = \delta_1^{-\gamma_1}$. It is now sufficient to remark that for every $m \in \{1, \dots, k-2\}$,

$$\left(\sum_{\ell=m+1}^{k-1} a_\ell \right)^{H-1+\varepsilon} \delta_1^{-\rho(k-m)} \leq \delta_1^{(\gamma_1(1-H-\varepsilon)-\rho)(k-m)}$$

and to choose for instance $\varepsilon = (1-H)/2$ and $\rho = \frac{\gamma_1(1-H)}{4} \wedge \frac{1}{2}$ so that

$$\gamma_1(1-H-\varepsilon) - \rho \geq \frac{\gamma_1(1-H)}{4} \wedge \frac{1}{2} > 0.$$

Let us now prove (4.64). We consider three cases:

Case 1: $k \geq 3$ and $m \in \{1, \dots, k-2\}$. $m \in \mathbb{N}$. In harmony with Lemma 4.6, we decompose $[\tau_{m-1}, \tau_m]$ in four intervals $[\tau_m^i, \tau_m^{i+1}]$, $i \in \{0, 1, 2, 3\}$, and also cut $\Lambda_{m,k}$ (which does not depend on k in this case) in four parts denoted by $\phi_{m,i}$:

$$\forall i \in \{0, 1, 2, 3\}, \quad \phi_{m,i}(s, t, W^j) = \int_{\tau_m^i}^{\tau_m^{i+1}} (t-r)^{H-\frac{1}{2}} - (s-r)^{H-\frac{1}{2}} dW_r^j.$$

By Lemma 4.5,

$$\frac{1}{t-s} |\phi_{m,i}(s, t, W^j)| \leq (t - \tau_m^i)^{H-\frac{3}{2}} |W_{\tau_m^i} - W_{\tau_m^{i+1}}| + \frac{1}{2} \int_{\tau_m^i}^{\tau_m^{i+1}} (s-r)^{H-\frac{5}{2}} |W_r - W_{\tau_m^{i+1}}| dr.$$

If $\tau_{k-1} \leq s < t$, since $\Delta\tau_l \geq a_l$ for all l , we get

$$t - \tau_m^i \geq \max\left(\sum_{\ell=m+1}^{k-1} a_\ell, \tau_m^{i+1} - \tau_m^i\right) \text{ and } \forall r \in [\tau_m^i, \tau_m^{i+1}], \quad s - r \geq \max\left(\sum_{\ell=m+1}^{k-1} a_\ell, 1 + \tau_m^{i+1} - r\right).$$

Thus, for every $\varepsilon \in (0, 1-H)$,

$$\sup_{(s,t), \tau_{k-1} \leq s < t \leq s+1} \frac{1}{t-s} |\phi_{m,i}(s, t, W^j)| \leq C \left(\sum_{\ell=m+1}^{k-1} a_\ell \right)^{H-1+\varepsilon} \Xi_{m,\varepsilon,i}$$

where

$$\Xi_{m,\varepsilon,i} = (\tau_m^{i+1} - \tau_m^i)^{-\frac{1}{2}-\varepsilon} |W_{\tau_m^i}^j - W_{\tau_m^{i+1}}^j| + \frac{1}{2} \int_{\tau_m^i}^{\tau_m^{i+1}} (1 + \tau_m^{i+1} - r)^{-\left(\frac{3}{2}+\varepsilon\right)} |W_r^j - W_{\tau_m^{i+1}}^j| dr.$$

The interesting point is that $\Xi_{m,\varepsilon,i}$ does not depend on k . Furthermore, by Lemma 4.6, for every $p > 1$, $\varepsilon \in (0, 1-H)$ and $i \in \{0, 1, 2, 3\}$,

$$\mathbb{E}[|\Xi_{m,\varepsilon,i}|^p | \mathcal{E}_m]^\frac{1}{p} \leq C$$

where C does only depend on p , ε and H . Set $\Xi_{m,\varepsilon} = \sum_{i=0}^3 \Xi_{m,\varepsilon,i}$. Summing up the previous controls (on i), we deduce from Hölder inequality that for every $p > 1$ and $q > 1$ such that $\frac{1}{p} + \frac{1}{q} = 1$,

$$\begin{aligned} 0 \leq \mathbb{E} \left[\sup_{(s,t), \tau_{k-1} \leq s < t \leq s+1} \frac{1}{t-s} |\Lambda_{m,k}(s,t,W^j)| |\mathcal{E}_k| \right] &\leq \mathbb{E} [|\Xi_{m,\varepsilon}|^p |\mathcal{E}_m|^{\frac{1}{p}} \left(\sum_{\ell=m+1}^{k-1} a_\ell \right)^{H-1+\varepsilon} \mathbb{P}(\mathcal{E}_k | \mathcal{E}_m)^{\frac{1}{q}-1}] \\ &\leq C \left(\frac{1}{\delta_1} \right)^{(1-\frac{1}{q})(k-m)} \left(\sum_{\ell=m+1}^{k-1} a_\ell \right)^{H-1+\varepsilon}. \end{aligned} \quad (4.66)$$

The result follows in this case by noticing that for every $\rho \in (0, 1)$, there exists $q \in (1, +\infty)$ such that $\rho = 1 - 1/q$.

Case 2: $k \geq 2$ and $m = k - 1$. It corresponds to the integral on the interval $[\tau_{k-2}, \tau_{k-1} - 1]$. The proof is almost identical using the controls

$$t - \tau_m^i \geq 1 + \tau_m^{i+1} \wedge (\tau_{k-1} - 1) - \tau_m^i \quad \text{and} \quad \forall r \in [\tau_m^i, \tau_m^{i+1} \wedge (\tau_{k-1} - 1)], \quad s - r \geq 1 + \tau_m^{i+1} \wedge (\tau_{k-1} - 1) - r.$$

We do not detail it.

Case 3: $k \geq 1$ and $m = 0$. It corresponds to the integral on the interval $(-\infty, \tau_0]$ if $k \geq 2$ and $(-\infty, \tau_0 - 1]$ if $k = 1$. For the sake of simplicity, we only consider the case $k \geq 2$. Note that on this interval $W^1 = W^2$. We then write W only. By Lemma 4.5 and the fact that $\lim_{M \rightarrow +\infty} M^{-1/2-\varepsilon} W_{-M} = 0$, we have

$$\frac{1}{t-s} |\Lambda_{0,k}(s,t,W)| \leq \frac{1}{2} \int_{-\infty}^{\tau_0} (s-r)^{H-\frac{5}{2}} |W_r - W_{\tau_0}| dr$$

so that

$$\sup_{\tau_{k-1} \leq s < t \leq \lfloor s \rfloor + 1} \frac{1}{t-s} |\Lambda_{0,k}(s,t,W)| \leq \frac{1}{2} \left(\sum_{m=1}^{k-1} a_m \right)^{H-1+\varepsilon} \int_{-\infty}^{\tau_0} (1 + \tau_0 - r)^{-\frac{3}{2}-\varepsilon} |W_r - W_{\tau_0}| dr$$

where $\varepsilon \in (0, 1 - H)$. Let $p \geq 1$. As remarked previously, one has no information about the joint law of τ_0 and W^j . We compensate this failure by a rough argument. Using Cauchy-Schwarz inequality,

$$\begin{aligned} \mathbb{E}_{\bar{\mu}} \left[\left(\int_{-\infty}^{\tau_0} (1 + \tau_0 - r)^{-\frac{1}{2}-\varepsilon} |W_r - W_{\tau_0}| dr \right)^p \right] \\ \leq \sum_{u=1}^{+\infty} \mathbb{E} \left[\left(\int_{-\infty}^u (1 + u - r)^{-\frac{3}{2}-\varepsilon} |W_r - W_u| dr \right)^{2p} \right]^{\frac{1}{2}} \mathbb{P}_{\bar{\mu}}(\tau_0 = u)^{\frac{1}{2}}. \end{aligned}$$

Thanks to the stationarity of the increments of the Brownian motion, we deduce from a change of variable that

$$\mathbb{E} \left[\left(\int_{-\infty}^u (1 + u - r)^{-\frac{3}{2}-\varepsilon} |W_r - W_u| dr \right)^{2p} \right] \leq \mathbb{E} \left[\left(\int_0^{+\infty} (1 + r)^{-\frac{3}{2}-\varepsilon} |W_r| dr \right)^{2p} \right] =: C_p.$$

Using that for every $p \geq 1$ and $\varepsilon > 0$,

$$\mathbb{E} \left[\sup_{r \geq 1} \left(\frac{|W_r|}{r^{\frac{1+\varepsilon}{2}}} \right)^{2p} \right] < +\infty,$$

we deduce that C_p is finite. It remains to show that $\sum_{u=1}^{+\infty} \mathbb{P}_{\bar{\mu}}(\tau_0 = u)^{\frac{1}{2}} < +\infty$. This property has already been proved in (4.37). \square

PROPOSITION 4.5. *Assume that Step 1 and 2 are carried out as described in Section 3. Let Ψ satisfy $\int \Psi(x_1)\bar{\mu}_1(dx_1) + \int \Psi(x_2)\bar{\mu}_2(dx_2) < +\infty$ and assume that τ_0 is defined in terms of this function as in (4.37). Assume there exists $\delta_1 > 0$ such that for every $m \geq 1$, $\mathbb{P}(\mathcal{E}_m | \mathcal{E}_{m-1}) \geq \delta_1 \in (0, 1)$ and that for every $k \geq 1$, a_k defined in (2.16) satisfies $a_k = \varsigma^k$ with $\varsigma > 1$. Then, $\mathbf{H}_2(\theta)$ holds for every $\theta \in (1/2, H)$.*

Proof. First, thanks to a change of variable and to the decomposition introduced at the beginning of the current subsection 4.3, we have

$$\begin{aligned} \sum_{u=1}^{\Delta\tau_k} \rho^{\Delta\tau_k-u} \|B\|_{\theta}^{\tau_{k-1}+u-1, \tau_{k-1}+u} &\leq \sum_{m=1}^k \sum_{u=\tau_{k-1}+1}^{\tau_k} \rho^{\tau_k-u} \|\Lambda_{m,k}^j\|_{\theta}^{u-1,u} \\ &+ \sum_{m=1}^3 \sum_{u=\tau_{k-1}+1}^{\tau_k} \rho^{\tau_k-u} \|\Gamma_m^j\|_{\theta}^{u-1,u} \end{aligned} \quad (4.67)$$

where we used the notations introduced in (4.49). Thus, the idea is to bound each term of the right-hand side. First, for every $m \in \{0, \dots, k-1\}$, for every $u \in \{\tau_{k-1}, \dots, \tau_k\}$,

$$\|\Lambda_{m,k}^j\|_{\theta}^{u,u+1} \leq \sup_{\tau_{k-1} \leq s < t \leq \lfloor s \rfloor + 1} \frac{1}{t-s} |\Lambda_{m,k}(s, t, W^j)|$$

Since the right-hand member does not depend on u , we deduce that for every $m \in \{0, \dots, k-1\}$

$$\sum_{u=\tau_{k-1}+1}^{\tau_k} \rho^{\tau_k-u} \|\Lambda_{m,k}^j\|_{\theta}^{u-1,u} \leq \sup_{\tau_{k-1} \leq s < t \leq \lfloor s \rfloor + 1} \frac{1}{t-s} |\Lambda_{m,k}(s, t, W^j)| \sum_{w=0}^{+\infty} \rho^w. \quad (4.68)$$

Thus, by Lemma 4.7, it follows that for every $m \in \{0, \dots, k-1\}$

$$\mathbb{E} \left[\sum_{u=\tau_{k-1}+1}^{\tau_k} \rho^{\tau_k-u} \|\Lambda_{m,k}^j\|_{\theta}^{u-1,u} | \mathcal{E}_k \right] \leq \frac{C}{1-\rho} \Delta^{k-m}$$

where $\Delta \in (0, 1)$. As a consequence,

$$\mathbb{E} \left[\sum_{m=1}^{k-1} \sum_{u=\tau_{k-1}+1}^{\tau_k} \rho^{\tau_k-u} \|\Lambda_{m,k}^j\|_{\theta}^{u-1,u} | \mathcal{E}_k \right] \leq \frac{C}{(1-\rho)(1-\Delta)}.$$

Keeping in mind inequality (4.67), it remains to bound, independently of k and K , the terms involving $\Lambda_{k,k}$ and Γ_m , $m = 1, 2, 3$. The strategy is different since these terms depend on the path between $\tau_{k-1} - 1$ and τ_k . Let us begin by $\Lambda_{k,k}$. By a change of variable,

$$\sum_{u=\tau_{k-1}+1}^{\tau_k} \rho^{\tau_k-u} \|\Lambda_{k,k}^j\|_{\theta}^{u-1,u} = \sum_{v=0}^{\Delta\tau_k-1} \rho^{\Delta\tau_k-1-v} Z_{k,v} \quad \text{where} \quad Z_{k,v} = \|\Lambda_{k,k}^j\|_{\theta}^{\tau_{k-1}+v, \tau_{k-1}+v+1}.$$

Note that $Z_{k,0} = 0$. On the event $\mathcal{A}_{k,\ell}$, one knows that $\Delta\tau_k$ is deterministic. Denote it by $\Delta(k, \ell)$. Decomposing the event \mathcal{E}_k , it follows that

$$\mathbb{E} \left[\sum_{u=\tau_{k-1}+1}^{\tau_k} \rho^{\tau_k-u} \|\Lambda_{k,k}^j\|_{\theta}^{u-1,u} | \mathcal{E}_k \right] = \sum_{\ell \geq 0} \sum_{v=1}^{\Delta(k, \ell)} \rho^{\Delta(k, \ell)-1-v} \mathbb{E}[Z_{k,v} | \mathcal{A}_{k,\ell}] \mathbb{P}(\mathcal{A}_{k,\ell} | \mathcal{E}_k).$$

Using that $\mathcal{A}_{k,\ell} \subset \mathcal{E}_k \subset \mathcal{E}_{k-1}$ and Cauchy-Schwarz inequality, we remark that

$$\mathbb{E}[Z_{k,v} | \mathcal{A}_{k,\ell}] \mathbb{P}(\mathcal{A}_{k,\ell} | \mathcal{E}_k) \leq \mathbb{E}[Z_{k,v}^2 | \mathcal{E}_k]^{\frac{1}{2}} \mathbb{P}(\mathcal{A}_{k,\ell} | \mathcal{E}_k)^{\frac{1}{2}} \leq \frac{\mathbb{E}[Z_{k,v}^4 | \mathcal{E}_{k-1}]^{\frac{1}{4}}}{\mathbb{P}(\mathcal{E}_k | \mathcal{E}_{k-1})^{\frac{1}{4}}} \mathbb{P}(\mathcal{A}_{k,\ell} | \mathcal{E}_k)^{\frac{1}{2}}. \quad (4.69)$$

But $\mathbb{P}(\mathcal{E}_k|\mathcal{E}_{k-1}) \geq \delta_1 > 0$ and using Lemma 3.2, we have for every $\ell \geq 2$,

$$\mathbb{P}(\mathcal{A}_{k,\ell}|\mathcal{E}_k) = \frac{\mathbb{P}(\mathcal{A}_{k,\ell}|\mathcal{E}_{k-1})}{\mathbb{P}(\mathcal{E}_k|\mathcal{E}_{k-1})} \leq \delta_1^{-1} \mathbb{P}(\mathcal{B}_{k,\ell-1}|\mathcal{E}_{k-1}) \mathbb{P}(\mathcal{B}_{k,\ell}^c|\mathcal{B}_{k,\ell-1}) \leq \delta_1^{-1} 2^{-\alpha\ell} \quad (4.70)$$

so that

$$\mathbb{E}\left[\sum_{u=\tau_{k-1}+1}^{\tau_k} \rho^{\tau_k-u} \|\Lambda_{k,k}^j\|_{\theta}^{u-1,u}|\mathcal{E}_k\right] \leq C_{\delta_1,\rho} \sup_{v \in \mathbb{N}} \mathbb{E}[Z_{k,v}^4|\mathcal{E}_{k-1}]^{\frac{1}{4}} \quad (4.71)$$

where $C_{\delta_1,\rho}$ is a finite constant depending only on δ_1 and ρ . Set $\varepsilon_{\theta} = (H - \theta)/2$. Using that $(W_{u+\tau_{k-1}-1} - W_{\tau_{k-1}-1})_{u \geq 0}$ is independent of $\mathcal{F}_{\tau_{k-1}-1}$, we obtain for every $v \in \mathbb{N}$, for every $k \in \mathbb{N}$,

$$\mathbb{E}[Z_{k,v}^4|\mathcal{E}_{k-1}] \leq \mathbb{E}[(F_v(\bar{W}_u, 0 \leq u \leq v))^4]$$

where \bar{W} is a standard Brownian Motion and $F_v : \mathcal{C}^{\frac{1}{2}-\varepsilon_{\theta}}([0, v], \mathbb{R}^d) \rightarrow \mathbb{R}$ defined by

$$F_v(w_u, 0 \leq u \leq v) = \sup_{v+1 \leq s < t \leq v+2} \frac{1}{t-s} \left| \int_0^v (t+1-r)^{H-\frac{1}{2}} - (s+1-r)^{H-\frac{1}{2}} dw_r \right|.$$

By Lemma 4.5, for every $v \geq 1$,

$$F_v(\bar{W}_u, 0 \leq u \leq v) \leq v^{H-\frac{3}{2}} |\bar{W}_v| + \frac{1}{2} \int_0^v (v+1-r)^{H-\frac{5}{2}} |\bar{W}_r - \bar{W}_v| dr.$$

Denote by $(\tilde{W}_u)_{u \in [0,1]}$ the rescaled Brownian motion defined by $\tilde{W}_u = \sqrt{v} \bar{W}_{uv}$. By a change of variable, for every $v \geq 1$,

$$F_v(\bar{W}_u, 0 \leq u \leq v) \leq |\tilde{W}_1| + \frac{1}{2} \int_0^1 \sqrt{v} (v+1-uv)^{H-\frac{5}{2}} |\tilde{W}_u - \tilde{W}_1| du.$$

Checking that

$$\forall v \geq 1, \quad \int_0^1 \sqrt{v} (v+1-uv)^{H-\frac{5}{2}} du \leq \left(\frac{3}{2} - H\right)^{-1},$$

one deduces that

$$\sup_{v \geq 1} \mathbb{E}[Z_{k,v}^4|\mathcal{E}_{k-1}] \leq \mathbb{E}\left[\left(|\tilde{W}_1| + \frac{2}{\frac{3}{2} - H} \sup_{u \in [0,1]} |\tilde{W}_u - \tilde{W}_1|\right)^4\right].$$

By plugging this inequality into (4.71), this concludes the study of $\Lambda_{k,k}$.

As concerns Γ_i , $i = 1, 2, 3$, one deduces with a similar strategy as before that it is enough to show that for $i = 1, 2, 3$,

$$\mathbb{E}[|G_i(\bar{W}_u, 0 \leq u \leq 2)|^4] \leq C, \quad (4.72)$$

where, setting $h = t - s$,

$$\begin{aligned} G_1(w) &= \sup_{1 \leq s < t \leq 2} \frac{1}{(t-s)^{\theta}} \left| \int_0^{s-h} (t-r)^{H-\frac{1}{2}} - (s-r)^{H-\frac{1}{2}} dw_r \right|, \\ G_2(w) &= \sup_{1 \leq s < t \leq 2} \frac{1}{(t-s)^{\theta}} \left| \int_{s-h}^s (s-r)^{H-\frac{1}{2}} dw_r \right| \quad \text{and} \\ G_3(w) &= \sup_{1 \leq s < t \leq 2} \frac{1}{(t-s)^{\theta}} \left| \int_{s-h}^t (t-r)^{H-\frac{1}{2}} dw_r \right|. \end{aligned}$$

By Lemma 4.5, we have

$$\begin{aligned} \int_0^{s-h} (t-r)^{H-\frac{1}{2}} - (s-r)^{H-\frac{1}{2}} dw_r &\leq |w_{s-h} - w_0|(t-s) \\ &\quad + (t-s)^{\theta} \int_0^{s-h} (s-h-r)^{H-\theta-\frac{3}{2}} |w_r - w_{s-h}| dr \end{aligned}$$

where in the second line, we used that for every $r \in [0, s - h]$, $s - r \geq s - h - r$ and $s - r \geq h$. We deduce that

$$G_1(w) \leq C(1 + \|w\|_{\frac{1}{2}-\varepsilon_\theta}^{0,2}) \quad \text{with} \quad \varepsilon_\theta = \frac{H - \theta}{2}. \quad (4.73)$$

and (4.72) follows for $i = 1$. By an integration by parts, one also checks that

$$\begin{aligned} \left| \int_{s-h}^s (s-r)^{H-\frac{1}{2}} dw_r \right| &\leq \|w\|_{\frac{1}{2}-\varepsilon_\theta}^{0,2} \left(h^{H-\varepsilon_\theta} + (H - \frac{1}{2}) \int_{s-h}^s (s-r)^{H-\frac{3}{2}+\frac{1}{2}-\varepsilon_\theta} dr \right) \\ &\leq Ch^{H-\varepsilon_\theta} \|w\|_{\frac{1}{2}-\varepsilon_\theta}^{0,2}. \end{aligned}$$

The previous control also holds for $G_3(w)$ and since $H - \varepsilon_\theta \geq \theta$, it follows that (4.72) is also true for $i = 2, 3$. \square

4.4 Condition (2.9)

PROPOSITION 4.6. *Let $\alpha \in (0, 1/2)$ and assume that for every $k \geq 1$ and $\ell \geq 0$, $\Delta_3(\ell, k) = c_3 a_k 2^{\beta \ell}$ with $\beta > (1 - 2\alpha)^{-1}$, $a_k = \zeta^k$ with an arbitrary $\zeta > 1$. Then, for each $K > 0$, there is a choice of c_3 such that, for every $k \geq 0$, Condition (2.9) is a.s. satisfied at time τ_k on the event $\{\tau_k < +\infty\}$. In other words, for every $k \geq 0$, $\mathbb{P}(\Omega_{\alpha, \tau_k}^1 | \tau_k < +\infty) = 1$.*

Proof. For every $k \geq 0$, set

$$u_k := \sup_{T \geq 0} \left(\int_0^{+\infty} (1+t)^{2\alpha} |(\mathcal{R}_T g_w^{\tau_k})(t)|^2 dt \right)^{\frac{1}{2}}.$$

Since g_w is null on $(-\infty, \tau_0]$, $u_0 = 0$. Following carefully the proof of Lemma 5.15 in [9] (see (5.8) therein for notation), we check that there exists $C > 0$ such that for every $k \geq 1$,

$$\begin{aligned} u_k - u_{k-1} &\leq C \left(\frac{\tau_k - \tau_k^3}{\tau_k^3 - \tau_{k-1}^3} \right)^{\alpha - \frac{1}{2}} \left(\int_0^{s_{k, \ell_k^* + 1} - \tau_{k-1}} (1+t)^{2\alpha} |g_w(\tau_{k-1} + t)|^2 dt \right)^{\frac{1}{2}} \\ &\leq C \left(\frac{c_2}{c_3} \zeta^{-k} 2^{-(\beta-1)\ell_k^*} \right)^{\frac{1}{2} - \alpha} \left(\int_0^{s_{k, \ell_k^* + 1} - \tau_{k-1}} (1+t)^{2\alpha} |g_w(\tau_{k-1} + t)|^2 dt \right)^{\frac{1}{2}} \end{aligned}$$

where we used (2.16) and (3.28) in the second inequality (By Remark 4.7, $c_2 = (C_K)^{\frac{1}{2\alpha}}$). But by Lemmas 3.1(b) and 3.2, for all $K > 0$ and $\alpha \in (0, H)$, there exists $C > 0$ such that

$$\left(\int_0^{s_{k, \ell_k^* + 1} - \tau_{k-1}} (1+t)^{2\alpha} |g_w(\tau_{k-1} + t)|^2 dt \right)^{\frac{1}{2}} \leq C(1 + 2^{\alpha(\ell_k^* + 1)}(\ell_k^* + 1)).$$

By the condition $\beta > (1 - 2\alpha)^{-1}$ which ensures that $(\beta - 1)(\frac{1}{2} - \alpha) > \alpha$, it follows that for every $K > 0$ and $\alpha \in (0, 1/2)$, there exists another constant $C > 0$ (depending on H, K, α and β) such that

$$\forall k \geq 0, \quad u_{k+1} - u_k \leq C \left(\frac{c_2}{c_3} \zeta^{-k} \right)^{\frac{1}{2} - \alpha}.$$

Choosing c_3 large enough in order that $C \sum_{k \geq 1} \left(\frac{c_2}{c_3} \zeta^{-k} \right)^{\frac{1}{2} - \alpha} \leq 1$ yields $\sup_{k \geq 0} u_k \leq 1$. \square

5 Proof of Theorem 2.1

Let $\alpha \in (0, 1/2)$. We enforce the assumptions of Proposition 3.1 and 4.2. Assume that X^1 and X^2 have initial distributions μ_0 and μ respectively, where μ denotes an invariant distribution. First, denoting by $\bar{\mu}$ its first marginal, we recall that $\int |x|^r \bar{\mu}(dx) < +\infty$ for any positive r (see Proposition 4.6 of [10] if b is Lipschitz continuous or Proposition 3 of [3] otherwise). It is therefore

enough to show that for any initial condition $\tilde{\mu}$ of (X^1, X^2) satisfying $\tilde{\mu}(|x_1|^r + |x_2|^r) < +\infty$ for some $r > 0$, for each $\varepsilon > 0$ there is $C_\varepsilon > 0$ such that $\mathbb{P}(\tau_\infty > t) \leq C_\varepsilon t^{-(\frac{1}{8}-\varepsilon)}$.

Set $k^* := \inf\{k \geq 1, \Delta\tau_k = +\infty\}$. We have

$$\mathbb{P}(\tau_\infty > t) = \mathbb{P}(\tau_0 + \sum_{k=1}^{+\infty} \Delta\tau_k 1_{k^* > k} > t) \leq \mathbb{P}(\tau_0 > \frac{t}{2}) + \mathbb{P}(\sum_{k=1}^{+\infty} \Delta\tau_k 1_{k^* > k} > \frac{t}{2}). \quad (5.74)$$

Taking Ψ such that $\Psi(x) \leq C(1 + |x|^r)$ (which is possible by Remark 4.8), by an argument similar to that of (4.39) we deduce the existence of $C > 0$ and $\gamma_0 > 0$ such that

$$\mathbb{P}(\tau_0 > \frac{t}{2}) \leq C \exp(-\gamma_0 t).$$

Now, let us focus on the second term on the right-hand side of (5.74) and let $p \in (0, \alpha/\beta) \subset (0, 1)$. By the Markov inequality and the elementary inequality $|u + v|^p \leq |u|^p + |v|^p$,

$$\begin{aligned} \mathbb{P}(\sum_{k=1}^{+\infty} \Delta\tau_k 1_{k^* > k} > \frac{t}{2}) &\leq \frac{C}{t^p} \sum_{k=1}^{+\infty} \mathbb{E}[|\Delta\tau_k|^p | 1_{\{k^* > k\}}] \\ &\leq \frac{C}{t^p} \sum_{k=1}^{+\infty} \mathbb{E}[\mathbb{E}[|\Delta\tau_k|^p | 1_{\{\Delta\tau_k < +\infty\}} | \mathcal{F}_{\tau_{k-1}}] 1_{\tau_{k-1} < +\infty}]. \end{aligned}$$

Let us bound deterministically the above conditional expectations. On the one hand, if Step 1 fails (including the case where $\omega \in \Omega_{K, \alpha, \tau_{k-1}}^c$), $\Delta\tau_k = 1 + c_3 \varsigma^k$ where $\varsigma > 1$ can be chosen arbitrarily. On the other hand, by Lemma 3.2, we have for every $\ell \geq 2$,

$$\mathbb{P}(\mathcal{A}_{k, \ell} | \mathcal{F}_{\tau_{k-1}} \cap \{\tau_{k-1} < +\infty\}) \leq 2^{-\alpha \ell}.$$

Since by construction, $\Delta\tau_k \leq C \varsigma^k 2^{\beta \ell}$ (with $\beta > (1 - 2\alpha)^{-1}$) on $\mathcal{A}_{k, \ell}$, this yields

$$\mathbb{E}[|\Delta\tau_k|^p | 1_{\{\Delta\tau_k < +\infty\}} | \mathcal{F}_{\tau_{k-1}} \cap \{\tau_{k-1} < +\infty\}] \leq C \varsigma^{kp} \sum_{\ell=1}^{+\infty} 2^{(\beta p - \alpha)\ell} \leq C \varsigma^{kp}.$$

Thus, for every $p \in (0, \alpha(1 - 2\alpha))$,

$$\mathbb{P}(\sum_{k=1}^{+\infty} \Delta\tau_k 1_{k^* > k} > \frac{t}{2}) \leq \frac{C}{t^p} \sum_{k=1}^{+\infty} \varsigma^{kp} \mathbb{P}(k^* > k - 1).$$

But

$$\mathbb{P}(k^* > k - 1) = \prod_{m=1}^{k-1} \mathbb{P}(\mathcal{E}_m | \mathcal{E}_{m-1}) = \prod_{m=1}^{k-1} (1 - \mathbb{P}(\mathcal{E}_m^c | \mathcal{E}_{m-1}))$$

and by Propositions 3.1 and 4.2 the latter applied with (for instance) $\varepsilon = 1/2$, we have for every $m \geq 1$,

$$\mathbb{P}(\mathcal{E}_m^c | \mathcal{E}_{m-1}) \geq \mathbb{P}(\Delta\tau_m = +\infty | \Omega_{K, \alpha, \tau_{m-1}}) \mathbb{P}(\Omega_{K, \alpha, \tau_{m-1}} | \mathcal{E}_{m-1}) \geq \frac{\delta_0}{2}$$

where δ_0 is a positive number depending on $K_{\frac{1}{2}}$. It follows that

$$\mathbb{P}(k^* > k - 1) \leq (1 - \frac{\delta_0}{2})^{k-1}.$$

As a consequence,

$$\sum_{k=1}^{+\infty} \varsigma^{kp} \mathbb{P}(k^* > k - 1) \leq \sum_{k=1}^{+\infty} \varsigma^{pk} (1 - \frac{\delta_0}{2})^{k-1} < +\infty$$

if ζ is chosen in such a way that $\zeta^p < (1 - \frac{\delta_0}{2})^{-1}$ (This is possible since ζ is an arbitrary number greater than 1). Finally, for every $\alpha \in (0, 1/2)$, for every $p \in \alpha(1 - 2\alpha)$, there exists $C > 0$ such that $\mathbb{P}(\tau_\infty > t) \leq Ct^{-p}$. To conclude the proof, it remains to optimize in α .

Acknowledgements: J.Fontbona acknowledges support of Fondecyt Grant 1110923, Basal-CONICYT Center for Mathematical Modeling (CMM), and Millenium Nucleus NC120062. F.Panloup acknowledges partial support of CNRS, INSA Toulouse and CMM to a six months visit at Santiago in 2013-2014 during which part of this work was carried out.

A Control of $(X_t^1)_{t \in [\tau_k, \tau_k+1]}$ under (K, α) -admissibility

We show the first point of the proof of Lemma 3.1. Let K and \tilde{K} denote some positive constants. Let $\omega \in \Omega_{K, \alpha, \tau_k}$ and assume that $\|W^1\|_{\frac{1}{2}-\varepsilon_\theta}^{\tau_k, \tau_k+1} \leq \tilde{K}$ with $\varepsilon_\theta = \frac{H-\theta}{2}$. First, we bound $\|B^1\|_{\theta}^{\tau_k, \tau_k+1}$. With the notations introduced at the beginning of Section 4.3,

$$\begin{aligned} \|B^1\|_{\theta}^{\tau_k, \tau_k+1} &\leq \sup_{\tau_k \leq s < t \leq \tau_k+1} \frac{1}{t-s} \left| \int_{-\infty}^{\tau_k-1} (t-r)^{H-\frac{1}{2}} - (s-r)^{H-\frac{1}{2}} dW_r^1 \right| \\ &\quad + \sum_{m=1}^3 \|\Gamma_m\|_{\theta}^{\tau_k, \tau_k+1}. \end{aligned}$$

The first right-hand term is bounded by K since $\omega \in \Omega_{K, \alpha, \tau_k}$. As concerns that of the second line, we deduce from the end of the proof of Proposition 4.5 (see *e.g.* (4.73)) that

$$\forall m \in \{1, 2, 3\}, \quad \|\Gamma_m\|_{\theta}^{\tau_k, \tau_k+1} \leq C(1 + \|W^1\|_{\frac{1}{2}-\varepsilon_\theta}^{\tau_k-1, \tau_k+1}).$$

Under the assumptions, $\|W^1\|_{\frac{1}{2}-\varepsilon_\theta}^{\tau_k-1, \tau_k} \leq K$ ((K, α) -admissibility) and $\|W^1\|_{\frac{1}{2}-\varepsilon_\theta}^{\tau_k, \tau_k+1} \leq \tilde{K}$. It follows that

$$\forall m \in \{1, 2, 3\}, \quad \|B^1\|_{\theta}^{\tau_k, \tau_k+1} \leq C_{K, \tilde{K}} \tag{A.75}$$

where $C_{K, \tilde{K}}$ is a finite deterministic constant which does not depend on k .

In order to conclude the proof, it is now enough to bound $\sup_{t \in [0, 1]} |X_{\tau_k+t}|$ with respect to $|X_{\tau_k}^1|$ and $\|B^1\|_{\theta}^{\tau_k, \tau_k+1}$. This point is a classical property of fractional driven SDE but we prove it for the sake of completeness. First, note that Steps 1, 2 and 3 of the proof of Proposition 4.4 still hold under the assumptions of Lemma . Set $F(x) = 1 + |x|^2$. Let $\tau_k \leq s < t \leq \tau_k + 1$ such that $c_0(1 + \|B^1\|_{\theta}^{\tau_k, \tau_k+1}) \leq 1/2$. By the change of variable formula,

$$\begin{aligned} F(X_t) &= F(X_s) + \int_s^t (\nabla F|b)(X_u) du + \int_s^t (\nabla F(X_u)|\sigma(X_u) dB_u^1) \\ &\leq F(X_s)(1 + C(t-s)) + \left| \int_s^t (\nabla F(X_u)|\sigma(X_u) dB_u^1) \right|, \end{aligned}$$

where in the second line, we used that $(\nabla F|b)(x) \leq CF(x)$ (since b is a sublinear function) and Step 3 of the proof of Proposition 4.4. The functions ∇F and σ being Lipschitz continuous and σ being also bounded, we obtain similarly to Step 4 of the proof of Proposition 4.4 (see (4.48)) that if $t-s \leq \eta$ defined by (4.47) (replacing 0 and 1 by τ_k and $\tau_k + 1$ respectively),

$$\left| \int_s^t (\nabla F(X_u)|\sigma(X_u) dB_u^1) \right| \leq C(t-s)^{\frac{1}{2}+\theta} F(X_s) + \tilde{\beta}.$$

Then, it follows that for every $\tau_k \leq s < t \leq \tau_k + 1$ such that $t-s \leq \eta$

$$F(X_t) \leq F(X_s)(1 + C(t-s)) + \tilde{\beta}.$$

An iteration of this inequality on the sequence $(\tau_k + \ell\eta)_{\ell \geq 1}$ yields

$$\begin{aligned} \forall \ell \in \dots, \lfloor \frac{1}{\eta} \rfloor, \quad F(X_{\tau_k + \ell\eta}) &\leq F(X_{\tau_k})(1 + C\eta)^\ell + \tilde{\beta} \sum_{u=0}^{\ell-1} (1 + C\eta)^u \\ &\leq \exp(C)(F(X_{\tau_k}) + \frac{\tilde{\beta}}{C\eta}) \leq \exp(C)(F(X_{\tau_k}) + \tilde{C}\tilde{\beta}(1 + \|B^1\|_\theta^{\tau_k, \tau_k+1})^{\frac{4}{2\theta-1}}) \end{aligned}$$

where in the last inequality, we used that $\eta \geq (1 + \|B^1\|_\theta)^{\tau_k, \tau_k+1}^{-\frac{1}{\theta}}$. Applying again Step 3 of the proof of Proposition 4.4 yields the existence of another constant C (which does not depend on k) such that

$$\sup_{t \in [\tau_k, \tau_k+1]} F(X_t) \leq C(F(X_{\tau_k}) + \tilde{\beta}(1 + \|B^1\|_\theta^{\tau_k, \tau_k+1})^{\frac{4}{2\theta-1}}). \quad (\text{A.76})$$

The result follows since, by (A.75), the right-hand side is bounded by a deterministic constant depending only on K, \tilde{K} and θ on the set $\Omega_{K, \alpha, \tau_k} \cap \{\|W^1\|_{\frac{1}{2}-\varepsilon_\theta}^{\tau_k, \tau_k+1} \leq \tilde{K}\}$.

B Proof of (4.34)

It is enough to prove that there exists $C > 0$ such that for every k and K , $\mathbb{E}[\varphi_{\tau_k, \varepsilon_\theta}(W^j(\omega)) | \mathcal{E}_k] \leq C$. The fact that

$$\mathbb{E}\left[\sup_{(s,t) \in \tau_k \leq s < t \leq \tau_k+1} \frac{1}{t-s} \left| \int_{-\infty}^{\tau_k-1} (t-r)^{H-\frac{1}{2}} - (s-r)^{H-\frac{1}{2}} dW_r^j \right| \middle| \mathcal{E}_k\right]$$

is bounded by a constant which does not depend on k follows from Lemma 4.7. More precisely, if $k \geq 1$ this property is a consequence of (4.65) combined with the fact that $\{(s, t), \tau_k \leq s < t \leq \tau_k + 1\} \subset \{(s, t), \tau_{k-1} \leq s < t \leq s + 1\}$. If $k = 0$, it corresponds to Case 3 of the proof of Lemma 4.7. For the second part, if $k \geq 1$, we deduce from Cauchy-Schwarz Inequality that

$$\mathbb{E}[\|W^j\|_{\frac{1}{2}-\varepsilon_\theta}^{\tau_k-1, \tau_k} | \mathcal{E}_k] \leq \mathbb{E}[(\|W^j\|_{\frac{1}{2}-\varepsilon_\theta}^{\tau_k-1, \tau_k})^2 | \mathcal{E}_{k-1}]^{\frac{1}{2}} \mathbb{P}(\mathcal{E}_k | \mathcal{E}_{k-1})^{-\frac{1}{2}}.$$

But using that $(W_j)_{t \in [\tau_{k-1}, \tau_k]}$ is independent of \mathcal{E}_{k-1} and that $\mathbb{P}(\mathcal{E}_k | \mathcal{E}_{k-1}) \geq \delta_1$, it follows that

$$\mathbb{E}[\|W^j\|_{\frac{1}{2}-\varepsilon_\theta}^{\tau_k-1, \tau_k} | \mathcal{E}_k] \leq C_\theta \delta_1^{-\frac{1}{2}}$$

where $C_\theta := \mathbb{E}[(\|W^j\|_{\frac{1}{2}-\varepsilon_\theta}^{0,1})^2] < +\infty$. This concludes the proof.

References

- [1] Ludwig Arnold. *Random dynamical systems*. Springer Monographs in Mathematics. Springer-Verlag, Berlin, 1998.
- [2] Serge Cohen and Fabien Panloup. Approximation of stationary solutions of Gaussian driven stochastic differential equations. *Stochastic Process. Appl.*, 121(12):2776–2801, 2011.
- [3] Serge Cohen, Fabien Panloup, and Samy Tindel. Approximation of stationary solutions to SDEs driven by multiplicative fractional noise. *Stochastic Process. Appl.*, 124(3):1197–1225, 2014.
- [4] Laure Coutin. Rough paths via sewing lemma. To appear in ESAIM PS, 2012.
- [5] Hans Crauel. Non-Markovian invariant measures are hyperbolic. *Stochastic Process. Appl.*, 45(1):13–28, 1993.
- [6] D. Down, S.P. Meyn, and R.L. Tweedie. Exponential and uniform ergodicity of markov processes. *The Annals of Probability*, 23:1671–1691, 1995.

- [7] María J. Garrido-Atienza, Peter E. Kloeden, and Andreas Neuenkirch. Discretization of stationary solutions of stochastic systems driven by fractional Brownian motion. *Appl. Math. Optim.*, 60(2):151–172, 2009.
- [8] Paolo Guasoni. No arbitrage under transaction costs, with fractional Brownian motion and beyond. *Math. Finance*, 16(3):569–582, 2006.
- [9] Martin Hairer. Ergodicity of stochastic differential equations driven by fractional Brownian motion. *Ann. Probab.*, 33(2):703–758, 2005.
- [10] Martin Hairer and Alberto Ohashi. Ergodic theory for SDEs with extrinsic memory. *Ann. Probab.*, 35(5):1950–1977, 2007.
- [11] Martin Hairer and Natesh S. Pillai. Regularity of laws and ergodicity of hypoelliptic SDEs driven by rough paths. *Ann. Probab.*, 41(4):2544–2598, 2013.
- [12] Jae-Hyung Jeon, Vincent Tejedor, Stas Burov, Eli Barkai, Christine Selhuber-Unkel, Kirstine Berg-Sørensen, Lene Oddershede, and Ralf Metzler. *In Vivo* anomalous diffusion and weak ergodicity breaking of lipid granules. *Phys. Rev. Lett.*, 106:048103, Jan 2011.
- [13] S. C. Kou. Stochastic modeling in nanoscale biophysics: subdiffusion within proteins. *Ann. Appl. Stat.*, 2(2):501–535, 2008.
- [14] Benoit B. Mandelbrot and John W. Van Ness. Fractional Brownian motions, fractional noises and applications. *SIAM Rev.*, 10:422–437, 1968.
- [15] Jonathan C. Mattingly. Exponential convergence for the stochastically forced Navier-Stokes equations and other partially dissipative dynamics. *Comm. Math. Phys.*, 230(3):421–462, 2002.
- [16] David Nualart and Aurel Răşcanu. Differential equations driven by fractional Brownian motion. *Collect. Math.*, 53(1):55–81, 2002.
- [17] David J. Odde, Elly M. Tanaka, Stacy S. Hawkins, and Helen M. Buettner. Stochastic dynamics of the nerve growth cone and its microtubules during neurite outgrowth. *Biotechnology and Bioengineering*, 50(4):452–461, 1996.
- [18] Sorin Rădulescu and Marius Rădulescu. An application of Hadamard-Lévy’s theorem to a scalar initial value problem. *Proc. Amer. Math. Soc.*, 106(1):139–143, 1989.
- [19] L. C. Young. An inequality of the Hölder type, connected with Stieltjes integration. *Acta Math.*, 67(1):251–282, 1936.
- [20] M. Zähle. Integration with respect to fractal functions and stochastic calculus. I. *Probab. Theory Related Fields*, 111(3):333–374, 1998.