

Nitrogen Cycling Driven By Organic Matter Export In The South Pacific Oxygen Minimum Zone

T. Kalvelage, G. Lavik, P. Lam, Sylvie Contreras, L. Arteaga, C. R. Loscher,

A. Oschlies, A. Paulmier, L. Stramma, M. M. M. Kuypers

► To cite this version:

T. Kalvelage, G. Lavik, P. Lam, Sylvie Contreras, L. Arteaga, et al.. Nitrogen Cycling Driven By Organic Matter Export In The South Pacific Oxygen Minimum Zone. Nature Geoscience, 2013, 6 (3), pp.228-234. 10.1038/ngeo1739 . hal-00998635

HAL Id: hal-00998635 https://hal.science/hal-00998635v1

Submitted on 3 Sep 2014

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Nitrogen cycling driven by organic matter export in the South Pacific oxygen minimum zone

Tim Kalvelage¹, Gaute Lavik¹, Phyllis Lam¹, Sergio Contreras^{1,4}, Lionel Arteaga², Caroline Löscher³, Andreas Oschlies², Aurélien Paulmier^{1,5,6}, Lothar Stramma², Marcel M.M. Kuypers¹

¹Max Planck Institute for Marine Microbiology, Bremen, Germany
²GEOMAR Helmholtz Centre for Ocean Research Kiel, Germany
³Institute for General Microbiology, Kiel, Germany
⁴Current address: Large Lakes Observatory, University of Minnesota Duluth, USA
⁵Current address: Laboratoire d'Etudes en Géophysique et Océanographie Spatiales, Toulouse, France
⁶Current address: Instituto del Mar del Perú (IMARPE), Esquina Gamarra y General Valle S/N Chucuito Callao, Peru

Keywords: marine nitrogen cycle, export production, organic matter, nitrogen loss, N-deficit, anammox, secondary nitrite maximum, oxygen minimum zone, eastern tropical South Pacific, Peruvian upwelling system

Oxygen minimum zones are major sites of oceanic nitrogen-loss, and have been expanding globally. Nitrogen-loss occurs mainly as the production of dinitrogen gas by denitrification or the so-called anammox process – the anaerobic ammonium oxidation with nitrite. Activity of anammox has been found more common in recent studies in the

- 5 eastern tropical South Pacific, one of the largest oxygen minimum zones worldwide. As anammox requires substrates from multiple co-occurring nitrogen transformations, regulation of nitrogen-loss has to involve factors controlling overall nitrogen cycling, but which exactly remains unclear. Here, we present the most comprehensive nitrogen budget assessment for all major nitrogen fluxes to date for the eastern tropical South
- 10 Pacific oxygen minimum zone. Extensive ¹⁵N-labelling experiments, nutrient measurements and export production modelling, together show that overall nitrogen cycling therein are tightly linked to the export of organic matter. Nitrogen-loss is most active over the productive shelf, fuelled by high rates of sinking organic matter and benthic ammonium release; then declines sharply offshore. These results highlight the
- 15 importance of coastal oxygen minimum zones in oceanic nitrogen balance, and offer an empirical relationship for parameterization in biogeochemical models to more realistically assess the effects of climate change on oceanic carbon and nitrogen cycling.

Coastal upwelling of nutrient-rich deep water fuels high surface productivity at the eastern boundaries of (sub)tropical oceans. The resultant export of organic matter stimulates strong microbial respiration in the subsurface. Combined with poor ventilation, permanently O₂-deficient waters called oxygen minimum zones (OMZs) develop at mid-depths^{1,2}. Ongoing, global expansion and intensification of OMZs will expectedly continue as anthropogenic pressures on marine environments grow^{3,4,5}.

Although constituting only ~1% ($O_2 \le 20 \ \mu mol \ kg^{-1}$) of global ocean volume⁶, OMZs have a profound impact on oceanic nitrogen (N) balance as they account for ~20-40% of global oceanic N-loss⁷. Ocean de-oxygenation might enlarge the ocean volume subject to Nloss⁸, exacerbate N-limitation of phytoplankton, and reduce the ocean's capacity to attenuate rising atmospheric carbon dioxide. Assessing the effects of expanding OMZs on the future ocean's nutrient balance, however, remains speculative, as biogeochemical models do not

reproduce present-day global patterns of N-loss^{9,10,11}. A major deficiency of those models appears to be the poor representation of coastal regions; whereas an increasing number of studies indicates that N-loss in shelf OMZs^{12,13,14}, coastal-offshore OMZ water mass exchange¹⁵ and OMZ-sediment interactions¹⁶ play more important roles on the overall N-

35 budget.

Based on the observed accumulations of nitrite (NO_2^-) and associated N-deficits, most N-loss in OMZ waters has traditionally been attributed to heterotrophic denitrification^{17,18,19}, the stepwise reduction of nitrate (NO_3^-) to gaseous dinitrogen (N_2) . Recent studies have, however, often failed to detect significant denitrifying activity in OMZs; rather, anammox

40 has more commonly been identified as a major N₂-forming pathway in these environments^{12,13,14,20}.

The regulation of N-loss activity including anammox is not fully understood. Anammox requires NH_4^+ and NO_2^- . Sources and sinks of both compounds have been identified in the OMZs, including aerobic NH_3 and NO_2^- oxidation, as well as anaerobic NO_3^-

reduction to NO₂⁻ and dissimilatory NO₃^{-/} NO₂⁻ reduction to NH₄⁺ (DNRA)^{14,15,21,22}.
 Surprisingly, O₂ sensitivity assays show that these processes in OMZ waters share a large overlapping range of O₂ concentrations (>0-20 µmol L⁻¹) in which they can co-occur, implying that within this range, controlling factors other than O₂ are more important^{23,24}.
 Enhanced autotrophic and heterotrophic N-cycling activity in the upper OMZ^{13,14,20,21}, and

25

50 generally elevated anammox rates usually measured in coastal versus offshore OMZs⁶, suggest that N-loss might ultimately be regulated by export production of organic matter.

To test this hypothesis, we conducted a large-scale survey of N-cycling rates, functional gene abundances, chlorophyll, nutrient and O₂ concentrations, as well as modelled export production, throughout the eastern tropical South Pacific (ETSP), one of the three

55 major OMZs in the world.

Dissolved Inorganic Nitrogen in the South Pacific Oxygen Minimum Zone

Consistent with past observations in the ETSP^{18,26}, pronounced secondary NO₂⁻ maxima were found in the offshore OMZ between 10°S and 18°S (supplementary Fig. 1),

- 60 extending up to 100's km westward with maximum concentrations of ~11 μ mol L⁻¹. Based on the spatial distribution of measured O₂, the lower OMZ boundary occurred at ~600 m on average near the Peruvian shelf (Figs S1-2). Henceforth, this is used as a depth cut-off to differentiate coastal OMZ stations, where the OMZ is in direct contact with sediments and benthic N-fluxes (<600m), from all others that are defined as offshore OMZ stations.
- Integrated over the thickness of the OMZ (defined by $O_2 \le 15 \ \mu mol \ L^{-1}$ where N-loss activity remains detectable in O_2 -sensitivity assays²⁴; Figs 1b, S2), NO_2^- concentrations reached >2 mol m⁻² in the offshore region (Fig. 1d). Concentrations of NH_4^+ were low (<0.25 $\mu mol \ L^{-1}$) throughout the OMZ, but could be $\ge 0.5 \ \mu mol \ L^{-1}$ over the shelf and near the upper OMZ boundary further offshore. Deeper in the offshore OMZ, plumes of elevated NH_4^+
- 70 concentrations (≤~3 µmol L⁻¹) sometimes occurred (supplementary Fig. 1), resulting in high depth-integrated values (Fig. 1e).

Offshore OMZs were characterized by severe N-deficits, expressed here as strongly negative N* with minima from -8 μ mol N L⁻¹ at 3.58°S down to -32 μ mol N L⁻¹ at 16°S. Depth-integrated values of N* (Fig. 1e) and NO₂⁻ were significantly correlated (Spearman

- 75 R=-0.61, p≤0.001). The southward intensification of both NO₂⁻ maxima and N* minima likely reflects the accumulated effects of time-integrated microbial activity in OMZ waters that advect poleward along the continental slope with the Peru-Chile Undercurrent ^{27,28}. Over the Peruvian shelf between 12°S and 14°S, extreme N-deficits (N* down to -60 µmol L⁻¹, Supplementary Fig. 1) were detected along with the presence of hydrogen sulphide (H₂S).
- 80 These stations are not further considered in the remaining discussions unless otherwise indicated, as the resident microbial communities and processes profoundly differ from typical OMZ scenarios (Schunck et al. submitted.).

Sources of Nitrite

Nitrite in the OMZs can be generated by NH₃ oxidation, the first step of nitrification; or by the reduction of nitrate to nitrite^{6,21,22}. Ammonia oxidation has been identified as an important NO₂⁻ source in the Peruvian OMZ, that is active under near-anoxic conditions^{21,23,29}. Our measured rates of NH₃ oxidation generally peaked at the base of the oxycline (~90 nmol N L⁻¹ d⁻¹), decreased to detection limit at the stations furthest offshore, and were not detectable in the core of the OMZ (Tables 1 and S2). The presence of both

archaeal and bacterial ammonia-oxidizers is verified by the detection of their biomarker functional genes encoding ammonia monooxygenase subunit A (Tables 1 and S3).

Integrated over the thickness of the OMZ, NO_2^- production via NH_3 oxidation increased from undetectable at the westernmost stations to $\leq 4.7 \text{ mmol } NO_2^- \text{ m}^{-2} \text{ d}^{-1}$ near the

95 coast (Fig. 2a; supplementary Table 1). For the entire OMZ volume examined (~ 5.5×10^5 km³), NH₃ oxidation is estimated to produce ~3.8 Tg N y⁻¹ of NO₂⁻, with 24% attributed to coastal OMZ (≤ 600 m) and 76% offshore (>600 m) (Fig. 3). Although significant rates have also been reported for the surface mixed layer in the ETSP³⁰, the mixed layer was not included in the current OMZ budget.

100 Overall, NH₃ oxidation accounted for only ~7% of the total NO₂⁻ production. The majority came from NO₃⁻ reduction to NO₂⁻, consistent with previous findings in the Peruvian, Namibian and Arabian Sea OMZs^{15,21,22}. Apart from its association with anammox, NO₃⁻ reduction to NO₂⁻ is the first step in denitrification and DNRA, and NO₃⁻ is the next preferred terminal electron acceptor after O₂ for the oxidation of organic matter. NO₃⁻

105 reduction was detected throughout the OMZ at all investigated stations; it reached a maximum (~1 μ mol N L⁻¹ d⁻¹) over the central shelf, but dropped to ~10 nmol N L⁻¹ d⁻¹ at the westernmost offshore stations (Table 1).

Depth-integrated rates showed similarly declining trend offshore (Fig. 2c; supplementary Table 1). Integration over the whole region yields an annual NO₃⁻ reduction of

110 ~49 Tg N, of which 29% occurs in the coastal OMZ and 71% offshore (Fig. 3). Like previous observations from the Peruvian²³ and the Arabian Sea¹⁵ OMZs, NO₃⁻ reduction significantly correlated with depth-integrated NO₂⁻ concentrations (Spearman R=0.71, p \leq 0.001) (Table 2), which indicates that NO₃⁻ reduction is a major contributor to the secondary NO₂⁻ maxima.

115 Sinks of Nitrite

Nitrite oxidation, the second step in nitrification, was most active in the upper OMZ throughout the ETSP. Its activity was detected deeper into the OMZ than NH₃ oxidation, consistent with earlier reports^{23,29}. Nitrite oxidation rates were highest (928 nmol N L⁻¹ d⁻¹) over the Peruvian shelf despite low O₂ levels (Table 1), and declined sharply to $\leq \sim 20$ nmol N

120 $L^{-1} d^{-1}$ along the furthest offshore transect. Although NO₂⁻ oxidation is believed to require O₂, this process has been detected at <1-2 µmol O₂ L⁻¹ in the Peruvian^{23,29} and Namibian OMZs²². O₂ sensitivity assays (~1-25 µmol L⁻¹) at two stations further demonstrated only a moderate attenuation by low O₂ (at most ~50% activity reduction at <1 µmol L⁻¹)

(supplementary Fig. 3), which agrees well with observations in the Namibian OMZ^{22} .

125 Clearly, NO₂⁻ oxidizers are well adapted to O₂-deficient environments.

 NO_2^- -supply from NH_3 oxidation, the first step of nitrification, is thought to constrain NO_2^- oxidation rates. Despite the significant correlation between NH_3 and NO_2^- oxidation rates (Spearman R=0.73, p≤0.001) (Table 2), NO_2^- oxidation in the ETSP OMZ exceeded those of NH_3 oxidation often by more than tenfold (Fig. 2a,b; supplementary Table 1,2).

130 Similar observations in the OMZs off Namibia²² and Peru^{23,29}, indicate a decoupling of the two steps of nitrification in O₂-deficient systems. A likely alternative NO₂⁻ source is NO₃⁻ reduction.

Based on modeled N-fluxes a NO₂⁻ "shunt", in which 45-74% of the NO₃⁻ reduced to NO₂⁻ by "denitrifying" micro-organisms is re-oxidized by aerobic NO₂⁻ oxidizers, has been proposed for the ETSP³¹. In agreement, our annual rates of NO₂⁻ oxidation for the coastal (7 Tg N y⁻¹) and offshore OMZ (23 Tg N y⁻¹) are equivalent to 51% and 65%, respectively, of NO₃⁻ reduction (Fig. 3). The strong correlation between the two processes (Spearman R=0.75, p≤0.001) (Table 2) indicates a close coupling between NO₂⁻ oxidation and NO₃⁻ reduction in the ETSP OMZ.

- 140 Meanwhile, only sporadic and low rates of DNRA ($\leq 1.3 \text{ nmol } \text{L}^{-1} \text{d}^{-1}$) were detected during our sampling period (Table 1). A general lack of detectable *nrfA*, a key functional gene encoding for the cytochrome $c \text{ NO}_2^{-1}$ reductase corroborates these results (Table 1). DNRA appears to exhibit a high degree of spatio-temporal variability, with similarly low rates measured on the Namibian shelf²², but with tenfold greater rates and *nrfA* gene
- abundance than observed in the OMZs off Peru²¹ and Oman¹⁴. Hence, we cannot exclude DNRA as a significant NO_x -sink for and NH_4^+ -source in the ETSP at other times.

Nitrogen-Loss Activities

At the time of our sampling, denitrification, expressed as the production of ${}^{30}N_2$ from

- ¹⁵NO_x, was generally non-detectable. Low rates of denitrification (~2-5 nmol L⁻¹ d⁻¹) were measured in three samples from the Peruvian shelf OMZ (Tables 1, S2). Substantially higher rates were detected in few samples containing measurable amounts of H₂S (Fig. 2e; supplementary Tables 1,2), suggesting a coupling with H₂S oxidation (ref. 32; Schunck et al. submitted.). In contrast to the conclusion drawn by a recent study³³, water-column
- 155 denitification was only of minor importance (<<1% total N-loss) for the overall N-budget in the ETSP OMZ (supplementary information).

N₂ production attributed to anammox was detected at all stations except the two furthest offshore, consistentent with previous studies in the ETSP^{13,20,34},. Anammox activity was often enhanced in the upper OMZ and markedly elevated in the bottom waters over the shelf and upper continental slope. Rates were highest (≤ 225 nmol N L⁻¹ d⁻¹) over the central shelf (10°S-16°S) and declined by two orders of magnitude westward (Table 1). The presence of anammox bacteria was verified by the detection of their characteristic hydrazine (N₂H₄) oxidoreductase genes (*hzo1* and 2) throughout the OMZ; whereas denitrifier-*nirS*, encoding for the cytochrome *cd*₁-containing NO₂⁻ reductase, was generally not detectable (Tables 2,

165 S3).

170

160

Depth-integrated anammox rates were >10 mmol N m⁻² d⁻¹ on the central shelf, similar to previous findings¹³, and <1 mmol N m⁻² d⁻¹ at the furthest offshore stations (Fig. 2d; supplementary Table 1). Altogether, anammox accounts for an annual N-loss of ~10 Tg in an area of 1.2×10^6 km², which is at the lower end of earlier estimates for the ETSP (9-26 Tg N v⁻¹)^{13,18,23,35}.

Flux measurements of dissolved inorganic nitrogen and N₂ made just prior to our sampling demonstrate that the sediments underlying the OMZ are additional sites of N-loss¹⁶. Combined with reaction-diffusion modeling, anammox and denitrification were shown to be

active N-sinks in the Peruvian coastal sediments. Based on the reported sedimentary NO_x⁻

175 fluxes and NO_x^- partitioning between anammox, denitrification, and DNRA¹⁶, we estimate a loss of 1 Tg N y⁻¹ from sediments in contact with the OMZ bottom waters (Fig. 3).

Conventionally, the accumulation of NO_2^- in OMZ waters has been interpreted as signs of active N-loss, and thus, is targeted by most field-sampling campaigns^{17,18,19,20,23,26,29}. Our data contradict this interpretation. Unlike NO_2^- , depth-integrated anammox rates did not

- 180 reveal any meridional trends, but decreased from shelf to offshore. Depth-integrated anammox rates and NO₂⁻ concentration were only moderately correlated (Spearman R=0.64, p<0.001) (Table 2). Furthermore, significant correlations between volumetric rates and NO₂⁻ concentrations were only observed for the shelf OMZ (Spearman R=0.72, p<0.001) and not offshore (Spearman, p>0.5). NO₂⁻ accumulation offshore probably resulted from a greater
- persistence of NO₃⁻ reduction to NO₂⁻ compared to other NO₂⁻-consuming processes in a poorly ventilated region, where the net NO₂⁻ gain was about five times higher compared to the coastal OMZ (11.4 and 2.2 Tg N y⁻¹, respectively).

Ongoing water column N-loss cannot be deduced simply from the intensity of N* minima, as shown by the lack of significant correlation (Spearman p>0.05) between

- anammox activity and N* (Table 2). While the depth-integrated N-deficit is strongest (most negative N*) offshore, anammox activity is highest over the shelf and upper continental slope. Though comprising only 10% of the area covered and merely 4% of the sampled OMZ volume, coastal OMZ waters contribute as much as 30% of the total N-loss (Fig. 3).
 Meanwhile, N-deficits in coastal OMZ waters amount to only 5% (4 Tg N) of the total N-
- 195 deficit (71 Tg N). Hence, the large N-deficit offshore most likely results from horizontal advection of N-deficient shelf waters²¹ that accumulate due to a long residence time in the offshore OMZ (~10 y based on N* and measured N-loss). This is analogous to to recent observations made in the Arabian Sea: substantial NO₂⁻ accumulation and low N-loss activity

in the central basin, compared to the rapid N-loss over the adjacent productive Omani 200 shelf^{14,15}.

Sources of Ammonium

N-loss driven by anammox requires NH_4^+ , which usually does not accumulate in OMZs. Ammonium concentrations can be kept low by a tight coupling between NH_4^+

205 production and consumption processes, while the NH_4^+ released at the reported remineralization rates may already be sufficient to fuel anammox. Major sources of NH_4^+ are water-column organic matter remineralization and sedimentary NH_4^+ release.

DNRA and organic matter ammonification are active benthic NH_4^+ sources off the coast of Peru. During two preceding cruises (M77-1 and 2) to the ETSP¹⁶, large NH_4^+ fluxes

210 (~0.5-4 mmol m² d⁻¹) from sediments into the overlying OMZ waters were measured on a cross-shelf transect at 11°S. The often enhanced anammox activity in the coastal OMZ bottom waters suggests a strong influence from NH₄⁺ diffusing out of the sediments^{13,36}. Assuming an average benthic NH₄⁺ flux of ~2 mmol m⁻² d⁻¹ and a typical anammox rate of ~4 mmol NH₄⁺ m⁻² d⁻¹ for the Peruvian coastal waters, the underlying sediments could supply
215 ~50% of the NH₄⁺ needed for the anammox rates observed. Clearly, additional NH₄⁺ sources are necessary to fulfil the remaining requirements for anammox, especially in offshore OMZ

waters, which are spatially decoupled from the sediments.

Based on the measured NO_3^- reduction rates, subsequent ammonification of Redfieldian organic matter generates 65% and 73% of the NH_4^+ needed for anammox in the

220 coastal and offshore OMZs, respectively (Fig. 3). These are likely underestimates, considering the observed preferential N-degradation of organic matter via NO_3^- respiration under suboxic conditions³⁷. Whether the reduction of NO_3^- is directly coupled to the

oxidation of organic matter, or indirectly via a recently proposed cryptic sulphur cycle³⁸, could not be discerned at this point.

- Remineralization of sinking organic matter and subsequent NH₄⁺ release is usually enhanced near the upper OMZ boundary, and would support the high anammox and NH₃ oxidation activity observed^{13,14,15,20,21,23}. On average, ~40% of their combined NH₄⁺ demands are supplied by NO₃⁻ reduction, with the remainder possibly coming from microaerobic organic matter remineralization²⁰. The activity of O₂-dependent nitrification at non-detectable O₂ concentrations in OMZs indicates that microaerobic respiration proceeds even at
- nanomolar O_2 levels, in accordance with an apparent half-saturation coefficient of <20 nmol L^{-1} previously reported for microaerobic respiration in these waters³⁹. High O_2 consumption rates, mainly attributable to heterotrophic respiration, and genes encoding for terminal respiratory oxidases with high O_2 -affinities were detected in the ETSP on the same
- 235 expedition (Kalvelage et al. unpubl.). While there are suggestions that O₂ is efficiently depleted down to the limits of microaerobic respiration in the OMZ core⁴⁰, regular intrusions of more oxygenated surface waters or mixing events, such as those related to eddies⁴¹, may sustain aerobic microbial activity in the upper OMZ.

240 Linking Surface Productivity and Sub-surface Nitrogen Cycling

Depth-integrated anammox rates correlated strikingly well with NO₃⁻ reduction, NO₂⁻ oxidation and NH₃ oxidation (Spearman R=0.88, 0.86 and 0.75, respectively; p \leq 0.001), indicating a common controlling factor for their concerted activity. Our data suggest that N-cycling processes in the OMZ are tightly coupled to the export of organic matter.

Export of organic matter at the base of the euphotic zone was estimated from net primary production $(NPP)^{42}$ and the ratio of export-to-total primary production (ef-ratio)⁴³. At the time of sampling, NPP was up to ~3 g organic C (C_{org}) m⁻² d⁻¹ near the coast and

decreased to <0.5 g C_{org} m⁻² d⁻¹ further offshore, values typical for the Peruvian upwelling system⁴⁴. Computed *ef*-ratios ranged from 0.16 (low-NPP sites) to 0.42 (high-NPP sites). The

- 250 resulting N-export production rates (converted from measured C:N=7.2 of surface particulate organic matter) were >10 mmol organic N (N_{org}) m⁻² d⁻¹ over the shelf and in the order of ~1 mmol N_{org} m⁻² d⁻¹ at the stations furthest offshore (Fig. 2f; supplementary Table 1). Export production was highly correlated to depth-integrated rates of anammox, NO₃⁻ reduction and NO₂⁻ oxidation (Spearman R=0.79, 0.75 and 0.60, respectively, p≤0.001) as well as NH₃
- 255 oxidation (Spearman R=0.56, p≤0.01) (Table 2). This suggests that the lateral distribution of N-cycling activity, including anammox, is mainly determined by the export of organic matter, which is the ultimate source of the required reactive substrates NH₄⁺ and NO₂⁻ in the OMZ.

Overall, we estimate NPP of 12 and 47 Tg N y⁻¹ in the coastal and offshore surface waters, respectively, which appear reasonable at a net lateral supply of 88 Tg NO_3^{-1} y⁻¹ to the

260 upwelling region (Fig. 3). The corresponding export fluxes are 4.4 and 9.9 Tg N_{org} y⁻¹. Taking organic matter sedimentation and export to the deep ocean into account, our results show that the export production to the OMZ is sufficient as an N-source to support the measured N-fluxes.

In summary, extensive sampling and experimentation throughout the ETSP OMZ shows that the activity of anammox and N-linked processes is highly correlated with export production. High productivity over the shelf and upper slope, as well as sedimentary NH₄⁺ release, drive high rates of tightly coupled N-cycling processes and thus N-loss via anammox in the shallow coastal OMZ compared to the offshore OMZ.

While the globally expanding OMZs might increase the oceanic volume conducive to 270 N-loss, N-loss would only continue to rise as long as there is sufficient nutrient supply for primary production in the euphotic zone, and nutrient supply is not hampered by intensified stratification (i.e. reduced upwelling) due to ocean warming. These positive and negative

feedbacks are important considerations for biogeochemical models, which at present do not adequately reproduce the observed spatial patterns of N-loss in OMZs. In light of our results,

275 the activities of N-loss via anammox appear to be directly linked to export production rates in biogeochemical models using the following empirical relationship: anammox = $0.7 \times N_{org}$ export (supplementary Fig. 4). This may facilitate a realistic assessment of the short- and long-term impacts of ocean de-oxygenation and changing productivity on N-cycling in OMZs, as well as their effects on neighbouring water masses.

280

Materials & Methods

Physico-chemical and N-cycling rate measurements

Large-scale distributions of chemical and biological variables were determined during the cruises M77-3 and 4 from December 2008 to February 2009 onboard R/V Meteor.

- Seawater was collected with either a conductivity-temperature-depth (CTD) rosette system fitted with 10-L Niskin bottles or a pump-CTD system (depth range: ~375 m). Continuous vertical profiles of chlorophyll-*a* were obtained fluorometrically and calibrated against discrete values derived from acetone extraction. Oxygen was measured with a Seabird sensor, a conventional amperometric microsensor and a highly sensitive STOX (Switchable Trace
- 290 amount Oxygen) sensor³⁹ (detection limit: 50 nmol L⁻¹). Dissolved inorganic N and PO₄³⁻ concentrations were analyzed using standard protocols^{45,46}. Nitrogen deficits were calculated as N* following Gruber & Sarmiento⁴⁷. Rates of microbial N-cycling (NH₃ and NO₂⁻ oxidation, NO₃⁻ reduction, anammox, denitrification and DNRA) were determined in short-term, time-series incubation experiments with various combinations of ¹⁵N-labeled and
- 295 unlabelled compounds as described in Füssel et al.²² and Holtapples et al.⁴⁸. Oxygen sensitivity assays for NO₂⁻ oxidation were conducted as previously described²². Consistent rates for anammox were calculated from the various ¹⁵N-incubation experiments

 $(^{15}NH_4^+\pm^{14}NO_2^-, ^{15}NO_2^-\pm^{14}NH_4^+, ^{15}NO_2^-\pm^{14}NH_4^+)$ for coastal OMZ stations; whereas more variability was associated with offshore OMZ stations. Although the possibility of substrate

- 300 stimulation due to ¹⁵N/¹⁴N-amendments cannot be fully eliminated, marine microbes including anammox and nitrifying bacteria^{13,22} are often associated with particles, and thus can experience substrate concentrations several orders of magnitude greater than the ambient water⁴⁹ such that our measured rates could also be substantially underestimated. In order to examine whether the export production is sufficient to support these measured rates of
- 305 various subsurface N-cycling processes and ultimately N-loss, the maximum potential rates for anammox from the various isotope-amendments (Supplementary Table 2) were used in budget calculations. Based on our combined rate measurements, nutrient inventories and subsequent modeling, the N-fluxes are sufficient to support all measured rates of N transformation. Hence, the here-presented measured rates may not be too far from reality.

310

315

Molecular ecological analyses

Water samples for nucleic-acids-based analyses were collected onto polyethersulfone membrane filters (0.2 μ m; Millipore) and immediately frozen at -80°C until further analysis. Nucleic acids were extracted using a Qiagen DNA/RNA All prep Kit following the manufacturers protocol with minor modifications⁵⁰. Functional genes for archaeal and bacterial (β -/ γ -proteobacterial) NH₃ oxidation (arch-*amoA* and β -/ γ -*am*oA, respectively), anammox (*hzo*1 and 2), denitrification (denitrifier-*nirS*) and DNRA (*nrfA*) were PCRamplified as described in Löscher et al.⁴⁹. Standards for quantitative PCRs were obtained from: *Nitrosococcus oceani* NC10 and *Nitrosomonas marina* NM22 and NM51 (γ - and β -

amoA, respectively), an environmental clone (GenBank accession number JF796147; archamoA), *Candidatus* "Scalindua profunda" (hzo1 and 2), *Pseudomonas aeruginosa* PAO1 (denitrifier-*nirS*) and *Escherichia coli* K12 (*nrfA*).

Modeling of export production

325 Export production was calculated from estimates of net primary production and the ratio of export production to total primary production (*ef*-ratio). Net primary production (NPP) at the time and location of our experimental stations was computed from measured chlorophyll-a concentrations and satellite-based (MODIS ocean color data) estimates of photosynthetic available radiation using the Vertically Generalized Production model⁴². *Ef*-

ratios were calculated from NPP and measured surface temperatures after Laws et al. 43 .

Author contributions

TK, GL and MK designed the study. TK, GL, SC and AP performed ¹⁵N-labeling

experiments. TK, GL and PL analyzed the data. CL carried out functional gene analyses.
 LA and AO modelled export production rates. LS provided CTD and ADCP data. TK, GL,
 PL and MK wrote the manuscript.

The authors declare no competing financial interests.

340

345

References

1.	Kamykowski, D. & Zentara, S. Hypoxia in the world ocean as recorded in the
	historical data set. Deep-Sea Res. 37, 1861-1874 (1990).

 Karstensen, J., Stramma, L. & Visbeck, M. Oxygen minimum zones in the eastern tropical Atlantic and Pacific oceans. *Progr. Oceanogr.* 77, 331-350 (2008).

- Diaz, R.J. & Rosenberg, R. Spreading Dead Zones and Consequences for Marine Ecosystems. *Science* 321, 926-929 (2008).
- Stramma, L. et al. Expanding Oxygen-Minimum Zones in the Tropical Oceans.
 Science 320, 655-658 (2008).
 - Keeling, R.F., Körtzinger, A. & Gruber, N. Ocean Deoxygenation in a Warming World. *Ann. Rev. Mar. Sci.* 463-493 (2009).
 - Lam, P. & Kuypers, M.M. Microbial Nitrogen Cycling Processes in Oxygen Minimum Zones. Ann. Rev. Mar. Sci. 3, 317-347 (2011).
- 355 Minimum Zones. Ann. Rev. Mar. Sci. **3**, 317-347 (2011).
 - Gruber, N. The Dynamics of the Marine Nitrogen Cycle and its Influence on Atmospheric CO2. *NATO ASI Series* 97-148 (2004).
 - Deutsch, C. et al. Climate-Forced Variability of Ocean Hypoxia. *Science* 333, 336-339 (2011).
- Moore, J.K. & Doney, S.C. Iron availability limits the ocean nitrogen inventory stabilizing feedbacks between marine denitrification and nitrogen fixation. *Glob. Biogeochem. Cyc.* 21, GB2001 (2007).
 - 10. Schmittner, A. et al. Future changes in climate, ocean circulation, ecosystems, and biogeochemical cycling simulated for a business-as-usual CO2 emission scenario until
- 365 year 4000 AD. *Glob. Biogeochem. Cyc.* **22**, GB1013 (2008).
 - Somes, C.J. et al. Simulating the global distribution of nitrogen isotopes in the ocean.
 Glob. Biogeochem. Cyc. 24, GB4019 (2010).
 - 12. Kuypers, M.M. et al. Massive nitrogen loss from the Benguela upwelling system through anaerobic ammonium oxidation. *PNAS* **102**, 6478-6483 (2005).
- Hamersley, M.R. et al. Anaerobic ammonium oxidation in the Peruvian oxygen minimum zone. *Limnol. Oceanogr.* 52, 923-933 (2007).

- 14. Jensen, M.M. et al. Intensive nitrogen loss over the Omani Shelf due to anammox coupled with dissimilatory nitrite reduction to ammonium. *ISME* **AOP** (2011).
- Lam, P. et al. Origin and fate of the secondary nitrite maximum in the Arabian Sea.
 Biogeosciences 8, 1565-1577 (2011).
- 16. Bohlen, L. et al. Benthic nitrogen cycling traversing the Peruvian oxygen minimum zone. *Geoch. Cosmoch. Acta* **75**, 6094-6111 (2011).
- Cline, J.D. & Richards, F.A. Oxygen Deficient Conditions and Nitrate Reduction in the Eastern Tropical North Pacific Ocean. *Limnol. Oceanogr.* 17, 885-900 (1972).
- Codispoti, L.A. & Packard, T.T. Denitrification Rates in the Eastern Tropical South-Pacific. J. Mar. Res. 38, 453-477 (1980).
 - Naqvi, S.W. Some aspects of the oxygen-deficient conditions and denitrification in the Arabian Sea. J. Mar. Res. 45, 1049-1072 (1987).
 - 20. Thamdrup, B. et al. Anaerobic ammonium oxidation in the oxygen-deficient waters

385 off northern Chile. *Limnol. Oceanogr.* **51**, 2145-2156 (2006).

- Lam, P. et al. Revising the nitrogen cycle in the Peruvian oxygen minimum zone.
 PNAS 106, 4752-4757 (2009).
- Füssel, J. et al. Nitrite oxidation in the Namibian oxygen minimum zone. *ISME* AOP (2011).
- 23. Lipschultz, F. et al. Bacterial transformations of inorganic nitrogen in the oxygendeficient waters of the Eastern Tropical South Pacific Ocean. *Deep-Sea Res.* 37, 1513-1541 (1990).
 - 24. Kalvelage, T. et al. Oxygen Sensitivity of Anammox and Coupled N-Cycle Processes in Oxygen Minimum Zones. *PLoS ONE* **6**, e29299 (2011).
- 395 25. Fuenzalida, R. et al. Vertical and horizontal extension of the oxygen minimum zone in the eastern South Pacific Ocean. *Deep-Sea Res. II* **56**, 992-1003 (2009).

- Codispoti, L.A. & Christensen, J.P. Nitrification, Denitrification and Nitrous Oxide Cycling in the Eastern Tropical South Pacific Ocean. *Marine Chemistry* 16, 277-300 (1985).
- Wooster, W.S., Chow, T.J. & Barrett, I. Nitrite Distribution in Peru Current Waters. J.
 Mar. Res. 23, 210-221 (1965).
 - Nelson, S.S. & Neshyba, S. On the southernmost extension of the Peru-Chile Undercurrent. *Deep-Sea Res.* 26A, 1387-1393 (1979).
 - 29. Ward, B.B., Glover, H.E. & Lipschultz, F. Chemoautotrophic Activity and
- 405 Nitrification in the Oxygen Minimum Zone off Peru. *Deep-Sea Res.* **36**, 1031-1051 (1989).
 - Fernández, C., Farías, L. & Alcaman, M.E. Primary production and nitrogen regeneration processes in surface waters of the Peruvian upwelling system. *Progr. Oceanogr.* 83, 159-168 (2009).
- 410 31. Anderson, J.J. et al. A model for nitrite and nitrate distributions in oceanic oxygen minimum zones. *Deep-Sea Res.* **29**, 1113-1140 (1982).
 - 32. Lavik, G. et al. Detoxification of sulphidic African shelf waters by blooming chemolithotrophs. *Nature* **457**, 581-584 (2008).
 - 33. Dalsgaard, T. et al. Anammox and denitrification in the oxygen minimum zone of the
- 415 eastern South Pacific. *Limnol. Oceanogr.* **57**, 1331-1346 (2012).
 - 34. Ward, B.B. et al. Denitrification as the dominant nitrogen loss process in the Arabian Sea. *Nature* 461, 78-81 (2009).
 - Deutsch, C. et al. Denitrification and N2 Fixation in the Pacific Ocean. *Glob. Biogeochem. Cyc.* 15, 483-506 (2001).
- 420 36. Dalsgaard, T. et al. N2 production by the anammox reaction in the anoxic water column of Golfo Dulce, Costa Rica. *Nature* **422**, 606-608 (2003).

- Van Mooy, B.A., Keil, R.G. & Devol, A.H. Impact of suboxia on sinking particulate organic carbon: Enhanced carbon flux and preferential degradation of amino acids via denitrification. *Geoch. Cosmoch. Acta* 66, 457-465 (2002).
- 425 38. Canfield, D.E. et al. A Cryptic Sulfur Cycle in Oxygen-Minimum Zone Waters off the Chilean Coast. *Science* **330**, 1375-1378 (2010).
 - Revsbech, N.P. et al. Determination of ultra-low oxygen concentrations in oxygen minimum zones by the STOX sensor. *Limnol. Oceanogr.: Methods* 7, 371-381 (2009).
 - 40. Thamdrup, B., Dalsgaard, T. & Revsbech, N.P. Widespread functional anoxia in the
- 430 oxygen minimum zone of the eastern South Pacific. *Deep-Sea Res. I* in Press (2012).
 - Bertrand, A., Ballón, M. & Chaigneau, A. Acoustic Observation of Living Organisms Reveals the Upper Limit of the Oxygen Minimum Zone. *PLoS ONE* 5, e10330 (2010).
 - 42. Behrenfeld, M.J. & Falkowski, P.G. Photosynthetic rates derived from satellite-based chlorophyll concentration. *Limnol. Oceanogr.* **42**, 1-20 (1997).
- 435 43. Laws, E.A. et al. Temperature effects on export production in the open ocean. *Glob. Biogeochem. Cyc.* 14, 1231-1246 (2000).
 - Pennington, J.T. et al. Primary production in the eastern tropical Pacific: A review.
 Progr. Oceanogr. 69, 285-317 (2006).
 - 45. Grasshoff, K. et al. *Methods of seawater analysis*. (Wiley: 1999).
- 440 46. Holmes, R.M. et al. A simple and precise method for measuring ammonium in marine and freshwater ecosystems. *Canadian Journal of Fisheries and Aquatic Sciences* 56, 1801-1808 (1999).
 - Gruber, N. & Sarmiento, J.L. Global Patterns of Marine Nitrogen Fixation and Denitrification. *Glob. Biogeochem. Cyc.* 11, 235-266 (1997).

- 445 48. Holtappels, M. et al. 15N-labeling experiments to dissect the contributions of heterotrophic denitrification and anammox to nitrogen removal in the OMZ waters of the ocean. *Methods Enzymol.* 486, 223-251 (2011).
 - 49. Stocker, R. Marine Microbes See a Sea of Gradients. *Science* **338**, 628-633 (2012).
 - 50. Loescher, C.R. et al. Production of oceanic nitrous oxide by ammonia-oxidizing
- 450 archaea. *Biogeosc. Disc.* **9**, 2095-2122 (2012).

Correspondence and requests for materials should be directed to: mkuypers@mpi-bremen.de 455

Acknowledgements

We wish to thank the Peruvian government for access to their territorial waters. We sincerely thank cruise leader Martin Frank as well as the crews of the cruises M77-3 and 4

460 onboard *R/V Meteor* for their support at sea. We are grateful for the technical and analytical assistance of Gabriele Klockgether, Inka Boosmann, Violeta Leon, Imke Grefe and Andreas Ellrott. We thank Niels Peter Revsbech, who generously provided the STOX sensors and Tim Ferdelmann for offering valuable comments to improve the article. This study was supported by the Deutsche Forschungsgemeinschaft through the Sonderforschungsbereich 754

465 "Climate-Biogeochemistry Interactions in the Tropical Ocean" and the Max Planck Society.

Table 1 | Abundance of selected N-functional genes and N-conversion rates in the ETSP during cruise M77-3.

Functional genes and N-conversion rates were not always determined at the same station and/or depths but with a comparable latitudinal and longitudinal as well as vertical resolution.

		N-functional gene abundances (10 ² copies mL ⁻¹)							N-conversion rates (nmol N L ⁻¹ d ⁻¹)					
		arch-amoA	β-amoA	γ-amoA	hzo1	hzo2	den-nirS*	nrfA	NH3 ox.	Anammox	Denitri- fication	DNRA	$NO_2^- ox.$	NO ₃ ⁻ red.
`oastal OMZ (≤600 m)	N:	63 (64)	8 (49)	0 (64)	8 (64)	42 (64)	0 (63)	0 (64)	27 (33)	33 (33)	3 (33)	7 (33)	27 (32)	27 (32)
	Range:	0.16-2773	0.05-1056	-	0.05-0.09	0.14-12.8	-	-	0.22-48.8	2.84-227	2.21-5.42	0.48-1.74	8.48-928	3.79-1010
	Mean:	676	135	-	0.07	4.45	-	-	8.24	43.4	4.19	1.14	172	203
0	Median:	90	5.0	-	0.06	3.77	-	-	3.40	21.2	4.94	1.10	65.4	101
ffshore OMZ (>600 m)	N:	67 (71)	2 (33)	0 (72)	4 (71)	43 (72)	2 (72)	1 (72)	17 (40)	33 (40)	0 (40)	3 (40)	27 (40)	25 (34)
	Range:	0.04-2332	0.15-1.36	-	0.01-0.09	0.06-14.7	0.06-1.98	0.11	0.51-88.8	0.71-46.9	-	0.33-1.31	4.58-186	4.53-77.4
	Mean:	352	0.75	-	0.08	3.15	1.02	0.11	20.9	6.14	-	0.82	40.6	32.1
0	Median:	89.5	0.75	-	0.08	1.51	1.02	0.11	5.79	3.01	-	0.83	30.2	22.3

N = number of samples in which N-functional genes/N-processes were detected; in parenthesis: number of samples analyzed. *denitrifier-nirS.

	NH ₃ oxidation	NO ₂ - oxidation	NO ₃ - reduction	Anammox	Export Production
$\mathbf{NH_4}^+$	0.51*	0.31	0.08	0.30	0.29
NO ₂ ⁻	0.46*	0.49*	0.71***	0.64**	0.10
N *	-0.08	-0.20	-0.05	-0.02	-0.02
Export Production	0.56*	0.60**	0.75***	0.79***	
Anammox	0.75***	0.86***	0.88***		
NO ₃ ⁻ reduction	0.49*	0.75***			
NO ₂ ⁻ oxidation	0.73***				

 Table 2 | Spearman rank correlation of depth-integrated nutrients and N-cycling rates as well as modelled export productions.

5

Presented values are correlation coefficients with significant values denoted by * ($p\leq0.05$), ** ($p\leq0.01$) and *** ($p\leq0.001$).

Figure legends

Figure 1 | Maps of sampling locations and nutrient distributions in the ETSP

OMZ. a, Sampling sites during M77-3 (•) and M77-4 (•) and ¹⁵N-experimental stations (•). **b**, vertical extent of the OMZ (in m) as defined by $O_2 \le 15 \mu \text{mol } \text{L}^{-1}$. **c-f**, concentrations of NO₃⁻, NO₂⁻, NH₄⁺ and N* (in mol m⁻²) integrated over the thickness of the OMZ. Red line in panel **a**. denotes the 600m-isobath that was used demarcate the coastal OMZ from offshore OMZ.

Figure 2 | Depth-integrated N-cycling rates in the ETSP OMZ. a,b, the two steps of the aerobic nitrification, NH₃ oxidation and NO₂⁻ oxidation. c, NO₃⁻ reduction to NO₂⁻. d,e, N-loss due to anammox as well as denitrification coupled to the oxidation of H₂S during a sulfidic event on the Peruvian shelf. f, modelled export of organic N from the euphotic zone to the OMZ. All rates in mmol N m⁻² d⁻¹.

Figure 3 N-fluxes and nutrient inventory of the ETSP OMZ. Black numbers indicate inventories of dissolved inorganic nitrogen (in Tg N). They were derived from the depth-integrated values over the OMZ thickness shown in Figs. 1-2, and then based on the 600 m seafloor depth cut-off for coastal versus offshore OMZs, the depth-integrated values were further integrated over the areal extents of the two types of OMZs. Fluxes (in Tg N y⁻¹) within the OMZ or across its boundaries are given in colour and white, respectively. A detailed description of the flux calculations is included in the supplementary information.





