



HAL
open science

A non symmetric extension of Amblard-Girard copulas

Emil Stoica

► **To cite this version:**

| Emil Stoica. A non symmetric extension of Amblard-Girard copulas. 2014. hal-00999399

HAL Id: hal-00999399

<https://hal.science/hal-00999399>

Preprint submitted on 3 Jun 2014

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

A non symmetric extension of Amblard-Girard copulas

Emil Stoica

Dragan European University of Lugo

Abstract

A copula is a function that completely describes the dependence structure between the marginal distributions. One of the most important parametric family of copulas is the Farlie-Gumbel-Morgenstern (FGM) family. In practical applications this copula has been shown to be somewhat limited and a symmetric extension of this family, known as the Amblard-Girard copula, has been introduced. We propose a new non symmetric extension of this family.

A copula is a function that completely describes the dependence structure. It contains all the information to link the marginal distributions to their joint distribution. To obtain a valid multivariate distribution function, it suffices to combine several marginal distribution functions with any copula function. Thus, for the purposes of statistical modeling, it is desirable to have a large collection of copulas at one's disposal. Many examples of copulas can be found in the literature, most are members of families with one or more real parameters.

One of the most important parametric family of copulas is the Farlie-Gumbel-Morgenstern (FGM) family defined as

$$C_{\theta}^{FGM}(u, v) = uv + \theta uv(1 - u)(1 - v), \quad u, v \in [0, 1], \quad (1)$$

where $\theta \in [-1, 1]$. The density function of FGM copulas is given by

$$\frac{\partial^2 C_{\theta}^{FGM}(u, v)}{\partial u \partial v} = \theta(2u - 1)(2v - 1) + 1,$$

for any $u, v \in [0, 1]$.

Members of the FGM family are symmetric, i.e., $C_{\theta}^{FGM}(u, v) = C_{\theta}^{FGM}(v, u)$ for all (u, v) in $[0, 1]^2$ and have the lower and upper tail dependence coefficients equal to 0.

The copula given in (1) is PQD for $\theta \in (0, 1]$ and NQD for $\theta \in [-1, 0)$. In practical applications this copula has been shown to be somewhat limited, for copula dependence parameter $\theta \in [-1, 1]$, Spearman's correlation $\rho \in [-1/3, 1/3]$ and Kendall's $\tau \in [-2/9, 2/9]$, for more details on copulas see, for example, [1].

To overcome this limited dependence, several authors proposed extensions of this family.

- Theoretical issues: [2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24]
- Applications: [25, 26, 27, 28, 29, 30]

Let C_1, C_2 be two copulas proposed by [9]:

$$C_i(u, v) = uv + \theta_i \phi_i(u) \phi_i(v)$$

with $i \in \{1, 2\}$. The parameter θ_i is called the dependence parameter while the function ϕ_i is called the generator. Let $C = C_1 \star C_2$ where \star is the copula-product, see for instance [1]. The density associated to C can be computed as:

$$\begin{aligned} c(u, v) &= \int_0^1 c_1(u, t) c_2(v, t) dt \\ &= \int_0^1 [1 + \theta_1 \phi_1'(u) \phi_1'(t)] [1 + \theta_2 \phi_2'(v) \phi_2'(t)] dt \\ &= 1 + \theta_1 \theta_2 \int_0^1 \phi_1'(t) \phi_2'(t) dt \phi_1'(u) \phi_2'(v), \end{aligned}$$

the cross-product terms being null in view of the constraint $\phi_i(0) = \phi_i(1) = 0$, for all $i \in \{1, 2\}$. Letting $\theta = \theta_1 \theta_2 \int_0^1 \phi_1'(t) \phi_2'(t) dt$, routine calculations show that

$$C(u, v) = uv + \theta \phi_1(u) \phi_2(v).$$

Therefore, C can be considered as a non-symmetric extension of [9].

References

- [1] Roger B Nelsen. *An introduction to copulas*. Springer, 1999.
- [2] E. Stoica. A stability property of farlie-gumbel-morgenstern distributions. <http://hal.archives-ouvertes.fr/hal-00861234>.
- [3] E. Stoica. Convex sums of farlie-gumbel-morgenstern distributions. <http://hal.archives-ouvertes.fr/hal-00862992>.
- [4] E. Stoica. Extensions of farlie-gumbel-morgenstern distributions: A review. <http://hal.archives-ouvertes.fr/hal-00864676>.
- [5] E. Stoica. A non symmetric extension of farlie-gumbel-morgenstern distributions. *Journal of Parametric and Non-Parametric Statistics*, 1, 2013. <http://jpanps.altervista.org/>.
- [6] E. Stoica. On copula efficiency. <http://hal.archives-ouvertes.fr/hal-00954223>.

- [7] Dong Yong-quan. Generation and prolongation of fgm copula. *Chinese journal of engineering mathematics*, 25:1137–1141, 2008.
- [8] Matthias Fischer and Ingo Klein. Constructing generalized fgm copulas by means of certain univariate distributions. *Metrika*, 65(2):243–260, 2007.
- [9] Cécile Amblard and Stéphane Girard. Symmetry and dependence properties within a semiparametric family of bivariate copulas. *Journal of Non-parametric Statistics*, 14(6):715–727, 2002.
- [10] Cécile Amblard and Stéphane Girard. Estimation procedures for a semi-parametric family of bivariate copulas. *Journal of Computational and Graphical Statistics*, 14(2), 2005.
- [11] Cécile Amblard and Stéphane Girard. A new extension of bivariate fgm copulas. *Metrika*, 70(1):1–17, 2009.
- [12] José Antonio Rodríguez-Lallena and Manuel Úbeda-Flores. A new class of bivariate copulas. *Statistics & probability letters*, 66(3):315–325, 2004.
- [13] Carles M Cuadras. Constructing copula functions with weighted geometric means. *Journal of Statistical Planning and Inference*, 139(11):3766–3772, 2009.
- [14] Fabrizio Durante. A new family of symmetric bivariate copulas. *Comptes Rendus Mathématique*, 344(3):195–198, 2007.
- [15] H Bekrizadeh, GA Parham, and MR Zadkarmi. A new class of positive dependent bivariate copula and its properties. In *2nd Workshop on Copula and its Applications*, page 12, 2012.
- [16] C Amblard and S Girard. A semiparametric family of symmetric bivariate copulas. *Comptes Rendus de l'Académie des Sciences-Series I-Mathematics*, 333(2):129–132, 2001.
- [17] JS Huang and Samuel Kotz. Modifications of the farlie-gumbel-morgenstern distributions. a tough hill to climb. *Metrika*, 49(2):135–145, 1999.
- [18] Norman L Johnson and Samuel Kott. On some generalized farlie-gumbel-morgenstern distributions. *Communications in Statistics-Theory and Methods*, 4(5):415–427, 1975.
- [19] Norman L Johnson and Samuel Kotz. On some generalized farlie-gumbel-morgenstern distributions-ii regression, correlation and further generalizations. *Communications in Statistics-Theory and Methods*, 6(6):485–496, 1977.
- [20] I Bairamov, S Kotz, and M Bekci. New generalized farlie-gumbel-morgenstern distributions and concomitants of order statistics. *Journal of Applied Statistics*, 28(5):521–536, 2001.

- [21] CD Lai and M Xie. A new family of positive quadrant dependent bivariate distributions. *Statistics & probability letters*, 46(4):359–364, 2000.
- [22] Ismihan Bairamov and Samuel Kotz. On a new family of positive quadrant dependent bivariate distributions. *International Mathematical Journal*, 3(11):1247–1254, 2003.
- [23] RB Nelsen, JJ Quesada-Molina, and JA Rodríguez-Lallena. Bivariate copulas with cubic sections. *Journal of Nonparametric Statistics*, 7(3):205–220, 1997.
- [24] JJ Quesada-Molina and JA Rodríguez-Lallena. Bivariate copulas with quadratic sections. *Journaltitle of Nonparametric Statistics*, 5(4):323–337, 1995.
- [25] Helene Cossette, Etienne Marceau, and Fouad Marri. On the compound poisson risk model with dependence based on a generalized farlie–gumbel–morgenstern copula. *Insurance: Mathematics and Economics*, 43(3):444–455, 2008.
- [26] Margaret Armstrong and Alain Galli. Sequential nongaussian simulations using the fgm copula. *Cerna Working Paper*, 2002.
- [27] Mathieu Bargès, Hélène Cossette, Stéphane Loisel, Etienne Marceau, et al. On the moments of the aggregate discounted claims with dependence introduced by a fgm copula. *Astin Bulletin*, 41(1):215–238, 2011.
- [28] Yoon-Sung Jung, Jong-Min Kim, and Jinhwa Kim. New approach of directional dependence in exchange markets using generalized fgm copula function. *Communications in Statistics-Simulation and Computation*®, 37(4):772–788, 2008.
- [29] Jong-Min Kim, Yoon-Sung Jung, Engin A Sungur, Kap-Hoon Han, Changyi Park, and Insuk Sohn. A copula method for modeling directional dependence of genes. *BMC bioinformatics*, 9(1):225, 2008.
- [30] D Long and R Krzysztofowicz. Farlie-gumbel-morgenstern bivariate densities: Are they applicable in hydrology? *Stochastic Hydrology and Hydraulics*, 6(1):47–54, 1992.