



HAL
open science

Economic values of body weight, reproduction and parasite resistance traits for a Creole goat breeding goal

Mélanie Gunia, Nathalie Mandonnet, Rémy R. Arquet, Gisèle Alexandre, Jean-Luc Gourdine, Michel Naves, Valérie Angeon, Florence Phocas

► **To cite this version:**

Mélanie Gunia, Nathalie Mandonnet, Rémy R. Arquet, Gisèle Alexandre, Jean-Luc Gourdine, et al.. Economic values of body weight, reproduction and parasite resistance traits for a Creole goat breeding goal. *Animal*, 2013, 7 (1), pp.22-33. 10.1017/S1751731112001413 . hal-01000063

HAL Id: hal-01000063

<https://hal.science/hal-01000063>

Submitted on 29 May 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Economic values of body weight, reproduction and parasite resistance traits for a Creole goat breeding goal

M. Gunia¹, N. Mandonnet^{1†}, R. Arquet², G. Alexandre¹, J-L. Gourdine¹, M. Naves¹, V. Angeon³ and F. Phocas⁴

¹Unité de Recherches Zootechniques, UR143, INRA Antilles Guyane, domaine de Duclos, 97170 Petit-Bourg, Guadeloupe, F.W.I.; ²PTEA, UE1294, INRA Antilles Guyane, domaine de Gardel, 97160 Le Moule Guadeloupe, F.W.I.; ³CEREGMIA, Université des Antilles et de la Guyane, Faculté de droit et d'économie, Campus de Schoelcher, BP 7209, 97275 Schoelcher Cedex, Martinique, F.W.I.; ⁴INRA Génétique Animale et Biologie Intégrative, UMR 1313, 78352 Jouy-en-Josas, France

(Received 27 September 2011; Accepted 1 May 2012; First published online 9 July 2012)

A specific breeding goal definition was developed for Creole goats in Guadeloupe. This local breed is used for meat production. To ensure a balanced selection outcome, the breeding objective included two production traits, live weight (BW11) and dressing percentage (DP) at 11 months (the mating or selling age), one reproduction trait, fertility (FER), and two traits to assess animal response to parasite infection: packed cell volume (PCV), a resilience trait, and faecal worm eggs count (FEC), a resistance trait. A deterministic bio-economic model was developed to calculate the economic values based on the description of the profit of a Guadeloupean goat farm. The farm income came from the sale of animals for meat or as reproducers. The main costs were feeding and treatments against gastro-intestinal parasites. The economic values were 7.69€ per kg for BW11, 1.38€ per % for FER, 3.53€ per % for DP and 3×10^{-4} € per % for PCV. The economic value for FEC was derived by comparing the expected profit and average FEC in a normal situation and in an extreme situation where parasites had developed resistance to anthelmintics. This method yielded a maximum weighting for FEC, which was -18.85€ per log(eggs per gram). Alternative scenarios were tested to assess the robustness of the economic values to variations in the economic and environmental context. The economic values of PCV and DP were the most stable. Issues involved in paving the way for selective breeding on resistance or resilience to parasites are discussed.

Keywords: breeding objective, economic value, goat, parasite resistance, fertility

Implication

This study provides the definition of the breeding goal for a local breed of goat used for meat production in the tropics. The breeding goal definition includes weight and reproduction as well as parasite resistance and resilience traits. This step of defining a breeding goal is crucial for the implementation of a breeding programme aimed at increasing goat meat production in Guadeloupe by giving priority to the indigenous breed of the island.

Introduction

Approximately 31 000 goats (Agreste, 2006) are reared for meat production in Guadeloupe (F.W.I., 16° Lat. N., 61° Long. W.). The local breed, named Creole, originates from a mixture of West African, European and Indian breeds imported to the island during colonisation. Naturally selected

for its ability to thrive in a tropical environment, this breed shows good reproduction and parasite resistance performance. However, the goat population in Guadeloupe does not fulfil the local demand for goat meat since production represents only 58% of the apparent consumption (Alexandre *et al.*, 2008). As an example, 1.4 tonnes of goat and sheep meat were imported in 2009 (Agreste, 2009).

Therefore, farmers, extension services and the guadeloupean organisation of goat breeders, called Cabricoop, have decided to organise a breeding programme for the Creole breed, designed with INRA expertise. Selection on adaptive as well as productivity traits has been advised for tropical areas (Shrestha and Fahmy, 2007) due to the specific constraints (climate, diseases, poor feeding) of the environment. As shown by a survey (Gunia *et al.*, 2010), farmers wish to improve the meat production of the Creole breed as well as its reproductive performance. Disease resistance traits should also be included in the breeding objective, because of the high prevalence of gastro-intestinal nematodes

† E-mail: nathalie.mandonnet@antilles.inra.fr

such as *Haemonchus* sp. and *Trichostrongylus* sp. in Guadeloupe. They are responsible for more than 80% of the mortality rate before weaning (Aumont *et al.*, 1997) and 30% loss in productivity in kids at slaughter age.

Defining the breeding goal is the preliminary crucial step before the setting up of a breeding programme (Hazel, 1943, Ponzoni and Newman, 1989). The breeding goal should encompass all traits affecting profitability, which is typically defined at the farm level. Economic values are derived by the partial differentiation of the profit function with respect to individual traits (Moav and Hill, 1966). Each economic value indicates the marginal increase in profit caused by the improvement of one physical unit of the corresponding trait of the breeding objective.

The aim of this study was threefold: first, derive the economic values of production, reproduction and parasite resistance traits for a Creole goat breeding scheme, second, analyse the sensitivity of these economic values to changes in the economic and the environmental context of goat meat production and third, identify the means of assisting farmers in appropriating and exploiting our findings.

Material and methods

Basic features of the model

A deterministic bio-economic model was developed to calculate the economic values of breeding goal traits relevant to Creole goat production. This model considered the profit of a Guadeloupean goat farm as a function of specific biological traits to be genetically improved. Deterministic models have been used widely for such aims, in goat (Bett *et al.*, 2007) or sheep (Kosgey *et al.*, 2003; Legarra *et al.*, 2007), as well as in cattle (Phocas *et al.*, 1998). The level of management modelled for the farm used in this study was above average relative to those farms registered in the Cabricoop breeding organisation. This model farm represented the management and farming practices observed in most Cabricoop farms (Gunia *et al.*, 2010). Some practices described in the model (such as three kiddings in 2 years and selective anthelmintic treatment of does) were not found in all Cabricoop farms. Hence, this goat farm model does not represent the diversity of practices in field data, but instead represents a technical level that could be achieved by most of the Cabricoop farms.

Performance recording has not yet been implemented by farmers. Therefore, the herd performances and production parameters were derived from the 30-year-old database of the Creole goat experimental herd at INRA-Gardel (Alexandre *et al.*, 1997; Mahieu *et al.*, 2008). The prices used in this study were the market prices of 2010. Only variable costs (which depend on the level of animal performance) and fixed cost per animal (which does not vary with the performance level) were considered in the cost function. The fixed costs per herd were not considered because they do not depend on the level of animal performance or the number of animals. Therefore, the profit function is a gross margin of the modelled farm without accounting for any salaries, subsidies, investments or housing taxes. Because genetic improvement should not be a substitute for technical progress, the farm management is




Figure 1 Flock composition of the modelled Creole goat herd for one year.

assumed to be at maximum technical efficiency to derive the economic values (Smith *et al.*, 1986). Equations of costs and income, as well as the profit function were calculated using Microsoft Excel 2002[®].

Description of herd management

A pasture-fed herd of 30 Creole does was simulated. The flock composition and dynamics for one year are shown in Figure 1. The corresponding flock structure parameters as well as the main production parameters are presented in Table 1.

The 30 does were mated three times in 2 years with a ratio of 15 does mated per natural service buck. Kids were reared by their mothers, before being weaned at 3 months of age. Female replacement was carried out within the herd, whereas new bucks were brought in from other farms. Each year, 7.8 young females were kept to renew the doe herd. The other females and males were sold at 11 months of age (except for one male, which was consumed by the farmer's family); three young bucks were sold alive for breeding, whereas the others were sold for slaughter. All kids were sold at 11 months of age, whatever their weight. All goats grazed on *Dichanthium* sp. pasture throughout the year. Commercial concentrate was given to does for 2 months before delivery and for 3 months after kidding and to kids from weaning to 11 months of age. Growth was assumed to be linear from weaning to 11 months and from 11 months to adult age.

Kids between 3 and 11 months of age and adult bucks were regularly drenched against gastro-intestinal nematodes. The main species found in Guadeloupe were *Haemonchus contortus*, *Trichostrongylus colubriformis* and *Oesophagostomum*

Table 1 Flock structure and production parameters in the Creole goat farm model

Production and management parameters	Unit	Mean
Flock structure parameters		
Fertility	%	82
Litter size	kids	2.3
Pre-weaning mortality rate	%	18
Post-weaning mortality rate	%	12
Annual culling rate of does	%	15
Annual culling rate of bucks	%	50
Mortality rate of does	%	11
Mortality rate of bucks	%	3
Production parameters		
Weaning weight (at 90 days)	kg	8.8
Weight at 11 months	kg	17.9
Mature weight of does (at 4 years)	kg	37
Weight of buck at 3 years	kg	35.4
Average daily gain (weaning to 11 months)	g/day	37
Average daily gain (11 months to mature weight) of does	g/day	17
Average daily gain (11 months to 3 years) of bucks	g/day	23
Dressing percentage	%	40
Average daily milk yield	l/day	1.02
Milk fat content	g/kg	56
Milk protein content	g/kg	37
Proportion of does with a PCV \geq 16	%	44.5
Faecal worm eggs count at 11 months	epg ^a	938

PCV = packed cell volume.

^aWorm eggs per gram of faeces.

columbianum. The drenching frequency was three times per year for adults and four times per year for kids. Does were drenched using the Famacha[®] method (Bath *et al.*, 1996). The Famacha[®] method is a measurement of the ocular conjunctivae colour, which allows selective drenching of only anaemic animals. In humid countries where *H. contortus*, a haematophagous nematode, is the primary parasitic pathogen, this method can be applied at the farm level to reduce the number of treatments administered. However, other parasites such as *T. colubriformis* cannot be monitored using this method. In Creole goats, a Famacha[®] score of 4 or 5 (anaemic animals) can be related to a packed cell volume (PCV) below or equal to 16. Using this method, 55.5% of the fertile Creole does were drenched each year and the drenched goats were dosed 1.54 times on average (Mahieu *et al.*, 2007). The does that failed to reproduce were not treated. They were less susceptible to gastro-intestinal nematode infection than pregnant and lactating does (Mandonnet *et al.*, 2005).

All animals were sprayed fortnightly with acaricide to prevent the occurrence of tick-borne diseases such as heartwater and dermatophilosis.

Breeding goal traits

The main traits in the breeding goal and their abbreviations are presented in Table 2. In order to ensure a balanced selection, the breeding objective consisted of two production

Table 2 Breeding goal traits for Creole goat

Trait	Unit	Abbreviation
Live weight at 11 months (mating or selling age)	kg	BW11
Fertility: number of doe kiddings per mating	%	FER
Dressing percentage at 11 months	%	DP
PCV of does	%	PCV
Faecal worm eggs count at 11 months	log(epg) ^a	FEC

BW11 = live weight at 11 months; FER = fertility; DP = dressing percentage at 11 months; PCV = packed cell volumes of does; FEC = faecal worm eggs count measured at 11 months of age.

^aNatural logarithm of worm eggs per gram of faeces.

traits, live weight (BW11) and dressing percentage (DP) at 11 months (the mating or selling age), and one reproduction trait: fertility (FER), which is the number of doe kiddings per mating. To assess animal response to parasite infection, two other traits were also considered. PCV, a resilience trait, is a measurement of the proportion of red cells in the blood used to diagnose anaemia caused by haematophagous parasites such as *H. contortus*. This trait was measured on does. Faecal worm eggs count (FEC) is a measurement of the number of gastro-intestinal parasite eggs found in the faeces. This measurement was used to assess the resistance of an animal and was measured on kids at 11 months of age. Mortality was not included in the breeding objective because the estimate of the genetic variability of this trait in Creole goats was null, probably due to the very low number of dead animals in the INRA herd (Mandonnet *et al.*, 2003). Litter size (LS) was deliberately not included in the breeding objective because in the model only the positive effects of LS could be taken into account easily, although negative effects are expected, such as increased labour for the farmer (especially if artificial milking is required) and lower kid weight.

Income and costs equations

Profit. The farm profit was calculated as the difference between income and costs:

$$\text{Profit} = \text{Income} - \text{costs} \quad (1)$$

The farm income came from the sale of animals: culled does and bucks sold for meat purposes as well as a few young bucks sold to other herds for breeding purposes. The farm costs were feed (grass and commercial pellets), treatments against gastro-intestinal parasites and ticks as well as identification (ear tags). The prices are presented in Table 3.

For the income and costs equations, five categories of animals were considered in the model: kidding does, non-kidding does, bucks, kids from 3 to 11 months of age (i.e. weaning to selling or mating age) and kids before weaning. The average number of animals in each category was expressed as a function of the number N_0 of mated does according to the following equation:

$$N_i = a_i \times N_0 \quad (2)$$

Economic values of body weight, reproduction and parasite resistance traits for a Creole goat breeding goal

Table 3 Prices in the income and costs equations of the Creole goat farm model

	Unit	Value
Carcase price of adults	€/kg of carcase	12.70
Carcase price of kids at 11 months	€/kg of carcase	11.96
Price of living animals at 11 months	€/kg of live weight	10.00
Anthelmintics cost	€/kg of live weight	0.03
Grass cost	€/kg of dry matter	0.10
Commercial concentrate cost	€/kg of dry matter	0.44
Acaricide cost	€/animal	0.04
Ear tag cost	€/kid	2.00

where i is the animal category (one kidding does, two non-kidding does, three bucks, four kids from 3 to 11 months of age, five kids before weaning) and a_i is a coefficient varying for each animal category:

$$a_1 = \text{FER};$$

$$a_2 = (1 - \text{FER});$$

$$a_3 = \text{breeding buck to breeding does ratio};$$

N_4 , the number of kids from 3 to 11 months of age, is the average of the number of kids weaned and the number of kids reaching 11 months of age.

$$a_4 = 0.5 \times [1.5 \times \text{FER} \times \text{LS} \times (1 - \text{pre-weaning mortality rate}) + 1.5 \times \text{FER} \times \text{LS} \times (1 - \text{pre-weaning mortality rate}) \times (1 - \text{post-weaning mortality rate})];$$

$$a_4 = 0.5 \times 1.5 \times \text{FER} \times \text{LS} \times (1 - \text{pre-weaning mortality rate}) \times (2 - \text{post-weaning mortality rate});$$

N_5 , the number of kids before weaning, is the average of the number of kids born and the number of kids weaned.

$$a_5 = 0.5 \times [1.5 \times \text{FER} \times \text{LS} + 1.5 \times \text{FER} \times \text{LS} \times (1 - \text{pre-weaning mortality rate})];$$

$$a_5 = 0.5 \times 1.5 \times \text{FER} \times \text{LS} \times (2 - \text{pre-weaning mortality rate}).$$

The coefficient 0.5 was used to calculate the average between the two groups of kids: weaned kids and 11-month-old kids for a_4 , and kids born and weaned for a_5 . The coefficient of 1.5 was used to take into account the three kiddings in 2 years to obtain an annual profit function.

Income. The income equations were:

$$\begin{aligned} \text{Income of meat} &= \text{Number of animals sold for meat} \\ &\quad \times \text{Live weight} \times \text{Dressing percentage} \\ &\quad \times \text{Price per kg of carcase weight} \end{aligned} \quad (3)$$

$$\begin{aligned} \text{Income of animals sold alive} &= \text{Number of animals sold alive} \times \text{Live weight} \\ &\quad \times \text{Price per kg of live weight} \end{aligned} \quad (4)$$

The selling price of a goat depended on its carcase or live weight. A fixed price per kg of live weight was also considered for young males sold alive as reproducers. For animals sold for slaughter, the price per kg of carcase weight was higher for the stock class where the average carcase weights are higher.

Carcase quality is not presently taken into account in Guadeloupe. Carcase price is not much higher than live weight price due to the differences in marketing network: official for goats sold to the slaughter house (controlled prices with taxes and fees deducted) and informal for live goats (with free prices).

Subsidies were ignored in the income equations because of their variable amounts and questionable persistence.

Costs. The cost equations were:

$$\begin{aligned} \text{Anthelmintics costs for does} &= N_1 \times \text{Proportion of does with PCV} \\ &\leq 16 \times \text{Live weight} \times \text{Price of anthelmintics} \\ &\quad \times \text{Number of treatments per year} \end{aligned} \quad (5)$$

$$\begin{aligned} \text{Anthelmintics costs for bucks and kids} &= N_i \times \text{Live weight} \times \text{Price of anthelmintics} \\ &\quad \times \text{Number of treatments per year; with } i = 3, 4 \end{aligned} \quad (6)$$

$$\begin{aligned} \text{Acaricide costs} &= N_i \times \text{Price of acaricide} \\ &\quad \times \text{Number of treatments per year; with } i = 0, 3, 4 \end{aligned} \quad (7)$$

$$\text{Identification costs} = N_5 \times \text{Price of ear tags} \quad (8)$$

The price of anthelmintics (expressed per kg of live weight) was the average of the market prices of different anthelmintics commonly used in Guadeloupe.

Feed costs were based on diets composed of *Dichantium sp* grass and commercial pellets if the grass was not sufficient to meet the animal feed requirements. Energy and protein needs as well as feed intake were calculated for each animal category. They were expressed as a function of live weight and average daily gain and were calculated using two different systems. INRA equations (Sauvant *et al.*, 2007) were used for doe diets. The requirements and feed intake were different for does whose mating was successful and does whose mating was unsuccessful. Increased feed requirements for lactating does were attributed to the does and not to the kids. For bucks and growing kids, the equations of metabolisable energy and metabolisable protein requirements considered were based on the equations of Sahlou *et al.* (2004) for indigenous goats. Dry matter (DM) intake was calculated from the equations of Luo *et al.* (2004). For all animal categories, the amount of grass required was increased by 10% from the calculated amount to take into account refusals. The cost of grass was the maintenance cost of the pasture. It included the cost of fencing, fertilisation, irrigation and labour (Fleury J., personal communication). The cost of pellets was the commercial price paid by Cabricoop farmers.

Derivation of the economic values of BW11, FER, DP and PCV in the base situation

The economic values were the marginal variation in annual profit per mated doe, due to the increase of one unit of the

trait, while maintaining the other traits constant at the population mean:

$$EV_t = (P_1 - P_0) / N_0 \quad (9)$$

with EV_t being the economic value of trait t , P_1 the profit after an increase of one unit of the trait t and P_0 the initial profit for $N_0 = 30$ does in the base situation. The economic values of the breeding goal traits were derived by 'finite differences' approximation of derivatives rather than by direct analytical derivatives of the profit function with respect to trait t , since the efficiency of this procedure has been proven by Phocas *et al.* (1998).

Standardised economic values were calculated by multiplying the economic values (expressed in euro per physical unit of the traits) by the genetic standard deviations of the traits. The units of the traits used in this study are presented in Table 2. These genetic standard deviations were 1.89 for BW11, 12.25 for FER, 1.60 for PCV and 0.38 FEC in Creole goats (Gunia *et al.*, 2011). For DP, no data were available for Creole goats or other breeds of goat in the literature; hence, the genetic standard deviation (1.22) given by Greeff *et al.* (2008) in sheep was used.

Derivation of the economic value for DP implies only the income equation for meat (3) because DP was not linked to any cost.

Derivation of the economic value for BW was based on the income equations (3) and (4), the feed cost equations (equations not detailed here) and the anthelmintic cost equations (5) and (6).

FER influences the number of kids, as shown in equation (2). Therefore, derivation of its economic value requires that changes in all income and costs concerning kids are taken into account. Changes in the feed costs of does also need to be accounted for because the diet differs between fertile and barren does.

According to equation (5), a change in profit ΔP will occur on increasing the proportion of does with $PCV \geq 16$ by 1%. In order to rescale this ΔP at the animal level to derive the economic value for PCV, an underlying normal distribution of PCV was assumed that allowed the derivation of the economic value for PCV by using the probability density function $\Phi(x)$ of a normal distribution:

$$EV_{PCV} = \Phi(x) \cdot \Delta P \quad (10)$$

$$\Phi(x) = \frac{1}{\sqrt{2\pi\sigma^2}} e^{-(x-\mu)^2/2\sigma^2} \quad (11)$$

with $\mu = 24.13$ being the mean of the PCV distribution, $\sigma = 4.97$ its phenotypic standard deviation and $x = 16$ the threshold value of PCV.

Independent calculation of the economic value of FEC when anthelmintics are ineffective

No obvious relationship was found between FEC and costs or revenues, which made it difficult to derive the economic value of this trait. Therefore, a somewhat different method was used to independently calculate the maximum economic

value of FEC when selection was focused only on this trait. The economic value of FEC was derived by considering the expected profit and average FEC in the base situation and in a situation in which anthelmintics were ineffective, because parasites had developed resistance to these products.

The equation used to derive the economic value of FEC was as follows:

$$EV_{FEC} = \frac{1}{N_0} \times \frac{\delta P}{\delta FEC} = \frac{1}{N_0} \times \frac{\delta P}{\delta(e^{FEC})} \times \frac{\delta(e^{FEC})}{\delta FEC} \\ \approx \frac{1}{N_0} \times \frac{P_A - P_B}{e^{FEC_A} - e^{FEC_B}} \times e^{FEC_B} \quad (12)$$

with EV_{FEC} being the economic value of FEC, N_0 the number of does, which was fixed and equal to 30, P the profit, e^{FEC} the exponential function of FEC, A the situation with ineffective anthelmintics and B the base situation. FEC was the only trait in the breeding goal, because the change in profit was not calculated by maintaining BW11, FER and PCV constant at their average level.

In the situation with ineffective anthelmintics, parasites were not controlled by any veterinary product. Mortality and FEC increased (by 55% to 60% for mortality and by almost 400% for FEC), whereas FER, LS and growth rate decreased by 1.2%, 3% and 35%, respectively. Drenching was discontinued due to its inefficiency. The modified parameters in this extreme scenario are presented in Table 4. They were based on the performance records at INRA-Gardel, in 1999, when the anthelmintics applied were no longer effective (Chevalier, 2001). Considerable loss in production had occurred during this period, with an increase in mortality of 55% to 60% and a decrease in the average daily gain of 35% for kids during the pre-weaning period and the post-weaning period until 11 months of age. No data were available on growth after 11 months or on the mortality of mature does and bucks. We therefore assumed that they showed the same reduction in growth and increase in mortality as that found in kids after weaning.

Sensitivity analysis

After deriving the economic values in the base scenario with a fixed number of does, six alternative situations were studied to analyse the sensitivity of profit and economic values to changes in the economic context. The six situations were as follows: fixed amount of grass, fixed amount of pellets, increased carcase price, increased pellets price, drought and expensive anthelmintic.

Fixed feed input scenarios. In the base situation described previously, the number of does was assumed to be fixed each year. Another strategy to define a breeding goal is to consider some physical or economical constraints that involve using a fixed amount of input instead of a fixed number of does. The following two situations were considered: a situation with a fixed amount of grass and a situation with a fixed amount of pellets. These two situations provided a good insight into the importance of the different

Table 4 Flock structure and production parameters in the Creole goat farm model in the situation with ineffective anthelmintics

Parameters	Unit	Mean
Flock structure parameters		
Fertility	%	81
Litter size	kids	2.23
Pre-weaning mortality rate	%	29
Post-weaning mortality rate	%	19
Mortality rate of does	%	17
Mortality rate of buck	%	4.7
Production parameters		
Weaning weight	kg	8.03
Weight at 11 months	kg	13.9
Mature weight of does (at 4 years)	kg	25.3
Weight of buck at 3 years	kg	26.3
Average daily gain (weaning to 11 months)	g/day	24
Average daily gain (11 months to mature weight) of does	g/day	11
Average daily gain (11 months to 3 years) of bucks	g/day	15
Faecal worm eggs count at 11 months	epg ^a	4588

^aWorm eggs per gram of faeces.

traits in two different feeding systems. Limitation in the amount of grass may occur when farmers cannot expand their pasture surface. On an island, this situation is highly probable. Financial difficulties (limitation in cash flow) could also prevent the farmer from increasing the quantity of pellets. In these two situations, the economic values were derived by changing the number of does after a marginal increase in a trait in order to fit the size of the quota (Groen, 1989). If the marginal increase of the trait led to an increase in the total amount of fodder (or pellets) consumed, the total number of does was reduced to maintain the amount of feed at its initial level. The economic values in the fixed input situations were calculated using the following equation:

$$EV_{ft} = (P_{ft} - P_0) / N_0 \quad (13)$$

where EV_{ft} is the economic value of the trait t in the situation with fixed input f , N_0 the number of mated does in the base situation, P_{ft} the profit after an increase of one unit of the trait t and adjustment of the number of does in the situation with fixed input f and P_0 the initial profit.

The new number of does changed to fit the quota and used in the income and costs equations for the calculation of P_{ft} was as follows:

$$N_{ft} = \frac{QT_f}{\sum_{i=1}^4 a_i Q_{ift}} \quad (14)$$

with N_{ft} being the new number of does after an increase of one unit of the trait t in the situation with fixed input f , QT_f the total quantity of feed f given to the whole herd in one year in the base situation before the increase of trait t , Q_{ift} the quantity of feed f given to the animal category i ($i = 1$ to 4) in one year after an increase of one unit of the trait t and a_i the coefficient varying for each animal category i already described in equation (2).

Slow increase in price. In this situation, we studied the consequences on the farm profit of realistic and likely increases in carcase weight price and concentrate price. We estimated what these prices would be in 2015 using a linear extrapolation of the prices of 2000 to 2010. Each price was increased while maintaining the other prices constant. The carcase price would be increased by 2.50€ per kg of carcase, which represents an increase of 20% of the initial price. The commercial pellet price would be increased by 0.048€ per kg of DM matter, which represents an increase of 10.9% of the initial price.

Extreme situations. We tested the stability of the farm profit and the economic values in two extreme situations: 'drought' and 'expensive anthelmintic'. These two situations were realistic, even if their probability of occurrence may be low.

In the 'drought' scenario, rainfall would decrease markedly in the driest part of the island at present, where most of the goat farms are situated. In this situation, farmers would have to buy hay instead of growing grass. We used the present price of hay (0.40€/kg of DM), even though this price could increase in the case of long-lasting drought. Hay is expensive in Guadeloupe because agricultural machines are imported from Metropolitan France at a high cost, their maintenance is expensive and the volumes of hay produced are limited, resulting in high production costs per hay bale.

In the 'expensive anthelmintic' scenario, the present anthelmintic would no longer be used, because of a resistance developed by parasites to these products. The only possibility of treatment would be the use of a new and expensive anthelmintic. Its price was fixed at 10 times the present price, that is, 0.3€ per kg of live weight.

The use of alternative scenarios provided us with the possibility of assessing the robustness of the economic values to variations in the economic and environmental contexts.

Results

Income, costs and profit

The income, costs and profit in different situations are presented in Table 5. In the base situation as well as in the situation with a fixed feed input, the profit of the farm was 90.29€/doe per year. The total profit per year for the 30-doe herd was therefore 2709€. In the situation used to calculate the economic value of FEC, with ineffective anthelmintics, profit was reduced by 73.36€. This reduction represents an 81% loss in profit, which clearly emphasises the cost of parasite infections. In the slow price rise situations, the change in profit varied. The 11% increase in the price of the pellets had a limited impact on profit, which only increased by 3.5%, whereas the 20% increase in carcass price had a major impact on profit, which increased by 40%. In extreme situations, profit decreased markedly. In the drought situation, the costs exceeded the income and the farmer lost 96€/doe per year. Buying hay would not be a good solution to compensate for the lack of grass. A 44% loss in profit was observed in the situation that required the use of expensive anthelmintics.

General consideration of economic values in alternative situations

The economic values for BW11, FER, DP and PCV in alternative situations as well as the economic value of FEC are presented in

Table 6. The economic values of BW11, FER, DP and PCV were all positive in the base situation as well as in the alternative situations. The economic value of FEC was the only negative one. Using standardised economic values facilitated a comparison of traits. In the base situation, FER had the highest standardised economic value, followed by BW11, which clearly indicated the importance of these two traits. The standardised economic value of FEC came in third position (in absolute value). However, this value should be considered with caution due to the calculation method used. It reflects more the maximum economic value of FEC than its current value in 2011. The economic value of DP was lower, whereas the economic value of PCV was very low, which is probably due to the underestimation of the costs associated with PCV. The highest economic values were found for the scenario with increased carcass price and the lowest for the drought scenario. The economic values of BW11 and FER were very sensitive to changes of situation, which was undoubtedly due to their relationship with the modified prices and costs, whereas the economic values of DP and PCV, which were not related to the value or the amount of concentrate and grass, remained unchanged.

Economic values with fixed feed input

In the situations with fixed feed input, the economic values of BW11 and FER decreased. In the situation with a fixed

Table 5 *Income, costs and profit (in €/doe per year) of a Creole goat farm in different situations*

Situation	Base or fixed feed input	Ineffective anthelmintics	Increased carcass price	Increased pellets price	Drought	Expensive anthelmintic
Income	193.12	98.25	229.38	193.12	193.12	193.12
Costs	102.83	81.31	102.83	106	289.06	142.72
Profit	90.29	16.93	126.55	87.12	-95.94	50.4

Table 6 *Economic values for each trait of the breeding goal for Creole goat in different situations*

Situation	Base	Ineffective anthelmintics	Fixed amount of grass	Fixed amount of pellets	Increased carcass price	Increased pellets price	Drought	Expensive anthelmintic
Economic values in trait unit ^a								
BW11 (kg)	7.69	–	5.1	6.74	9.52	7.65	2.63	6.26
FER (%)	1.38	–	0.87	0.28	1.8	1.34	0.34	0.91
DP (%)	3.53	–	3.53	3.53	4.27	3.53	3.53	3.53
PCV (%)	3×10^{-4}	–	3×10^{-4}	3×10^{-4}	3×10^{-4}	3×10^{-4}	3×10^{-4}	3×10^{-4}
FEC log(epg) ^b	–	-18.85	–	–	–	–	–	–
Standardised economic values ^d								
BW11	14.56	–	9.65	12.77	18.03	14.5	4.99	11.86
FER	16.9	–	10.68	3.37	22.01	16.43	4.2	11.16
DP	4.32	–	4.32	4.32	5.23	4.32	4.32	4.32
PCV	4.7×10^{-4}	–	4.7×10^{-4}	4.7×10^{-4}	4.7×10^{-4}	4.7×10^{-4}	4.7×10^{-4}	4.7×10^{-3}
FEC	–	-7.12	–	–	–	–	–	–

BW11 = live weight at 11 months; FER = fertility; DP = dressing percentage at 11 months; PCV = packed cell volumes of does; FEC = faecal worm eggs count measured at 11 months of age.

^aEconomic values expressed in €/trait unit per doe per year.

^bNatural logarithm of worm eggs per gram of faeces.

^cEconomic values in €/trait unit per doe per year multiplied by the genetic standard deviation of the trait. The standardised economic values provided the gain in profit caused by the improvement of one genetic standard deviation of the trait.

amount of pellets, the economic values of FER decreased considerably (by 80%), whereas BW11 was less affected (12% loss). In contrast, in the situation with a fixed amount of grass, the economic values of FER and BW11 decreased almost equally (by 37% for FER and by 34% for BW11).

An increase in FER of 1% led to an increase in the number of pregnant and lactating does as well as the number of kids. Therefore, the total amount of grass and pellets consumed increased. The increase in the amount of pellets provided was especially high, due to the higher proportion of pregnant and suckling does, which required pellets in addition to grass. The reduction in the number of does to fit the quota was therefore higher in the situation with a fixed amount of pellets than in the situation with a fixed amount of grass. In the fixed amount of pellets scenario, the new profit (P_{ft}) and therefore the economic value of FER decreased more markedly than in the fixed amount of grass scenario.

An increase in BW11 led to an increase in the average weights of kids and does, whose energy requirements increased accordingly. For kids, the amounts of both grass and pellets increased. For does, the feed intake capacity increased and allowed them to fulfil their additional energy requirements with grass instead of concentrate. The total amount of pellets consumed by does was even slightly reduced. Hence, an increase in BW11 of 1 kg had more consequences on the total amount of grass than on the total amount of pellets. The reduction in the number of does to fit the quota is therefore higher in the situation with a fixed amount of grass, which decreased the new profit (P_{ft}) and therefore the economic value of BW11 more drastically than in the situation with a fixed amount of pellets.

Economic values with a slow increase in price

In the situation with carcass prices increased by 20%, the economic values of BW11, FER and DP increased by 21% to 30%. In the scenario where pellet prices increased by 11%, the economic value of FER decreased by 3%, BW11 remained almost the same and the others did not vary. The economic value of BW11 remained stable because an increase of this trait did not alter significantly the total amount of pellets consumed, as explained in the previous paragraph.

Economic values in extreme situations

The extreme situations modified the economic values considerably. They all decreased, except for the economic value of PCV, which multiplied by 10 in the situation with the use of expensive anthelmintics. This economic value was still lower than the economic values of BW11 and FER, which were reduced by 18% and 34%, respectively. In the drought scenario, the standardised economic values of BW11, FER and DP were all in the same range. The increased price of hay reduced the economic values of BW11 by 66% and those of FER by 75%, bringing them down to the same level as the economic value of DP.

Discussion

Profit and economic values under alternative situations

This research sought to use profit equations in a bio-economic model to derive the economic values of the traits

of the breeding objective. As shown in a survey (Gunia *et al.*, 2010), goat breeding is a supplementary professional activity. The small size of herds (about 30 does) and the 2709€ profit per year are not sufficient to maintain this activity alone. Extreme situations jeopardise the future of the goat breeding activity by substantially decreasing a farmer's profit.

The differences in the timing and the frequency of expression of the different traits were not accounted for by using discounted gene flow principles. The gene flow principles are based on the number of expected improved progeny when following the genes of an improved reproducer. When accounting for each of the selection paths in a closed selection nucleus, each selection path makes the same genetic contribution in the long term. Because all costs and expenses are expressed by breeding doe over all age classes, the right number of expressions of improved parents for each trait is accounted for at a given time period at birth of their progeny. Then, the only issue is the fact that one may wish to discount the expressions of improved traits to account for their different patterns of expressions over time. In our case, all traits are expressed with a very short period of time for a given progeny cohort (between 11 months and 16 months). Therefore, there is not a long delay between the expression of production and functional traits. Considering discounting would only have a significant impact on the relative economic weights if high discount rates are considered. However, imposing high discount rates yields a short-term perspective, which conflicts with the inherently long-term nature of genetic improvement. Without discounting, the weight of the resilience trait is almost set to zero in the breeding goal. Discounting will only reinforce the very small economic value of resilience trait *v.* production and resistant traits. Therefore, we propose to test the robustness of the economic weights and regularly update them, rather than to account for discounted gene flow in the derivation of breeding goal.

Therefore, running alternative scenarios facilitated the assessment of the robustness of the economic values to changes of situation. The economic values under different constraints were not equivalent in Creole goats. As summarised by Groen (1989), different economic values are obtained in different perspectives (fixed number of animals, fixed input) when the value of production factors differs between alternative uses. The equivalence of the economic value in different perspectives could be obtained with the use of three conditions, but were not applicable in our case. They are as follows: one-product situation or income (or other products considered as negative costs), equilibrium in a purely competitive industry and all costs being variable per unit of product. The fixed number of does or fixed amount of grass situations are probably the most appropriate perspectives in Guadeloupe breeding systems.

Choice of traits and comparison of their economic values

The traits included in the breeding objective were close to the traits desired by farmers, as reported in a survey (Gunia *et al.*, 2010). Creole goats are an indigenous breed, whose

meat production is important to farmers. Its maternal qualities are also important, especially for farmers using Creole does as a maternal line for crossbreeding.

Fertility had the highest standardised economic value in the base situation. This trait was the most important, even if the replacement costs were not included in the derivation of its economic value. Age was the main reason for culling and does that failed to reproduce were generally kept for one or two other cycles of reproduction before culling.

LS was not included in the breeding objective due to the difficulties in modelling the negative effects of an increase of this trait. Moreover, farmers want to maintain the high LS of Creole goats (2.3 kids per doe), without increasing it. In studies on sheep breeds used for meat production (Wolfova *et al.*, 2009; Byrne *et al.*, 2010), LS was one of the traits with the highest standardised economic value. As observed by Amer *et al.* (1999a), the economic value of LS depended on its current average level and on the intensity of the production systems. These authors modelled the proportion of singles, twins, triplets and quadruplets as flock prolificacy increased. The economic value reduced as LS increased and the proportion of triplets or quadruplet increased. They found an optimal LS of 2.2 or 2.3 lambs per ewe lambing, which is the current prolificacy level of Creole does. Therefore, evolution of LS will be monitored and in case of necessity, future selection could aim at stabilising LS by including this trait in a restricted selection index (Itoh and Yamada, 1987).

Desirability of selecting Creole goats for parasite resistance and resilience

In Guadeloupe, as in many tropical countries, gastrointestinal parasites are endemic. Their infections are responsible for a high mortality rate, reduced growth rate, reduced reproductive performance and general loss of productivity (Perry *et al.*, 2002). Our results show that goat farm profit is reduced by 81% when parasite infections are no longer controlled by anthelmintics.

However, even if these results emphasise the benefits of using anthelmintics to control parasitism, these anthelmintics are not a sustainable solution because firstly, they have a cost, which in some rare cases cannot be compensated for by reduced loss (Perry *et al.*, 2002), secondly, residues of these products can be found in meat (Love, 1999) and thirdly, parasites develop resistance to the treatments (Kaplan, 2004). Kaplan (2004) suggested the possibility that in some places, there would not be any efficient anthelmintic remaining. Non-chemical alternative approaches have been developed for more integrated parasite control for small ruminants (Jackson and Miller, 2006; Torres-Acosta and Hoste, 2008) and for Creole goats (Mahieu *et al.*, 2009). These methods include (among others) rotational grazing, improved nutrition, selective drenching and selection of resistant or resilient animals (Baker and Gray, 2004). Selection is a long-term process with a persistent effect, in contrast to the other short-term measures aimed at reducing FEC (Eady *et al.*, 2003). Resistant animals, with low FEC, can

suppress the establishment of parasites or parasitic burden. Resilient animals are those that are capable of surviving and producing when facing a parasitic challenge. They can resist the effects of infection rather than resisting the infection itself (Bisset and Morris, 1996).

Feasibility of selecting Creole goats for parasite resistance and resilience

Faecal egg count, the density of worm eggs excreted in the faeces of the animal, is the traditional and convenient trait used to select animals resistant to parasites (Bishop and Morris, 2007). These animals contaminate the pasture less, which has a strong impact on the epidemiology of worm infection and reduces the need for treatments (Eady *et al.*, 2003). Another measurement, PCV, has also been used as a proxy for resilience (Baker *et al.*, 2001) when *H. contortus* is predominant. When animals are submitted to *H. contortus* infections, the genetic correlation between PCV and FEC is negative (Baker *et al.*, 2001; Mandonnet *et al.*, 2001). Hence, resistant animals are also resilient, at least under the challenge of this parasite (Bisset and Morris, 1996).

Attempts to select sheep and goats for resistance/resilience while comparing their performance with susceptible or control lines have proven successful (Bisset and Morris, 1996; Vagenas *et al.*, 2002). Commercial programmes that include resistance to nematodes in sheep have already been implemented in Australia, New Zealand (Woolaston and Baker, 1996) and the United Kingdom (Jackson and Miller, 2006). Previous studies in Creole goats have shown that it is possible to select this breed for resistance and resilience to parasites (Mandonnet *et al.*, 2001) by selecting on their FEC or PCV.

Benefits of selecting goats for resistance and resilience

Limited knowledge is available on goats in tropical countries under *H. contortus* infection. The main results produced are in sheep under infection from other parasite species. The selection of small ruminants on parasite resistance is interesting because it seems that parasites cannot develop resistance against the mechanisms that their host has developed (Kemper *et al.*, 2009). It is also the method with the strongest effect on FEC reduction (Eady *et al.*, 2003), which means reduced egg output leading to reduced pasture larval contamination, and therefore subsequent larval challenge (Bishop and Stear, 2003). The main benefits are thus in terms of the epidemiology of the parasite. The original hypothesis was that selecting for low FEC would increase live weight. However, this predicted increase in productivity has not often been attained (Greer, 2008). Even if gains in productivity are hypothetical, it seems reasonable to assume some reduction in the number of necessary anthelmintic treatments, as occurred in Merino sheep selected for low FEC (Kahn *et al.*, 2003). However, selecting animals on FEC would not be equivalent to drenching them, because adult worms would not be eliminated from the digestive tract.

Selection on resilience is a method that does not apply any selection pressure on parasites, which can complete their life

cycle unimpeded. Therefore, resilient animals will still contaminate the pasture. However, they can better cope with the challenge in terms of productivity (Bisset and Morris, 1996). In contrast to selection on resistance, production benefits can be expected from selection on resilience. Nevertheless, it is difficult to predict the actual benefit in terms of productivity, reduced number of treatments or potential change in FEC. Selection on resilience has received less attention in the literature than selection on resistance and the results vary according to the nematode species, the traits used to assess resilience (drench requirement, growth under challenge, PCV) and, more generally, the environment and breeding conditions.

In Creole goats, there were no significant genetic correlations between FEC and production traits, but there were favourable genetic correlations between PCV of does and weight and fertility (Gunia *et al.*, 2011). The genetic correlation of FEC measured on kids at 11 months and PCV of does was not significantly different from zero, whereas the genetic correlations between FEC and PCV measured at the same age were strongly negative (Mandonnet *et al.*, 2001; Mandonnet *et al.*, 2006). Selecting Creole goats on FEC will obviously reduce FEC in the herd and therefore pasture contamination. Productivity benefits will probably be limited, but it seems possible that drenching requirements will be reduced. Selecting on PCV will help maintain better herd health and will probably lead to some gain in production. The need for treatments will be reduced. However, it is not clear whether FEC will decrease significantly. These two traits would clearly be complementary for improving resistance and resilience in goats.

Deriving economic values for parasite resistance and resilience
As reported by Kominakis and Theodoropoulos (1999), assigning an economic value to FEC is not straightforward. Previous attempts in the scientific literature for FEC in sheep relied mostly on the desired gain method (Woolaston, 1994; Kominakis and Theodoropoulos, 1999; Gicheha *et al.*, 2005), even if this method has been criticised (Gibson and Kennedy, 1990). An attempt to derive an economic value for FEC in sheep in New Zealand and Australia was made by Amer *et al.* (1999b). This economic value takes into account a reduction in parasite epidemiology and reduced contamination in commercial flocks. It links changes in flock average FEC to changes in the adult worm burden. These changes in worm burden are related to the costs of reduced body weight and fleece weight and the costs of increased mortality. The economic values for FEC varied from -0.02 NZ\$ to -0.03 NZ\$ per 1% increase in flock average FEC in ewes and in lamb, respectively, for one year in New Zealand, and from -0.13 A\$ to 0.04 A\$ per adult and per lamb, respectively, in summer-dry areas in Australia for one year. They are low, and even if comparison is hazardous due to differences in currencies and breeding systems, they seem closer to the economic value of PCV found in Creole goats than to the maximum economic value attributed to FEC in Creole goats.

In this study, the economic value of FEC is derived considering an extreme situation, when parasites are resistant

to anthelmintics and when infection levels are not controlled by any alternative method. This weighting is a maximum value and does not reflect the economic value at the present time. Nevertheless, it provides a good insight into the long-term benefits of parasite resistance when anthelmintics will no longer be effective. Selection on reduced FEC will become increasingly more important as the drenching frequency reduces when shifting from a preventive to a curative approach. Analysis of the database on production traits and FEC for Creole goats during normal years (when parasites were controlled by anthelmintics) did not show any clear relationship between FEC and production traits. The effects of change in pasture contamination have not yet been evaluated for Creole goats, but it seems that an increase of FEC of one unit during the normal period does not lead to any change in productivity. The high weighting of FEC is therefore the result of a non-linear phenomenon, when a sudden increase in FEC results in considerable production loss.

The economic value of PCV only accounts for the costs of anthelmintic treatments that are still relatively inexpensive, which explains the low value obtained. The costs and performance recording of PCV may be considered more expensive than the potential benefit of improving this trait. However, this value does not take into account the potential gain in productivity that could be achieved from an improvement in animal PCV.

Transferring scientific knowledge into farming practices

The two economic values for resistance and resilience raise important issues. Which point of view should be favoured (short or long term)? How could selection on these traits be implemented on-farm? Farmers are not fully aware of the importance of resistance to parasites (Gunia *et al.*, 2010) and rely strongly on anthelmintics. If their awareness of the anthelmintics issue is not improved, they will probably not want to invest in selection on resistance or resilience.

A preliminary step could be to attribute weighting to FEC according to the desired gain method, to aim at maintaining this trait without degrading it. At first, PCV could be retained in the breeding objective, even without being recorded. Simulation will be run to ensure that this trait does not deteriorate.

An easier way, perhaps more suitable for farmers, could be to select animals on Famacha[®] mark (measurement of the ocular conjunctivae colour). This trait is moderately heritable (Riley and Van Wyk, 2009), very easy to record and highly correlated with PCV. If the Famacha[®] method is widely adopted by farmers (presently, this method is still new and has only been adopted by a few), it would probably make more sense to farmers to select on a directly visible trait. This trait could be used as an indicator trait for PCV or could even replace PCV in the breeding objective. However, further research would be required to study the feasibility of selection including this new trait in Creole goats.

Selection of Creole goats for resistance and resilience is feasible and highly desirable. Therefore, FEC and PCV should be included in the breeding goal and their weights should be

high enough to prevent deterioration of these traits. In the future, selection on Famacha[®] mark could be implemented to encourage farmers in appropriating and exploiting our findings.

Conclusion

Our study presents a balanced breeding goal for production and reproduction as well as parasite resistance and resilience traits. Defining such a breeding goal is a crucial step in setting up a breeding programme for Creole goats. Different alternatives are proposed for the integration of resistance and resilience. With further research, selection on both resistance and resilience can be implemented in Guadeloupe and will pave the way for a tropically adaptative selective breeding of goats.

Acknowledgements

We acknowledge 'la Région Guadeloupe' and the European Community (FEOGA) for the financial support provided, Donald White (ABIES doctoral school) for the English proofreading, as well as Régis Alexandre (Chambre d'Agriculture), Rangit Manicom (Iguavie), Marylene Madassamy (Cabricoop), Stephan Asselin de Beauville (Iguavie) and Jérôme Fleury (INRA) for their information and advice about field data, Harry Archimède and Daniel Sauvart (INRA) for the help on feeding equations, Maurice Mahieu (INRA) and Jean Marie-Felix Cherdieu d'Alexis (Veterinarian) for information about anthelmintics and integrated methods for parasite control, Fabien Stark (EPLEFPA de la Guadeloupe) and all INRA-URZ colleagues for the relevant advice and ideas given.

References

Agreste. 2006. Effectifs d'animaux, Guadeloupe. Retrieved July 10, 2011, from <http://www.agreste.agriculture.gouv.fr/IMG/pdf/R97108D06.pdf>

Agreste. 2009. Mémento agricole, Guadeloupe. Retrieved July 10, 2011, from http://www.agreste.agriculture.gouv.fr/IMG/pdf_D97110C02.pdf

Alexandre G, Aumont G, Fleury J, Mainaud JC and Kandassamy T 1997. Zootechnical performances of Creole goats in Guadeloupe (French West Indies). A twenty-year survey in an experimental farm. INRA Productions Animales 10, 7–20.

Alexandre G, Asselin de Beauville S, Shitalou E and Zebus MF 2008. An overview of the goat meat sector in Guadeloupe: conditions of production, consumer preferences, cultural functions and economic implications. Livestock Research for Rural Development 20, Article 14. Retrieved August 1, 2011, from <http://www.lrrd.org/lrrd20/1/alex20014.htm>

Amer PR, McEwan JC, Dodds KG and Davis GH 1999a. Economic values for ewe prolificacy and lamb survival in New Zealand sheep. Livestock Production Science 58, 75–90.

Amer PR, Woolaston RR, Eady SJ and McEwan JC 1999b. Economic values for sheep internal parasite resistance traits in New Zealand and Australia. Proceedings of the Association for the Advancement of Animal Breeding and Genetics 13, 504–507.

Aumont G, Pouillot R, Simon R, Hostache G, Varo H and Barré N 1997. Digestive parasitism of small ruminants in the French West Indies. INRA Productions Animales 10, 79–89.

Baker RL and Gray GD 2004. Appropriate breeds and breeding schemes for sheep and goats in the tropics. In Worm Control for Small Ruminants in Tropical Asia (ed. RA Sani, GD Gray and RL Baker), Monograph No. 113, pp. 63–75. Australian Centre for International Agriculture Research (ACIAR), Canberra, Australia.

Baker RL, Audho JO, Aduda EO and Thorpe W 2001. Genetic resistance to gastro-intestinal nematode parasites in Galla and Small East African goats in the sub-humid tropics. Animal Science 73, 61–70.

Bath GF, Malan FS and Van Wyk JA 1996. The "FAMACHA[®]" Ovine Anaemia Guide to assist with the control of haemonchosis. Proceedings of the 7th Annual Congress of the Livestock Health and Production Group of the South African Veterinary Association, Port Elizabeth, p. 5.

Bett RC, Kosgey IS, Bebe BO and Kahi AK 2007. Breeding goals for the Kenya Dual Purpose goat. I. Model development and application to smallholder production systems. Tropical Animal Health and Production 39, 477–492.

Bishop SC and Stear MJ 2003. Modeling of host genetics and resistance to infectious diseases: understanding and controlling nematode infections. Veterinary Parasitology 115, 147–166.

Bishop SC and Morris CA 2007. Genetics of disease resistance in sheep and goats. Small Ruminant Research 70, 48–59.

Bisset SA and Morris CA 1996. Feasibility and implications of breeding sheep for resilience to nematode challenge. International Journal for Parasitology 26, 857–868.

Byrne TJ, Amer PR, Fennessy PF, Cromie AR, Keady TWJ, Hanrahan JP, McHugh MP and Wickham BW 2010. Breeding objectives for sheep in Ireland: a bio-economic approach. Livestock Science 132, 135–144.

Chevalier M 2001. Restauration de la sensibilité au Levamisole de *Trichostrongylus colubriformis* par substitution de population parasitaire dans un élevage caprin de Guadeloupe F.W.I. Thèse pour le diplôme d'Etat de Docteur Vétérinaire, Ecole Nationale Vétérinaire de Nantes, France.

Eady SJ, Woolaston RR and Barger IA 2003. Comparison of genetic and nongenetic strategies for control of gastrointestinal nematodes of sheep. Livestock Production Science 81, 11–23.

Gibson JP and Kennedy BW 1990. The use of constrained selection indexes in breeding for economic merit. Theoretical and Applied Genetics 80, 801–805.

Gicheha MG, Kosgey IS, Bebe BO and Kahi AK 2005. Economic values for resistance to gastrointestinal helminths in meat sheep in Kenya. Journal of Animal Breeding and Genetics 122, 165–171.

Greeff JC, Safari E, Fogarty NM, Hopkins DL, Brien FD, Atkins KD, Mortimer SI and van der Werf JHJ 2008. Genetic parameters for carcase and meat quality traits and their relationships to liveweight and wool production in hogget Merino rams. Journal of Animal Breeding and Genetics 125, 205–215.

Greer AW 2008. Trade-offs and benefits: implications of promoting a strong immunity to gastrointestinal parasites in sheep. Parasite Immunology 30, 123–132.

Groen AF 1989. Cattle breeding goals and production circumstances. PhD, Department of Farm Management and Department of Animal Breeding, Wageningen Agricultural University, Wageningen, The Netherlands.

Gunia M, Phocas F, Arquet R, Alexandre G and Mandonnet N 2011. Genetic parameters for body weight, reproduction and parasite resistance traits in the Creole goat. Journal of Animal Science 89, 3443–3451.

Gunia M, Mandonnet N, Arquet R, de la Chevrotière C, Naves M, Mahieu M and Alexandre G 2010. Production systems of Creole goat and their implications for a breeding programme. Animal 4, 2099–2115.

Hazel LN 1943. The genetic basis for constructing selection indexes. Genetics 28, 476–490.

Itoh Y and Yamada Y 1987. Comparisons of selection indexes achieving predetermined proportional gains. Genetics Selection Evolution 19, 69–81.

Jackson F and Miller J 2006. Alternative approaches to control – Quo vadit? Veterinary Parasitology 139, 371–384.

Kahn LP, Knox MR, Walkden-Brown SW and Lea JM 2003. Regulation of the resistance to nematode parasites of single- and twin-bearing Merino ewes through nutrition and genetic selection. Veterinary Parasitology 114, 15–31.

Kaplan RM 2004. Drug resistance in nematodes of veterinary importance: a status report. Trends in Parasitology 20, 477–481.

Kemper KE, Elwin RL, Bishop SC, Goddard ME and Woolaston RR 2009. *Haemonchus contortus* and *Trichostrongylus colubriformis* did not adapt to long-term exposure to sheep that were genetically resistant or susceptible to nematode infections. International Journal for Parasitology 39, 607–614.

Kominakis A and Theodoropoulos G 1999. Selection of dairy sheep in Greece for parasitic nematode resistance: defining the aggregate genotype and evaluating selection schemes. Animal Science 69, 535–542.

Kosgey IS, van Arendonk JAM and Baker RL 2003. Economic values for traits of meat sheep in medium to high production potential areas of the tropics. Small Ruminant Research 50, 187–202.

Economic values of body weight, reproduction and parasite resistance traits for a Creole goat breeding goal

- Legarra A, Ramon M, Ugarte E and Perez-Guzman MD 2007. Economic weights of fertility, prolificacy, milk yield and longevity in dairy sheep. *Animal* 1, 193–203.
- Love S 1999. Residues and worm control in sheep and goats. Wool and sheepmeat services. Annual Conference 1999, University of New England, Armidale, NSW, Australia, 15–16 November, 1999, pp. 145–147.
- Luo J, Goetsch AL, Nsahlai IV, Moore JE, Galyean ML, Johnson ZB, Sahlu T, Ferrell CL and Owens FN 2004. Voluntary feed intake by lactating, Angora, growing and mature goats. *Small Ruminant Research* 53, 357–378.
- Mahieu M, Arquet R, Kandassamy T, Mandonnet N and Hoste H 2007. Evaluation of targeted drenching using Famacha[®] method in Creole goat: reduction of anthelmintic use, and effects on kid production and pasture contamination. *Veterinary Parasitology* 146, 135–147.
- Mahieu M, Archimede H, Fleury J, Mandonnet N and Alexandre G 2008. Intensive grazing system for small ruminants in the Tropics: The French West Indies experience and perspectives. *Small Ruminant Research* 77, 195–207.
- Mahieu M, Arquet R, Fleury J, Coppy O, Marie-Magdeleine C, Boval M, Archimede H, Alexandre G, Bambou JC and Mandonnet N 2009. Integrated control of gastrointestinal parasitism in grazing small ruminants in the humid tropics. 16^{emes} Rencontres autour des Recherches sur les Ruminants, Paris, les 2 et 3 Decembre 2009, pp. 265–268.
- Mandonnet N, Aumont G, Fleury J, Arquet R, Varo H, Gruner L, Bouix J and Khang JVT 2001. Assessment of genetic variability of resistance to gastrointestinal nematode parasites in Creole goats in the humid tropics. *Journal of Animal Science* 79, 1706–1712.
- Mandonnet N, Ducrocq V, Arquet R and Aumont G 2003. Mortality of Creole kids during infection with gastrointestinal strongyles: a survival analysis. *Journal of Animal Science* 81, 2401–2408.
- Mandonnet N, Bachand A, Mahieu A, Arquet R, Baudron F, Abinne-Molza L, Varo H and Aumont G 2005. Impact on productivity of peri-parturient rise in fecal egg counts in Creole goats in the humid tropics. *Veterinary Parasitology* 134, 249–259.
- Mandonnet N, Menendez-Buxadera A, Arquet R, Mahieu M, Bachand M and Aumont G 2006. Genetic variability in resistance to gastro-intestinal strongyles during early lactation in Creole goats. *Animal Science* 82, 283–287.
- Moav R and Hill WG 1966. Specialised sire and dam lines. IV. Selections within lines. *Animal Production* 8, 375–390.
- Perry BD, Randolph TF, McDermott JJ, Sones KR and Thornton PK 2002. Investing in animal health research to alleviate poverty. ILRI (International Livestock Research Institute), Nairobi, Kenya.
- Phocas F, Bloch C, Chapelle P, Becherel F, Renand G and Menissier F 1998. Developing a breeding objective for a French purebred beef cattle selection programme. *Livestock Production Science* 57, 49–65.
- Ponzoni RW and Newman S 1989. Developing breeding objectives for Australian beef-cattle production. *Animal Production* 49, 35–47.
- Riley DG and Van Wyk JA 2009. Genetic parameters for FAMACHA (c) score and related traits for host resistance/resilience and production at differing severities of worm challenge in a Merino flock in South Africa. *Veterinary Parasitology* 164, 44–52.
- Sahlu T, Goetsch AL, Luo J, Nsahlai IV, Moore JE, Galyean ML, Owens FN, Ferrell CL and Johnson ZB 2004. Nutrient requirements of goats: developed equations, other considerations and future research to improve them. *Small Ruminant Research* 53, 191–219.
- Sauvant D, Giger-Reverdin S and Meschy F 2007. Alimentation des caprins. Alimentation des bovins, ovins et caprins. Editions Quae-INRA, Versailles, France, pp. 139–152.
- Shrestha JNB and Fahmy MH 2007. Breeding goats for meat production 3. Selection and breeding strategies. *Small Ruminant Research* 67, 113–125.
- Smith C, James JW and Brascamp EW 1986. On the derivation of economic weights in livestock improvement. *Animal Production* 43, 545–551.
- Torres-Acosta JFJ and Hoste H 2008. Alternative or improved methods to limit gastro-intestinal parasitism in grazing sheep and goats. *Small Ruminant Research* 77, 159–173.
- Vagenas D, Jackson F, Russel AJF, Merchant M, Wright IA and Bishop SC 2002. Genetic control of resistance to gastro-intestinal parasites in crossbred cashmere-producing goats: responses to selection, genetic parameters and relationships with production traits. *Animal Science* 74, 199–208.
- Wolfova M, Wolf J and Milerski M 2009. Calculating economic values for growth and functional traits in non-dairy sheep. *Journal of Animal Breeding and Genetics* 126, 480–491.
- Woolaston RR 1994. Preliminary evaluation of strategies to breed Merinos for resistance to roundworms. Proceedings of the 5th World Congress on Genetics Applied to Livestock Production, Guelph, Canada, pp. 281–284.
- Woolaston RR and Baker RL 1996. Prospects of breeding small ruminants for resistance to internal parasites. *International Journal for Parasitology* 26, 845–855.