

1Table S1. Theoretical models comparing at least two strategies for delaying the evolution of resistance to two or more pesticides or
 2drugs. C = *Combination*, PA = *Periodic application*, M = *Mosaic* and RA = *Responsive alternation*.

Pesticide	Reference	Strategies	Relative efficacy for delaying resistance
or drug		compared	
Insecticide			
s	[1]	C ^a , PA, RA	C > PA > RA
	[2]	C, M, RA	C > M > RA
	[3]	C, PA, RA	C > PA = RA
	[4]	C ^a , PA	C > PA
	[5]	C, PA, M	C > PA > M
	[6]	C, RA	C > RA
	[7]	PA, M, RA	PA = M = RA
	[8]	C, PA, M	C > PA > M
	[9]	C, RA	conditional ^b
<i>Bacillus</i>	[10]	C ^a , PA, M, RA	conditional ^b for C but PA = M > RA
<i>thuringien</i>	[11]	C, PA	C > PA
<i>sis toxins</i>	[12]	C, RA	C ≥ RA

Herbicides	[13]	C, PA, RA	$C > PA = RA$
	[14]	C, PA	$C > PA$
	[15]	C, PA, RA	$C > PA > RA$
Antibiotics	[16]	PA, M	$M \geq PA$
	[17]	C, PA, M	$C > M \geq PA$
	[18]	C, RA	$C > RA$
	[19]	PA, M	$M > PA$
	[20]	PA, M	conditional ^b
Other	[21]	C, PA, M	$C > M \geq PA$
	[22]	C, PA	$C > PA$
pesticides	[23]	C, PA, RA	$C > PA = RA$
	[24]	C, PA, M	$PA = M > C$
or drugs	[25]	C, RA	$C > RA$
	[26]	PA, M	$M \geq PA$
Unspecifie	[27]	C, RA	conditional ^b
	[28]	C, RA	$C \geq RA$
d			
pesticides	[29]	C, PA	$C \geq PA$
or drugs			

3^a *Combination* is considered as a *half-dose Combination* of the pesticides or drugs; *full-dose*

4*Combination* is compared with *half-dose Combination* in [10].

5^b The ranking of the strategies depends on the setting for one or several input or output parameters.

Table S2. Empirical studies comparing at least two strategies for delaying the evolution of resistance to two or more pesticides or drugs. C = Combination, PA = Periodic application, M = Mosaic and RA = Responsive alternation.

Pesticide or drug	Reference	Strategies compared	Relative efficacy for delaying resistance	Type of study	Species	Pitfalls ^c		Specific settings ^d	
						PE	IC	HF	AR
Insecticide						√	√	√	√
s	[30]	C, PA, RA	C = PA > RA	Laboratory	<i>Blatella germanica</i>				
	[31]	C ^a , PA, RA	C = PA = RA	Field	<i>Bemisia tabaci</i>	√	√	√	
					<i>Culex</i>	√	√	√	
	[32]	PA, RA	PA > RA	Laboratory	<i>quinquefasciatus</i>				
	[33]	C ^a , PA, RA	PA > C > RA	Field	<i>Scirtothrips citri</i>	√	√	√	√
	[34]	C, PA, RA	C > PA > RA	Laboratory	<i>Musca domestica</i>	√	√	√	√
	[35]	PA, RA	PA > RA	Field	<i>Musca domestica</i>	√	√	√	
	[36]	C, PA, RA	C = PA > RA	Laboratory	<i>Haematobia irritans</i>	√	√	√	√

	[37]	M, RA	M > RA	Laboratory	<i>Musca domestica</i>		√	√	
	[38]	C, PA, RA	C = PA = RA	Laboratory	<i>Musca domestica</i>	√	√	√	√
	[39]	C, PA, RA	C = PA > RA	Laboratory	<i>Bemisia argentifolii</i>	√	√	√	√
<i>Bacillus</i>				Greenhouse				√	
<i>thuringien</i>	[40]	PA, M	PA > M	e	<i>Plutella xylostella</i>				
<i>sis toxins</i>				Greenhouse		√	√		
	[41]	C, RA, M	C > RA > M	e	<i>Plutella xylostella</i>				
	[42]	C, RA	C > RA	Laboratory	<i>Plodia interpunctella</i>	√	√	√	√
					<i>Culex</i>	√	√		√
	[43]	C, RA	C > RA	Laboratory	<i>quinquefasciatus</i>				
Herbicides	[44]	C, PA	C > PA	Field	<i>Thlaspi arvense</i>			√	√
					<i>Pseudomonas</i>	√		√	
Antibiotics	[45]	PA, M	PA > M	ICU ^b	<i>aeruginosa</i>				
					<i>Acinetobacter</i>	√	√	√	
	[46]	PA, M	M > PA	ICU ^b	<i>baumannii</i>				

8^a *Combination* is considered as a *half-dose Combination* of the pesticides or drugs.

9^b Intensive care unit.

10^c Pitfalls: PE = problems in experimental design (lack of replicates, no randomization, confounding factors etc.) and IC = incorrect

11 comparisons.

12^d Specific settings: HF = high frequencies of resistance before selection and AR = absence of refuges when testing *Combination*.

13References

- 141 Argentine, J. *et al.* (1994) Computer simulation of insecticide resistance management
15 strategies for control of Colorado potato beetle (Coleoptera: Chrysomelidae). *J. Agr.*
16 *Entomol.* 11, 137-155
- 172 Curtis, C.F. (1985) Theoretical models of the use of insecticide mixtures for the
18 management of resistance. *Bull. Entomol. Res.* 75, 259-265
- 193 Curtis, C.F. *et al.* (1993) Are there effective resistance management strategies for vectors
20 of human disease? *Biol. J. Linn. Soc.* 48, 3-18
- 214 Knippling, E.F. and Klassen, W. (1984) Influence of insecticide use patterns on the
22 development of resistance to insecticides - a theoretical study. *Southw. Entomol.* 9, 351-
23 368
- 245 Lenormand, T. and Raymond, M. (1998) Resistance management: the stable zone
25 strategy. *Proc. R. Soc. Lond. B* 265, 1985-1990
- 266 Mani, G.S. (1985) Evolution of resistance in the presence of two insecticides. *Genetics*
27 109, 761-783
- 287 Mani, G.S. (1989) Evolution of resistance with sequential application of insecticides in
29 time and space. *Proc. R. Soc. Lond. B* 238, 245-276
- 308 Roush, R.T. (1989) Designing resistance management programs - how can you choose.
31 *Pestic. Sci.* 26, 423-441

- 329 Tabashnik, B.E. and McGaughey, W.H. (1994) Resistance risk assessment for single and
33 multiple insecticides - responses of Indian meal moth (Lepidoptera, Pyralidae) to *Bacillus*
34 *thuringiensis*. *J. Econ. Entomol.* 87, 834-841
- 3510 Caprio, M.A. (1998) Evaluating resistance management strategies for multiple toxins in
36 the presence of external refuges. *J. Econ. Entomol.* 91, 1021-1031
- 3711 Caprio, M.A. (2001) Source-sink dynamics between transgenic and non-transgenic
38 habitats and their role in the evolution of resistance. *J. Econ. Entomol.* 94, 698-705
- 3912 Roush, R.T. (1998) Two-toxin strategies for management of insecticidal transgenic crops:
40 can pyramiding succeed where pesticide mixtures have not? *Philos. Trans. R. Soc. Lond.*
41 *B* 353, 1777-1786
- 4213 Diggle, A.J. *et al.* (2003) Herbicides used in combination can reduce the probability of
43 herbicide resistance in finite weed populations. *Weed Res.* 43, 371-382
- 4414 Neve, P. *et al.* (2003) Simulating evolution of glyphosate resistance in *Lolium rigidum* II:
45 past, present and future glyphosate use in Australian cropping. *Weed Res.* 43, 418-427
- 4615 Powles, S.B. *et al.* (1997) Herbicide resistance: impact and management. In *Advances in*
47 *Agronomy*, 58, 57-93, Elsevier Academic Press Inc
- 4816 Bergstrom, C. *et al.* (2004) Ecological theory suggests that antimicrobial cycling will not
49 reduce antimicrobial resistance in hospitals. *Proc. Natl. Acad. Sci. USA* 101, 13285-13290

- 5017 Bonhoeffer, S. *et al.* (1997) Evaluating treatment protocols to prevent antibiotic
51 resistance. *Proc. Natl. Acad. Sci. USA* 94, 12106-12111
- 5218 D'Agata, E.M. *et al.* (2008) The impact of different antibiotic regimens on the emergence
53 of antimicrobial-resistant bacteria. *PLoS ONE* 3, e4036
- 5419 Lo, M. *et al.* (2002) Comparison of antimicrobial cycling and mixing using stochastic
55 mathematical models. In *42nd Interscience Conference on Antimicrobial Agents and*
56 *Chemotherapy*, American Society for Microbiology, Washington, DC, USA.
- 5720 Reluga, T.C. (2005) Simple models of antibiotic cycling. *Math. Med. Biol.* 22, 187-208
- 5821 Sun, H.R. *et al.* (2010) Qualitative analysis of models with different treatment protocols
59 to prevent antibiotic resistance. *Math. Biosci.* 227, 56-67
- 6022 Barnes, E. *et al.* (1995) Worm control and anthelmintic resistance: adventures with a
61 model. *Parasitol. today* 11, 56-63
- 6223 Smith, G. (1990) A mathematical model for the evolution of anthelmintic resistance in a
63 direct life cycle nematode parasite. *Int. J. Parasitol.* 20, 913-921
- 6424 Dobson, R.J. *et al.* (1987) A genetic model describing the evolution of levamisole
65 resistance in *Trichostrongylus colubriformis*, a nematode parasite of sheep. *J. Math. Appl.*
66 *Med. Biol.* 4, 279-293
- 6725 Curtis, C.F. and Otoo, L.N. (1986) A simple model of the build-up of resistance to
68 mixtures of anti-malarial drugs. *Trans. R. Soc. Trop. Med. Hyg.* 80, 889-892

- 6926 Boni, M. *et al.* (2008) Benefits of using multiple first-line therapies against malaria. *Proc.*
70 *Natl. Acad. Sci. USA* 105, 14216-14221
- 7127 Birch, C.P.D. and Shaw, M.W. (1997) When can reduced doses and pesticide mixtures
72 delay the build-up of pesticide resistance? A mathematical model. *J. Appl. Ecol.* 34, 1032-
73 1042
- 7428 Comins, H.N. (1986) Tactics for resistance management using multiple pesticides. *Agr.*
75 *Ecosys. Envir.* 16, 129-148
- 7629 Jaffe, K. *et al.* (1997) Dynamics of the emergence of genetic resistance to biocides among
77 asexual and sexual organisms. *J. Theor. Biol.* 188, 289-299
- 7830 Burden, G.S. *et al.* (1960) Development of chlordane and malathion resistance in the
79 German cockroach. *J. Econ. Entomol.* 53, 1138-1139
- 8031 Castle, S.J. *et al.* (2002) Field evaluation of different insecticide use strategies as
81 resistance management and control tactics for *Bemisia tabaci* (Hemiptera: Aleyrodidae).
82 *Bull. Entomol. Res.* 92, 449-460
- 8332 Georghiou, G. (1983) Management of resistance in arthropods. In *Pest Resistance to*
84 *Pesticides* (Georghiou, G., and Saito, T., eds), pp. 769–792, Plenum
- 8533 Immaraju, J.A. *et al.* (1990) Field-evaluation of insecticide rotation and mixtures as
86 strategies for citrus thrips (Thysanoptera, Thripidae) resistance management in California.
87 *J. Econ. Entomol.* 83, 306-314

- 8834 MacDonald, R.S. *et al.* (1983) Effect of 4 spray regimes on the development of
89 permethrin and dichlorvos resistance, in the laboratory, by the housefly (Diptera,
90 Muscidae). *J. Econ. Entomol.* 76, 417-422
- 9135 MacDonald, R.S. *et al.* (1983) Development of resistance to permethrin and dichlorvos by
92 the housefly (Diptera, Muscidae) following continuous and alternating insecticide use on
93 4 farms. *Can. Entomol.* 115, 1555-1561
- 9436 McKenzie, C.L. and Byford, R.L. (1993) Continuous, alternating, and mixed insecticides
95 affect development of resistance in the horn fly (Diptera, Muscidae). *J. Econ. Entomol.*
96 86, 1040-1048
- 9737 Pimentel, D. and Bellotti, A.C. (1976) Parasite-host population systems and genetic
98 stability. *Am. Nat.* 110, 877-888
- 9938 Pimentel, D. and Burgess, M. (1985) Effects of single versus combinations of insecticides
100 on the development of resistance. *Environ. Entomol.* 14, 582-589
- 10139 Prabhaker, N. *et al.* (1998) Evaluation of insecticide rotations and mixtures as resistance
102 management strategies for *Bemisia argentifolii* (Homoptera: Aleyrodidae). *J. Econ.*
103 *Entomol.* 91, 820-826
- 10440 Zhao, J.Z. *et al.* (2010) Testing insecticide resistance management strategies: mosaic
105 versus rotations. *Pest Manag. Sci.* 66, 1101-1105

- 10641 Zhao, J.Z. *et al.* (2003) Transgenic plants expressing two *Bacillus thuringiensis* toxins
 107 delay insect resistance evolution. *Nat. Biotechnol.* 21, 1493-1497
- 10842 McGaughey, W.H. and Johnson, D.E. (1992) Indian meal moth (Lepidoptera, Pyralidae)
 109 resistance to different strains and mixtures of *Bacillus thuringiensis*. *J. Econ. Entomol.* 85,
 110 1594-1600
- 11143 Georghiou, G.P. and Wirth, M.C. (1997) Influence of exposure to single versus multiple
 112 toxins of *Bacillus thuringiensis* subsp. *israelensis* on development of resistance in the
 113 mosquito *Culex quinquefasciatus* (Diptera: Culicidae). *Appl. Env. Microbiol.* 63, 1095-
 114 1101
- 11544 Beckie, H. and Reboud, X. (2009) Selecting for weed resistance: herbicide rotation and
 116 mixture. *Weed Technol.* 23, 363-370
- 11745 Martinez, J.A. *et al.* (2006) Comparison of antimicrobial cycling and mixing strategies in
 118 two medical intensive care units. *Crit. Care Med.* 34, 329-336
- 11946 Sandiumenge, A. *et al.* (2006) Impact of diversity of antibiotic use on the development of
 120 antimicrobial resistance. *J. Antimicrob. Chemother.* 57, 1197-1204