



# Acclimation of leaf nitrogen to vertical light gradient at anthesis in wheat is a whole-plant process that scales with the size of the canopy

Delphine Moreau, Vincent Allard, Oorbessy Gaju, Jacques Le Gouis, M. John Foulkes, Pierre Martres

## ► To cite this version:

Delphine Moreau, Vincent Allard, Oorbessy Gaju, Jacques Le Gouis, M. John Foulkes, et al.. Acclimation of leaf nitrogen to vertical light gradient at anthesis in wheat is a whole-plant process that scales with the size of the canopy. *Plant Physiology*, 2012, 160 (3), pp.1479-1490. 10.1104/pp.112.199935 . hal-01001606

**HAL Id: hal-01001606**

**<https://hal.science/hal-01001606>**

Submitted on 22 Jan 2021

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# Acclimation of Leaf Nitrogen to Vertical Light Gradient at Anthesis in Wheat Is a Whole-Plant Process That Scales with the Size of the Canopy<sup>1[W][OA]</sup>

Delphine Moreau<sup>2</sup>, Vincent Allard, Oorbessy Gaju, Jacques Le Gouis, M. John Foulkes, and Pierre Martre\*

Institut National de la Recherche Agronomique, Unité Mixte Recherche 1095 Genetics, Diversity, and Ecophysiology of Cereals, F-63039 Clermont-Ferrand, cedex 02, France (D.M., V.A., J.L.G., P.M.); Blaise Pascal University, Unité Mixte Recherche 1095 Genetics, Diversity, and Ecophysiology of Cereals, F-63170 Aubiere, France (D.M., V.A., J.L.G., P.M.); and Division of Plant and Crop Sciences, School of Biosciences, University of Nottingham, Leicestershire LE12 5RD, United Kingdom (O.G., M.J.F.)

Vertical leaf nitrogen (N) gradient within a canopy is classically considered as a key adaptation to the local light environment that would tend to maximize canopy photosynthesis. We studied the vertical leaf N gradient with respect to the light gradient for wheat (*Triticum aestivum*) canopies with the aims of quantifying its modulation by crop N status and genetic variability and analyzing its ecophysiological determinants. The vertical distribution of leaf N and light was analyzed at anthesis for 16 cultivars grown in the field in two consecutive seasons under two levels of N. The N extinction coefficient with respect to light ( $b$ ) varied with N supply and cultivar. Interestingly, a scaling relationship was observed between  $b$  and the size of the canopy for all the cultivars in the different environmental conditions. The scaling coefficient of the  $b$ -green area index relationship differed among cultivars, suggesting that cultivars could be more or less adapted to low-productivity environments. We conclude that the acclimation of the leaf N gradient to the light gradient is a whole-plant process that depends on canopy size. This study demonstrates that modeling leaf N distribution and canopy expansion based on the assumption that leaf N distribution parallels that of the light is inappropriate. We provide a robust relationship accounting for vertical leaf N gradient with respect to vertical light gradient as a function of canopy size.

In cereals, as in many crop species, nitrogen (N) nutrition is a major determinant in the elaboration of grain yield and quality (Lemaire and Millard, 1999; Lawlor, 2002; Hikosaka, 2005). N is involved in both meristematic and photosynthetic activities, with consequences on plant architecture and carbon acquisition and in fine on grain yield and protein concentration. Beside the total amount of N absorbed by the crop, the allocation of N among plant organs plays a key role in determining crop productivity and quality (Grindlay, 1997; Dreccer et al., 1998; Hikosaka, 2005).

Light interception and leaf N content are the two main factors governing carbon assimilation at the leaf scale (Evans, 1989). For various species, both light and leaf N attenuate with cumulative leaf area index counted from the top of the canopy (Field, 1983; Hirose and Werger, 1987). Leaf N vertical gradients have been regarded as an adaptive response to the local light environment, maximizing canopy photosynthesis and N utilization efficiency (Hirose and Werger, 1987; Hikosaka et al., 1994; Drouet and Bonhomme, 1999), as N is largely contained in the assimilatory enzyme Rubisco. Theoretical studies indicated that leaf N maximizes canopy photosynthesis when it parallels the light gradient (i.e. when the light [ $K_L$ ] and N [ $K_N$ ] extinction coefficients are equal), considering that the leaf N gradient is “optimal” in accordance with the “optimization theory” (Field, 1983; Hirose and Werger, 1987; Anten et al., 1995b).

Factors other than the photosynthetic photon flux density (PPFD) might be responsible for the observed leaf N distribution. For instance, the acropetal gradients of leaf age (Hikosaka et al., 1994; Hikosaka, 2005) and light composition (Rousseaux et al., 1999) are known to strengthen the leaf N gradient. However, the impact of each of these factors has been shown to be much less than that of the PPFD gradient (Werger and Hirose, 1991; Pons and de Jong-van Berkel, 2004), although for the grass species *Brachypodium pinnatum* other factors than light might be involved (Pons et al., 1993). At the

<sup>1</sup> This work was supported by the Biotechnology and Biological Sciences Research Council of the United Kingdom and the French National Institute for Agricultural Research through the “NUE traits” project (grant no. IN-BB-06).

<sup>2</sup> Present address: Institut National de la Recherche Agronomique, Unité Mixte Recherche 1347 Agroécologie, F-21065 Dijon, France.

\* Corresponding author; e-mail pierre.martre@clermont.inra.fr.

The author responsible for distribution of materials integral to the findings presented in this article in accordance with the policy described in the Instructions for Authors (www.plantphysiol.org) is: Pierre Martre (pierre.martre@clermont.inra.fr).

[W] The online version of this article contains Web-only data.

[OA] Open Access articles can be viewed online without a subscription.

www.plantphysiol.org/cgi/doi/10.1104/pp.112.199935

molecular level, the process could be driven by the import of compounds such as cytokinins transported in the transpiration stream (Pons et al., 2001; Boonman et al., 2007). Although the actual N distribution usually follows the light gradient, in all studies it is less steep than the calculated optimal N profile maximizing canopy photosynthesis (Pons et al., 1989; Yin et al., 2003). Possible reasons for this discrepancy have been discussed in detail by Kull (2002). Sink-source relations and in particular the demand for N could modulate the light-leaf N relationship (Dreccer et al., 1998), but conflicting results have been reported regarding the effect of N availability on the light-leaf N relationship. While some authors found no effect of N availability (Sinclair and Shiraiwa, 1993; Milroy et al., 2001), others found that the N gradient relative to light (i.e.  $K_L/K_N$ ) was steeper under low N (Hikosaka et al., 1994; Grindlay et al., 1995; Lötscher et al., 2003) or that the response of the light-leaf N relationship to N availability depended on the developmental stage (Dreccer et al., 2000). Interspecific differences in the light-leaf N relationship have also been reported and were related to differences in phenotypic plasticity (Aerts, 1996) or plant architecture (leaf stature and branching pattern; Anten et al., 1995a; Lötscher et al., 2003).

Since canopy photosynthesis is dependent upon the leaf N gradient, it has been suggested that the pattern of leaf N distribution could be responsible for part of the genetic variability associated with the negative correlation between grain yield and protein concentration reported for various crop species (Dreccer et al., 1998). In wheat (*Triticum aestivum*), N accumulated before anthesis contributes 30% to 70% of grain N (Mi et al., 2000; Kichey et al., 2007). The efficiency of N translocation from the lower to the upper leaves may increase with the steepness of the N gradient, with only a negligible effect on canopy carbon gain integrated over the whole grain-filling period. This hypothesis is consistent with experimental studies based on a range of genotypes showing that, at a given grain yield level, grain protein concentration is positively related to the efficiency of N translocation either from the lower to the upper leaves (Wang et al., 2005) or from the leaves to the grains (Monaghan et al., 2001; Jukanti and Fischer, 2008). Only a few studies have investigated the intraspecific variability of the light-N relationship at the intraspecific level (Shiraiwa and Sinclair, 1993; Bindraban, 1999; Bertheloot et al., 2008; van Oosterom et al., 2010). For wheat, published analyses of the genetic variability of the light-leaf N relationship were limited to only two to five genotypes, and no genetic differences were found (Bindraban, 1999; Bertheloot et al., 2008).

This paper focuses on the genetic variability of the vertical leaf N gradient with respect to light for wheat. Three main issues were investigated. What is the effect of N supply on the vertical distribution of leaf N? Does the distribution of leaf N with respect to light differ among genotypes? If the adjustment of leaf N to the light gradient varies with both the genotype and N supply, could this genetic and environmental variability

have a unique ecophysiological determinant (driving variable)?

These questions were addressed using 16 genotypes (Supplemental Table S1) covering a wide range of variation for N use efficiency (i.e. grain dry mass yield per unit of available mineral N from the soil and fertilizer), for grain protein concentration (Le Gouis et al., 2000; Foulkes et al., 2006; Gaju et al., 2011) and for the deviation from the negative correlation between grain yield and protein concentration (Oury et al., 2003). The 16 genotypes were grown in the field under two conditions of N supply (N<sup>-</sup> and N<sup>+</sup> for low- and high-N treatments, respectively) in order to modulate crop N status at Clermont-Ferrand (CF) in France in two consecutive seasons (experiments CF07 and CF08). In addition, four of the 16 cultivars representing the variability observed for N utilization and N uptake efficiency were grown in the field under two conditions of N supply at Sutton Bonington (SB) in the United Kingdom in one season (experiment SB07). The distribution of leaf N was analyzed at anthesis. The first reason for this is that the distribution of both light and leaf N within the canopy is relatively stable from this phenological stage until almost the end of grain filling (Bertheloot et al., 2008). Whereas the canopy green area index (GAI) decreases dramatically during the grain-filling period, the structure of the canopy affecting light interception does not change significantly during that period. Both the vertical light and N distributions down the canopy are unchanged during most of the grain-filling period; therefore, the  $K_N$ -to- $K_L$  ratio is constant during that period (Bertheloot et al., 2008). Similarly, Archontoulis et al. (2011) showed that  $K_N$ -to- $K_L$  ratio is not modified during the vegetative and reproductive stages for field-grown sunflower (*Helianthus annuus*) crops. Therefore, as most of the final grain yield results from carbon assimilated after anthesis (Bidingier et al., 1977; Gebbing and Schnyder, 1999), the N distribution at anthesis is very relevant in terms of carbon assimilation and grain yield in wheat. A second reason is that the number and potential size of grains are determined around anthesis, which therefore appears as a critical stage in the formation of grain yield. A better understanding of the ecophysiological determinants of leaf N gradient at this phenological stage could consequently be crucial for improving wheat productivity and quality (Dreccer et al., 1998).

## RESULTS

### Canopy Size and N Uptake at Anthesis

All the traits under study were affected by the year of experiment at CF (Table I). That is, lamina N and GAI were higher in 2008 than in 2007. Correspondingly, crops were denser and a larger percentage of the incident PPFD was intercepted by the canopy in 2008 (62% on average) than in 2007 (89% on average; Fig. 1). By contrast, average main shoot leaf lamina N mass per unit leaf area ( $N_{LA}^{sh}$ ) was lower in 2008 than in 2007

**Table I.** GAI, main shoot leaf lamina N, and average  $N_{LA}^{sh}$  for wheat

Crops were grown in the field at CF during the 2006–2007 (CF07) and 2007–2008 (CF08) growing seasons and at SB during the 2006–2007 growing season (SB07). For each experiment, the minimum, maximum, and average values across the 16 (at CF) or four (at SB) cultivars analyzed are reported. Data are means  $\pm$  SE for two treatments  $\times$  three independent replicates. *P* values of the main effects and interactions from split-plot ANOVAs at CF are reported.

Cultivars and Statistical Data	GAI			Lamina N			$N_{LA}^{sh}$		
	CF07	CF08	SB07	CF07	CF08	SB07	CF07	CF08	SB07
	$m^2$ green tissue $m^{-2}$ ground			$g$ N $m^{-2}$ ground			$g$ N $m^{-2}$ lamina		
Alchemy	3.17 $\pm$ 0.54	5.14 $\pm$ 1.48	–	2.43 $\pm$ 0.83	4.18 $\pm$ 2.39	–	1.90 $\pm$ 0.29	1.57 $\pm$ 0.39	–
Beaver	3.05 $\pm$ 0.59	5.98 $\pm$ 1.88	–	2.22 $\pm$ 1.05	4.60 $\pm$ 2.32	–	1.94 $\pm$ 0.44	1.59 $\pm$ 0.41	–
Consort	2.96 $\pm$ 0.62	5.78 $\pm$ 2.00	–	1.80 $\pm$ 0.83	4.64 $\pm$ 2.48	–	2.14 $\pm$ 0.44	1.59 $\pm$ 0.47	–
Paragon	3.62 $\pm$ 0.90	5.90 $\pm$ 1.59	–	2.03 $\pm$ 0.77	3.64 $\pm$ 1.68	–	1.71 $\pm$ 0.25	1.39 $\pm$ 0.45	–
Rialto	2.82 $\pm$ 0.51	4.51 $\pm$ 1.63	8.30 $\pm$ 1.74	1.80 $\pm$ 0.83	3.58 $\pm$ 1.85	8.62 $\pm$ 3.72	1.98 $\pm$ 0.35	1.77 $\pm$ 0.44	2.02 $\pm$ 0.43
Robigus	3.13 $\pm$ 0.62	5.90 $\pm$ 1.41	–	2.48 $\pm$ 0.76	4.10 $\pm$ 2.45	–	1.76 $\pm$ 0.33	1.39 $\pm$ 0.45	–
Savannah	2.96 $\pm$ 0.64	5.24 $\pm$ 1.50	8.99 $\pm$ 1.37	2.43 $\pm$ 1.09	4.21 $\pm$ 1.84	7.95 $\pm$ 2.97	2.16 $\pm$ 0.48	1.61 $\pm$ 0.31	1.82 $\pm$ 0.35
Soissons	3.54 $\pm$ 0.76	6.03 $\pm$ 1.61	–	2.42 $\pm$ 0.69	3.44 $\pm$ 1.26	–	1.97 $\pm$ 0.40	1.53 $\pm$ 0.29	–
Arche	3.51 $\pm$ 0.66	7.08 $\pm$ 1.45	–	2.24 $\pm$ 0.85	4.33 $\pm$ 1.80	–	1.48 $\pm$ 0.24	1.35 $\pm$ 0.31	–
CF9107	3.33 $\pm$ 0.66	6.06 $\pm$ 1.78	–	2.68 $\pm$ 0.90	3.91 $\pm$ 1.57	–	1.74 $\pm$ 0.20	1.47 $\pm$ 0.35	–
CF99102	3.14 $\pm$ 0.72	5.46 $\pm$ 1.39	9.41 $\pm$ 1.94	2.20 $\pm$ 0.52	3.47 $\pm$ 1.52	8.56 $\pm$ 3.40	1.74 $\pm$ 0.24	1.47 $\pm$ 0.36	1.74 $\pm$ 0.33
Perfector	3.23 $\pm$ 0.81	6.07 $\pm$ 1.89	8.42 $\pm$ 1.33	2.27 $\pm$ 1.00	4.47 $\pm$ 2.11	8.42 $\pm$ 2.56	1.71 $\pm$ 0.27	1.53 $\pm$ 0.33	2.01 $\pm$ 0.37
Quebon	2.69 $\pm$ 0.71	5.84 $\pm$ 1.64	–	1.78 $\pm$ 0.72	4.49 $\pm$ 2.04	–	1.91 $\pm$ 0.39	1.63 $\pm$ 0.38	–
Recital	3.01 $\pm$ 0.73	5.96 $\pm$ 1.28	–	1.96 $\pm$ 0.74	3.77 $\pm$ 1.39	–	1.49 $\pm$ 0.16	1.65 $\pm$ 0.37	–
Renan	3.55 $\pm$ 0.77	7.09 $\pm$ 1.17	–	2.46 $\pm$ 0.85	3.66 $\pm$ 0.93	–	1.67 $\pm$ 0.18	1.61 $\pm$ 0.42	–
Toisonador	3.22 $\pm$ 0.80	5.33 $\pm$ 1.42	–	2.21 $\pm$ 1.17	4.30 $\pm$ 1.87	–	1.75 $\pm$ 0.35	1.56 $\pm$ 0.35	–
Minimum	2.82	4.51	8.30	1.78	3.44	7.95	1.48	1.39	1.74
Maximum	3.55	7.08	9.41	2.68	4.60	8.62	2.16	1.77	2.02
Average	3.18	5.85	8.78	2.14	4.07	8.39	1.82	1.54	1.90
<i>P</i> values									
N	<0.001			<0.001			<0.001		
Year	<0.001			<0.001			<0.001		
Genotype	<0.001			0.003			<0.001		
N $\times$ year	<0.001			<0.001			<0.001		
N $\times$ genotype	0.470			<0.001			0.004		
Year $\times$ genotype	<0.001			<0.001			<0.001		
N $\times$ year $\times$ genotype	0.491			0.182			0.004		

(Table I). All traits varied with genotype (Table I). The traits were also greatly influenced by N treatments. For both years, greater N supply increased GAI, main shoot lamina N, and  $N_{LA}^{sh}$ . For all the traits under study, the year  $\times$  genotype and the year  $\times$  N interactions were significant, indicating that the differences between years were more or less marked according to the cultivar and to the N treatment. Except for GAI, the genotype  $\times$  N interaction was significant, indicating that the magnitude of the response to N treatment varied across cultivars.

Similar conclusions could be drawn from the study of the four cultivars in the three experiments (CF07, CF08, and SB07; Table I). Hence, for the three experiments, the two N treatments strongly modulated crop growth and N economy. Altogether, the different site/year/N treatment/cultivar combinations generated a wide range of variations of canopy structure and total canopy N at anthesis.

### Vertical Light Distribution

$K_L$  is commonly estimated from stratified light measurements through the canopy, but this method is very time consuming. Therefore, at CF07, we assessed whether  $K_L$  could be estimated without significant bias using a simplified method in which PPFD is measured above the canopy and at ground level only, assuming


that light decreases exponentially with cumulative GAI (Shearman et al., 2005). For all the N treatment/cultivar combinations, the decrease of light as a function of cumulative GAI was well described by an exponential function (all  $r^2 > 0.982$  and  $P < 0.001$ ; Fig. 1). The average measurement error variance of  $K_L$  was similar for the two methods, averaging 0.0041 ( $n = 12$ ).  $K_L$  values estimated using the two methods were tightly correlated ( $r = 0.949$ ,  $P < 0.001$ , degrees of freedom = 31; Supplemental Fig. S1). The slope of the relationship was not different from 1 ( $P = 0.65$ , slope = 0.929, 95% confidence interval = 0.828–1.044), and the intercept was not different from 0 ( $P = 0.21$ , intercept = 0.048). Since the two methods gave similar results, we used  $K_L$  values estimated with the simplified method in the three experiments.

$K_L$  differed neither between years nor between N treatments (Table II). However,  $K_L$  differed among cultivars, and the year  $\times$  genotype interaction was significant. Averaging across the N treatments,  $K_L$  ranged from 0.30 (cv Consort) to 0.52 (cv Recital) at CF07 and from 0.37 (cv Recital) to 0.54 (cv Rialto) at CF08 (Table II).

### Vertical Leaf N Distribution with Respect to Vertical Light Distribution

In all cases, the decline of leaf lamina N mass per unit leaf area ( $N_{LA}$ ) down the canopy was related to

**Figure 1.** Cumulative GAI counted from the top of the canopy ( $F$ ) versus relative PPFD ( $I/I_0$ ) at anthesis for 16 cultivars of wheat (A–P) grown in the field under N– and N+ at CF during the 2006–2007 (CF07) and 2007–2008 (CF08) growing seasons and for four cultivars grown at SB during the 2006–2007 growing season (SB07). Data are means  $\pm$  SE for three independent replicates. Lines were fitted to the data using the  $K_L$  reported in Table II.



the accumulated GAI, but the relationship depended on the site, year, N treatment, and cultivar (Fig. 2). According to the site/year/N treatment/cultivar combination,  $N_{LA}$  declined from the flag leaf to the lowest leaf in the canopy by 48% to 83%.

For N–, the relationship between the relative photosynthetic leaf N and the relative incident PPFD could be described satisfactorily by Equation 5 for each site/year/cultivar combination (Fig. 3). The observed  $N_{LA}$  of the flag leaf ( $N_{LA}^f$ ) and that estimated using Equation 4 were not different ( $P = 0.754$ ; Supplemental Fig. S2A). For N+, in most cases, the values obtained by Equation 4 indicated a higher decrease of  $N_{LA}$  at the top of the canopy than observed values, but below the flag leaf it described well the  $N_{LA}$  gradient.  $N_{LA}^f$  estimated using Equation 4 was higher than observed values, as the intercept of the estimated versus observed relation differed from zero ( $P = 0.023$ ; Supplemental Fig. S2B). This relationship was characterized by the N extinction coefficient with respect to light ( $b$ ).  $b$  was modified by the cultivar, and averaged across the N treatments  $b$  varied between 1.23 (cv Paragon) and 2.20 (cv Consort) at CF07 and between 0.66 (cv Alchemy) and 0.98 (cv Récital) at CF08 (Table II). Values were higher for CF07 than for CF08. The year  $\times$  genotype interaction was significant for  $b$ . Whatever the genotype,  $b$  was higher for N– (averaging 1.40 across the cultivars and years) than for N+ (averaging 0.96 across the cultivars and years), indicating that the N gradient

was systematically steeper under limiting N supply. The differences between N treatments were consistent for all the cultivars (nonsignificant genotype  $\times$  N treatment interaction). Similar conclusions could be drawn from the study of the four cultivars studied at SB.

#### Relationship between Vertical Leaf N Distribution, Crop N Status, and Canopy Size

When all the site/year/N treatment/cultivar combinations were considered, shoot N concentration explained only 36% of  $b$  variations (Fig. 4A). A nitrogen nutrition index (NNI) can be determined to quantify crop N status; this index has been shown to be a better indicator of crop N status than shoot N concentration because it is well connected with the physiological regulation of N uptake at the canopy level (Lemaire et al., 2008). The relationship between  $b$  and NNI was thus analyzed (Fig. 4B). As expected, NNI explained a larger part (52%) of  $b$  variations than shoot N concentration. There was no relationship between  $b$  and  $N_{LA}^{sh}$  ( $P = 0.15$ ) or  $N_{LA}^f$  ( $P = 0.86$ ).

The relationship between  $b$  and GAI was then analyzed in order to assess whether the variability associated with the vertical N distribution, both among cultivars and between N treatments, could be related to differences in canopy size. When the different site/year/N treatment/cultivar combinations were considered,  $b$  decreased with

**Table II.** Canopy  $K_L$  and  $b$  for 16 cultivars of wheat

Crops were grown in the field at CF during the 2006–2007 (CF07) and 2007–2008 (CF08) growing seasons and at SB during the 2006–2007 growing season (SB07). Data are means  $\pm$  SE for two treatments  $\times$  three independent replicates.  $P$  values of the main effects and interactions from split-plot ANOVAs for the 16 cultivars at CF are reported.

Cultivars and Statistical Data	$K_L$			$b$		
	CF07	CF08	SB07	CF07	CF08	SB07
	$m^2 \text{ ground } m^{-2} \text{ green tissue}$			dimensionless		
Alchemy	0.33 $\pm$ 0.04	0.49 $\pm$ 0.03	–	1.22 $\pm$ 0.09	0.64 $\pm$ 0.11	–
Beaver	0.34 $\pm$ 0.04	0.41 $\pm$ 0.03	–	1.84 $\pm$ 0.23	0.68 $\pm$ 0.11	–
Consort	0.30 $\pm$ 0.04	0.41 $\pm$ 0.06	–	1.85 $\pm$ 0.20	0.83 $\pm$ 0.13	–
Paragon	0.47 $\pm$ 0.08	0.51 $\pm$ 0.05	–	1.11 $\pm$ 0.10	0.70 $\pm$ 0.08	–
Rialto	0.48 $\pm$ 0.03	0.54 $\pm$ 0.07	0.47 $\pm$ 0.11	1.15 $\pm$ 0.15	0.85 $\pm$ 0.07	0.45 $\pm$ 0.05
Robigus	0.44 $\pm$ 0.06	0.39 $\pm$ 0.02	–	1.26 $\pm$ 0.12	0.76 $\pm$ 0.12	–
Savannah	0.33 $\pm$ 0.04	0.41 $\pm$ 0.03	0.45 $\pm$ 0.05	1.67 $\pm$ 0.24	0.75 $\pm$ 0.11	0.51 $\pm$ 0.08
Soissons	0.39 $\pm$ 0.09	0.38 $\pm$ 0.06	–	1.72 $\pm$ 0.22	0.94 $\pm$ 0.17	–
Arche	0.42 $\pm$ 0.05	0.38 $\pm$ 0.03	–	1.42 $\pm$ 0.11	0.76 $\pm$ 0.08	–
CF9107	0.47 $\pm$ 0.04	0.42 $\pm$ 0.10	–	1.29 $\pm$ 0.13	0.80 $\pm$ 0.06	–
CF99102	0.48 $\pm$ 0.04	0.46 $\pm$ 0.04	0.51 $\pm$ 0.07	1.25 $\pm$ 0.11	0.69 $\pm$ 0.05	0.37 $\pm$ 0.04
Perfector	0.43 $\pm$ 0.06	0.39 $\pm$ 0.04	0.39 $\pm$ 0.11	1.33 $\pm$ 0.12	0.68 $\pm$ 0.09	0.40 $\pm$ 0.08
Quebon	0.51 $\pm$ 0.07	0.44 $\pm$ 0.04	–	1.49 $\pm$ 0.16	0.74 $\pm$ 0.08	–
Recital	0.52 $\pm$ 0.05	0.37 $\pm$ 0.02	–	1.41 $\pm$ 0.09	0.96 $\pm$ 0.05	–
Renan	0.42 $\pm$ 0.06	0.38 $\pm$ 0.02	–	1.41 $\pm$ 0.16	0.84 $\pm$ 0.08	–
Toisonдор	0.46 $\pm$ 0.04	0.45 $\pm$ 0.03	–	1.28 $\pm$ 0.15	0.82 $\pm$ 0.07	–
Minimum	0.30	0.37	0.39	1.11	0.64	0.37
Maximum	0.52	0.54	0.51	1.85	0.96	0.51
Average	0.42	0.43	0.46	1.42	0.78	0.43
$P$ values						
N	0.205			<0.001		
Year	0.617			<0.001		
Genotype	<0.001			<0.001		
N $\times$ year	0.603			0.003		
N $\times$ genotype	0.881			0.174		
Year $\times$ genotype	<0.001			<0.001		
N $\times$ year $\times$ genotype	0.826			0.465		

increasing GAI according to a unique power function that accounted for 79% of  $b$  variations (Fig. 4C):

$$b = \beta \times \text{GAI}^\alpha \quad (1)$$

No correlation was found between  $K_L$  and GAI ( $r = 0.014$ ,  $P = 0.904$ ); hence, the relationship between  $b$  and GAI was due to a correlation between  $K_N$  and GAI ( $r = 0.92$ ,  $P > 0.001$ ). As a result of the isometric overall relationship between  $b$  and GAI and the exponential decay of the light down the canopy,  $b$  was linearly related to the reciprocal of  $\ln$  of the canopy transmittance (Supplemental Fig. S3).

The allometric relation between  $b$  and GAI was statistically significant for 14 of the 16 cultivars (Table III; Supplemental Fig. S4), and the scaling coefficient ( $\beta$ ) differed among the cultivars, ranging from 2.16 (cv Rialto) to 5.41 (cv Perfector). No difference was found for the scaling exponent ( $\alpha$ ) of the  $b$ -GAI relationship, and there was no shift along the common slope (Table III), but the parameters  $\alpha$  and  $\beta$  were correlated ( $r = -0.854$ ,  $P < 0.001$ ).

## DISCUSSION


Previous reports based on several species studied under controlled conditions suggested that plant internal

N availability could be determinant in the development of the vertical leaf N gradient (Anten et al., 1995b; Lötscher et al., 2003). This suggested that factors other than light could play a key role in the determination of the vertical leaf N gradient with respect to light. Therefore, we analyzed the vertical leaf N gradient in relation to different ecophysiological variables for 16 cultivars of wheat grown under two conditions of N supply in three field experiments. To the best of our knowledge, this is the first study examining the genetic variability related to the vertical leaf N gradient, with respect to the light gradient, at the intraspecific level. Determining whether the genetic variability in this trait can be related to ecophysiological variables had never been tested before. For a staple crop like wheat, such findings could have important implications for plant breeding and modeling.

## The Vertical Distribution of Light Varied Among Cultivars

The vertical light gradient is the main environmental factor governing the development of the vertical leaf N gradient in canopies (Werger and Hirose, 1991; Pons and de Jong-van Berkel, 2004). In this study, the distribution of light was described by the  $K_L$ . It is broadly acknowledged that the estimation of  $K_L$ , which assumes

**Figure 2.** Cumulative GAI counted from the top of the canopy ( $F$ ) versus  $N_{LA}$  at anthesis for 16 cultivars of wheat (A–P) grown in the field under N– and N+ at CF during the 2006–2007 (CF07) and 2007–2008 (CF08) growing seasons and for four cultivars grown at SB during the 2006–2007 growing season (SB07). Data are means  $\pm$  SE for three independent replicates. Lines were fitted to the data by reduced major axis regression using logarithmic transformation of Equation 4. Symbols and lines are as in Figure 1.



that light attenuates exponentially within the canopy, is a simplification that may not describe the actual light environment, due to sunflecks, diffuse radiations, or variations of leaf angle with depth (Sinclair and Shiraiwa, 1993; Kull and Jarvis, 1995; Johnson et al., 2010). Beyond these concerns, we assessed  $K_L$  using a simplified method: instead of performing stratified light measurements at different depths within the canopy, light was measured above the canopy and at ground level only. This methodological choice, motivated by the large number of cultivars under study, was shown to provide reliable estimations of  $K_L$ . Several authors have stressed that the development of high-throughput methods for field-based plant phenotyping represents a major bottleneck for analyzing the genetic variability and determinism of plant functional traits (Montes et al., 2007). The method used in this study is well adapted to analyze the genetic variability of  $K_L$ , an important canopy-level trait, and therefore represents an important methodological advance.


N treatments did not modify  $K_L$ . The stability of  $K_L$  for different conditions of N supply reported in this study is consistent with previous studies (Robertson and Giunta, 1994; Muirinen and Peltonen-Sainio, 2006). By contrast, significant differences among cultivars were found. The values of  $K_L$  reported here are lower than those reported in some other investigations, where stem and ear surface areas were calculated as their

projected surface area (O'Connell et al., 2004; Shearman et al., 2005) or where only leaf surface area was considered to calculate GAI (Muirinen and Peltonen-Sainio, 2006). However, the range of genetic variation for  $K_L$  observed in this study is in good agreement with the literature (for review, see Martre et al., 2007). Thus, while increasing N supply increased GAI, the intensity with which light attenuated with increasing cumulative GAI was independent of N supply and canopy size but varied with cultivar. The large year  $\times$  genotype interaction component nevertheless precludes unambiguous cultivar ranking for this trait.

#### The Ratio of Vertical Leaf N to Light Gradients at Anthesis Scales with the Size of the Canopy

A power relationship was observed between vertical N and light distributions for each site/year/N treatment/cultivar combination. However, the adjustment of the leaf N gradient to the light gradient (reflected by the value of  $b$ ) varied according to the cultivar. In addition, it differed with the N treatment: whatever the cultivar, the leaf N gradient relative to the light gradient was steeper at lower N supply. Different descriptors related to crop N status and canopy size were considered.  $b$  decreased with increasing shoot N concentration according to a trend that was similar to that reported by Lötscher et al.






**Figure 3.** Relative PPFD incident on the leaf ( $I_i/I_0$ ) versus relative photosynthetic  $N_{LA}$   $[(N_{LA} - n_b)/(N_{LA}^0 - n_b)]$  at anthesis for 16 cultivars of wheat (A–P) grown in the field under N– and N+ at CF during the 2006–2007 (CF07) and 2007–2008 (CF08) growing seasons and for four cultivars grown at SB during the 2006–2007 growing season (SB07). For  $[(N_{LA} - n_b)/(N_{LA}^0 - n_b)]$ , data are means  $\pm$  SE for three independent replicates. For  $I_i/I_0$ , data at each leaf rank were estimated a posteriori using the fitted curve between  $I_i/I_0$  and the height in the canopy (data not shown) and the height of each leaf rank. Lines were fitted to the data by reduced major axis regression using logarithmic transformation of Equation 5. Symbols are as in Figure 1.

(2003) for several monocot and dicot species, but most of the variability of  $b$  remained unexplained by shoot N concentration. NNI is a better descriptor of crop N status than shoot N concentration (Lemaire et al., 2008) and, as hypothesized by Lötscher et al. (2003), it accounted for a larger part of the variability of  $b$  than shoot N concentration. A key result of this study was that  $b$  decreased with GAI, indicating that the leaf N gradient to the light gradient became progressively more uniform when canopy size increased. Under N+, most of the cultivars accumulated more N at the top of the canopy (mainly in the flag leaf) than expected by an exponential function of relative light. Nevertheless, this finding did not challenge the observed relationship between  $b$  and GAI. Indeed, this relationship remained valid even if we excluded the canopies for

which luxury N storage existed at the top of the canopy or the flag leaf for all the canopies (data not shown). In addition, there was no correlation between  $K_L$  and GAI. Therefore, the relationship between  $b$  and GAI stemmed from the relationship between  $K_N$  and GAI.

Whether  $b$  should be considered as a function of GAI, or rather GAI as a function of  $b$ , is a priori not straightforward. Indeed, if there is a strong vertical attenuation of leaf N, then the crop would not be able to maintain a high GAI, since the leaf at bottom of the canopy would start to senesce due to N shortage. In other words, GAI would decrease as a consequence of increasing  $b$ . In our study, the number of leaves per main shoot at anthesis was nearly constant, ranging from four to five leaves. Therefore, the variations of GAI were not caused by differences in leaf number per



**Figure 4.**  $b$  versus shoot N concentration (A), crop NNI (B), and GAI (C) at anthesis for 16 cultivars of wheat grown in the field under N– and N+ at CF during the 2006–2007 (CF07) and 2007–2008 (CF08) growing seasons and for four cultivars grown at SB during the 2006–2007 growing season (SB07). Data are means  $\pm$  SE for three independent replicates. Solid lines indicate reduced major axis regression fitted to the data after logarithmic transformation. DM, Dry mass.



**Table III.** Summary statistics of the linear regression analysis of  $b$  versus GAI at anthesis for 16 wheat cultivars

Reduced major axis intercepts (coefficient) and slopes (exponents) and fitted axis means were calculated using the log-log version of the function equation  $b = \beta \times \text{GAI}^\alpha$ . CI, Confidence intervals. Data are plotted in Supplemental Figure S3.

Cultivars and Statistical Data	Exponent		Coefficient		Fitted Axis Mean	GAI at Which $b = 1$	$r^2$	P Value
	$\alpha$	95% CI	$\beta$	95% CI				
	$\text{m}^2 \text{ ground m}^{-2} \text{ green tissue}$		$\text{dimensionless}$					
Alchemy	-1.28	-1.76 to 0.06	4.59	2.57 to 8.12	0.25	3.30	0.99	0.006
Beaver	-1.17	-1.69 to 0.06	5.36	2.88 to 10.00	0.32	4.22	0.98	0.008
Consort	-0.95	-1.28 to 0.04	3.88	2.63 to 5.75	0.37	4.16	0.99	0.005
Paragon	-0.90	-1.20 to 0.04	3.23	2.19 to 4.68	0.24	3.66	0.99	0.004
Rialto	-0.76	-1.47 to 0.13	2.16	-0.93 to 5.01	0.19	2.74	0.74	0.027
Robigus	-0.86	-1.67 to 0.09	3.27	1.35 to 7.94	0.28	3.98	0.95	0.028
Savannah	-1.01	-1.37 to 0.08	4.14	2.45 to 6.92	0.19	4.05	0.95	0.001
Soissons	-0.95	-1.81 to 0.10	4.89	1.82 to 12.9	0.32	5.28	0.95	0.025
Arche	-0.87	-1.29 to 0.06	4.17	2.34 to 7.41	0.26	5.21	0.98	0.009
CF9107	-0.73	-2.36 to 0.16	2.78	0.57 to 13.5	0.27	4.04	0.77	0.123
CF99102	-0.95	-1.16 to 0.05	3.17	2.29 to 4.36	0.15	3.39	0.98	<0.001
Perfector	-1.25	-2.13 to 0.19	5.42	1.62 to 18.20	0.15	3.85	0.84	0.010
Quebon	-0.76	-1.02 to 0.03	2.62	1.91 to 3.63	0.29	3.57	0.99	0.005
Recital	-0.53	-0.85 to 0.04	2.36	1.62 to 3.47	0.32	5.10	0.97	0.013
Renan	-0.74	-2.09 to 0.15	3.32	0.76 to 14.79	0.25	5.06	0.84	0.086
Toison d'or	-0.82	-1.78 to 0.10	2.98	1.12 to 7.94	0.29	3.82	0.92	0.041
P values	0.336		<0.001		0.998			
Common parameter	-0.920	-1.017 to -0.847						

main shoot but rather by variation in individual leaf size and the number of main shoots per square meter. We can thus conclude that under our experimental conditions, the variations in  $b$  were caused by the variation in GAI.

This study is concordant with previous studies (Pons et al., 1989; Kull, 2002) challenging the optimization theory (Hirose and Werger, 1987). In our study, GAI covered a wide range of variation (from approximately 2 to 12  $\text{m}^2 \text{ m}^{-2}$ ), but under normal growth conditions, GAI of a mature canopy is generally higher than 4  $\text{m}^2 \text{ m}^{-2}$ , and according to our results,  $b$  is then lower than 1. These values are in good agreement with all available studies showing that actual leaf N distribution is more uniform ( $b < 1$ ) than the calculated optimal distribution, which maximizes canopy photosynthesis ( $b = 1$ ; Chen et al., 1993; Evans, 1993; Hollinger, 1996; Anten et al., 1998). As pointed out by Farquhar (1989), the parallelism between the vertical distributions of light and N ( $b = 1$ ) was found in models that do not take into account the costs of the construction and maintenance of photosynthesis. It can be assumed that these costs are proportional to GAI and that  $b$  scales with GAI so that the N distribution maximizes canopy photosynthesis independently of GAI. Our results strongly suggest that the acclimation of the leaf N gradient to the light gradient is a whole-plant process that depends notably on the size of the canopy, as suggested by Kull (2002).

Both theoretical and experimental results showed that the relative response of canopy photosynthesis to changes in  $b$  increases with the size of the canopy (Hirose and Werger 1987; Schieving et al., 1992; Sands, 1995). In our study, different stands covering a wide range of GAI were used to analyze the ecophysiological determinants of  $b$ . Under our growing conditions,

commercial wheat crops have a maximum GAI ranging between 4 and 6  $\text{m}^2 \text{ m}^{-2}$  (Foulkes et al., 1998); thus, according to Equation 1,  $b$  may range from 1.06 to 0.67. According to Hirose and Werger (1987), such variations in  $b$  would change canopy photosynthesis up to 10% to 15%. This rough estimation of the potential effect of changes in  $b$  on canopy photosynthesis has to be put into perspective using a mechanistic dynamic simulation model of whole-canopy photosynthesis over the whole growing season. A modeling analysis would also allow quantifying the effect of the genetic variability of  $\beta$  of the  $b$ -GAI relationship reported in this study on canopy photosynthesis and final grain crop biomass and grain yield.

Anyway, modeling leaf N distribution and canopy expansion based on the assumption that leaf N distribution is always optimal to maximize photosynthesis according to the optimization theory (i.e.  $b = 1$ ) is definitely inappropriate (Johnson et al., 2010). The relation between  $b$  and GAI is robust for a wide range of cultivars. Yet, empirically, it provides a good basis for improving our representation of N distribution within whole-plant models. Our work also puts emphasis on the importance of accurate modeling of GAI, because it determines both the extent of the light-intercepting area and the distribution of N among leaves (through  $b$  modulation) and therefore the photosynthesis per leaf area unit.

In summary, the analysis of wheat canopies grown in a wide range of site/year/N treatment/cultivar combinations provided a large variability of the vertical leaf N distribution at anthesis and allowed the three questions framed in the introduction to be answered. Regarding the effect of N supply on the distribution of leaf N, we showed that the distribution of leaf N, with respect to the light distribution, was steeper

when N supply decreased. Regarding the second question, when all the cultivars in each site/year/N treatment combination were considered, we found a wide range of  $b$  values, but most of these variations reflected a phenotypic plasticity rather than a genetic variability for  $b$ , as the genotype  $\times$  environment interaction was significant. Finally, we found that GAI was a unique ecophysiological determinant of both genetic and environmental variations of  $b$ , and we can conclude that the vertical distribution of leaf N scales with the size of the canopy. This finding is in good agreement with previous reports showing that the acclimation of the leaf N gradient to the light gradient is a whole-plant process that depends notably on the size of the canopy (Kull, 2002). The  $\beta$  of this allometric relationship differed among cultivars, suggesting that cultivars could be differently adapted to low-productivity environments.

## MATERIALS AND METHODS

### Plant Material and Growing Conditions

Sixteen genotypes of wheat (*Triticum aestivum*) from the United Kingdom and France elite germplasm (Supplemental Table S1) were studied in a 2-year field experiment carried out at CF (45°47' N, 3°10' E, 329 m elevation) during the 2006–2007 (experiment CF07) and 2007–2008 (experiment CF08) growing seasons (named 2007 and 2008, respectively). The genotypes comprised UK- and French-bred cultivars/advanced lines representing feed wheat (Alchemy, Arche, Beaver, Consort, Paragon, Rialto, Robigus, and Savannah) and bread-making wheat (CF9107, CF99102, Perfector, Quebon, Recital, Renan, Soissons, and Toisonador). All cultivars had winter growth habit, except the spring cultivar Paragon. Four of the cultivars (Rialto, Savannah, CF99102, and Perfector), representing the variability observed for N utilization and N uptake efficiency (Gaju et al., 2011), were also grown in 2007 (in experiment SB07) at the University of Nottingham in SB (52°50' N, 1°14' W, 52 m elevation). Soil characteristics for the different experiments are reported in Supplemental Table S2. At CF, crops were sown at a density of 250 seeds  $m^{-2}$  on November 10, 2006 and November 6, 2007. At SB, crops were sown at a density of 300 seeds  $m^{-2}$  on October 4, 2006. At both sites, the experimental design was a split plot in which N treatments were randomized on main plots, cultivars were randomized on the subplots, and each treatment combination was replicated three times. Subplot size was  $24 \times 1.65$  m at SB and  $7 \times 1.5$  m at CF. Interrow spacing was 0.17 and 0.125 m at CF and SB, respectively.

Weather conditions during the crop growth seasons are summarized in Supplemental Figure S5. From sowing to anthesis of cv Rialto, which has a rate of development in the middle of the range of the 16 cultivars (Supplemental Table S1), daily mean air temperature averaged 9.7°C and 7.9°C at CF07 and CF08, respectively, and was 9.1°C at SB07. During the same period, accumulated weekly radiation averaged 68 and 74 MJ  $m^{-2} week^{-1}$  at CF in 2007 and 2008, respectively, and was 53 MJ  $m^{-2} week^{-1}$  at SB. At CF07, the crop received 156 mm of rainfall during the vegetative period, which was lower than the long-term mean. Rainfall was also lower than the long-term mean in January and February 2008. At SB, rainfall was lower than the long-term mean in April and May 2007, during which the crops received 58 mm. At CF, anthesis was recorded between May 4 and 22, 2006, and between May 15 and June 1, 2007, depending on the cultivar (Supplemental Table S1). At SB, it was recorded on June 3 and 4, 2007 depending on the cultivar.

In all experiments, N+ and N− treatments were applied. For N+, the rates of N fertilization were determined using the balance-sheet method to optimize grain yield (Rémy and Hébert, 1977). N was applied as ammonium nitrate granules (34.5% N) in three (SB) to four (CF) splits. At CF, in both years, 4, 8, 8, and 4 g N  $m^{-2}$  were applied when cv Rialto, reached growth stage 21 (GS21; Zadoks et al., 1974), GS31, GS37, and GS61, respectively. At SB, 6, 9, and 6 g N  $m^{-2}$  were applied at GS21, GS31, and GS39. The amount of N applied under N− was adjusted in each site-season according to the soil mineral N measured in February (Supplemental Table S2) with the aim of providing approximately 10 g N  $m^{-2}$  from the combined soil mineral N and fertilizer N, corresponding to a moderate to severe N limitation sufficient to reduce grain yield by

approximately 30% compared with N+ conditions. At CF, in both years, 4 g N  $m^{-2}$  was applied in one split when cv Rialto reached GS31, while at SB, no N was applied in the N− treatment. All other crop inputs, including weed, disease, and pest control and potassium, phosphate, and sulfur fertilizers, were applied at levels to prevent non-N nutrients, weeds, diseases, and pests from limiting yield. At SB, the plant growth regulator chlormequat was applied at GS31. The major growth stages were determined for each cultivar on five to 10 tagged plants per plot every 2 to 3 d according to Tottman (1987).

### Plant Sampling

When each cultivar reached GS61, plant material in a 0.5- $m^2$  area per subplot was sampled by cutting at ground level. The total fresh mass of the samples was determined, and a 25% subsample (by fresh mass) was randomly selected. The main and secondary shoots were separated and counted. Shoots were classified as main when the top of their ear was in the top 0.15-m layer of the canopy. Thirty main shoots were randomly selected. The distances from the base of the shoot to individual culm leaf ligules and to the base and top of the ear were measured on 10 main shoots. Then, the 30 main shoots were dissected into individual leaf laminae, stem (including leaf sheath), and ear. The fresh mass of the secondary shoots of the subsamples was determined, and subsamples of secondary shoot (approximately 20%) were randomly selected. Their fresh mass was determined before separating them into laminae (pooled), stem (including leaf sheath), and ear. For both main and secondary shoots, green and dead/nonphotosynthetic tissues were analyzed separately as described below.

### Green Surface Area Determination

The projected surface areas of green laminae, stems, and ears were determined using a Li-3100 area meter (LI-COR). For the main shoots, the lamina surface area of each phytomer was determined, while for the secondary shoots, the total lamina surface area was determined. Actual stem and ear surface areas were calculated as their projected surface area multiplied by  $\pi/2$  (Lang, 1991). Canopy GAI ( $m^2$  green tissue  $m^{-2}$  ground) was calculated as the sum of lamina, stem, and ear surface area of the main and secondary shoots. GAI between successive main shoot leaf ligules was calculated as the sum of half of the laminae and stem surface area of the main and secondary shoots (plus ear peduncle and ear surface for the upper layer) located below and above each leaf ligule. The laminae and stem surface areas of secondary shoots were distributed to each layer in proportion to the laminae and stem surface areas of primary shoots.

### Light Measurements and $K_L$ Calculations

The vertical distribution of PPFD was measured 1 d before plant sampling using a 90-cm-long linear ceptometer (LP-80 AccuPAR; Decagon Devices) equipped with an external PPFD sensor. Simultaneous measurements were taken diagonally across the rows and above the canopy. At CF07, measurements were taken every 10 cm from the top of the canopy to the ground level in three different areas of each subplot (Bertheloot et al., 2008). The attenuation of PPFD down the canopy was described according to the Beer-Lambert's law (Monsi and Saeki, 2005):

$$I = I_0 \times \exp(-K_L \times F) \quad (2)$$

where  $F$  ( $m^2$  green tissue  $m^{-2}$  ground) is the cumulative GAI from the top of the canopy and  $I_0$  and  $I$  ( $\mu mol m^{-2} s^{-1}$ ) are the PPFD values on a horizontal level above and within the canopy at depth  $F$ , respectively. For GAI higher than 0.2  $m^2 m^{-2}$ , leaf clumping is very limited in wheat (Demarez et al., 2008); therefore, unbiased estimation of  $K_L$  could be done without taking into account the foliage aggregation.

The PPFD at the height of each culm leaf ligule was estimated from an exponential function equation fitted to the PPFD values plotted against the height from the ground level. Then,  $K_L$  was estimated by fitting Equation 2 to the estimated log-transformed PPFD values at the ligule heights plotted against log-transformed  $F$ . At CF08 and SB07, PPFD was measured above the canopy and at ground level only after confirming the validity of this simplified method of PPFD measurements using CF07 data (Supplemental Fig. S1) and independent data sets including canopies with GAI values ranging from 4.6 to 9.7  $m^2 m^{-2}$  (data not shown).

PPFD was measured at SB07 using a 100-cm-long linear ceptometer (SunScan System SS1; Δ-T Devices). Ten measurements were taken per subplot.  $K_L$  was

then calculated from Equation 2 (Shearman et al., 2005). All the PPFD measurements were restricted to 10 AM to 2 PM (solar time).

The PPFD incident on the leaf surface ( $I_i$ ;  $\mu\text{mol m}^{-2} \text{s}^{-1}$ ) at depth  $F$  in the canopy was calculated as follows (Thornley and Johnson, 2000):

$$I_i = I_0 \frac{K_L}{1-m} \times \exp(-K_L \times F) \quad (3)$$

where  $m$  (dimensionless) is the PPFD transmittance of the leaf laminae, set at 0.08 (Masoni et al., 1996).  $I_i$  was calculated after logarithmic transformation of Equation 3. The error due to the use of only the light extinction of the canopy in Equation 3 is most likely negligible (5%–15% at most for genotypes with an erect leaf posture).

### Main Shoot Dry Mass, N Concentration, Specific $N_{LA}$ , Canopy $K_N$ , and NNI

Lamina, stem, and ear dry mass of the main and secondary shoots was determined separately after oven drying at 80°C for 48 h. Samples were then milled, and their total N concentration (N mass per unit dry mass) was determined with the Dumas combustion method (Association of Official Agricultural Chemists method no. 7.024) using a FlashEA 1112 N/Protein Analyzer (Thermo Electron) at CF and a NA-2000 elemental analyzer (Fisons) at SB. Specific  $N_{LA}$  (g N  $\text{m}^{-2}$  lamina) was calculated by dividing the mass of green lamina N by the green lamina surface area. Average  $N_{LA}$  of main shoot leaves was calculated by dividing the total lamina N by the total laminae surface area.

Vertical leaf lamina N distribution through the canopy can be related to  $F$  as follows:

$$N_{LA} = (N_{LA}^f - n_b) \times \exp(-K_N \times F) + n_b \quad (4)$$

where  $n_b$  (g N  $\text{m}^{-2}$  leaf lamina) is the  $N_{LA}$  at which light-saturated photosynthesis is nil;  $K_N$  ( $\text{m}^2 \text{ground m}^{-2}$  green tissue) is the N extinction coefficient; and  $N_{LA}^f$  approximates the  $N_{LA}$  of the flag leaf. For wheat, after heading,  $N_{LA}^f$  may significantly depart from the  $N_{LA}$  at the top of the canopy because of the presence of the ear. Previous works indicated that optimal N distribution can be given by the Lagrange multiplier (Field, 1983; Hirose and Werger, 1987) when maximal photosynthesis is proportional to light availability across leaf layers (Farquhar, 1989). Equation 3 was then derived by Anten et al. (1995b), who showed that N distribution is optimal when convexity and the initial slope of the light-response curve of photosynthesis are constant across leaves.

After eliminating  $F$  from Equations 3 and 4,  $N_{LA}$  can be related to PPFD:

$$N_{LA} = (N_{LA}^f - n_b) \times \left( \frac{I_i}{I_{i0}} \right)^b + n_b \quad (5)$$

where  $b$  (dimensionless) is the coefficient of leaf N distribution with respect to relative PPFD incident on a leaf and is equal to the ratio of  $K_N$  to  $K_L$  and  $I_{i0}$  ( $\mu\text{mol m}^{-2} \text{s}^{-1}$ ) is the PPFD incident on the leaf at midpoint (in terms of GAI) of the flag leaf lamina.  $b$  and  $N_{LA}^f$  were estimated after logarithmic transformation of Equation 5. In wheat,  $n_b$  ranges from 0.25 to 0.55 g N  $\text{m}^{-2}$  (Araus and Tapia, 1987; Evans, 1993; Lawlor, 1995; Dreccer et al., 2000; Sudo et al., 2003; Vos et al., 2005; Del Pozo et al., 2007). In a preliminary analysis, we conducted a sensitivity analysis in which  $n_b$  was varied from 0.25 to 0.55 g N  $\text{m}^{-2}$  with intervals of 0.05 g N  $\text{m}^{-2}$ . This sensitivity analysis showed that  $b$  tended to increase linearly with  $n_b$ . However, the values of  $b$  estimated when either a fixed value of  $n_b$  was used or both  $b$  and  $n_b$  were estimated simultaneously for each cultivar were closely correlated with a slope not significantly different from 1 and  $n_b$  was not significantly different among cultivars (data not shown). Therefore,  $n_b$  was set at the nominal value of 0.4 g N  $\text{m}^{-2}$ . The  $N_{LA}$  of a layer was related to  $I_i/I_{i0}$  at midpoint of that layer (Milroy et al., 2001) estimated using Equation 2 and to  $K_L$  estimated from PPFD measurements taken above the canopy and at ground level. According to Anten et al. (1995b), an optimal  $N_{LA}$  distribution maximizing canopy photosynthesis is given when  $K_L$  equals  $K_N$  and thus  $b$  equals 1.  $b$  is a key parameter defining the acclimation of leaf photosynthesis to the light environment (Sands, 1995). The value of  $b$  increases with the nonuniformity of the N distribution (increasing difference between the  $N_{LA}$  values at the top and bottom of the canopy).

Based on the critical dilution curve of wheat (Justes et al., 1994), a NNI was calculated to quantify crop N status. Although this curve is empirical, it is based on solid theoretical grounds (Lemaire and Gastal, 1997). Climatic conditions can affect growth and N uptake differently, but the NNI incorporates

these effects in terms of crop N needs (for a thorough discussion of N dilution in crops, see Lemaire et al., 2008; Gonzalez-Dugo et al., 2010). For a given shoot dry mass (DM; g  $\text{m}^{-2}$ ), NNI was calculated as the ratio between the actual and critical shoot N concentrations ( $N_c$ ; g N  $\text{g}^{-1}$  dry mass) defined by the critical dilution curve (Justes et al., 1994):

$$N_c = 40.96 \times \text{DM}^{-0.442} \quad (6)$$

A NNI value close to 1 indicates an optimal crop N status, and a value lower (or higher) than 1 indicates N deficiency (or N excess).

### Statistics

All regression analyses were done in R-2.12.2 for Windows (R Development Core Team, 2007). The two methods for estimating  $K_L$  were compared by reduced major axis regression (Warton et al., 2006) with the package SMATR for R (Falster et al., 2006). All ANOVAs were performed with Statgraphics Plus 4.1 for Windows (Statistical Graphics). ANOVA procedures for a split-plot design was used at CF for the 16 cultivars. Replications were regarded as random effects, while N treatment and genotype were fixed effects. For ANOVA across the three experiments and the four cultivars, site and year were regarded as random effects. Statistical differences were judged at  $\alpha = 0.05$ .

### Supplemental Data

The following materials are available in the online version of this article.

**Supplemental Figure S1.** Comparison of the two methods used to estimate the canopy  $K_L$ .

**Supplemental Figure S2.** Estimated versus observed flag leaf lamina N mass per unit leaf area.

**Supplemental Figure S3.** Relationship between the ratio of  $K_N$  to  $K_L$  and the reciprocal of the ln of the canopy transmittance.

**Supplemental Figure S4.** Coefficient of leaf N distribution with respect to relative light versus GAI at anthesis for 16 cultivars of wheat.

**Supplemental Figure S5.** Weather data for the three growing seasons considered in this study.

**Supplemental Table S1.** Cultivar name, country of origin, registration year, height, and anthesis date.

**Supplemental Table S2.** Soil characteristics.

### ACKNOWLEDGMENTS

We thank J. Messaoud, B. Bonnemoy, J.L. Joseph, and P. Lemaire (all Institut National de la Recherche Agronomique CF) and John Alcock, Matthew Tovey, and Dr. Stella Hubbart (all University of Nottingham) for their technical assistance as well as Dr. B. Andrieu (Institut National de la Recherche Agronomique Grignon) and Dr. Tom Sinclair (University of Florida) for helpful comments on an earlier version of the manuscript.

Received May 8, 2012; accepted September 13, 2012; published September 14, 2012.

### LITERATURE CITED

- Aerts R (1996) Nutrient resorption from senescing leaves of perennials: are there general patterns? *J Ecol* **84**: 597–608
- Anten NPR, Schieving F, Medina E, Werger MJA, Schuffelen P (1995a) Optimal leaf area indices in C3 and C4 mono and dicotyledonous species at low and high nitrogen availability. *Physiol Plant* **95**: 541–550
- Anten NPR, Schieving F, Werger MJA (1995b) Patterns of light and nitrogen distribution in relation to whole canopy carbon gain in C3 and C4 mono and dicotyledonous species. *Oecologia* **101**: 504–513
- Anten NPR, Werger MJA, Medina E (1998) Nitrogen distribution and leaf area indices in relation to photosynthetic nitrogen use efficiency in savanna grasses. *Plant Ecol* **138**: 63–75
- Araus JL, Tapia L (1987) Photosynthetic gas exchange characteristics of wheat flag leaf blades and sheaths during grain filling: the case of a

- spring crop grown under Mediterranean climate conditions. *Plant Physiol* **85**: 667–673
- Archontoulis SV, Vos J, Yin X, Bastiaans L, Danalatos NG, Struik PC** (2011) Temporal dynamics of light and nitrogen vertical distributions in canopies of sunflower, kenaf and cynara. *Field Crops Res* **122**: 186–198
- Bertheloot J, Martre P, Andrieu B** (2008) Dynamics of light and nitrogen distribution during grain filling within wheat canopy. *Plant Physiol* **148**: 1707–1720
- Bidinger F, Musgrave RB, Fischer RA** (1977) Contribution of stored preanthesis assimilate to grain yield in wheat and barley. *Nature* **270**: 431–433
- Bindraban PS** (1999) Impact of canopy nitrogen profile in wheat on growth. *Field Crops Res* **63**: 63–77
- Boonman A, Prinsen E, Gilmer F, Schurr U, Peeters AJM, Voesenek LACJ, Pons TL** (2007) Cytokinin import rate as a signal for photosynthetic acclimation to canopy light gradients. *Plant Physiol* **143**: 1841–1852
- Chen JL, Reynolds JF, Harley PC, Tenhunen JD** (1993) Coordination theory of leaf nitrogen distribution in a canopy. *Oecologia* **93**: 63–69
- Del Pozo A, Perez P, Gutierrez D, Alonso A, Morcuende R, Martinez-Carrasco R** (2007) Gas exchange acclimation to elevated CO<sub>2</sub> in upper-sunlit and lower-shaded canopy leaves in relation to nitrogen acquisition and partitioning in wheat grown in field chambers. *Environ Exp Bot* **59**: 371–380
- Demarez V, Duthoit S, Baret F, Weiss M, Dedieu G** (2008) Estimation of leaf area and clumping indexes of crops with hemispherical photographs. *Agric Meteorol* **148**: 644–655
- Dreccer MF, Slafer GA, Rabbinge R** (1998) Optimization of vertical distribution of canopy nitrogen: an alternative trait to increase yield potential in winter cereals. *J Crop Prod* **1**: 47–77
- Dreccer MF, Van Oijen M, Schapendonk AHCM, Pot CS, Rabbinge R** (2000) Dynamics of vertical leaf nitrogen distribution in a vegetative wheat canopy: impact on canopy photosynthesis. *Ann Bot (Lond)* **86**: 821–831
- Drouet JL, Bonhomme R** (1999) Do variations in local leaf irradiance explain changes to leaf nitrogen within row maize canopies? *Ann Bot (Lond)* **84**: 61–69
- Evans JR** (1989) Photosynthesis and nitrogen relationships in leaves of C<sub>3</sub> plants. *Oecologia* **78**: 9–19
- Evans JR** (1993) Photosynthetic acclimation and nitrogen partitioning within a lucerne canopy. 1. Canopy characteristics. *Aust J Plant Physiol* **20**: 55–67
- Falster DS, Warton DI, Wright IJ** (2006) SMATR: standardised major axis tests and routines, version 2.0. <http://www.bio.mq.edu.au/ecology/SMATR/> (October 12, 2012)
- Farquhar GD** (1989) Models of integrated photosynthesis of cells and leaves. *Philos Trans R Soc Lond B Biol Sci* **323**: 357–367
- Field C** (1983) Allocating leaf nitrogen for the maximization of carbon gain: leaf age as a control on the allocation program. *Oecologia* **56**: 341–347
- Foulkes J, Holdsworth MJ, Kerr S, Kightly S, Barraclough PB, Hawkesford MJ, Shewry PR** (2006) A study of the scope for the application of crop genomics and breeding to increase nitrogen economy within cereal and rapeseed based food chains. Final report for project AR0714. Department for Environment, Food and Rural Affairs, London
- Foulkes MJ, Sylvester-Bradley R, Scott RK** (1998) Evidence for differences between winter wheat cultivars in acquisition of soil mineral nitrogen and uptake and utilization of applied fertilizer nitrogen. *J Agric Sci* **130**: 29–44
- Gaju O, Allard V, Martre P, Snape JW, Heumez E, LeGouis J, Moreau D, Bogard M, Griffiths S, Orford S, et al** (2011) Identification of traits to improve the nitrogen-use efficiency of wheat genotypes. *Field Crops Res* **123**: 139–152
- Gebbing T, Schnyder H** (1999) Pre-anthesis reserve utilization for protein and carbohydrate synthesis in grains of wheat. *Plant Physiol* **121**: 871–878
- Gonzalez-Dugo V, Durand J-L, Gastal F** (2010) Water deficit and nitrogen nutrition of crops: a review. *Agron Sustain Dev* **30**: 529–544
- Grindlay DJC** (1997) Towards an explanation of crop nitrogen demand based on the optimization of leaf nitrogen per unit leaf area. *J Agric Sci* **128**: 377–396
- Grindlay DJC, Sylvester-Bradley R, Scott RK** (1995) The relationship between canopy green area and nitrogen in the shoot. In G Lemaire, IJ Burns, eds, *Diagnostic Procedures for Crop N Management*, Vol 82. INRA Editions, Poitiers, France, pp 53–60
- Hikosaka K** (2005) Leaf canopy as a dynamic system: ecophysiology and optimality in leaf turnover. *Ann Bot (Lond)* **95**: 521–533
- Hikosaka K, Terashima I, Katoh S** (1994) Effects of leaf age, nitrogen nutrition and photon flux density on the distribution of nitrogen among leaves of a vine (*Ipomoea tricolor* Cav.) grown horizontally to avoid mutual shading of leaves. *Oecologia* **97**: 451–457
- Hirose T, Werger MJA** (1987) Maximizing daily canopy photosynthesis with respect to the leaf nitrogen allocation pattern in the canopy. *Oecologia* **72**: 520–526
- Hollinger DY** (1996) Optimality and nitrogen allocation in a tree canopy. *Tree Physiol* **16**: 627–634
- Johnson IR, Thornley JHM, Frantz JM, Bugbee B** (2010) A model of canopy photosynthesis incorporating protein distribution through the canopy and its acclimation to light, temperature and CO<sub>2</sub>. *Ann Bot (Lond)* **106**: 735–749
- Jukanti AK, Fischer AM** (2008) A high-grain protein content locus on barley (*Hordeum vulgare*) chromosome 6 is associated with increased flag leaf proteolysis and nitrogen remobilization. *Physiol Plant* **132**: 426–439
- Justes E, Mary B, Meynard JM, Machet JM, Thelier-Huche L** (1994) Determination of a critical nitrogen dilution curve for winter wheat crops. *Ann Bot (Lond)* **74**: 397–407
- Kichey T, Hirel B, Heumez E, Dubois F, Le Gouis J** (2007) In winter wheat (*Triticum aestivum* L.), post-anthesis nitrogen uptake and remobilisation to the grain correlates with agronomic traits and nitrogen physiological markers. *Field Crops Res* **102**: 22–32
- Kull O** (2002) Acclimation of photosynthesis in canopies: models and limitations. *Oecologia* **133**: 267–279
- Kull O, Jarvis PG** (1995) The role of nitrogen in a simple scheme to scale-up photosynthesis from leaf to canopy. *Plant Cell Environ* **18**: 1174–1182
- Lang ARG** (1991) Application of some of Cauchy's theorems to estimation of surface areas of leaves, needles and branches of plants, and light transmittance. *Agric Meteorol* **55**: 191–212
- Lawlor DW** (1995) Photosynthesis, productivity and environment. *J Exp Bot* **46**: 1449–1461
- Lawlor DW** (2002) Carbon and nitrogen assimilation in relation to yield: mechanisms are the key to understanding production systems. *J Exp Bot* **53**: 773–787
- Le Gouis J, Béghin D, Heumez E, Pluchard P** (2000) Genetic differences for nitrogen uptake and nitrogen utilisation efficiencies in winter wheat. *Eur J Agron* **12**: 163–173
- Lemaire G, Gastal F** (1997) N uptake and distribution in plant canopies. In G Lemaire, ed, *Diagnosis of the Nitrogen Status in Crops*. Springer-Verlag, Berlin, pp 3–43
- Lemaire G, Jeuffroy M-H, Gastal F** (2008) Diagnosis tool for plant and crop N status in vegetative stage: theory and practices for crop N management. *Eur J Agron* **28**: 614–624
- Lemaire G, Millard P** (1999) An ecophysiological approach to modelling resource fluxes in competing plants. *J Exp Bot* **50**: 15–28
- Lötscher M, Stroh K, Schnyder H** (2003) Vertical leaf nitrogen distribution in relation to nitrogen status in grassland plants. *Ann Bot (Lond)* **92**: 679–688
- Martre P, Semenov MA, Jamieson PD** (2007) Simulation analysis of physiological traits to improve yield, nitrogen use efficiency and grain protein concentration in wheat. In JHJ Spiertz, PC Struik, HH Van Laar, eds, *Scale and Complexity in Plant Systems Research, Gene-Plant-Crop Relations*. Springer, Dordrecht, The Netherlands, pp 181–201
- Masoni A, Ercoli L, Mariotti M** (1996) Spectral properties of leaves deficient in iron, sulfur, magnesium, and manganese. *Agron J* **88**: 937–943
- Mi G, Tang L, Zhang F, Zhang J** (2000) Is nitrogen uptake after anthesis in wheat regulated by sink size? *Field Crops Res* **68**: 183–190
- Milroy SP, Bange MP, Sadras VO** (2001) Profiles of leaf nitrogen and light in reproductive canopies of cotton (*Gossypium hirsutum*). *Ann Bot (Lond)* **87**: 325–333
- Monaghan JM, Snape JW, Chojceki AJS, Kettlewell PS** (2001) The use of grain protein deviation for identifying wheat cultivars with high grain protein concentration and yield. *Euphytica* **122**: 309–317
- Monsi M, Saeki T** (2005) On the factor light in plant communities and its importance for matter production. 1953. *Ann Bot (Lond)* **95**: 549–567
- Montes JM, Melchinger AE, Reif JC** (2007) Novel throughput phenotyping platforms in plant genetic studies. *Trends Plant Sci* **12**: 433–436
- Muurinen S, Peltonen-Sainio P** (2006) Radiation-use efficiency of modern and old spring cereal cultivars and its response to nitrogen in northern growing conditions. *Field Crops Res* **96**: 363–373
- O'Connell MG, O'Leary GJ, Whitfield DM, Connor DJ** (2004) Interception of photosynthetically active radiation and radiation-use efficiency of

- wheat, field pea and mustard in a semi-arid environment. *Field Crops Res* **85**: 111–124
- Oury FX, Berard P, Brancourt-Hulmel M, Depatureaux C, Doussinault G, Galic N, Giraud A, Heumez E, Lecomte C, Pluchard P, et al** (2003) Yield and grain protein concentration in bread wheat: a review and a study of multi-annual data from a French breeding program. *J Genet Breed* **57**: 59–68
- Pons TL, de Jong-van Berkel YEM** (2004) Species-specific variation in the importance of the spectral quality gradient in canopies as a signal for photosynthetic resource partitioning. *Ann Bot (Lond)* **94**: 725–732
- Pons TL, Jordi W, Kuiper D** (2001) Acclimation of plants to light gradients in leaf canopies: evidence for a possible role for cytokinins transported in the transpiration stream. *J Exp Bot* **52**: 1563–1574
- Pons TL, Schieving F, Hirose T, Werger MJA** (1989) Optimization of leaf nitrogen allocation for canopy photosynthesis in *Lysimachia vulgaris*. In H Lambers, ML Cambridge, H Konings, TL Pons, eds, *Causes and Consequences of Variation in Growth Rate and Productivity of Higher Plants*. SPB Academic Publishing, The Hague, The Netherlands, pp 175–186
- Pons TL, Van Rijnbeck H, Scheurwater I, Van Der Verf A** (1993) Importance of the gradient in photosynthetically active radiation in a vegetation stand for leaf nitrogen allocation in two monocotyledons. *Oecologia* **95**: 416–424
- R Development Core Team** (2007) *A Language and Environment for Statistical Computing*. R Foundation for Statistical Computing, Vienna
- Rémy JC, Hébert J** (1977) Le devenir des engrais dans le sol. *Compte Rendu de l'Académie d'Agriculture de France* **63**: 700–710
- Robertson MJ, Giunta F** (1994) Responses of spring wheat exposed to pre-anthesis water stress. *Aust J Agric Res* **45**: 19–35
- Rousseaux MC, Hall AJ, Sanchez RA** (1999) Light environment, nitrogen content, and carbon balance of basal leaves of sunflower canopies. *Crop Sci* **39**: 1093–1100
- Sands PJ** (1995) Modelling canopy production. I. Optimal distribution of photosynthetic resources. *Aust J Plant Physiol* **22**: 593–601
- Schieving F, Werger MJA, Hirose T** (1992) Canopy structure, nitrogen distribution and whole canopy photosynthetic carbon gain in growing and flowering stands of tall herbs. *Vegetatio* **102**: 173–181
- Shearman VJ, Sylvester-Bradley R, Scott RK, Foulkes MJ** (2005) Physiological processes associated with wheat yield progress in the UK. *Crop Sci* **45**: 175–185
- Shiraiwa T, Sinclair TR** (1993) Distribution of nitrogen among leaves in soybean canopies. *Crop Sci* **33**: 804–808
- Sinclair TR, Shiraiwa T** (1993) Soybean radiation-use efficiency as influenced by nonuniform specific leaf nitrogen distribution and diffuse radiation. *Crop Sci* **33**: 808–812
- Sudo E, Makino A, Mae T** (2003) Differences between rice and wheat in ribulose-1,5-bisphosphate regeneration capacity per unit of leaf-N content. *Plant Cell Environ* **26**: 255–263
- Thornley JHM, Johnson IR** (2000) *Plant and Crop Modelling: A Mathematical Approach to Plant and Crop Physiology*. Blackburn Press, Caldwell, NJ
- Tottman DR** (1987) The decimal code for the growth stages of cereals, with illustrations. *Ann Appl Biol* **110**: 441–454
- van Oosterom E, Borrell AK, Chapman SC, Broad IJ, Hammer GL** (2010) Functional dynamics of leaf nitrogen balance of sorghum. I. N-balance during pre-anthesis. *Field Crops Res* **115**: 19–28
- Vos J, van der Putten PEL, Birch CJ** (2005) Effect of nitrogen supply on leaf appearance, leaf growth, leaf nitrogen economy and photosynthetic capacity in maize (*Zea mays* L.). *Field Crops Res* **93**: 64–73
- Wang ZJ, Wang JH, Zhao CJ, Zhao M, Huang WJ, Wang CZ** (2005) Vertical distribution of nitrogen in different layers of leaf and stem and their relationship with grain quality of winter wheat. *J Plant Nutr* **28**: 73–91
- Warton DI, Wright IJ, Falster DS, Westoby M** (2006) Bivariate line-fitting methods for allometry. *Biol Rev Camb Philos Soc* **81**: 259–291
- Werger MJA, Hirose T** (1991) Leaf nitrogen distribution and whole canopy photosynthetic carbon gain in herbaceous stands. *Plant Ecol* **97**: 11–20
- Yin X, Lantinga EA, Schapendonk AHCM, Zhong X** (2003) Some quantitative relationships between leaf area index and canopy nitrogen content and distribution. *Ann Bot (Lond)* **91**: 893–903
- Zadoks JC, Chang TT, Konzak CF** (1974) A decimal code for the growth stages of cereals. *Weed Res* **14**: 415–421