

Delayed generalized muscle atrophy following unilateral hindlimb casting in old rats is prevented by a whey or a high protein diet but not a free leucine-enriched diet

Hugues H. Magne, Isabelle Savary-Auzeloux, Carole Migné, Marie-Agnès Peyron, Lydie Combaret, Didier Remond, Dominique Dardevet

▶ To cite this version:

Hugues H. Magne, Isabelle Savary-Auzeloux, Carole Migné, Marie-Agnès Peyron, Lydie Combaret, et al.. Delayed generalized muscle atrophy following unilateral hindlimb casting in old rats is prevented by a whey or a high protein diet but not a free leucine-enriched diet. 1 st Montréal International Symposium of GRAPA: from Humans to Cells (MHC), Groupe de Recherche en Activité Physique Adaptée (GRAPA). FRA., May 2013, Montréal, Canada. hal-01002245

HAL Id: hal-01002245 https://hal.science/hal-01002245v1

Submitted on 3 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Delayed generalized muscle atrophy following unilateral hindlimb casting in old rats is prevented by a whey or a high protein diet but not a free leucine-enriched diet.

Hugues Magne ^{1, 2}, Isabelle Savary-Auzeloux ^{1, 2}, Carole Migné ^{1,2}, Marie-Agnès Peyron ^{1, 2}, Lydie Combaret ^{1, 2}, Didier Rémond ^{1, 2} and Dominique Dardevet ^{1,2}

- (1) Clermont Université, Université d'Auvergne, Unité de Nutrition Humaine, BP 10448, F-63000 CLERMONT-FERRAND
- (2) INRA, UMR 1019, UNH, CRNH Auvergne, F-63000 CLERMONT-FERRAND

Introduction

During ageing, limb immobilization induces general and delayed muscle atrophy during the rehabilitation period. Understanding the underlying mechanisms and limit this generalized muscle atrophy is then a major issue in elderlies.

Objective

We investigated the impact of dietary supplementations on generalized and delayed muscle atrophy induced by immobilization and subsequent recovery in old-adult-rats.

Methods

185 old-adult-rats were subjected to unilateral hindlimb casting immobilization for 8 days (I0-I8) and allowed to recover for 10-40 days (R10, R20, R30, R40). Animals were fed during the recovery period a control diet (C) or the same diet supplemented with leucine (LEU), whey (WHEY) or high-protein diet (HIGH-PROT). Muscle mass, muscle protein breakdown (MPB) and protein synthesis (MPS) were measured in non-immobilized *gastrocnemius* muscles. Data were analyzed by ANOVA.

Results

Muscle mass was unchanged in the non-immobilized limb from I0 to R20 but atrophied thereafter (-15.5%), traducing a general atrophy. MPB was unaltered in this limb but MPS became resistant to food intake compared to pair-fed non-immobilized-controls. In LEU group, *gastrocnemius* atrophied similarly despite a better stimulation of postprandial MPS (+20%). In the WHEY and HIGH-PROT groups, no atrophy was observed in the the non-immobilized-*gastrocnemius* throughout the experimental period (no difference at R40vsI0).

Discussion/conclusion

LEU supplementation was able to stimulate MPS in the non-immobilized limb via the mTOR-signalling-pathway, but it remained inefficient to prevent the muscle atrophy in the non-immobilized limb. By contrast, WHEY or HIGH PROT diets were able to prevent atrophy of the non-immobilized limb by sustaining MPS during the entire rehabilitation.