



Soluble milk proteins promote the recovery of muscle functional properties after an immobilization

Vincent V. Martin, Sébastien S. Ratel, Julien J. Siracusa, Charlotte C. Baudry, Pascale P. Le Ruyet, Lydie Combaret, Isabelle I. Auzeloux, Christelle C. Guillet, Dominique Dardevet

► To cite this version:

Vincent V. Martin, Sébastien S. Ratel, Julien J. Siracusa, Charlotte C. Baudry, Pascale P. Le Ruyet, et al.. Soluble milk proteins promote the recovery of muscle functional properties after an immobilization. 1 st Montréal International Symposium of GRAPA : from Humans to Cells (MHC), GRavitation and AstroParticle Physics., May 2013, Montreal, Canada. hal-01002246

HAL Id: hal-01002246

<https://hal.science/hal-01002246>

Submitted on 3 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Soluble milk proteins promote the recovery of muscle functional properties after an immobilization

V. Martin¹, S. Ratel¹, J. Siracusa¹, C. Baudry², P. Le Ruyet², L. Combaret³, I. Savary-Auzeloux³, C. Guillet³ & D. Dardevet³

¹Clermont Université, Université Blaise Pascal, Laboratoire des Adaptations Métaboliques à l'Exercice en conditions Physiologiques et Pathologiques (EA 3533), Clermont-Fd

²Lactalis Recherche et Développement, Retiers

³Clermont Université, Université d'Auvergne, INRA, Unité de Nutrition Humaine (UNH, UMR 1019), Clermont-Fd

INTRODUCTION

It has been shown that soluble milk proteins are more efficient in the recovery of muscle mass after a period of immobilization. However, their effect on muscle functional properties remains unknown.

OBJECTIVE

To evaluate the effect of milk proteins quality on the recovery of muscle functional properties after an immobilization.

METHODS

After an initial (I0) evaluation of the contractile properties of the plantar flexors (isometric torque and concentric power), the ankle of 30 adult rats was immobilized by casting for 8 days. After cast removal, rats were fed either with a diet with 13% of casein (CAS); or 13% of native soluble milk proteins (NAT); or 13% of proteins from whey (WHEY). During the recovery period, contractile properties were evaluated 7 (R7), 21 (R21) and 42 days (R42) after cast removal.

RESULTS

The immobilization induced an homogeneous depression of isometric torque and concentric power in all the experimental groups. At R21, isometric torque had fully recovered in WHEY and NAT, but was still significantly depressed in CAS, where complete recovery occurred at R42. No significant difference of power recovery kinetics was observed between CAS, WHEY and NAT at R21. However, power was increased in NAT at R42 as compared to I0 (+ 14.7 ± 26.0% ; P<0.001), but not in CAS and WHEY.

DISCUSSION/CONCLUSION

Soluble milk proteins clearly speed up the recovery of isometric torque as compared to casein. However, in dynamic condition, the increase of concentric power above baseline values is only observed in the group fed with native soluble milk proteins.