



HAL
open science

A conserved molecular basis for photoperiod adaptation in two temperate legumes

James L. J. L. Weller, Lim Chee L. C. Liew, Valerie F. G. V. F. G. Hecht,
Vinodan V. Rajandran, Rebecca E. R. E. Laurie, Stephen S. Ridge, Bénédicte
B. Wenden, Jacqueline K. J. K. Vander Schoor, Odile O. Jaminon, Christelle
C. Blassiau, et al.

► To cite this version:

James L. J. L. Weller, Lim Chee L. C. Liew, Valerie F. G. V. F. G. Hecht, Vinodan V. Rajandran, Rebecca E. R. E. Laurie, et al.. A conserved molecular basis for photoperiod adaptation in two temperate legumes. *Proceedings of the National Academy of Sciences of the United States of America*, 2012, 109 (51), pp.21158 - 21163. 10.1073/pnas.1207943110 . hal-01004100

HAL Id: hal-01004100

<https://hal.science/hal-01004100>

Submitted on 29 May 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

A conserved molecular basis for photoperiod adaptation in two temperate legumes

James L. Weller^{a,1}, Lim Chee Liew^{a,2}, Valérie F. G. Hecht^a, Vinodan Rajandran^a, Rebecca E. Laurie^b, Stephen Ridge^a, Bénédicte Wenden^{c,3}, Jacqueline K. Vander Schoor^a, Odile Jaminon^d, Christelle Blassiau^e, Marion Dalmais^f, Catherine Rameau^c, Abdelhafid Bendahmane^f, Richard C. Macknight^b, and Isabelle Lejeune-Hénaut^d

^aSchool of Plant Science, University of Tasmania, Hobart, Tasmania 7001, Australia; ^bDepartment of Biochemistry, University of Otago, Dunedin 9054, New Zealand; ^cInstitut Jean-Pierre Bourgin, Institut National de la Recherche Agronomique (INRA) Unité Mixte de Recherche (UMR) 1318, INRA-AgroParis Tech, F-78000 Versailles, France; ^dINRA, UMR 1281, Stress Abiotiques et Différenciation des Végétaux cultivés, 80203 Péronne, France; ^eUMR 1281-Stress Abiotiques et Différenciation des Végétaux cultivés, Université des Sciences et Technologies de Lille, 59650 Villeneuve d'Ascq, France; and ^fUnité de Recherche en Génomique Végétale, UMR 1165 ERL8196 INRA-Université d'Evry Val d'Essonne-Centre National de la Recherche Scientifique, F-91057 Evry, France

Edited by Maarten Koornneef, Max Planck Institute for Plant Breeding Research, Cologne, Germany, and approved November 7, 2012 (received for review May 13, 2012)

Legumes were among the first plant species to be domesticated, and accompanied cereals in expansion of agriculture from the Fertile Crescent into diverse environments across the Mediterranean basin, Europe, Central Asia, and the Indian subcontinent. Although several recent studies have outlined the molecular basis for domestication and eco-geographic adaptation in the two main cereals from this region, wheat and barley, similar questions remain largely unexplored in their legume counterparts. Here we identify two major loci controlling differences in photoperiod response between wild and domesticated pea, and show that one of these, *HIGH RESPONSE TO PHOTOPERIOD* (*HR*), is an ortholog of *EARLY FLOWERING 3* (*ELF3*), a gene involved in circadian clock function. We found that a significant proportion of flowering time variation in global pea germplasm is controlled by *HR*, with a single, widespread functional variant conferring altered circadian rhythms and the reduced photoperiod response associated with the spring habit. We also present evidence that *ELF3* has a similar role in lentil, another major legume crop, with a distinct functional variant contributing to reduced photoperiod response in cultivars widely deployed in short-season environments. Our results identify the factor likely to have permitted the successful prehistoric expansion of legume cultivation to Northern Europe, and define a conserved genetic basis for major adaptive changes in flowering phenology and growth habit in an important crop group.

crop adaptation | *Pisum sativum* | *Lens culinaris*

Many of the world's earliest agricultural systems were based around crops from two important groups: cereals and legumes. Although molecular and genetic analyses have led to considerable progress in understanding the genetic changes underlying domestication and adaptation in several cereal crops, similar efforts in legumes are in general much less advanced. Among the legumes domesticated in the world's oldest farming culture in the Neolithic Near East, the temperate long-day (LD) species lentil (*Lens culinaris* Medik.), pea (*Pisum sativum* L.), and chickpea (*Cicer arietinum* L.) all persist as crops of global economic importance. Of these crops, pea has the widest distribution, the most diverse phenology, and is the best understood molecularly, and offers prospects for a detailed exploration of molecular events important in early cultivation and spread (1, 2).

P. sativum is now generally viewed as a complex species that includes a wide variety of cultivated and wild forms with pink, purple, or white flowers (1). Wild *P. sativum* lines are characterized by dehiscent pods and a rough, thick seed coat, and include both tall, climbing forms distributed around the Mediterranean (*P. sativum* var. *elatius*) and shorter forms restricted to the Near East (*P. sativum* var. *humile*), which intergrade in their areas of overlap. Cytogenetic differences and analyses of genetic diversity support the view that the majority of cultivated peas originated from a distinct gene pool within var. *humile* (1), although recent molecular studies also highlight the likely genomic contribution

from other wild forms, and emphasize the importance of introgression and recombination within the complex (2, 3). Domesticated variants of *P. sativum* include garden or "vegetable" pea (var. *sativum*) and field pea (var. *arvense*) grown for dry seed or as a forage crop. A distinct taxon (*P. sativum* var. *abyssinicum*) found in highland regions of Ethiopia and southern Yemen is believed to represent an independent domesticate with a substantial genomic contribution from a second, related species, the yellow/orange-flowered *Pisum fulvum* Sibth. & Sm (1, 3).

Control of flowering time is widely acknowledged as an important feature of plant adaptation, and there has been much recent effort directed toward understanding the molecular basis for flowering time adaptation in both wild and domesticated species (4). One common feature of flowering-time adaptation is the relaxation of mechanisms that operate to delay flowering under unfavorable conditions, through loss of function in genes controlling responsiveness to photoperiod or vernalization. These changes reduce the length of the growth cycle, permitting a shift from winter to spring cropping in temperate regions, and enabling expansion to areas where the growing season is limited by short duration or water availability. Similar changes also underlie expansion of temperate crops to low latitudes where the short photoperiods and lack of vernalizing temperatures would otherwise preclude flowering. Wild *Pisum* in its native range displays a typical winter habit in which plants germinate in autumn, overwinter in the vegetative state, and flower in response to increasing day-length in spring (5). Anecdotal reports of experiments in controlled conditions also suggest that wild *P. sativum* lines generally only flower when grown under LD photoperiods (6). This obligate or near-obligate requirement for LDs suits pea to a winter cropping cycle and has been retained in certain forage cultivars (7). However, the majority of cultivated pea accessions from higher latitudes have a quantitative LD response and are grown as a spring crop (7).

In this study, we undertook an initial genetic analysis of the differences in flowering and photoperiod responsiveness between

Author contributions: J.L.W., V.F.G.H., C.R., and I.L.-H. designed research; J.L.W., L.C.L., V.F.G.H., V.R., R.E.L., S.R., B.W., J.K.V.S., O.J., C.B., M.D., and R.C.M. performed research; A.B. contributed new reagents/analytic tools; J.L.W., L.C.L., V.F.G.H., V.R., R.E.L., S.R., B.W., J.K.V.S., M.D., C.R., R.C.M., and I.L.-H. analyzed data; and J.L.W. wrote the paper.

The authors declare no conflict of interest.

This article is a PNAS Direct Submission.

Data deposition: The sequence reported in this paper has been deposited in the GenBank database (accession nos. [JN983406](https://doi.org/10.1073/pnas.1207943110) and [JN983407](https://doi.org/10.1073/pnas.1207943110)).

¹To whom correspondence should be addressed. E-mail: jim.weller@utas.edu.au.

²Present address: Melbourne School of Land and Environment, University of Melbourne, Parkville, Victoria 3010, Australia.

³Present address: Institut National de la Recherche Agronomique, UR419, Unité de Recherche sur les Espèces Fruitières, Centre de Bordeaux, F-33140 Villenave-d'Ornon Cedex, France.

This article contains supporting information online at www.pnas.org/lookup/suppl/doi:10.1073/pnas.1207943110/-DCSupplemental.

wild and domesticated pea. We identified two major loci controlling these differences, and showed that one of them likely corresponds to an ortholog of the circadian clock gene *EARLY FLOWERING 3 (ELF3)*. We further demonstrate that variation in this gene is also associated with photoperiod-insensitive early flowering in the related crop species lentil.

Results

We first sought to gain an overview of natural variation for flowering and photoperiod responsiveness in pea, in a selection of lines representing a broad range of genetic diversity (3, 8) (Table S1). All lines flowered within a narrow developmental window under LD, but much wider variation was observed under short days (SD), where *P. fulvum*, var. *abyssinicum* and most domesticated lines flowered, but lines of var. *elatius* and *humile*, several landraces, and several winter cultivars did not (Fig. 1A). This finding suggests that ancestral *P. sativum* was an obligate LD plant, consistent with the winter habit observed for wild forms in their natural range (5, 6), and suggests that evolution of early-flowering types in which environmental constraints to flowering are genetically relaxed has been a key feature of postdomestication spread.

We next examined the genetic basis for changes in SD flowering under domestication in *P. sativum* in a cross between NGB5839, an isogenic dwarf derivative of the spring cultivar Torsdag, and the wild line JI1794, a representative accession of the northern race of *P. sativum* var. *humile* proposed as the major wild contributor to the domesticated pea gene pool (1). Like many spring cultivars, NGB5839 and its progenitor cv. Torsdag carry recessive alleles at the *HIGH RESPONSE TO PHOTOPERIOD (HR)* locus that confer early flowering in SD (1). We examined the association between node of flower initiation and markers for a range of flowering-related genes. As expected, we identified significant association with markers at the top of linkage group III in the region of *HR*, but also with markers in the middle of linkage group VI, and further genotyping of markers in these two regions defined two quantitative trait loci (QTL) (Fig. 1B and Table S2). These loci showed contrasting patterns of inheritance,

with the early flowering conferred by domesticated alleles showing recessive inheritance in the case of *QTL3* but dominant inheritance at *QTL6* (Fig. 1C). The loci also showed significant interaction, with extreme delay of flowering under SD only seen in plants homozygous for the JI1794 allele at *QTL6* and carrying at least one JI1794 allele at *QTL3* (Fig. 1C). We conclude that wild alleles at both loci are required for full expression of the winter habit. *QTL3* and *QTL6* together explained 88% of the observed variance for node of flower initiation, as estimated by two-way ANOVA based on peak markers *MAX1* and *RNAhel* (Fig. 1C). In addition, single marker tests failed to provide evidence for significant effects of other known flowering loci, including *LF* (9), members of the *FT* family (10), or *SN* (11) (Table S3). Interestingly, markers for both *QTL3* and *QTL6* were also associated with variation in shoot branching, which was reduced in an additive manner by homozygosity for the domesticated alleles (Fig. 1C).

Because NGB5839 carries recessive alleles at the *HR* locus, we considered that *QTL3* might be largely or entirely equivalent to *HR*. Previous studies have characterized the effects of *HR* on responses to photoperiod and light quality, and in a more applied setting, on winter frost damage in field-grown plants (7, 11). To further characterize the physiological effects of *HR*, we generated near-isogenic *HR* and *hr* lines in which dominant *HR* alleles were introgressed into the NGB5839 background from the line WL1771 (Wellensiek's Dominant) (12) through six successive backcrosses. Consistent with previous studies (11, 13), the *hr* allele was associated with early flowering and reduced branching in plants grown under SD and reduced sensitivity to the light quality of extended photoperiods (Fig. 2A). In *HR* plants, flowering was more effectively promoted by extensions with light of low red (R):far-red (FR) ratio than of high R:FR ratio, whereas in *hr* plants both types of photoperiod extension were equally effective (Fig. 2A). One factor that influences responses to both light quality and photoperiod is the circadian clock, and we previously noted that rhythmic expression of several circadian clock genes showed unusual light-dependent damping in NGB5839 (14).




Fig. 1. Adaptation to photoperiod in pea is controlled by two major-effect QTL. (A) Survey of photoperiod-regulated flowering in *Pisum*. Plants received an 8-h photoperiod of natural daylight (SD) extended with low-irradiance ($10 \mu\text{mol}\cdot\text{m}^{-2}\cdot\text{s}^{-1}$) white light (LD) from mixed fluorescent and incandescent sources. Data are mean \pm SE for $n = 4$. Lines not flowering in SD conditions are indicated by a "+" symbol and had produced a minimum of 55 vegetative nodes before termination of the experiment 180 d after sowing. All lines are *P. sativum* var. *sativum* unless indicated. Lines carrying the *hr* (6C) mutation are shaded in pink, and the two lines used for subsequent genetic analysis are indicated by black arrowheads. (B) Location of QTL controlling SD flowering on linkage groups III and VI in the F_2 of a domesticated (NGB5839) \times wild (var. *humile*; JI1794) cross. The one-LOD and two-LOD confidence intervals around the peak are indicated by dark red and pale red shading, respectively. (C) Genotype means \pm SE for interaction of *QTL3* and *QTL6* in the control of flowering and other developmental traits. Genotypes at *QTL3* and *QTL6* were inferred from the genotype of peak markers *MAX1* (*QTL3*) and *RNAhel* (*QTL6*), with the wild (JI1794) and domesticated (NGB5839) alleles indicated by the suffixes -w and -d, respectively. Significance levels (*** $P < 0.001$; ns, $P > 0.05$) and proportion of variance explained for the individual locus effects and their interaction (int.) were determined by two-way ANOVA and indicated to the right of each plot.

origins were identical over the 3.8-kb genomic *ELF3* sequence apart from the *hr* mutation, and this sequence was also shared by 250 of 251 *hr* lines in the CRB collection. The *hr* mutation and its corresponding *HR* haplotype are thus widespread across domesticated germplasm and represent a highly differentiated form of the *ELF3* gene. Other sequence differences seem unlikely to have a major influence on flowering, in view of the fact that all 26 predicted amino acid substitutions in our *ELF3* sequence dataset affected poorly conserved residues and are unlikely to be functionally significant (Fig. S2), and furthermore that there was no significant effect of haplogroup on flowering under either SD or LD (Fig. S3).

Genetic variation for flowering time and photoperiod responsiveness has been documented in other temperate LD legumes, including lentil (e.g., ref. 28) and chickpea (29). In both cases, early flowering and reduced photoperiod sensitivity has been an important adaptation to cropping in South Asia, where the season is limited by short day-lengths or terminal drought (29, 30). To examine the possibility that *ELF3* orthologs might also underlie variation for phenology in other legumes, we examined the genetic control of flowering in crosses between early- and late-flowering lentil lines. We resolved a major locus controlling SD flowering (Fig. 5 A and B) in a cross between the photoperiod responsive line cv. Northfield (ILL5588) and the early-flowering line ILL6005, which is a derivative of cv. Precoz, an early-flowering line from Argentina (30). This locus likely corresponds to the *SN* locus previously defined in crosses with cv. Precoz (28). We




Fig. 3. A mutation in *ELF3* ortholog is the likely basis for the *hr* spring phenotype. (A) Details of the *PsELF3* 5C/6C polymorphism. (B) Complementation of flowering and petiole phenotypes of the *Arabidopsis elf3-1* mutant by the 5C (*HR*) but not the 6C (*hr*) form of *35S::PsELF3*, under 8-h SD conditions. Representative plants are shown for two independent transformants for each construct. (C) Association of the 5C/6C polymorphism with photoperiod responsiveness in a selection of *P. sativum* germplasm. Plants received an 8-h photoperiod of natural daylight (SD) extended with low-irradiance ($10 \mu\text{mol}\cdot\text{m}^{-2}\cdot\text{s}^{-1}$) white light (LD) from mixed fluorescent and incandescent sources. Data are mean \pm SE for $n = 64$ (5C) and $n = 20$ (6C).




Fig. 4. Sequence diversity in the *HR* gene. Neighbor-joining tree representing genetic distances among haplotypes identified in a 3.8-kb region of the *HR* gene in 110 diverse *Pisum* lines (*P. sativum* var. *sativum* except where indicated). Node support (%) was obtained from 10,000 bootstrap replicates. Numbering of haplotypes corresponds to Table S1 and Fig. S2, with bold letters indicating distinct haplotype groups. Haplotypes present in more than one line are indicated by filled triangles with size proportional to the number of lines represented, with the single haplotype containing the *hr* mutation designated by an open triangle. Haplotypes including lines that flowered in SD conditions, despite carrying an apparently functional form of *HR*, are indicated by an asterisk.

assessed the genetic relationship of this locus to candidate photoperiod response genes and showed that it was tightly linked to the lentil *ELF3* ortholog (Fig. 5B), effectively excluding the possibility that it is an ortholog of pea *SN*, which is located in pea linkage group VII in a distinct part of the legume comparative map (20, 21). Sequencing of *ELF3* genomic DNA and cDNA in ILL6005 and cv. Precoz revealed a translationally silent G-to-A substitution in the last nucleotide of exon 3 (Fig. 5C) causing missplicing and skipping of exon 3. This change is predicted to cause a frame-shift in translation of exon 4 and termination after four missense amino acids. (Fig. 5D). These results suggest that this lentil locus is an ortholog of pea *HR*, and provides evidence that *HR* may play a similar role in photoperiod adaptation in lentil as in pea.

Discussion

The genetic basis for adaptive changes in flowering time has been extensively analyzed in *Arabidopsis* and many cereal crops, but is less well understood in other species, including legumes, which are a large and important crop group and include many of the world's oldest domesticated species. In this study we characterized natural variation for photoperiod responsiveness in the temperate LD species pea, including a broad range of domesticated and wild material. We show that wild forms have an obligate requirement for exposure to long photoperiods to flower, which is also seen in a wide range of primitive domesticated types. In contrast, many other domesticated lines of *P. sativum* from diverse origins do flower in SD, indicating that the ability to flower early under restrictive photoperiods has been an important factor in the expansion of pea cultivation. In crosses between a domestic and a wild line, we distinguished major contributions from two loci corresponding to the previously studied *HR* locus and




Fig. 5. Mutation in an *ELF3* ortholog is also associated with early flowering in lentil. (A) Differing photoperiod responsiveness in *Lens culinaris* lines ILL5588 and ILL6005. Plants received an 8-h photoperiod of natural daylight extended with 2-h (SD) or 16-h (LD) low-irradiance ($10 \mu\text{mol}\cdot\text{m}^{-2}\cdot\text{s}^{-1}$) white light from mixed fluorescent and incandescent sources. Data are mean \pm SE for $n = 9$ –10. (B) Cosegregation of flowering time under SD with a marker for *LcELF3* in the F_2 generation of cross ILL6005 \times ILL5588. The flowering time ranges of the two parental lines are shown as horizontal bars. (C) Details of the mutation in *LcELF3* genomic DNA. Early-flowering segregants carry a translationally silent G-to-A substitution in the last nucleotide of exon 3. (D) Details of splicing defect. PCR with the indicated primers (small arrowheads) revealed a 52-bp deletion in the *LcELF3* mRNA from ILL6005, corresponding to skipping of exon 3. This was verified by sequencing and results in a frame-shift in translation of exon 4 and termination after four missense amino acids.

a novel locus, *QTL6*. We identified *ELF3* as a strong positional candidate for *HR* and showed that *HR* influences several physiological characteristics also controlled by *ELF3* in *Arabidopsis*, including photoperiod responsiveness, sensitivity to low R:FR (31), and light regulation of circadian gene expression rhythms (15) (Fig. 2). Furthermore, we identified sequence polymorphism in pea *ELF3* that eliminated its function in transgenic *Arabidopsis* (Fig. 3B), was strongly associated with photoperiod responsiveness across a diverse set of pea germplasm (Fig. 3C), and was tightly linked to the *HR* locus. Together, these results strongly implicate *ELF3* as *HR*, but the alternative possibility that aspects of the *hr* phenotype derive from other nearby genes cannot be definitively excluded.

Identification of *HR* as a likely *ELF3* ortholog also prompted us to examine in detail the sequence diversity at the *HR* locus and its relationship to flowering time and photoperiod adaptation. In combined resequencing and TILLING analyses of over 500 genetically diverse pea accessions, we found evidence for only a single functional variant in the *HR* gene. The widespread nature of this *hr* mutation and the fact that it occurs within a closely related but distinct group of haplotypes suggest that the mutation originated relatively recently within an already differentiated lineage, and has undergone rapid dispersal. Archaeological evidence for the early history of pea cultivation suggests that following a relatively rapid westward spread from the inferred domestication center into Mediterranean Europe, expansion to

the north occurred only after a significant lag (32, 33). We speculate that this early westward lineage may be represented by haplogroup A, and the *hr* mutation may have arisen within this lineage, permitting summer cropping and thereby enabling expansion to areas with colder winters. More recently, reincorporation of functional *HR* alleles from winter forage pea cultivars into a variety of other end-use types is an important factor in the strategy for breeding winter forms of these crops (23) and the identification of a putative causal mutation should facilitate this process.

The observation that domesticated alleles at *QTL6* promote flowering under SD to an extent equivalent to the *hr* mutation (Fig. 1C) suggests the possibility of a second, independent route to a reduction in photoperiod response and the spring habit. The existence of *HR*-independent flowering variation within each of the three major *HR* haplogroups (Figs. 1A and Fig. S2) also indicates the likely contribution of additional genetic factors to photoperiod adaptation across the broader *Pisum* germplasm, which might include *QTL6*. In addition, although the previously characterized *LF/TFL1c* gene (9) did not influence SD flowering in our mapping population (Table S3), other studies of *LF* function and interaction indicate recessive *lf* alleles can confer earlier flowering in photoperiod-responsive genetic backgrounds under SD (11), and suggest *LF* as another candidate for some of the variation in SD flowering we observe. Interestingly, we found moderate to strong bootstrap support for association of var. *elatius* lines with each of the three var. *sativum* *HR* haplotype groups (Fig. 4), with haplogroup A in particular associated with var. *elatius* lines from Italy and Greece. This finding is consistent with the possibility that distinct lineages of var. *sativum* originated from different subsets of wild germplasm, and that early flowering arose independently in these lineages. The distinct form *P. sativum* var. *abyssinicum* is found in Ethiopia and Yemen and has been suggested as a probable independent domesticate with a clear genomic contribution from the second wild *Pisum* species, *P. fulvum*. We found that although var. *abyssinicum* lines carry a functional *HR* gene, they can all flower in SD, as expected from their distribution at latitudes below 10° N. Interestingly, *P. fulvum* itself (which also carries a predicted functional *HR* gene) is also able to flower in SD, which could be understood as an adaptation to a shorter effective season in its natural habitat compared with wild *P. sativum* forms (34). In view of the suggested relationship between *P. fulvum* and *P. sativum* var. *abyssinicum*, it will be interesting in future to examine whether their early-flowering phenotypes might have the same molecular basis and are related to the *HR*-independent early-flowering seen within var. *sativum*.

Evolution of early flowering under domestication has been studied in a number of other crop groups, most notably the cereals. Interestingly, these studies have identified a number of cases of convergence, with adaptation in different species conferred by changes in orthologous genes. Studies of photoperiod adaptation genes in wheat and barley have identified two main genetic routes to the spring growth habit: through loss-of-function *vm2* mutations in the grass-specific *ZCCT* transcription factors that act as repressors of flowering, or through semidominant mutations causing deregulated expression of the flower-promoting *Ppd1/PRR37* response regulator genes (35, 36). Evidence also suggests the importance of *PRR37* for photoperiod response in the warm-season cereals sorghum and rice (37, 38). The convergent nature of these adaptive changes implies the potential for similar convergence in other plant groups, and we examined this possibility in another legume species, lentil, which shows significant variation for flowering time and photoperiod responsiveness (28, 39). Linkage, sequencing and expression analyses show that a locus with a large effect on SD flowering is associated with a functionally significant polymorphism in the lentil *HR* ortholog. The origins of the recessive early allele are obscure but it seems to have arisen relatively recently in an Argentinian *macrosperma* landrace, and has subsequently provided an important means of broadening the genetic base of South Asian *microsperma* material (30). These results implicate altered *ELF3* function as a convergent route for evolution of photoperiod-hypersensitive

early-flowering in two different temperate legumes in different contexts. Taken together with recent reports that *ELF3* orthologs are also involved in natural variation for flowering in rice, barley, and *Arabidopsis* (31, 40–42), our results also suggest *HR* orthologs and functionally associated clock genes including *ELF4* and *LUX* (25) as possible candidates underlying genetic variation for phenology in a wider range of legumes. More generally, our characterization of *HR* and *QTL6* loci provides prospects for better understanding flowering-time adaptation, identifying new molecular targets in breeding for phenology and abiotic stress tolerance, and probing the early history of cultivation in this important group of crop plants.

Methods

Plant material was obtained from the *Pisum* germplasm collections at the John Innes Centre (Norwich, United Kingdom) and the Australian Temperate Field Crops Collection (Horsham, VIC, Australia). Lentil lines were provided by W. Erskine (International Center for Agricultural Research in the Dry Areas, Aleppo, Syria). Plants for expression experiments were grown in growth cabinets at 20 °C under 150 $\mu\text{mol}\cdot\text{m}^{-2}\cdot\text{s}^{-1}$ white light from cool-white fluorescent tubes. All other plants were grown under an 8-h photoperiod of natural daylight either with (LD) or without (SD) an 8-h extension of 10 $\mu\text{mol}\cdot\text{m}^{-2}\cdot\text{s}^{-1}$ white light from compact fluorescent tubes (LDH; R:FR = 4.8), 40 W incandescent globes (LDL; R:FR = 0.6), or a combination of both of these light sources (LD; R:FR = 1.2). The propensity to branch was quantified as the ratio of the length of lateral branches to the increase in total plant

height over a specified interval. QTL analysis was performed on a population of 92 F_2 plants from a cross between NGB5839 and JI1794 grown in SD, using JoinMap 4 and MapQTL 6 (Kyazma). Details of gene-based markers used are given in Table S5. Full-length pea and lentil *ELF3* genes and cDNAs were isolated using PCR techniques, genome walking (GenomeWalker universal kit; Clontech), and rapid amplification of cDNA ends (SMART RACE cDNA amplification kit; Clontech) using specific primers (Table S6). Harvested tissue for expression experiments consisted of both leaflets from the uppermost fully expanded leaf. RNA extraction, reverse-transcription, and real-time PCR analysis were performed as described in ref. 14. Construct preparation and *Arabidopsis* transformation were also carried out as previously described (14) and several independent transformants per construct were characterized through several generations under both LD and SD conditions. Distance and parsimony-based methods were used for phylogenetic analyses in PAUP*4.0b10 (<http://paup.csit.fsu.edu>).

ACKNOWLEDGMENTS. We thank N. Ellis, J. Burstin, W. Erskine, N. Weeden, I. Murfet, R. Redden, F. Sussmilch, and A. de Saint-Germain for access to materials, background information, and data; B. Potts and J. Freeman for advice on statistical and quantitative trait locus analysis; I. Cummings, T. Winterbottom, and M. Oates for technical assistance; N. Ellis, S. Abbo, and colleagues from the University of Tasmania for helpful comments on the manuscript; and I. Murfet for support in the initial stages of the lentil work. This work was supported by Australian Research Council Grant DP0878723 (to J.L.W.); New Zealand Foundation for Research Science and Technology Grant C10X0704 (to R.C.M.); an Institut National de la Recherche Agronomique Missions des Relations Internationales grant (to B.W.); and by Génomplante (project SNEPA, ANR06-GPLA019).

- Zohary D, Hopf M (2000) *Domestication of Plants in the Old World* (Oxford Univ Press, New York).
- Vershinin AV, Allnutt TR, Knox MR, Ambrose MJ, Ellis TH (2003) Transposable elements reveal the impact of introgression, rather than transposition, in *Pisum* diversity, evolution, and domestication. *Mol Biol Evol* 20(12):2067–2075.
- Jing R, et al. (2010) The genetic diversity and evolution of field pea (*Pisum*) studied by high throughput retrotransposon based insertion polymorphism (RBIP) marker analysis. *BMC Evol Biol* 10:44.
- Alonso-Blanco C, et al. (2009) What has natural variation taught us about plant development, physiology, and adaptation? *Plant Cell* 21(7):1877–1896.
- Abbo S, Shtienberg D, Lichtenzveig J, Lev-Yadun S, Gopher A (2003) The chickpea, summer cropping, and a new model for pulse domestication in the ancient near east. *Q Rev Biol* 78(4):435–448.
- Makasheva RK (1984) *The Pea* (A.A. Balkema, Rotterdam).
- Lejeune-Hénaut I, et al. (1999) Floral initiation in field-grown forage peas is delayed to a greater extent by short photoperiods, than in other types of European varieties. *Euphytica* 109:201–211.
- Zong X, et al. (2009) Analysis of a diverse global *Pisum* sp. collection and comparison to a Chinese local *P. sativum* collection with microsatellite markers. *Theor Appl Genet* 118(2):193–204.
- Foucher F, et al. (2003) *DETERMINATE* and *LATE FLOWERING* are two *TERMINAL FLOWER1/CENTRORADIALIS* homologs that control two distinct phases of flowering initiation and development in pea. *Plant Cell* 15(11):2742–2754.
- Hecht V, et al. (2011) The pea *GIGAS* gene is a *FLOWERING LOCUS T* homolog necessary for graft-transmissible specification of flowering but not for responsiveness to photoperiod. *Plant Cell* 23(1):147–161.
- Murfet IC (1985) *CRC Handbook of Flowering*, ed Halevy A (CRC Press, Boca Raton), pp 97–126.
- Murfet IC (1971) Flowering in *Pisum*. Three distinct phenotypic classes determined by the interaction of a dominant early and a dominant late gene. *Heredity* 26(2): 243–257.
- Reid JB (1982) Flowering in peas: Effect of the gene *Hr* on spectral sensitivity. *Crop Sci* 22(2):266–268.
- Liew LC, et al. (2009) *DIE NEUTRALIS* and *LATE BLOOMER 1* contribute to regulation of the pea circadian clock. *Plant Cell* 21(10):3198–3211.
- McWatters HG, Bastow RM, Hall A, Millar AJ (2000) The *ELF3* zeitnehmer regulates light signalling to the circadian clock. *Nature* 408(6813):716–720.
- Hazen SP, et al. (2005) *LUX ARRHYTHMO* encodes a Myb domain protein essential for circadian rhythms. *Proc Natl Acad Sci USA* 102(29):10387–10392.
- Allen T, et al. (2006) *Arabidopsis FHY3* specifically gates phytochrome signaling to the circadian clock. *Plant Cell* 18(10):2506–2516.
- Salathia N, et al. (2006) *FLOWERING LOCUS C*-dependent and -independent regulation of the circadian clock by the autonomous and vernalization pathways. *BMC Plant Biol* 6:10.
- Michaels SD, Bezerra IC, Amasino RM (2004) *FRIGIDA*-related genes are required for the winter-annual habit in *Arabidopsis*. *Proc Natl Acad Sci USA* 101(9):3281–3285.
- Hecht V, et al. (2005) Conservation of *Arabidopsis* flowering genes in model legumes. *Plant Physiol* 137(4):1420–1434.
- Bordat A, et al. (2011) Translational genomics in legumes allowed placing in silico 5460 unigenes on the pea functional map and identified candidate genes in *Pisum sativum* L. *G3 (Bethesda)* 1(2):93–103.
- Murfet IC (1973) Flowering in *Pisum. Hr*, a gene for high response to photoperiod. *Heredity* 31(12):157–164.
- Lejeune-Hénaut I, et al. (2008) The flowering locus *Hr* colocalizes with a major QTL affecting winter frost tolerance in *Pisum sativum* L. *Theor Appl Genet* 116(8):1105–1116.
- Yu JW, et al. (2008) *COP1* and *ELF3* control circadian function and photoperiodic flowering by regulating *GI* stability. *Mol Cell* 32(5):617–630.
- Nusinow DA, et al. (2011) The *ELF4-ELF3-LUX* complex links the circadian clock to diurnal control of hypocotyl growth. *Nature* 475(7356):398–402.
- Deulvot C, et al. (2010) Highly-multiplexed SNP genotyping for genetic mapping and germplasm diversity studies in pea. *BMC Genomics* 11:468.
- Triques K, et al. (2008) Mutation detection using ENDO1: Application to disease diagnostics in humans and TILLING and Eco-TILLING in plants. *BMC Mol Biol* 9:42.
- Sarker A, Erskine W, Sharma B, Tyagi MC (1999) Inheritance and linkage relationships of days to flower and morphological loci in lentil (*Lens culinaris* Medikus subsp. *culinaris*). *J Hered* 90(2):270–275.
- Millan T, et al. (2006) Chickpea molecular breeding: New tools and concepts. *Euphytica* 147(1–2):81–103.
- Erskine W, et al. (1998) A bottleneck in lentil: Widening its genetic base in south Asia. *Euphytica* 101(2):207–211.
- Jiménez-Gómez JM, Wallace AD, Maloof JN (2010) Network analysis identifies *ELF3* as a QTL for the shade avoidance response in *Arabidopsis*. *PLoS Genet* 6(9):6.
- Colledge S, Conolly J (2006) *The Origins and Spread of Domestic Plants in Southwest Asia and Europe* (Left Coast Press, Walnut Creek, CA).
- Purugganan MD, Fuller DQ (2009) The nature of selection during plant domestication. *Nature* 457(7231):843–848.
- Abbo S, et al. (2011) Experimental growing of wild pea in Israel and its bearing on Near Eastern plant domestication. *Ann Bot (Lond)* 107(8):1399–1404.
- Yan L, et al. (2004) The wheat *VRN2* gene is a flowering repressor down-regulated by vernalization. *Science* 303(5664):1640–1644.
- Beales J, Turner A, Griffiths S, Snape JW, Laurie DA (2007) A pseudo-response regulator is misexpressed in the photoperiod insensitive Ppd-1a mutant of wheat (*Triticum aestivum* L.). *Theor Appl Genet* 115(5):721–733.
- Murakami M, Matsushika A, Ashikari M, Yamashino T, Mizuno T (2005) Circadian-associated rice pseudo response regulators (OsPRRs): Insight into the control of flowering time. *Biosci Biotechnol Biochem* 69(2):410–414.
- Murphy RL, et al. (2011) Coincident light and clock regulation of pseudoresponse regulator protein 37 (PRR37) controls photoperiodic flowering in sorghum. *Proc Natl Acad Sci USA* 108(39):16469–16474.
- Erskine W, Ellis RH, Summerfield RJ, Roberts EH, Hussain A (1990) Characterization of responses to temperature and photoperiod for time to flowering in a world lentil collection. *Theor Appl Genet* 80:193–199.
- Zakhrabekova S, et al. (2012) Induced mutations in circadian clock regulator *Mat-a* facilitated short-season adaptation and range extension in cultivated barley. *Proc Natl Acad Sci USA* 109(11):4326–4331.
- Faure S, et al. (2012) Mutation at the circadian clock gene *EARLY MATURITY 8* adapts domesticated barley (*Hordeum vulgare*) to short growing seasons. *Proc Natl Acad Sci USA* 109(21):8328–8333.
- Saito H, et al. (2012) *Ef7* encodes an *ELF3*-like protein and promotes rice flowering by negatively regulating the floral repressor gene *Ghd7* under both short- and long-day conditions. *Plant Cell Physiol* 53(4):717–728.