



HAL
open science

Environmental assessment of bioethanol production from lignocellulosic crops (2nd generation)

Karine K. Dufosse, Benoit Gabrielle, Jean-Louis Drouet

► To cite this version:

Karine K. Dufosse, Benoit Gabrielle, Jean-Louis Drouet. Environmental assessment of bioethanol production from lignocellulosic crops (2nd generation). Proceedings, Conference Ecotech & tools, Nov 2011, Montpellier, France. 1 p. hal-01004311

HAL Id: hal-01004311

<https://hal.science/hal-01004311>

Submitted on 3 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Environmental assessment of bioethanol production from lignocellulosic crops

Authors: K. Dufossé B. Gabrielle, J.-L. Drouet

INRA, AgroParisTech, Environment and arable crops Res. Unit, Grignon, France.

Novelty: A multi-scale approach using agro-ecosystem and landscape models is developed to calculate the emissions of greenhouse gases (GHG), especially N₂O, related to crop and landscape management. This approach is used to improve the Life Cycle Assessment (LCA) of second generation biofuels.

Context: 20% of fuel have to be produced from renewable origin in 2020, and meet a 50% decrease of GHG emission compared to fossil fuel (RED Directive, 2009/28/CE). Besides, agriculture represents 10 to 15% of GHG emissions in France and especially 65% of N₂O emissions which have a huge global warming potential (310 times higher than CO₂, according to IPCC 2007). Therefore, it is essential to carefully develop, evaluate and use relevant methods to assess the environmental balance of lignocellulosic crops (2nd generation biofuels).

Even if they all demonstrate a net reduction in GHG emissions, LCA of 2nd generation biofuels do not present any common methodology (Cherubini and Stromman, 2011). Main differences during the crop cycle are due to the generic factors used to calculate GHG emissions and the lack of consideration for land use change. A multi-scale approach is required to fulfill a complete LCA. Indeed, GHG emissions are strongly bound with local pedo-climatic conditions and technological options (especially N application rates). This approach is used to estimate direct GHG emissions, downstream indirect GHG emissions and emissions due to land use change.

Scope: The LCA is carried out for a lignocellulosic bio-ethanol industrial unit, located in France, supplied by a mixture of feedstocks: annual and dedicated crops (triticale and fiber sorghum), perennial crops (miscanthus and switchgrass), forest and crop residues (straw) and possibly short rotation coppice (SRC). The study focuses mainly on a supply area around the industrial unit located in the Champagne-Ardenne region. However, we aim at developing generic methods.

Material and methods: The agro-ecosystem model CERES-EGC (Gabrielle *et al.*, 2006) is used to simulate GHG emissions at plot scale (i.e., a few hectares). This model calculates direct emissions in relation to local soil and climate conditions.

Downstream indirect emissions are simulated from the NitroScape model (Duret *et al.* 2011), that accounts for hydrological and atmospheric transfers of reactive nitrogen between landscape elements (e.g., plots, farm buildings). That model works at a scale of typically a small watershed or a few (20-30) square kilometers and make it possible to estimate uncertainties made when considering only direct emissions.

The CERES-EGC model will be possibly used at regional scale and combined with prospective scenarios of land use change from conventional crops towards lignocellulosic crops.

Results and discussion: The methodological approach for assessing LCA is currently under development. Primary results are expected for the end of 2011 (November). They should present LCA simulated at farm gate for a mixture of feedstock and integrated at plot and regional scales.

Conclusion: This method is useful to reduce uncertainties in estimates of GHG emission, one of the steps that have important impact in biofuel LCA. Moreover it will be applied in the French project FUTUROL to assess the sustainability of a project of bio-ethanol production plant.

References

- Cherubini F. and Stromman A. H. (2011), Life Cycle assessment of bioenergy systems: State of the art and future challenges, *Bioresource Technology*, 102 (2), 437-451.
- Directive, 2009/28/CE of the European Parliament and of the Council, OJ L 140/16.
- Duret S., Drouet J.L., Durand P., Hutchings N.J., Theobald M.R., Salmon-Monviola J., Dragosits U., Maury O., Sutton M.A., and Cellier P. (2010), NitroScape: A model to integrate nitrogen transfers and transformations in rural landscapes, *Environmental Pollution*, In press, available on line 2 July 2011.
- Gabrielle B., Laville P., Hénault C., Nicoulaud B. and Germon J.C. (2006), Simulation of N₂O emissions from wheat-cropped soils using CERES, *Nutrient Cycling in Agroecosystems*, 74 (2), 133-146.
- IPCC (2007), The Physical Science Basis, 4th Assessment Report, Chapt. 7, table 7.7, Cambridge University Press, Cambridge, UK, and New York, USA.