



HAL
open science

Food sensory characteristics: their unconsidered roles in the feeding behaviour of domestic ruminants

Angélique Favreau-Peigné, René Baumont, Cécile Ginane

► To cite this version:

Angélique Favreau-Peigné, René Baumont, Cécile Ginane. Food sensory characteristics: their unconsidered roles in the feeding behaviour of domestic ruminants. *Animal*, 2013, 7 (5), pp.806-813. 10.1017/S1751731112002145 . hal-01004443

HAL Id: hal-01004443

<https://hal.science/hal-01004443>

Submitted on 28 May 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Food sensory characteristics: their unconsidered roles in the feeding behaviour of domestic ruminants

A. Favreau-Peigné^{1,2,3,4†}, R. Baumont^{1,2} and C. Ginane^{1,2}

¹INRA, UMR1213 Herbivores, F-63122 Saint-Genès-Champanelle, France; ²VetAgroSup, F-63370 Lempdes, France; ³INRA, UMR791 Modélisation Systémique Appliquée aux Ruminants, F-75005 Paris, France; ⁴AgroParisTech, UMR Modélisation Systémique Appliquée aux Ruminants, F-75005 Paris, France

(Received 5 March 2012; Accepted 24 August 2012; First published online 6 December 2012)

When domestic ruminants are faced with food diversity, they can use pre-ingestive information (i.e. food sensory characteristics perceived by the animal before swallowing the food) and post-ingestive information (i.e. digestive and metabolic consequences, experienced by the animal after swallowing the food) to evaluate the food and make decisions to select a suitable diet. The concept of palatability is essential to understand how pre- and post-ingestive information are interrelated. It refers to the hedonic value of the food without any immediate effect of post-ingestive consequences and environmental factors, but with the influence of individual characteristics, such as animal's genetic background, internal state and previous experiences. In the literature, the post-ingestive consequences are commonly considered as the main force that influences feeding behaviour whereas food sensory characteristics are only used as discriminatory agents. This discriminatory role is indeed important for animals to be aware of their feeding environment, and ruminants are able to use their different senses either singly or in combination to discriminate between different foods. However, numerous studies on ruminants' feeding behaviour demonstrate that the role of food sensory characteristics has been underestimated or simplified; they could play at least two other roles. First, some sensory characteristics also possess a hedonic value which influences ruminants' intake, preferences and food learning independently of any immediate post-ingestive consequences. Further, diversity of food sensory characteristics has a hedonic value, as animals prefer an absence of monotony in food sensory characteristics at similar post-ingestive consequences. Second, some of these food sensory characteristics become an indicator of post-ingestive consequences after their initial hedonic value has acquired a positive or a negative value via previous individual food learning or evolutionary processes. These food sensory characteristics thus represent cues that could help ruminants to anticipate the post-ingestive consequences of a food and to improve their learning efficiency, especially in complex environments. This review then suggests that food sensory characteristics could be of importance to provide pleasure to animals, to increase palatability of a food and to help them learn in complex feeding situations which could improve animal welfare and productivity.

Keywords: hedonic value, herbivore, food learning, palatability, sensory cues

Implications

Food sensory characteristics have a more important role than generally considered in the feeding behaviour of domestic ruminants. Producers could thus adapt their feeding management to influence preferences and palatability of food. Then, productivity (e.g. by increasing the acceptability of a new food) and animal welfare (e.g. by inducing pleasant stimuli) could be improved.

Introduction

At pasture, ruminants are faced with varied feeding situations, from monocultures to highly diversified permanent pastures. At pasture, they encounter a great variability of

food items that differ in quality and availability, depending on the vegetal species present, their phenological stages and their general state (height, humidity, etc.). In housed systems, ruminants are faced to a lesser extent with food diversity since their diets are generally constituted by one or two types of forages and concentrates. Even when these different components are offered in a complete mixed diet, food diversity can play a role because of animals' abilities to select food (Dumont, 1997).

As soon as they have choice opportunities, ruminants have to make decisions to select an adequate diet in quality and quantity. Animals can rely on two types of available information to evaluate foods: (i) pre-ingestive information, that is, food sensory characteristics perceived by the animal before swallowing the food, and (ii) post-ingestive information, that is, digestive and metabolic consequences felt by

[†] E-mail: angelique.favreau@agroparistech.fr

the animal after swallowing the food. This information is then integrated by the animal to evaluate food's palatability.

Palatability is a key concept to understand how pre- and post-ingestive information interact, but it encounters many definitions in the literature (Baumont, 1996; Forbes, 2010). In Merriam-Webster dictionary, a palatable food is described as agreeable to the palate or taste. Greenhalgh and Reid (1971) suggested that palatability reflects those characteristics of a food that invoke a sensory response in the animal. For Rogers (1990), palatability is a hypothetical construct which is needed to account for the hedonic value of food sensory characteristics; palatability is influenced by innate factors and can also be modified by learning. Studies concerning the food learning theory have proved that ruminants, as other animals, can associate sensory characteristics and post-ingestive consequences during food learning processes (Provenza, 1995; Forbes and Provenza, 2000). They can learn both positive (Villalba and Provenza, 1997; Duncan and Young, 2002) and negative (Zahorik *et al.*, 1990) consequences when presented experimentally with simple conditioning procedures (i.e. one pre- and post-ingestive association at a time). Palatability is also influenced by the genetic background of the animal and its internal state (i.e. alliesthesia; Cabanac, 1979; Brondel and Cabanac, 2007), as well as by environmental conditions and social context (Forbes, 2010). As a result, the concept of palatability integrates various components: (i) an 'intrinsic or pre-conditioned palatability', that is, innate like or dislike attributable to the sensory properties of the food as the result of evolution and/or *in utero* pre-conditioning, (ii) a 'learned or conditioned palatability', that is, response to food based on prior associations between its sensory characteristics and its post-ingestive consequences and (iii) an 'organismic or reconditioned palatability', that is, acclimatization to a new state, including changes in the physiology of the animal (e.g. deprivation), in the nutritive value of the food or in some aspects of the environment (Kissileff, 1990; Forbes, 2010). As there are many methodological difficulties in separating the different components of the whole palatability concept, the term palatability will hereafter refer to the sensory pleasantness of food without any immediate effect of post-ingestive consequences and environmental factors, but with the probable effects of internal-state signals, genetic background and previous food learning.

The concept of palatability, which refers to the hedonic value of food sensory characteristics, appears to be inter-related with the post-ingestive consequences (Provenza, 1995) or even totally calibrated by them (Garcia, 1989). In studies on food preferences and aversions, it is normal to consider the post-ingestive consequences as the main force that influences preferences, and to relegate the pre-ingestive stimuli to the simple status of discriminatory agents. Sensory characteristics are actually important for animals to discriminate between various food items as this review will show, focusing on the different senses involved in this essential task.

However, sensory motivation to eat can override post-ingestive signals in some cases (Baumont *et al.*, 1990a), sensory characteristics can induce preferences in the absence of any immediate post-ingestive consequences (e.g. umami taste; Gherardi and Black, 1991), and ruminants show difficulties to learn in complex experimental feeding situation where the food sensory characteristics are just used as discriminating tools. It thus seems that the role of sensory characteristics in ruminants' feeding behaviour have been underestimated and is more important than previously described in the literature. As a consequence, this review aims to emphasize some of the other roles of food sensory characteristics in the feeding behaviour of domestic ruminants.

Sensory characteristics: a way to discriminate between foods

The unique role that is consensually assigned to food sensory characteristics is the one that allows animals to be aware of their feeding environment and to discriminate between different food items using their different senses, either singly or in combination. Animals will use one sense or another depending on its distance with the food. Sight and smell are supposed to help animals in making a first sensory evaluation of food without any direct contact. Touch and taste are in use as soon as there is a direct contact between the lips and mouth of the animal and the plant. Taste and smell are, however, often mixed together resulting in a global stimulation of the oro-pharyngeal area, called flavour.

The sense of sight is supposed to be mainly used to orientate herbivores at pasture by a discrimination process at a large scale (Arnold, 1966a). At a smaller scale, sight can also allow them to discriminate between objects of different brightness (Bazely and Ensor, 1989) and colours (Kendrick and Baldwin, 1986), swards differing in height (Bazely, 1990), plant species (i.e. clover and ryegrass, Edwards *et al.*, 1997) and different clover polymorphisms by recognizing white leaf marks (Cahn and Harper, 1976). Furthermore, Arnold (1966a) showed that the diets selected by blinkered sheep were of lower quality than those of control sheep. Thus, sight is used by sheep to discriminate and identify foods, and this affects their diet selection to some extent.

The sense of touch is also of importance since grazing animals generally select against rough, harsh, sticky or spiny material (Baumont, 1996; Vallentine, 2001). Conversely, goats can overcome physical defences of certain trees because they have hard mouth parts that are unaffected by spines and thorns (Cheeke and Dierenfeld, 2010). Krueger *et al.* (1974) determined that touch appeared primarily to supplement the sense of taste in selection of plant species, but plants selected, at least in part, on the basis of touch were all categorized as 'low' in coarseness (e.g. mountain knotweed, mountain sorrel). Physical characteristics and the sense of touch play a role in the acceptability of some food items as it is the case for various strains of *Bromus* grasses (Arnold, 1966b).

Taste is the last discriminating sense used by the animal before swallowing a food. It refers to the five basic tastes, namely the sweet, salty, bitter and sour tastes, incremented recently by the umami taste (Lindemann, 2001). Various behavioural methods have been used to demonstrate that ruminants perceive and discriminate among tastes (Ginane *et al.*, 2011). For instance, two-bottle choice tests have shown that sheep, goats and cattle are sensitive to sweet, salty, sour and bitter tastes (Goatcher and Church, 1970c; Krueger *et al.*, 1974). These sensitivities, measured as the lowest concentration discriminated by the animals, differed both between species (e.g. cattle > goats > sheep for the sweet taste) and between tastes within species (i.e. sweet < salty < sour < bitter for all studied ruminant species) (Goatcher and Church, 1970a and 1970c). Sensitivities for bitter taste vary with the grazer/browser dichotomy: grazers (domestic ruminants who eat grass, e.g. cattle and sheep) are more sensitive to bitter taste than browsers (domestic ruminants who also eat forbs, shrubs and trees, e.g. goats; Glendinning, 1994). It was supposed that browsers have evolved to be less sensitive to bitter taste because they commonly encounter toxic plants that have a bitter taste (Glendinning, 1994). The umami taste, induced by monosodium glutamate, is perceived by sheep since (i) it increased intake of sham-fed animals (Colucci and Grovum, 1993) and (ii) it induced preferences when added to hay compared with the same untreated hay (Gherardi and Black, 1991). Recent studies, using either a non-caloric flavour or monosodium glutamate to induce the umami taste, showed that lambs are able to perceive umami taste in a feed, as they increased their preference for this feed (Favreau *et al.*, 2010a; Villalba *et al.*, 2011; Bach *et al.*, 2012).

Compared with taste, the sense of smell is supposed to be of lower importance in diet selection (Krueger *et al.*, 1974). As it is difficult to differentiate between animals' appreciation of smell (from volatile components of the food) and taste (from soluble components) because they both mix and form a flavour once the food enters the mouth (Forbes and Mayes, 2002), smell is often considered to supplement the sense of taste. Natural or artificial flavours are frequently used in conditioning experiments by adding them to a common food so as to create distinct foods from the animals' sensory perception (Provenza *et al.*, 1996; Favreau *et al.*, 2010b). Many experiments have thus shown that sheep discriminate between different artificial flavours, such as between orange and aniseed (Kyriazakis *et al.*, 1997), onion and coconut (Villalba *et al.*, 1999) or between onion flavour at different concentrations (Villalba and Provenza, 2000). Sheep and goats were also shown to discriminate many different flavours (maple, strawberry, onion, etc.) from the same untreated food (Robertson *et al.*, 2006). Only a few experiments have used natural odours, such as plant aromatic extract or natural compounds (e.g. tannic acid, cedarwood oil, etc.), and these showed that ruminants are able to discriminate them by comparing intakes of normal *v.* anosmic sheep (Arnold *et al.*, 1980) or by submitted goats to cafeteria trials (Rosa *et al.*, 2002).

Sensory characteristics: a way to get pleasure

From neuroscientists' point of view, palatability corresponds to the 'liking' or hedonic component of the brain reward system (Berridge, 1996), which deals with affective responses to the food and induces pleasure (Berridge and Kringelbach, 2008).

How to assess the palatability of food sensory characteristics?

A non-invasive and simple way to assess the hedonic value of food sensory characteristics is to observe the animals' behavioural responses to these sensory characteristics, without any immediate effect of post-ingestive consequences.

This can be achieved by (i) measuring short-term preferences and/or intake since it can be assumed that the post-ingestive consequences of the food have not been felt by the animal yet (Favreau *et al.*, 2010c), (ii) by using an ingredient able to modify the sensory properties of the food without adding any post-ingestive consequence (e.g. a calorie-free sweetener such as saccharin; Hellekant *et al.*, 1994), (iii) by offering the different food items simultaneously since it increases the difficulty for the animal to identify which food item is responsible for which consequence (Duncan and Young, 2002; Favreau *et al.*, 2010b), and/or (iv) by checking, with a conditioning procedure, that the intake of the added sensory modifier does not cause any post-ingestive consequence at the dose rate involved in the study (Gherardi and Black, 1991; Favreau *et al.*, 2010a).

Two methods using fistulated animals could also be used to assess the hedonic value of food sensory characteristics by avoiding any influence of immediate post-ingestive consequences on animals' behavioural responses to the food. The first one is the sham-feeding method which allows the oro-pharyngeal stimulation of the animal while ingesting the food but prevents the food ingested by the animal from entering into the rumen thanks to the presence of an oesophageal fistula (Grovum and Chapman, 1988; Colucci and Grovum, 1993). The second method consists in controlling both the oro-pharyngeal stimuli and the post-ingestive consequences of the diet. It can be achieved by offering one food to be orally consumed (o) while the same or another food is introduced in the rumen of the animal via a rumen fistula (r) in an equal amount to that of the food orally consumed. The palatability of the two tested food (A and B) could then be assessed by comparing feeding behaviour in reversed situations (i.e. comparison of Ao/Br and Bo/Ar situations), where the food sensory characteristics are different, but the diet composition and the post-ingestive consequences are similar (Greenhalgh and Reid, 1971; Favreau *et al.*, 2010c).

Operant conditioning can also be a way to assess food palatability. It consists of positive or negative reinforcements applied to the animal after it performs the appropriate behaviour. The animal thus decides to act or not in order to gain a reward or to avoid a punishment. Then, the amount of work an animal is prepared to perform in order to obtain food is

used to assess its palatability. Sheep were thus trained to press a panel to receive a food reward which allowed Hutson and Van Mourik (1981) to show that the highest level of response was obtained for barley and wheat over peas, lucerne, lupins and oats. Animals were also trained to walk in order to obtain a good-quality food while a poor-quality food is freely available; the longer distance an animal is prepared to walk, the more the good-quality food is preferred (Dumont *et al.*, 1998). Some studies used the walking distance index to determine how food palatability is increased by food restriction (Schütz *et al.*, 2006; Verbeek *et al.*, 2011).

Does the palatability of food sensory characteristics affect intake and feeding pattern?

Using two good-quality forages (lucerne and grass hays) with different sensory characteristics, daily dry matter intake did not differ when controlling for post-ingestive consequences (Favreau *et al.*, 2010c; Figure 1). Greenhalgh and Reid (1971) found similar results when offering two good-quality forages (dried grass and meadow hay), whereas they demonstrated that sensory characteristics influenced the daily intake of a low-quality forage (oat straw) compared with a good-quality forage (dried grass; Figure 1).

These differences could be explained by an insufficient sensory contrast between the better-quality forages used in the first two studies which cannot influence daily intake regardless of post-ingestive consequences. In return, when the contrast is stronger, and even if previous knowledge about the respective post-ingestive consequences of the foods has surely participated in shaping their palatability, it appears that foods' palatability can affect daily intake.




Figure 1 Voluntary intake (mean \pm s.e.) measured in three studies involving the same methodology consists in offering one food at the trough to be orally consumed *ad libitum* (indicated on the upper part of the bars) while the same or another food is introduced in the rumen of the animal via a rumen fistula (indicated in the lower part of the bars). The first experiment (Greenhalgh and Reid, 1971) used a good-quality forage (dried grass = Dg, 13.9% of CP in dry matter) and a low-quality forage (oat straw = Os, 3.9% CP). The second (Greenhalgh and Reid, 1971) and third (Favreau *et al.*, 2010c) experiments used two good-quality forages (meadow hay = Mh, 7.7% CP v. dried grass = Dg, 13.9% CP; lucerne hay = Lh, 15.6% CP v. grass hay = Gh, 7.2% CP, respectively).

Food palatability can also influence intake in the short term. It has been shown that satiated sheep ate a substantial meal after a second distribution of a sufficiently palatable hay; this meal was then associated with an increase in rumen fill of up to 10% of the daily maximum observed without this second distribution (Baumont *et al.*, 1990b). The hedonic stimulation induced by the new distribution thus overrode the satiety signals due to rumen fill. In the same way, moderate energy preloads, combined with manipulations of palatability during a test meal, resulted in overconsumption in humans, thus demonstrating that the hedonic value of food sensory characteristics decreased the ability of short-term satiety cues to stop intake (Yeomans *et al.*, 2001). In sheep, when post-ingestive consequences were controlled, the intake of lucerne hay was higher than that of grass hay just after food distribution, indicating that the motivation to eat was then higher than the negative fill effect of the hay introduced in the rumen (Favreau *et al.*, 2010c). Finally, the physical form (pellets or cubes) and the addition of monosodium glutamate (i.e. umami taste) increased the palatability of straw (a poor-quality food) which induced higher sham intakes in the short term (Colucci and Grovum, 1993). On the contrary, a low palatability can decrease the short-term intake of a food in cattle (palm kernel expeller; Spörndly and Åsberg, 2006).

Does the palatability of food sensory characteristics affect preferences?

As soon as ruminants have choice opportunities, they express preferences. These preferences are influenced by the effect of post-ingestive consequences (Provenza, 1995), but some results suggest that the relative hedonic value of food sensory characteristics is involved as well. Sensory characteristics were indeed able to induce food preferences by their own, that is, without any immediate positive post-ingestive reinforcement as it is the case with umami taste (Grovum and Chapman, 1988; Gherardi and Black, 1991). The umami taste, induced by monosodium glutamate, seems to have a positive hedonic value in sheep, as it induced strong and constant preferences for the treated food over the same untreated food, regardless of the doses (Gherardi and Black, 1991; Favreau *et al.*, 2010a). In cattle, umami induced such a preference in young animals (e.g. in animals weaned at 3 weeks of age, and from 3 to 6 weeks of age; Waldern and van Dyk, 1971). In the same way, when different flavours are sprayed on the same food (i.e. to ensure similar post-ingestive consequences) and compared in short-term choice tests with the same unflavoured food, animals often express clear preferences. This was the case for sheep preferring an aniseed-flavoured orchard grass hay, at a rate of 80%, compared with an orange-flavoured orchard grass hay (Favreau *et al.*, 2010b), suggesting that the aniseed flavour has a higher hedonic value than the orange flavour in sheep. However, this result is inconsistent with those of other studies in which sheep did not express any preference between the two flavours (Kyriazakis *et al.*, 1997; Arsenos *et al.*, 2000). Such variability was also frequently reported in

the literature concerning sweet taste (see Ginane *et al.*, 2011 for review). Individual variability in preferences between flavours or tastes in the absence of immediate post-ingestive influence are also widely encountered in different experiments (Goatcher and Church, 1970a and 1970b; Robertson *et al.*, 2006). In the absence of immediate post-ingestive consequences that usually calibrate food preferences, it appears logical to encounter a great individual variability since it is known that individual characteristics, such as animal's physiological and emotional states (Brondel and Cabanac, 2007 and Tamashiro *et al.*, 2007, respectively) and previous experiences (Provenza, 1995) *in utero* (Simitzis *et al.*, 2008) or early in life (Miller-Cushon and DeVries, 2010), can affect palatability. Experiences *in utero* and early in life cause neurological, morphological and physiological changes because of the interactions between genome, environment and post-ingestive consequences; this participates in the uniqueness of each individual (Provenza *et al.*, 2003). These differences in preferences can lead to variability in food selection in the field; as a consequence, it can result in less grazing pressure on individual vegetation species, which can represent an evolutionary advantage for social herbivore species (Goatcher and Church, 1970b).

The hedonic value of food sensory diversity

When ruminants are offered a choice between foods that differ in how long they have not been eaten, they generally prefer the one that was not consumed recently. This was true for sheep preferring the flavour presented first within a conditioning period over that presented second (Arsenos *et al.*, 2000), for ewes preferring the opposite species (i.e. white clover or ryegrass) to the one they had previously grazed (Newman *et al.*, 1992; Parsons *et al.*, 1994), and for heifers temporarily increasing their preference for the hay that had not been offered during the previous period (Ginane *et al.*, 2002). The authors suggested that the preference for the 'new' hay could be explained by (i) the animal's need to select a balanced diet that best meets its homeostatic requirements or (ii) the animals' search for rarity for functional purposes, such as maintaining gut flora diversity or (iii) an attractive effect of novelty, considered as a search for diversity. On that last point, Scott and Provenza (1998) demonstrated that a variety of flavours is preferred relative to monotony by lambs when food nutrient content was constant between the different food items available. Similarly, sheep always preferred in the short term the hay they were not used to eating, whatever its nature (lucerne or grass hay), even when the post-ingestive consequences associated with animals' diet were similar (Favreau *et al.*, 2010c). This preference for diversity suggests the existence of a hedonic dimension for food sensory characteristics. This is consistent with the concept of sensory specific satiety, which assumes that the hedonic value of food sensory characteristics will decrease as it is consumed (Rolls, 1986). This is also consistent with recent findings showing that lambs receiving a monotonous diet had higher cortisol levels 1 h after food presentation than lambs receiving a diverse diet,

indicating a stressing effect of monotony (Villalba *et al.*, 2012). Ruminants thus appear to search for diversity for their pleasure as well as for their needs. As a consequence, the absence of food diversity, encountered when animals are conducted outdoors on monoculture or fed indoors with the same diet over several weeks or months, may have negative implications on animal welfare.

Does the palatability of food sensory characteristics affect food learning?

As mentioned above, sheep showed a higher initial preference for aniseed-flavoured orchard grass hay compared with orange-flavoured orchard grass hay (Favreau *et al.*, 2010b). After conditioning, the preference for the flavoured hay associated with the negative post-ingestive consequence was still influenced by these initial preferences, that is, sheep negatively conditioned on aniseed expressed a lesser avoidance than sheep negatively conditioned on orange (0.53 ± 0.04 v. 0.25 ± 0.03 , respectively; Favreau *et al.*, 2010b). Similar results were found with a higher initial hedonic preference for fenugreek flavour over garlic flavour which then influenced the conditioned responses of sheep (Arsenos *et al.*, 2000). Therefore, it appears that the initial palatability of a food influences its evaluation by sheep in a learning context. It then affects the animals' propensity to make an associative learning between sensory characteristics and post-ingestive consequences. It may be easier for animals to associate a food initially preferred with a positive than with a negative consequence.

Sensory characteristics: a way to indicate the nature of the post-ingestive consequences of the food

The condition for a food sensory characteristic to become an informative signal is that animals are able to give it a meaning via evolution and/or by using individual learning processes. When animals encounter a new food and have no knowledge about it, they hesitate or even refuse to eat it. This neophobic response is considered as adaptive, as it helps animals to avoid eating harmful foods at the first encounter, and give them the opportunity to learn about their post-ingestive consequences (Birch, 1999). At the individual learning level, it has been largely demonstrated that ruminants, such as sheep (du Toit *et al.*, 1991; Burritt and Provenza, 1992), cattle (Ralphs and Provenza, 1999) and goats (Duncan and Young, 2002), are able to associate food sensory characteristics with post-ingestive consequences when the experimental learning procedure is simple (i.e. one association between sensory characteristics and post-ingestive consequences to learn at a time), and modify their diet selection accordingly. However, animals show difficulties to learn when the experimental learning procedure becomes more complex and closer to natural situations, which could be achieved experimentally either by offering several different foods simultaneously during the conditioning procedure (Duncan and Young, 2002), by increasing the number of consequences associated with each food (Ginane *et al.*, 2005) or by modifying

the length of feeding bouts and the rate of nutrient delivery (Duncan *et al.*, 2007). In these studies designed to test food-learning abilities in ruminants, food sensory characteristics can only be used as discriminating tools because they are meaningless for the animals at the very first association, that is, they do not indicate to the animal the post-ingestive consequences associated with the food. Diverse sensory characteristics of plants could, however, act as crude indicators of their nutritional and toxic values in the field. For instance, several secondary plant compounds have a bitter taste and are toxic to herbivores (Garcia and Hankins, 1975; Rogosic *et al.*, 2008), and a young green plant is generally more nutritious than a mature plant (Kühnle and Müller, 2011). Further, other species use food sensory characteristics as indicators of their potential toxic or nutritive values, as shown with birds avoiding invertebrates' aposematic colouring (Aronsson and Gamberale-Stille, 2008; Skelhorn *et al.*, 2008).

Edwards *et al.* (1997) demonstrated that sheep could use visual and smelling cues (in turves of perennial ryegrass or white clover) to find rewards (highly preferred pelleted food). Sheep expressed an aversion for a familiar food when added with a cue, that is, a novel odour, which was previously associated with toxicosis; this was true for both artificial flavour (coconut) and natural odour (the odour of *Astragalus bisulcatus*; Provenza *et al.*, 2000). In that study, it appears that the sensory characteristic has acquired a meaning, indicating a probable toxicity to the animal which, as a consequence, reduced its preference for the food associated with that cue. Conversely, sensory characteristics can be a positive cue. Food neophobia can be overcome by adding a familiar flavour or odour on a novel food (Launchbaugh *et al.*, 1997; Van Tien *et al.*, 1999). Familiarity can be considered as a cue indicating that the food is safe (Burritt and Provenza, 1989).

Further, a recent study demonstrated that the use of food sensory characteristics (flavours experimentally associated with negative or positive post-ingestive consequences) as indicators of the value of the food allowed sheep to improve their learning efficiency in a subsequent complex learning task compared with naïve sheep (Favreau *et al.*, 2010b). This also underlines their ability to generalize their knowledge about flavours to different situations varying in complexity and confirms previous results (Launchbaugh and Provenza, 1994; Ginane and Dumont, 2006). So sheep can rely on sensory cues, which provide an essential help in learning about food especially in complex environments. Howery *et al.* (2000) demonstrated that cattle can learn to associate visual cues (traffic barricades and traffic cones) with disparate food qualities and use this information to forage more efficiently in environments varying in complexity (fixed or variable locations of foods in the field).

Plants encountered in the field by animals differ in many sensorial characteristics and notably in their taste profiles; some of them are supposed to reveal the value of the food. Thus, they constitute an interesting natural cue to study the potential role of food sensory characteristics as an indicator of post-ingestive consequences. Hence, several plant secondary




Figure 2 Diagram of the interrelated roles of food sensory characteristics and post-ingestive consequences in the feeding behaviour of domestic ruminants and its control. The post-ingestive consequences play a major role in ruminants' feeding behaviour because of their fill effect and/or their nutritive/toxic effect. All sensory characteristics are discriminating characteristics (i.e. they help animals to distinguish between different food items). Some of them possess a hedonic value that influences food learning, intake and preferences at least on the short term. Then, some of them have acquired a positive or a negative value through individual learning or evolutionary processes. This value could play an indicator role that allows ruminants to anticipate the post-ingestive consequences associated with the food consumed, and then improve their food learning efficiency especially in complex environment.

compounds that are toxic to herbivores have a bitter taste (Garcia and Hankins, 1975; Rogosic *et al.*, 2008), whereas sweet and umami tastes are supposed to be associated with energy (Swithers and Davidson, 2008) and protein (Naim *et al.*, 1991) contents, respectively. A recent study demonstrated that the umami taste would have a positive value, expressed by a high initial preference, whereas the bitter taste would have a negative value as shown by the reluctance of sheep to increase their preference for this taste once it is associated with positive consequences (Favreau *et al.*, 2010a). Thus, herbivores seem to attribute *a priori* values to natural basic tastes. Consequently, it is suggested that ruminants could use primary tastes as cues to anticipate food consequences and then to forage both more efficiently and more safely, although this still needs to be tested.

Conclusion

This review brings to light that the roles of food sensory characteristics have been previously underestimated or simplified, and underlines three important roles of food sensory characteristics on ruminants' feeding behaviour (Figure 2). First, all sensory characteristics can be considered as discriminating characteristics that allow animals to distinguish between different food items. Second, some of these sensory characteristics possess a hedonic value that influences ruminants' intake, preferences and food learning independently of any immediate post-ingestive consequences. Third, for some of them, the initial hedonic value has acquired a positive or a negative value either via previous individual

food learning or via evolutionary processes. This value then makes possible for a sensory characteristic to become an indicator that allows ruminants to anticipate the post-ingestive consequences of a food and to improve their learning efficiency, which could be of interest for animals, especially in complex environments.

Perspectives and significance

This review suggests some avenues to follow in order to improve animal welfare and productivity by taking into account these multiple roles of food sensory characteristics. For instance, food sensory characteristics could be used to improve palatability of poor quality or new foods, so as to increase their acceptability, at least in the short term. Further, animals seem to search for an absence of monotony in food sensory characteristics. This could represent an easy way to enrich their environment and improve their welfare. Sensory cues could also be created by animal managers to help animals anticipate the post-ingestive consequences of some toxic plants so that they avoid them in the field.

Acknowledgements

The authors are grateful to INRA and the Auvergne Region for their funding.

References

Arnold GW 1966a. The special senses in grazing animals. I – Sight and dietary habits in sheep. *Australian Journal of Agricultural Research* 17, 523–529.

Arnold GW 1966b. The special senses in grazing animals. II – Smell, taste, and touch and dietary habits in sheep. *Australian Journal of Agricultural Research* 17, 531–542.

Arnold GW, De Boer ES and Boundy CAP 1980. The influence of odour and taste on the food preferences and food intake of sheep. *Australian Journal of Agricultural Research* 31, 571–587.

Aronsson M and Gamberale-Stille G 2008. Domestic chicks primarily attend to colour, not pattern, when learning an aposematic coloration. *Animal Behaviour* 75, 417–423.

Arsenos G, Hills J and Kyriazakis I 2000. Conditioned feeding responses of sheep towards flavoured foods associated with casein administration: the role of long delay learning. *Animal Science* 70, 157–169.

Bach A, Villalba JJ and Ipharraguerre IR 2012. Interactions between mild nutrient imbalance and taste preferences in young ruminants. *Journal of Animal Science* 90, 1015–1025.

Baumont R 1996. Palatability and feeding behaviour in ruminants. A review. *Annales de Zootechnie* 45, 385–400.

Baumont R, Malbert CH and Ruckebusch Y 1990a. Mechanical stimulation of rumen fill and alimentary behaviour in sheep. *Animal Production* 50, 123–128.

Baumont R, Segulier N and Dulphy JP 1990b. Rumen fill, forage palatability and alimentary behaviour in sheep. *Journal of Agricultural Science* 115, 277–284.

Bazely DR 1990. Rules and cues used by sheep foraging in monocultures. In *Behavioural mechanisms of food selection* (ed. RN Hugues), pp. 343–367. Springer-Verlag, Berlin Heidelberg.

Bazely DR and Ensor CV 1989. Discrimination learning in sheep with cues varying in brightness and hue. *Applied Animal Behaviour Science* 23, 293–299.

Berridge KC 1996. Food reward: brain substrates of wanting and liking. *Neuroscience & Biobehavioral Reviews* 20, 1–25.

Berridge KC and Kringelbach M 2008. Affective neuroscience of pleasure: reward in humans and animals. *Psychopharmacology* 199, 457–480.

Birch LL 1999. Development of food preferences. *Annual Review of Nutrition* 19, 41–62.

Brondel L and Cabanac M 2007. Alliesthesia in visual and auditory sensations from environmental signals. *Physiology & Behavior* 91, 196–201.

Burritt EA and Provenza FD 1989. Food aversion learning: ability of lambs to distinguish safe from harmful foods. *Journal of Animal Science* 67, 1732–1739.

Burritt EA and Provenza FD 1992. Lambs form preferences for nonnutritive flavors paired with glucose. *Journal of Animal Science* 70, 1133–1136.

Cabanac M 1979. Sensory pleasure. *The quarterly review of biology* 54, 1–29.

Cahn MG and Harper JL 1976. The biology of the leaf mark polymorphism in *Trifolium repens* L. *Heredity* 37, 327–333.

Cheeke PR and Dierenfeld ES 2010. *Comparative animal nutrition and metabolism*. CABI, Cambridge, UK.

Colucci PE and Grovum WL 1993. Factors affecting the voluntary intake of food by sheep. 6. The effect of monosodium glutamate on the palatability of straw diets by sham-fed and normal animals. *British Journal of Nutrition* 69, 37–47.

du Toit JT, Provenza FD and Nassis A 1991. Conditioned taste aversions: how sick must a ruminant get before it learns about toxicity in foods? *Applied Animal Behaviour Science* 30, 35–46.

Dumont B 1997. Diet preferences of herbivores at pasture. *Annales de Zootechnie* 46, 105–116.

Dumont B, Dutronc A and Petit M 1998. How readily will sheep walk for a preferred forage? *Journal of Animal Science* 76, 965–971.

Duncan AJ and Young SA 2002. Can goats learn about foods through conditioned food aversions and preferences when multiple food options are simultaneously available? *Journal of Animal Science* 80, 2091–2098.

Duncan AJ, Elwert C, Villalba JJ, Yearsley J, Pouloupoulou I and Gordon IJ 2007. How does pattern of feeding and rate of nutrient delivery influence conditioned food preferences? *Oecologia* 153, 617–624.

Edwards GR, Newman JA, Parsons AJ and Krebs JR 1997. Use of cues by grazing animals to locate food patches: an example with sheep. *Applied Animal Behaviour Science* 51, 59–68.

Favreau A, Baumont R, Ferreira G, Dumont B and Ginane C 2010a. Do sheep use umami and bitter tastes as cues of post-ingestive consequences when selecting their diet? *Applied Animal Behaviour Science* 125, 115–123.

Favreau A, Baumont R, Duncan AJ and Ginane C 2010b. Sheep use preingestive cues as indicators of postingestive consequences to improve food learning. *Journal of Animal Science* 88, 1535–1544.

Favreau A, Ginane C and Baumont R 2010c. Feeding behaviour of sheep fed lucerne v. grass hays with controlled post-ingestive consequences. *Animal* 4, 1368–1377.

Forbes JM 2010. *Palatability: principles, methodology and practice for farm animals*. CAB Reviews: Perspectives in Agriculture, Veterinary Science, Nutrition and Natural Resources 5, 1–15.

Forbes JM and Provenza FD 2000. Integration of learning and metabolic signals into a theory of dietary choice and food intake. In *Ruminant physiology: digestion, metabolism, growth and reproduction* (ed. PB Cronje), pp. 3–19. CAB International, Wallingford.

Forbes JM and Mayes RW 2002. Food choice. In *Sheep nutrition* (ed. M Freer and H Dove), pp. 51–69. CAB international, Wallingford.

Garcia J 1989. Food for Tolman: cognition and cathexis in concert. In *Aversion, avoidance, and anxiety: perspectives on aversively motivated behavior* (ed. T Archer and LG Nilsson), pp. 45–85. Lawrence Erlbaum Associates, Hillsdale, NJ, USA.

Garcia J and Hankins WG 1975. The evolution of bitter and the acquisition of toxiphobia. In *Olfaction and taste: 5th International Symposium* (ed. D Denton and J Coghlan), pp. 39–45. Academic Press, New York.

Gherardi SG and Black JL 1991. Effect of palatability on voluntary feed intake by sheep. I – Identification of chemicals that alter the palatability of a forage. *Austrian Journal of Agricultural Research* 42, 571–584.

Ginane C and Dumont B 2006. Generalization of conditioned food aversions in grazing sheep and its implication for food categorization. *Behavioral Processes* 73, 178–186.

Ginane C, Baumont R and Favreau-Peigné A 2011. Perception and hedonic value of basic tastes in domestic ruminants. *Physiology & Behavior* 104, 666–674.

Ginane C, Baumont R, Lassalas J and Petit M 2002. Feeding behaviour and intake of heifers fed on hays of various quality, offered alone or in a choice situation. *Animal Research* 51, 177–188.

Ginane C, Duncan AJ, Young SA, Elston DA and Gordon IJ 2005. Herbivore diet selection in response to simulated variation in nutrient rewards and plant secondary compounds. *Animal Behaviour* 69, 541–550.

- Glendinning JI 1994. Is the bitter taste rejection always adaptive? *Physiology & Behavior* 56, 1217–1227.
- Goatcher WD and Church DC 1970a. Taste responses in ruminants. III. Reactions of pygmy goats, normal goats, sheep and cattle to sucrose and sodium chloride. *Journal of Animal Science* 31, 364–372.
- Goatcher WD and Church DC 1970b. Taste responses in ruminants. IV. Reactions of pygmy goats, normal goats, sheep and cattle to acetic acid and quinine hydrochloride. *Journal of Animal Science* 31, 373–382.
- Goatcher WD and Church DC 1970c. Review of some nutritional aspects of the sense of taste. *Journal of Animal Science* 31, 973–981.
- Greenhalgh JFD and Reid GW 1971. Relative palatability to sheep of straw, hay and dried grass. *British Journal of Nutrition* 26, 107–116.
- Grovum WL and Chapman HW 1988. Factors affecting the voluntary intake of food by sheep. 4. The effect of additives representing the primary tastes on sham intakes by oesophageal-fistulated sheep. *British Journal of Nutrition* 59, 63–72.
- Hellekant G, Hård Af Segerstad C and Roberts TW 1994. Sweet taste in the calf. III. Behavioral responses to sweeteners. *Physiology & Behavior* 56, 555–562.
- Howery LD, Bailey DW, Ruyle GB and Renken WJ 2000. Cattle use visual cues to track food locations. *Applied Animal Behaviour Science* 67, 1–14.
- Hutson GD and Van Mourik SC 1981. Food preferences of sheep. *Australian Journal of Experimental Agriculture and Animal Husbandry* 21, 575–582.
- Kendrick KM and Baldwin BA 1986. The activity of neurones in the lateral hypothalamus and zona incerta of the sheep responding to the sight or approach of food is modified by learning and satiety and reflects food preference. *Brain Research* 375, 320–328.
- Kissileff HR 1990. Some suggestions on dealing with palatability – response to Ramirez. *Appetite* 14, 162–166.
- Krueger WC, Laycock WA and Price DA 1974. Relationship of taste, smell, sight, and touch to forage selection. *Journal of Range Management* 27, 258–262.
- Kühnle A and Müller C 2011. Relevance of visual and olfactory cues for host location in the mustard leaf beetle *Phaedon cochleariae*. *Physiological Entomology* 36, 68–76.
- Kyriazakis I, Papachristou TG, Duncan AJ and Gordon IJ 1997. Mild conditioned food aversions developed by sheep towards flavors associated with plant secondary compounds. *Journal of Chemical Ecology* 23, 727–746.
- Launchbaugh KL and Provenza FD 1994. The effect of flavor concentration and toxin dose on the formation and generalization of flavor aversions in lambs. *Journal of Animal Science* 72, 10–13.
- Launchbaugh KL, Provenza FD and Werkmeister MJ 1997. Overcoming food neophobia in domestic ruminants through addition of a familiar flavor and repeated exposure to novel foods. *Applied Animal Behaviour Science* 54, 327–334.
- Lindemann B 2001. Receptors and transduction in taste. *Nature* 413, 219–225.
- Miller-Cushon EK and DeVries TJ 2010. Effect of early feed type exposure on diet-selection behavior of dairy calves. *Journal of Dairy Science* 94, 342–350.
- Naim M, Ohara I, Kare MR and Levinson M 1991. Interaction of MSG taste with nutrition – perspectives in consummatory behavior and digestion. *Physiology & Behavior* 49, 1019–1024.
- Newman JA, Parsons AJ and Harvey A 1992. Not all sheep prefer clover: diet selection revisited. *Journal of Agricultural Science* 119, 275–283.
- Parsons AJ, Newman JA, Penning PD, Harvey A and Orr RJ 1994. Diet preference of sheep: effects of recent diet, physiological state and species abundance. *Journal of Animal Ecology* 63, 465–478.
- Provenza FD 1995. Postingestive feedback as an elementary determinant of food preference and intake in ruminants. *Journal of Range Management* 48, 2–17.
- Provenza FD, Kimball BA and Villalba JJ 2000. Roles of odor, taste, and toxicity in the food preferences of lambs: implications for mimicry in plants. *Oikos* 88, 424–432.
- Provenza FD, Scott CB, Phy TS and Lynch JJ 1996. Preference of sheep for foods varying in flavors and nutrients. *Journal of Animal Science* 74, 2355–2361.
- Provenza FD, Villalba JJ, Dziba LE, Atwood SB and Banner RE 2003. Linking herbivore experience, varied diets, and plant biochemical diversity. *Small Ruminant Research* 49, 257–274.
- Ralps MH and Provenza FD 1999. Conditioned food aversions: principles and practices, with special reference to social facilitation. *Proceedings of the Nutrition Society* 58, 813–820.
- Robertson E, Gordon IJ and Pérez-Barbería FJ 2006. Preferences of sheep and goats for straw pellets treated with different food-flavouring agents. *Small Ruminant Research* 63, 50–55.
- Rogers PJ 1990. Why a palatability construct is needed. *Appetite* 14, 167–170.
- Rogosic J, Estell RE, Ivankovic S, Kezic J and Razov J 2008. Potential mechanisms to increase shrub intake and performance of small ruminants in mediterranean shrubby ecosystems. *Small Ruminant Research* 74, 1–15.
- Rolls BJ 1986. Sensory-specific satiety. *Nutrition Reviews* 44, 93–101.
- Rosa Gd, Moio L, Napolitano F, Grasso F, Gubitosi L and Bordi A 2002. Influence of flavor on goat feeding preferences. *Journal of Chemical Ecology* 28, 269–281.
- Schütz K, Davison D and Matthews L 2006. Do different levels of moderate feed deprivation in dairy cows affect feeding motivation? *Applied Animal Behaviour Science* 101, 253–263.
- Scott LL and Provenza FD 1998. Variety of foods and flavors affects selection of foraging location by sheep. *Applied Animal Behaviour Science* 61, 113–122.
- Simitzis PE, Bizelis JA, Deligeorgis SG and Feggeros K 2008. Effect of early dietary experiences on the development of feeding preferences in semi-intensive sheep farming systems – a brief note. *Applied Animal Behaviour Science* 111, 391–395.
- Skelhorn J, Griksaitis D and Rowe C 2008. Colour biases are more than a question of taste. *Animal Behaviour* 75, 827–835.
- Spöndly E and Åsberg T 2006. Eating rate and preference of different concentrate components for cattle. *Journal of Dairy Science* 89, 2188–2199.
- Swithers SE and Davidson TL 2008. A Role for sweet taste: calorie predictive relations in energy regulation by rats. *Behavioral Neuroscience* 122, 161–173.
- Tamashiro KL, Hegeman MA, Nguyen MMN, Melhorn SJ, Ma LY, Woods SC and Sakai RR 2007. Dynamic body weight and body composition changes in response to subordination stress. *Physiology & Behavior* 91, 440–448.
- Vallentine JF 2001. *Grazing management*. Academic Press, San Diego, USA.
- Van Tien D, Lynch JJ, Hinch GN and Nolan JV 1999. Grass odor and flavor overcome feed neophobia in sheep. *Small Ruminant Research* 32, 223–229.
- Verbeek E, Waas JR, McLeay L and Matthews LR 2011. Measurement of feeding motivation in sheep and the effects of food restriction. *Applied Animal Behaviour Science* 132, 121–130.
- Villalba JJ and Provenza FD 1997. Preference for wheat straw by lambs conditioned with intraruminal infusions of starch. *British Journal of Nutrition* 77, 287–297.
- Villalba JJ and Provenza FD 2000. Roles of flavor and reward intensities in acquisition and generalization of food preferences: do strong plant signals always deter herbivory? *Journal of Chemical Ecology* 26, 1911–1922.
- Villalba JJ, Provenza FD and Rogosic J 1999. Preference for flavored wheat straw by lambs conditioned with intraruminal infusions of starch administered at different times after straw ingestion. *Journal of Animal Science* 77, 3185–3190.
- Villalba JJ, Bach A and Ipharraguerre IR 2011. Feeding behaviour and performance of lambs are influenced by flavor diversity. *Journal of Animal Science* 89, 2571–2581.
- Villalba JJ, Catanese F, Provenza FD and Distel RA 2012. Relationships between early experience to dietary diversity, acceptance of novel flavors, and open field behavior in sheep. *Physiology & Behavior* 105, 181–187.
- Waldern DE and van Dyk RD 1971. Effect of monosodium glutamate in starter rations on feed consumption and performance of early weaned calves. *Journal of Dairy Science* 54, 262–265.
- Yeomans MR, Lee MD, Gray RW and French SJ 2001. Effects of test-meal palatability on compensatory eating following disguised fat and carbohydrate preloads. *International Journal of Obesity* 25, 1215–1224.
- Zahorik DM, Houpt KA and Swartzman-Andert J 1990. Taste-aversion learning in three species of ruminants. *Applied Animal Behaviour Science* 26, 27–39.