



HAL
open science

Symmetries

Hugo Bacard

► **To cite this version:**

| Hugo Bacard. Symmetries. 2014. hal-01018837v1

HAL Id: hal-01018837

<https://hal.science/hal-01018837v1>

Preprint submitted on 6 Jul 2014 (v1), last revised 8 Jul 2014 (v2)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Symmetries

Hugo V. Bacard *

Western University

July 5, 2014

Abstract

Classical mathematics are founded within set theory, but sets don't have *symmetries*. We conjecture that if we allow sets with symmetries, then many problems such as *Mirror symmetry* or *Homological mirror symmetry* can be explained. One way to do this is to embed sets to higher categories and especially into higher groupoids as already envisioned by Grothendieck, and more recently by Voevodsky. We simply outline this idea in these notes.

«Tout est intéressant !»

Carlos Simpson

1 Symmetries in maths

Given a set $X = \{a, b, c, \dots\}$ such as the natural numbers $\mathbb{N} = \{0, 1, \dots, p, \dots\}$, there is a standard procedure that amounts to regard X as a category with only identity morphisms. This is the *discrete functor* that takes X to the category denoted by $Disc(X)$ where the hom-sets are given by $\text{Hom}(a, b) = \emptyset$ if $a \neq b$, and $\text{Hom}(a, b) = \{Id_a\} = 1$ if $a = b$.

But in category theory, there is also a procedure called *opposite or dual*, that takes a general category \mathcal{C} to its opposite \mathcal{C}^{op} . And the problem is that if we restrict this procedure

*E-mail address: hbacard@uwo.ca

to categories such as $Disc(X)$, there is no way to distinguish $Disc(X)$ from $Disc(X)^{op}$. And this is what we mean by *sets don't show symmetries*. In the program of Voevodsky, we can interpret this by saying that:

'The identity type is not good for sets, instead we should use the Equivalence type'

The problem is that so far we've used set theory with this lack of symmetries, as foundations for mathematics. And some *symmetry phenomenons* occur as we progress in maths, and sometimes we're unable to figure out why exactly.

Grothendieck [7] has already seen this when he moved from sheaves of sets, to sheaves of groupoid (stacks), because he wanted to allow objects to have symmetries (automorphisms). If we look at the Giraud-Grothendieck picture on nonabelian cohomology [6], then what happens is an extension of coefficients $\mathcal{U} : \mathbf{Set} \hookrightarrow \mathbf{Cat}$. But this embedding is too big as we mentioned in [3]. Rather we should consider first the comma category $\mathbf{Cat} \downarrow \mathcal{U}$, whose objects are functors $C \rightarrow Disc(X)$. And then we should consider the full subcategory consisting of functors $C \xrightarrow{\sim} Disc(X)$ that are equivalences of categories. This will force C to be a groupoid, that looks like a set. And we call such $C \xrightarrow{\sim} Disc(X)$ a *Quillen-Segal \mathcal{U} -object*.

This category of Quillen-Segal objects should be called the *category of sets with symmetries*. Following Grothendieck's point of view, we've denoted by $\mathbf{Cat}_{\mathcal{U}}[\mathbf{Set}]$ the comma category, and think of it as *categories with coefficients or coordinates in sets*. This terminology is justified by the fact that the functor $\mathcal{U} : \mathbf{Set} \hookrightarrow \mathbf{Cat}$ is a geometric morphism of (higher) topos. The category of set with symmetries is like the homotopy neighborhood of this point, similar to a one-point going to a disc or any contractible object. The advantage of the Quillen-Segal formalism is the presence of a Quillen model structure on $\mathbf{Cat}_{\mathcal{U}}[\mathbf{Set}]$ such that the fibrant objects are Quillen-Segal objects ([2, Theorem 8.2], [3, Theorem 1.2]).

In standard terminology this means that if we embed a set X in \mathbf{Cat} as $Disc(X)$, and take an 'injective resolution' of it, then we get an equivalence of groupoids $P \xrightarrow{\sim} Disc(X)$, and P has symmetries. Concretely what happen is just a factorization of the identity (type) $\text{Id} : Disc(X) \rightarrow Disc(X)$ as a cofibration followed by a trivial fibration:

$$Disc(X) \hookrightarrow P \xrightarrow{\sim} Disc(X).$$

The first morphism is also automatically an equivalence.

We regard in [3] this process of embedding $\mathbf{Set} \hookrightarrow QS\{\mathbf{Cat}_{\mathcal{U}}[\mathbf{Set}]\}$ as a *minimal homotopical enhancement*. The idea is that there is no good notion of homotopy (weak equivalence) in \mathbf{Set} , but there are at least two notions in \mathbf{Cat} : equivalences of categories and the equivalences of classifying spaces *à la* Grothendieck-Kan-Quillen-Segal-Thomason.

This last class of weak equivalences is important for what we believe happens with *mirror phenomenons*. We isolated the discussion in the next paragraph.

2 Mirrors

Given a compact Kähler manifold Y , we know that the cohomology groups $H^*(Y, \mathbb{C})$ have a Hodge decomposition $H^{p,q}$ (see [5] [10, 11]). Now because we have Poincaré duality, and the comparisons between singular and De Rham cohomologie, we know that any other space Z that has the same homotopy type as Y will have the same cohomology groups. Consequently they will share the same *Hodge diamond*, thus its symmetries.

This means that the symmetry of the Hodge diamond is mostly attached to the homotopy type of Y . This is not surprising anymore because it can already be seen from the equivalence of the De Rham cohomology which is analytic, and the Betti cohomology which is something purely simplicial. In fact, it can also be seen from the (smooth) homotopy invariance of De Rham cohomology.

We believe that this symmetry can be understood using the Quillen-Segal formalism as follows. Given Y , let's consider $Y_{top} \in \mathbf{Top}$. Recall that we have a Quillen equivalence $\mathcal{U} : \mathbf{Top} \rightarrow \mathbf{sSet}_Q$, where $\mathcal{U} = \mathit{Sing}$ is the singular functor whose left adjoint is the geometric realization. When we consider the comma category $\mathbf{sSet}_{Q\mathcal{U}}[\mathbf{Top}] = \mathbf{sSet} \downarrow \mathcal{U}$, we are literally creating in French a “*trait d'union*”, between the two categories. And when we consider the subcategory of Quillen-Segal objects, then our result [3, Theorem 1.2] says that we have a triangle that descends to a triangle of equivalences between the homotopy categories. In fact there is a much better statement.

$$\begin{array}{ccc}
 (\mathbf{sSet}_Q \downarrow \mathcal{U}) & & \\
 \downarrow & \searrow^s & \\
 \mathbf{Top} & \xrightarrow{\mathcal{U}} & \mathbf{sSet}_Q
 \end{array}$$

It turns out that if we apply [3, Theorem 1.2] to the same functor but we choose the Joyal model structure \mathbf{sSet}_J , we get the *Homotopy hypothesis* (see [1]).

A fibrant replacement of Y in the model category $\mathbf{sSet}_{Q\mathcal{U}}[\mathbf{Top}]$, is a trivial fibration $\mathcal{F} \xrightarrow{\sim} \mathcal{U}(Y)$, where \mathcal{F} is fibrant in \mathbf{sSet}_Q , that is a Kan complex. But a Kan complex is exactly an ∞ -groupoid. ∞ -Groupoids generalizes groupoids, and still are category-like. In particular we can take their opposite (or dual), just like we consider the opposite category \mathcal{C}^{op} of a usual category, as outlined in the beginning.

Conjecture 2.1. *Given Y as above, we can think of the mirror of Y as the opposite ∞ -groupoid \mathcal{F}^{op} . A good approximation of \mathcal{F}^{op} can be obtained by the schematization functor à la Toën applied to simplicial set (quasicategory) underlying \mathcal{F}^{op} .*

We can take as model for \mathcal{F} the fundamental ∞ -groupoid $\Pi_\infty(Y)$.

Toën schematization functor can also be obtained from the Quillen-Segal formalism ap-

plied to the embedding

$$\mathcal{U} : Sh(\mathbf{Var}(\mathbb{C})) \hookrightarrow sPresh(\mathbf{Var}(\mathbb{C})),$$

where on the right hand side we consider the model category of simplicial presheaves à la Jardine-Joyal.

3 Conjectures

We now list some conjectures. We use the same notations as [3].

Conjecture 3.1. 1. As mentioned above, we can enhance $Top \hookrightarrow \mathbf{sSet}_J[Top]$ by looking at the Quillen-Segal objects. Then given Y as before, then the mirror of Y should correspond to thhe¹ opposite ∞ -groupoid $\Pi_\infty(Y)$. Indeed by a theorem of Toën [9], we know that $\mathbb{Z}/2 = Gal(\mathbb{C}/\mathbb{R})$ acts on $\mathbf{ho}(\mathbf{sSet}_J)$, where 0 is the identity and 1 is the opposite-category construction. It seems that it's not surprising that $\mathbb{Z}/2$ appears in supersymmetry.

2. We can enhance $\mathbf{Vect}_{\mathbb{C}} \hookrightarrow (\infty, n) \mathbf{Cat}[\mathbf{Vect}_{\mathbb{C}}]$ with the relative pushout product. Taking $(\infty, n) \mathbf{Cat}[\mathbf{Vect}_{\mathbb{C}}]$ as coefficient for TQFT, should explain why TQFT are classified by fully dualizable objects. Indeed Toën's theorem has been generalized by Barwick and Schommer-Pries [4]. There is an action of $(\mathbb{Z}/2)^n$ on $\mathbf{ho}[(\infty, n) \mathbf{Cat}]$, that correspond to the different opposite construction for 1-morphisms, to n -morphisms. This action should explain the n -dualizable objects.

3. We can enhance $dg\text{-}\mathbf{Cat} \hookrightarrow (\infty, 2) \mathbf{Cat}[dg\text{-}\mathbf{Cat}]$ and similarly we have an action of $(\mathbb{Z}/2)^2$ on the homotopy category $\mathbf{ho}[(\infty, 2) \mathbf{Cat}]$. This action should explain the proof of Deligne's conjecture given by Tamarkin [8]. It should also agree with Tamarkin's answer to Drinfeld's question:

What do dg-categories form ?

4. We can enhance $\mathbf{Mod}_k \hookrightarrow (\infty, 1) \mathbf{Cat}[\mathbf{Mod}_k]$ with the relative pushout product. It should be interesting to let cohomology theories in algebraic geometry take their coefficient in this enhancement.

Remark 3.2. If we follow our philosophy, it's not surprising that there is no direct link between a variety Y and its mirror Y^{op} . Because there is no direct link in general between a classical category \mathcal{C} and its opposite \mathcal{C}^{op} , unless \mathcal{C} is Tannakian or at least has duals. It would be interesting to understand the statement of the Hodge conjecture in terms of ∞ -groupoid that are fixed by the homotopy action of $\mathbb{Z}/2 = Gal(\mathbb{C}/\mathbb{R})$.

¹'thhe' is the homotopy version of 'the' (Drinfeld)

References

- [1] D. Ara. On the homotopy theory of Grothendieck ∞ -groupoids. *ArXiv e-prints*, June 2012.
- [2] H. V. Bacard. Quillen-Segal objects and structures: an overview. *ArXiv e-prints*, June 2014.
- [3] H. V. Bacard. Understanding higher structures through Quillen-Segal objects. *ArXiv e-prints*, July 2014.
- [4] C. Barwick and C. Schommer-Pries. On the Unicity of the Homotopy Theory of Higher Categories. *ArXiv e-prints*, November 2011.
- [5] Pierre Deligne. Théorie de Hodge. III. *Inst. Hautes Études Sci. Publ. Math.*, (44):5–77, 1974.
- [6] Jean Giraud. *Cohomologie non abélienne*. Springer-Verlag, Berlin, 1971. Die Grundlehren der mathematischen Wissenschaften, Band 179.
- [7] Alexander Grothendieck. *Revêtements étales et groupe fondamental. Fasc. II: Exposés 6, 8 à 11*, volume 1960/61 of *Séminaire de Géométrie Algébrique*. Institut des Hautes Études Scientifiques, Paris, 1963.
- [8] D. Tamarkin. What do DG categories form? *ArXiv Mathematics e-prints*, June 2006.
- [9] Bertrand Toën. Vers une axiomatisation de la théorie des catégories supérieures. *K-Theory*, 34(3):233–263, 2005.
- [10] Claire Voisin. *Hodge theory and complex algebraic geometry. I*, volume 76 of *Cambridge Studies in Advanced Mathematics*. Cambridge University Press, Cambridge, 2002. Translated from the French original by Leila Schneps.
- [11] Claire Voisin. *Hodge theory and complex algebraic geometry. II*, volume 77 of *Cambridge Studies in Advanced Mathematics*. Cambridge University Press, Cambridge, 2003. Translated from the French by Leila Schneps.