

## State of the art of genomics for selection

Didier Boichard, Francois Guillaume, A. Baur, Pascal Croiseau, M.N. Rossignol, Marie Yvonne Boscher, T. Druet, L. Genestout, L. Journaux, Vincent Ducrocq, et al.

## ► To cite this version:

Didier Boichard, Francois Guillaume, A. Baur, Pascal Croiseau, M.N. Rossignol, et al.. State of the art of genomics for selection. Workshop of ICAR 2011 "New technologies and new challenges for breeding and herd management", Jun 2011, Bourg-en-Bresse, France. 1 p. hal-01019343

## HAL Id: hal-01019343 https://hal.science/hal-01019343

Submitted on 6 Jun 2020

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

## State of the art of genomics for selection

*D. Boichard*<sup>(1)</sup>, F. Guillaume<sup>(1,2)</sup>, A. Baur<sup>(3)</sup>, P. Croiseau<sup>(1)</sup>, M.N. Rossignol<sup>(4)</sup>, M.Y. Boscher<sup>(4)</sup>, T. Druet<sup>(5)</sup>, L. Genestout<sup>(4)</sup>, L. Journaux<sup>(3)</sup>, V. Ducrocq<sup>(1)</sup>, S. Fritz<sup>(3)</sup>.

<sup>(1)</sup> INRA ; <sup>(2)</sup> Institut de l'Elevage; <sup>(3)</sup> UNCEIA; <sup>(4)</sup> Labogena, France ; <sup>(5)</sup> Liège University, Belgium

Selection could be based on DNA polymorphism information of each animal. Nowadays, this information is obtained through chips providing individual genotyping results for many thousand SNP simultaneously. In the future, sequencing will become more affordable and provide a complete description of each individual genome. This polymorphism information could be used in genetic evaluation. The prediction is based on the genotype-phenotype relationships observed in a reference population and its accuracy depends on two parameters, the accuracy of SNP estimates (depending on the sample size and heritability) and linkage disequilibrium between SNP and QTL. Several genomic evaluation methods have been proposed, with varying underlying assumptions. The most popular, GBLUP, uses SNP information mainly to measure true relationships between animals. Other approaches, particularly Bayesian methods, try to detect the most predictive SNP. The French model for dairy cattle uses haplotypes instead of single SNPs, in order to maximize LD between markers and QTL. For each trait, 300-700 QTL characterized by 3-5 SNP are included in a QTL-BLUP. Large reference populations have been built by genotyping many progeny tested bulls and by sharing information in consortia such as the EuroGenomics initiative. Two major questions, the evaluation for new traits and the across population evaluation, need to be solved for a generalized use of genomic selection in all species and breeds.