



**HAL**  
open science

## Wood formation is altered in poplar under ozone and/or elevated carbon dioxide

Nicolas Richet, Dany Afif, Françoise Huber, Koffi Tozo, Pierrick Priault, Jacques Banvoy, Brigitte Pollet, Catherine Lapierre, Pascale Maillard, Patrick Gross, et al.

### ► To cite this version:

Nicolas Richet, Dany Afif, Françoise Huber, Koffi Tozo, Pierrick Priault, et al.. Wood formation is altered in poplar under ozone and/or elevated carbon dioxide. 26. New Phytologist Symposium Bioenergy trees, May 2011, Nancy, France. pp.43. hal-01019732

**HAL Id: hal-01019732**

**<https://hal.science/hal-01019732>**

Submitted on 3 Jun 2020

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

## 28. Wood formation is altered in poplar under ozone and/or elevated carbon dioxide

**NICOLAS RICHEL<sup>1</sup>, DANY AFIF<sup>1</sup>, FRANCOISE HUBER<sup>2</sup>, KOFFI TOZO<sup>3</sup>, PIERRICK PRIAULT<sup>1</sup>, JACQUES BANVOY<sup>1</sup>, BRIGITTE POLLET<sup>4</sup>, CATHERINE LAPIERRE<sup>4</sup>, PASCALE MAILLARD<sup>1</sup>, PATRICK GROSS<sup>1</sup>, PIERRE DIZENGREMEL<sup>1</sup>, PATRICK PERRE<sup>2</sup>, MIREILLE CABANE<sup>1</sup>**

<sup>1</sup>Nancy-Université, INRA, UMR 1137 Ecologie et Ecophysiologie Forestières, Boulevard des Aiguillettes, B.P. 70239, F-54506 Vandœuvre lès Nancy, France. <sup>2</sup>AgroParisTech, UMR 1092 LERFOB, ENGREF, 14 rue Girardet, F-54042 Nancy cedex, France ; INRA, UMR 1092 LERFOB, ENGREF, 14 rue Girardet, F-54042 Nancy cedex, France ; <sup>3</sup>Département de botanique, Faculté des sciences, Université de Lomé, B.P. 1515 Lomé, Togo. (4) AgroParisTech, UMR 1318, 78850 Thiverval-Grignon, France; INRA, UMR 1318, 78850 Thiverval-Grignon, France.

The industrial development led to an increase in the concentration of atmospheric carbon dioxide but also resulted in an increase of tropospheric ozone concentrations. Ozone has been suggested to cause the greatest amount of damage to vegetation as compared to other gaseous pollutants. On the opposite, elevated carbon dioxide is usually observed to enhance tree photosynthesis and growth.

We investigated the effects of elevated carbon dioxide and/or ozone on wood formation in young poplars. Trees were bent in order to induce tension wood formation on the upper side of the stem and opposite wood on the lower side. Under ozone, lignin and cellulose biosynthesis decreased probably due to lower availability in carbon skeletons. Cellulose was more affected than lignin resulting in a reduction of cellulose to lignin ratio in wood. Wood anatomy and density was also modified by ozone. Tension wood was generally more altered than opposite wood. All these results showed a strong impact of ozone on wood formation and wood quality and suggested a coordinated regulation of lignin and cellulose synthesis. Elevated carbon dioxide had a low impact on wood formation but the combination with ozone resulted in an attenuation of ozone effects.

## 29. Shoot Development In Short Rotation Coppice Willow.

**J. SALMON<sup>1</sup>, S. WARD<sup>2</sup>, S. HANLEY<sup>1</sup>, O. LEYSER<sup>2</sup>, A. KARP<sup>1</sup>**

<sup>1</sup>Centre for Bioenergy and Climate Change, Rothamsted Research, Harpenden, Herts., AL5 2JQ, UK; <sup>2</sup>Sainsbury Laboratory Cambridge University, Bateman Street, Cambridge, CB2 1LR, UK

Central to the suitability of willows for biomass production is the vigorous re-growth that occurs following the removal of apical dominance by coppicing. We are beginning to decipher the regulation of this process with respect to bud and shoot behaviour. A dedicated field trial was established in April 2008 comprising six genotypes contrasting in shoot number, stem diameter, and yield. After first coppice a subset of stools was extracted and examined for bud formation, position and number, revealing variation between genotypes in the proportion of buds that sprouted. This process is currently being repeated two years post-coppice, and followed in two mapping populations.

Preliminary QTL analysis placed two *More AXillary branching (MAX)* genes within a locus for shoot development in the K8 mapping population. The K8 alleles and alleles from diverse willow species have been cloned and used for transformation rescue of *Arabidopsis max* mutants. Variation in the degree of mutant rescue between allelic variants suggests functional allelic differences that could contribute to differences in coppicing phenotypes. Eleven new mapping populations have been generated with the aim of affirming, and discovering new, bud and shoot regulation QTLs.

This project aims to improve biomass willows in terms of increased yield and the production of biomass products tailored to a range of energy production processes.