



HAL
open science

A never-decisive and anonymous criterion for optimal growth models

Alain Ayong Le Kama, Ha-Huy Thai, Cuong Le Van, Katheline Schubert

► **To cite this version:**

Alain Ayong Le Kama, Ha-Huy Thai, Cuong Le Van, Katheline Schubert. A never-decisive and anonymous criterion for optimal growth models. *Economic Theory*, 2014, 55 (2), pp.281-306. 10.1007/s00199-013-0759-x . hal-01044568

HAL Id: hal-01044568

<https://hal.science/hal-01044568v1>

Submitted on 17 Nov 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

A Never-decisive and Anonymous Criterion for Optimal Growth Models *

Alain Ayong Le Kama Thai Ha-Huy
EconomiX, University Paris Ouest EPEE, University of Evry

Cuong Le Van
IPAG Business School, CNRS, Paris School of Economics

Katheline Schubert
Paris School of Economics, University Paris 1

April 16, 2013

Abstract

We address in this paper the question of the existence of a Social Welfare Function that would be sustainable and would allow to obtain solutions to optimal growth models. We define sustainability by two new axioms called Never-decisiveness of the present and Never-decisiveness of the future. We first show that a SWF which has Never-decisiveness properties cannot be defined on a ball of l_{∞}^+ . We must (i) restrict to the set of utility streams for which the value of the SWF is finite and (ii) introduce additional assumptions in order to obtain the Never-decisiveness properties. Our main result in this paper is therefore to show that the undiscounted utilitarian criterion is an anonymous and never-decisive criterion for optimal growth models. We consider the set of utilities of consumptions which are generated by a specific technology, namely a technology with decreasing returns for high levels of capital and restrict ourselves to good programmes, i.e. any programme for which intertemporal utility is well defined.

JEL classification codes: D60, D70, D90, Q0

Keywords: Anonymity, intergenerational equity, natural resources, never-decisiveness of the future, never-decisiveness of the present, no-dictatorship

*The authors thank an anonymous referee and the associate editor of this journal for very helpful and valuable comments and suggestions. They acknowledge the support of the French National Research Agency (ANR) under the CLEANER project (ANR_NT09_505778).

of the future, no-dictatorship of the present, optimal growth models, social welfare function, sustainability.

1 Introduction

Optimal growth theory is largely built around the discounted utilitarian approach, but the debate between authors subscribing to discounted utilitarianism and authors rejecting it has always been vivid. Ramsey [1928] himself, in a much-cited statement, held that “*discount later enjoyments in comparison with earlier ones [is] a practice which is ethically indefensible and arises merely from the weakness of imagination*”. This debate is particularly passionate when the sustainability of growth is concerned, because the process of discounting forces a fundamental asymmetry between present and future generations, particularly those in the distant future, and so appears to be in contradiction with the intergenerational equity concern underlying the search for sustainability.

Several alternative social welfare functions (SWF hereafter) have been proposed in the optimal growth literature. The Rawlsian or maximin criterion (Rawls [1971], Solow [1974]) maximizes the utility of the least well off generation. The undiscounted utilitarian criterion (Ramsey [1928], Koopmans [1965]) maximizes the infinite sum of the (negative) differences between actual utilities and their upper bound (the bliss), with a zero utility discount rate. With the overtaking criterion (Gale [1967]), one utility stream is said to be better than another if from some date on the first one is greater than the second one. Whereas the usual discounted utilitarian criterion assumes a constant discount rate, criteria with non constant and declining discount rates, hyperbolic for instance, have also been used (see e.g. Heal [1998]).

In the context of each particular growth model, the shape of the optimal growth paths obtained under these various criteria is intimately dependent on the properties of each criterion. For instance, in the canonical Dasgupta–Heal–Solow model of growth with exhaustible resources, optimal consumption asymptotically vanishes with the discounted utilitarian criterion, whereas it grows forever with the undiscounted utilitarian criterion, and remains constant with the maximin criterion (see e.g. Dasgupta and Heal [1979]). The properties leading to these different paths are related to discounting, but also to the degree of intertemporal substitutability of utilities embodied in the criterion; more deeply, these properties are con-substantial with the axiomatic foundations of the criteria, as we have been knowing since Koopmans [1960].

The axiomatic approach founding social welfare criteria considers the set of intertemporal utility streams generated by a general set of bounded

consumptions, and defines a preference relation between them. This relation is ideally required to satisfy two main axioms, the Pareto axiom and the Anonymity axiom. Pareto guarantees efficiency. Broadly speaking, it demands that the preference relation is sensitive to the well-being of each generation. Anonymity guarantees equity, that is an equal treatment of all generations, in the sense that any finite permutation of utilities does not change the evaluation of the utility stream. Besides, preferences are often required to be independent¹ and to satisfy some continuity, transitivity and completeness properties. Unfortunately, Basu and Mitra [2003] show that when the social welfare criterion is represented by a function, then it cannot satisfy simultaneously the Pareto and Anonymity axioms when instantaneous utilities are uniformly bounded: it is impossible to represent by a numerical function a preference relation embodying both the efficiency and equity requirements.

For practical purposes then, the point is to determine which axiom(s) we are ready to drop, depending on the context. Obviously, the discounted utilitarian criterion does not satisfy Anonymity. It seems very counterintuitive to drop Anonymity as far as sustainability is concerned. The undiscounted utilitarian criterion does not satisfy completeness: is not a well defined real valued function on all l_∞ ², and cannot therefore define a complete order on l_∞ . The overtaking criterion also it is not a well defined function of l_∞ , since it cannot rank a pair of utility streams of l_∞ for which neither the first overtakes the second, nor the second overtakes the first. This criterion does not also satisfy the axiom of "Strong Relative Anonymity", introduced by Asheim, d'Aspremont and Banerjee [2010a], where 'strong' reflects that all infinite permutations are considered. The maximin criterion does not satisfy the independence property. Nevertheless, neither completeness nor independence seem crucial for sustainability.

Chichilnisky [1996] drops the Anonymity axiom. But her purpose is to capture the idea of sustainable growth, and so she proposes a weaker concept of equity embodied in two new axioms: No-dictatorship of the present and No-dictatorship of the future. A social welfare function is said to give *a dictatorial role to the present* if it disregards the utilities of all generations from some generation on. Conversely, a social welfare function gives *a dictatorial role to the future* if it is only sensitive to the utilities of the generations coming after some generation. She exhibits, for the set of uniformly bounded

¹See Definition 1 below.

² $l_\infty = \{(y_g)_{g=1, \dots, \infty} : y_g \in \mathbb{R}, \sup_g |y_g| < \infty\}$.

consumptions, a social welfare criterion satisfying, besides the Pareto and independence requirements, these two axioms for sustainability. It is a mixed criterion, composed of one discounted utilitarian part taking into account the interests of the present, and a part taking into account only the interests of the future.

This approach is very appealing. But its drawback is that most of the time the Chichilnisky's criterion does not allow to obtain explicit optimal growth paths with optimal control techniques. Chichilnisky [1996] does not apply her criterion to growth models but confines herself to the axiomatic approach. Applications are presented in Chichilnisky [1997] and Heal [1998], in the framework of canonical optimal growth models with exhaustible or renewable resources. When the discounted utilitarian part of the criterion embodies a constant discount rate, they claim that they obtain, using non-standard techniques, a solution in the case of exhaustible resources, but they show that no solution exists in the model with renewable resources. Only when the discount rate is decreasing are they able to obtain a solution to the renewable resource model. Figuières and Tidball [2012] focus on the inexistence result of the case with a constant discount rate, and show that they can find in the model with renewable resources "near optimal" growth paths.

More recently, Alvarez-Cuadrado and Long [2009] have proposed another mixed criterion named the mixed Bentham-Rawls criterion, weighted average of a discounted utilitarian part as in Chichilnisky's criterion, and a maximin part. They show that Chichilnisky does not obtain this criterion because it does not satisfy the independence property she imposes, property that they do not find compelling.

We address in this paper the question of the existence of a continuous SWF which is sustainable, in the sense that this continuous SWF is anonymous, Pareto, independent, and allows us to obtain solutions to optimal growth models. We introduce a new concept of sustainability embodied in two new axioms we call Never-decisiveness of the present and Never-decisiveness of the future. These axioms are inspired by the No-dictatorship axioms. We show that in the set of utilities bounded uniformly by a constant, there exists, up to a positive scalar, a unique SWF which is continuous, anonymous, Pareto, independent. This SWF satisfies our two Never-decisiveness axioms. Moreover, we can use this SWF to solve the discrete time version of the original Ramsey model (Ramsey [1928]). Actually, the two Never-decisiveness axioms give another axiomatic foundation to

the Gale's overtaking criterion (Gale [1967]). More precisely, we show that, under the assumptions introduced above and some others (see below), the overtaking criterion for optimal growth models satisfies Anonymity, Pareto axioms and the two Never-decisiveness axioms for sustainability.

We come back to the seminal definition of social welfare functions through the Yosida Theorem. We consider the SWF, W , which are restrictions to l_∞^+ of continuous, linear functions on l_∞ . From the Yosida Theorem, such a SWF has a unique decomposition: $W(\mathbf{x}) = \sum_{t=0}^{\infty} a_t x_t + \phi(\mathbf{x})$, where $\mathbf{x} \in X$, a set of infinite sequences, $\sum_{t=0}^{\infty} |a_t| < \infty$ and ϕ is *purely finitely additive*. This SWF satisfies the No-dictatorship axioms introduced by Chichilnisky [1996] for sequences of utility streams. We show that, in l_∞ , a continuous and independent SWF satisfies the No-dictatorship axioms if, and only if, it satisfies the Never-decisiveness axioms. Notice that this criterion is a non-constructible criterion (see Lauwers [2012]).

Now consider optimal growth models with decreasing returns. In these models, the feasible consumption paths are uniformly bounded by the same quantity. So, without any loss of generality, we can assume that the set of sequences X is $[0, 1]^\infty$. With this restriction, we show that there exists no SWF W (as defined above) from X into \mathbb{R}_+ , with $a_t > 0, \forall t, \sum_{t=0}^{\infty} a_t = 1$, ϕ purely finitely additive, which satisfies the two Never-decisiveness axioms. This result comes from the assumption $\sum_{t=0}^{\infty} a_t = 1$. We therefore remove it and assume $\sum_{t=0}^{\infty} a_t = +\infty^3$. The criterion is not complete anymore. Then, we easily show that the SWF W satisfies the two Never-decisiveness axioms on the set of utility sequences for which the value of the social welfare function is finite.

Asheim, Mitra and Tungodden [2012] and Asheim and Mitra [2010] propose the Sustainable Discounted Utilitarianism. As Chichilnisky does, they drop the Anonymity axiom. They propose to replace it by the axiom of Hammond Equity for the Future introduced by Asheim and Tungodden [2004], which entails that the social welfare is sensitive to the interests of the present generation only when the present is worse off than the future. It may not be the case with our criterion (see Example in Appendix 1).

Basu and Mitra [2007] propose and characterize a set of new social welfare criteria called the overtaking Social Welfare Relations, allowing to ensure an equal treatment of all generations. They show that the overtaking SWRs can be characterized in terms of four axioms. Moreover, in a one-dimension Ramsey growth model, they actually show that the optimal

³For simplicity, we also assume that $\forall t, 0 < \underline{a} \leq a_t \leq \bar{a}$.

path for the Gale overtaking criterion coincides with the optimal path for overtaking SWRs. From Dana and Le Van [1990], one can see that this path is actually the optimal path for the SWF obtained from Anonymity, Pareto axioms and our Never-decisiveness axioms. Observe that Asheim, d'Aspremont and Banerjee [2010a] show that Basu and Mitra welfare criterion is also incomplete. In other words, it may be effective in the sense of selecting a small set of optimal or maximal elements for a given class of feasible infinite utility streams. They establish a result that generalize the time-invariant overtaking criterion to satisfy stationarity and an anonymity criterion called Strong Relative Anonymity.

To sum up, for the issue of sustainability and intergenerational equity, our main result is to show that the undiscounted utilitarian criterion is a non-dictatorial and anonymous criterion for optimal growth models. It is also upper-semi continuous for the product topology, under a set of assumptions which can be summarized as follows: instead of l_∞^+ as set of utilities, we just consider the set of utilities of consumptions which are generated by a specific technology, namely a technology with decreasing returns for high levels of capital; by doing so, we restrict ourselves to good programmes in the sense of Gale [1967], Brock [1970a], Arkin and Evstigneev [1987], Dana and Le Van [1990] and Dana and Le Van [1993], when the state variables are in a finite dimension, and Zaslavski [2007] when these variables belong to an abstract compact metric space, *i.e.* any programme for which intertemporal utility is well defined.⁴

The paper is organized as follows. In Section 2 we present the Never-decisiveness criteria when the state space is respectively l_∞^+ or $[0, 1]^\infty$. We show that for an optimal growth model with decreasing returns for high value of capital stock where the utilities of the streams of consumptions typically belong to $[0, 1]^\infty$, there exists no SWF which is never-decisive. We then restrict the state space to the sets of utilities for which the value of the SWF is finite. In Section 3, we show that, actually, the Gale's overtaking criterion, restricted to the set of good programmes, is anonymous and never-decisive⁵.

⁴It is well-known that the undiscounted utilitarian criterion is not complete on l_∞^+ . In order to decrease incompleteness, Lauwers [2012] introduces a stronger notion of anonymity called maximal anonymity. Unfortunately, the criterion he obtains is a non-constructible object if we impose it to be also Pareto.

⁵Gale [1967], Dana and Le Van [1990] and Dana and Le Van [1993] require that the technology exhibits decreasing returns. Here, we drop this assumption. We show that the turnpike result in Gale [1967] also holds with non-convex technologies. Zaslavski [2007] also drops the convexity assumption but assumes the asymptotic turnpike property holds.

Section 4 shows that using the overtaking criterion, one can obtain optimal solutions to growth models. Some examples of models are also presented in this section. In particular, we present an economy with a convex-concave production function (see Dechert and Nishimura [1983]). In Dechert and Nishimura [1983], the proof of existence of a poverty trap is given, but its precise value is not easy to compute. As a by product, we obtain here the explicit value of this poverty trap. We also use our criterion to solve two growth models, respectively with exhaustible and renewable resources.

2 Never-decisiveness criterion

We introduce here a new concept, called Never-decisiveness, based on Chichilnisky [1996] concept of No-dictatorship. We show that the two concepts are equivalent in l_∞ . As we are interested in finding solutions of optimal growth models, we restrict ourselves to the space of uniformly bounded consumptions. Then we know that the Chichilnisky criterion may not allow us to obtain solutions. We show how our criterion overcomes this problem.

2.1 Preliminaries

We first recall the concepts of No-dictatorship of the present and No-dictatorship of the future given by Chichilnisky [1996].

Let X denote a set of infinite sequences. A criterion W is a non-decreasing function from X into \mathbb{R} .

A criterion W exhibits **No Dictatorship of the Present** if there exist $\mathbf{x} \in X$, $\mathbf{y} \in X$ which satisfy $W(\mathbf{x}) > W(\mathbf{y})$, and for any N , there exist $k \geq N$, $(z_t)_{t \geq k+1}$, $(s_t)_{t \geq k+1}$ such that

$$W(x_0, \dots, x_k, z_{k+1}, \dots, z_{k+t}, \dots) \leq W(y_0, \dots, y_k, s_{k+1}, \dots, s_{k+t}, \dots).$$

A criterion W exhibits **No Dictatorship of the Future** if there exist $\mathbf{x} \in X$, $\mathbf{y} \in X$ which satisfy $W(\mathbf{x}) > W(\mathbf{y})$, and for any N , there exist $k \geq N$, $(z_t)_{t=0, \dots, k}$, $(s_t)_{t=0, \dots, k}$ such that

$$W(z_0, \dots, z_k, x_{k+1}, \dots, x_{k+t}, \dots) \leq W(s_0, \dots, s_k, y_{k+1}, \dots, y_{k+t}, \dots).$$

We now define the concepts of Never-decisiveness of the present and Never-decisiveness of the future.

A criterion W exhibits **Never-decisiveness of the Present** if for any $\mathbf{x} \in X$, $\mathbf{y} \in X$ which satisfy $W(\mathbf{x}) > W(\mathbf{y})$, then, for any N , there exist

$k \geq N$, $(z_t)_{t \geq k+1}$, $(s_t)_{t \geq k+1}$ such that

$$W(x_0, \dots, x_k, z_{k+1}, \dots, z_{k+t}, \dots) \leq W(y_0, \dots, y_k, s_{k+1}, \dots, s_{k+t}, \dots).$$

A criterion W exhibits **Never-decisiveness of the Future** if for any $\mathbf{x} \in X$, $\mathbf{y} \in X$ which satisfy $W(\mathbf{x}) > W(\mathbf{y})$, then, for any N , there exist $k \geq N$, $(z_t)_{t=0, \dots, k}$, $(s_t)_{t=0, \dots, k}$ such that

$$W(z_0, \dots, z_k, x_{k+1}, \dots, x_{k+t}, \dots) \leq W(s_0, \dots, s_k, y_{k+1}, \dots, y_{k+t}, \dots).$$

The following definition generalizes Definition 7 in Chichilnisky [1996, p.251]:

Definition 1 *Let X be a nonempty subset of l_∞ and W be a criterion from X into \mathbb{R} . The criterion W is independent if there exists a linear function l on l_∞ such that for any $\mathbf{x} \in l_\infty$, any $\mathbf{y} \in l_\infty$, $W(\mathbf{x}) = W(\mathbf{y}) \Leftrightarrow l(\mathbf{x} - \mathbf{y}) = 0$.*

From Chichilnisky [1996, p.251], when $X = l_\infty$, W can be represented by a linear function on l_∞ . Now, when X is a subset of l_∞ , we say that W is independent if it is affine.

We say that the SWF W is sustainable if it is complete (i.e. well defined on the whole space l_∞^+), increasing and satisfies the two No Dictatorship axioms.

We consider the SWF which are restrictions to l_∞^+ of continuous, linear functions on l_∞ . From Yosida theorem, such a SWF W has a unique decomposition

$$W(\mathbf{x}) = \sum_{t=0}^{\infty} a_t x_t + \phi(\mathbf{x})$$

where $\sum_{t=0}^{\infty} |a_t| < \infty$ and ϕ is purely finitely additive. If W is increasing then $a_t > 0$ for all t , and ϕ is increasing.

In the following Proposition, when $X = l_\infty$, sustainability is equivalent to Never-decisiveness:

Proposition 1 *Let W be a continuous independent SWF from l_∞ into \mathbb{R}_+ . Then W is sustainable if, and only if, it satisfies Never-decisiveness of the future and Never-decisiveness of the present.*

Proof: 1. From Chichilnisky [1996], Theorem 2, if W is sustainable then it has the decomposition:

$$W(\mathbf{x}) = \sum_{t=0}^{\infty} a_t x_t + \phi(\mathbf{x}) \tag{1}$$

where $\sum_{t=0}^{\infty} |a_t| < \infty$ and ϕ is purely finitely additive. Moreover $a_t > 0, \forall t$ and $\phi \neq 0$. We can assume $\sum_{t=0}^{\infty} a_t = 1$ and $\phi(1, 1, \dots, 1, \dots) > 0^6$.

2. Let us consider a SWF W as in (1). We will prove that it is never-decisive.

(a) **Never-decisiveness of the present**

Let $W(\mathbf{x}) > W(\mathbf{y})$ with $\mathbf{x} \in \ell_{\infty}, \mathbf{y} \in \ell_{\infty}$. Choose $\lambda > 0$ large enough s.t. $\phi(\lambda, \lambda, \dots) = \lambda\phi(1, 1, \dots) > \sum_{t=0}^{\infty} a_t x_t - \sum_{t=0}^{\infty} a_t y_t$ and $\lambda > \sup_t |y_t|$. For any N large enough, we have $\phi(\lambda, \lambda, \dots) = \lambda\phi(1, 1, \dots) > \sum_{t=0}^N a_t x_t - \sum_{t=0}^{\infty} a_t y_t$. Take such an N . For any $k \geq N$, define $z_t = 0, \forall t \geq k+1, z'_t = \lambda, \forall t \leq k$. Then $\phi(y_0, \dots, y_k, z'_{k+1}, \dots) = \phi(\lambda, \lambda, \dots) = \lambda\phi(1, 1, \dots)$. Then

$$\begin{aligned} W(x_0, \dots, x_k, z_{k+1}, \dots) &= \sum_{t=0}^k a_t x_t < \sum_{t=0}^{\infty} a_t y_t + \lambda\phi(1, 1, \dots) \\ &< \sum_{t=0}^k a_t y_t + \sum_{t=k+1}^{\infty} a_t z'_t + \phi(y_0, \dots, y_k, z'_{k+1}, \dots) \\ &= W(y_0, \dots, y_k, z'_{k+1}, \dots). \end{aligned}$$

(b) **Never-decisiveness of the future**

Let $W(\mathbf{x}) > W(\mathbf{y})$ with $\mathbf{x} \in \ell_{\infty}, \mathbf{y} \in \ell_{\infty}$. Let $\lambda > 0$ satisfy $\lambda + \phi(\mathbf{y}) > \phi(\mathbf{x})$. There exists N s.t. for any $k \geq N, \lambda \sum_{t=0}^k a_t + \phi(\mathbf{y}) > \phi(\mathbf{x}) + \sum_{t=k+1}^{\infty} a_t x_t$, since $\sum_{t=0}^{\infty} a_t = 1$ and $\sum_{t=T}^{\infty} a_t x_t \rightarrow 0$, when $T \rightarrow +\infty$. For any $k \geq N$, define $z_t = 0$ for $t = 0, \dots, k, z'_t = \lambda$ for $t = 0, \dots, k$. Then

$$\begin{aligned} W(z_0, \dots, z_k, x_{k+1}, \dots) &= \sum_{t=k+1}^{\infty} a_t x_t + \phi(\mathbf{x}) < \lambda \sum_{t=0}^k a_t + \phi(\mathbf{y}) \\ &= \sum_{t=0}^k a_t z'_t + \phi(\mathbf{y}) = W(z'_0, \dots, z'_k, y_{k+1}, \dots). \end{aligned}$$

■

3. Obviously, if W is a Never-decisive social welfare function then it is sustainable.

2.2 The case where $X \subseteq [0, 1]^{\infty}$

We mentioned at the beginning of the paper that in optimal growth models with decreasing returns the feasible consumption paths are uniformly bounded. So, without loss of generality, we can first assume $X = [0, 1]^{\infty}$.

⁶This last assumption is the one used by Chichilnisky [1996].

Proposition 2 *The SWF W from X into \mathbb{R}_+ :*

$$W(\mathbf{x}) = \sum_{t=0}^{\infty} a_t x_t + \phi(\mathbf{x})$$

with $a_t > 0, \forall t, \sum_{t=0}^{\infty} a_t = 1, \phi$ positive, purely finitely additive, satisfies the two No dictatorship properties.⁷

Proof: a) **No Dictatorship of the present**

Let \mathbf{x} satisfy $x_t = \zeta$ for any t with $\zeta < \min\{1, \phi(1, 1, \dots)\}$. Let $y_t = 0, \forall t$. Take $\gamma = \mathbf{0}$ and $\sigma_t = 1, \forall t$. Then, for any K ,

$$\begin{aligned} W(x_0, \dots, x_K, \gamma_{K+1}, \dots) &= \zeta \sum_{t=0}^K a_t + \phi(x_0, \dots, x_K, 0, \dots, 0, \dots) \\ &= \zeta \sum_{t=0}^K a_t + \phi(\mathbf{0}) = \zeta \sum_{t=0}^K a_t \\ &< \zeta < \sum_{K+1}^{\infty} a_t \sigma_t + \phi(1, 1, \dots) \\ &= W(y_0, \dots, y_K, \sigma_{K+1}, \dots) \end{aligned}$$

b) **No Dictatorship of the future** Let $x_0 = 1, x_t = 0$ for $t \geq 1$, and let all components of \mathbf{y}, \mathbf{z} and \mathbf{s} be equal to 0. Then, since $a_0 > 0$ and $x_0 = 1 > 0 = y_0, W(\mathbf{x}) > W(\mathbf{y})$, but

$$W(z_0, z_1, \dots, z_K, y_{K+1}, \dots) = W(s_0, \dots, s_K, x_{K+1}, \dots)$$

for all $K \geq 0$. Essentially, the property that W is increasing in all components implies No Dictatorship of the Future. ■

However, in Appendix 2, we show that the basic one sector Ramsey model with a criterion defined by (1), where x_t is the utility at period t of the representative consumer, has no solution if the marginal productivity at the origin of the production function is larger than 1. When this marginal productivity is less than one, the criterion defined by (1) and the one defined by the discounted sum of utilities give the same solution to the Ramsey model.

The following proposition shows that there exists no independent, continuous (for the l_∞ topology) SWF which is never-decisive in $[0, 1]^\infty$.

⁷In Chichilnisky [1996], we have a proof for l_∞ . Here, we give a proof for $[0, 1]^\infty$.

Proposition 3 *There exists no continuous independent SWF W from X into \mathbb{R}_+ :*

$$W(\mathbf{x}) = \sum_{t=0}^{\infty} a_t x_t + \phi(\mathbf{x})$$

with $a_t > 0, \forall t, \sum_{t=0}^{\infty} a_t = 1, \phi$ positive, purely finitely additive, which satisfies the two Never-decisiveness properties.

Proof: *Claim 1.* If $\phi(1, 1, \dots) \geq 1$, then W does not satisfy the Never-decisiveness of the future axiom.

Indeed, let $W(\mathbf{x}) > W(\mathbf{y})$ where $x_t = 1, \forall t$ and $y_t = 0, \forall t$. Let N be the first index s.t. $a_N > 0$. Then $\forall k \geq N, \forall (z_0, \dots, z_k) \in [0, 1]^k, \forall (z'_0, \dots, z'_k) \in [0, 1]^k$,

$$\begin{aligned} W(z_0, \dots, z_k, x_{k+1}, \dots, x_{k+t}) &= \sum_{t=0}^k a_t z_t + \sum_{t=k+1}^{\infty} a_t x_t + \phi(1, 1, \dots) \\ &\geq \phi(1, 1, \dots) \\ &\geq \sum_{t=0}^{\infty} a_t \\ &\geq \sum_{t=0}^k a_t z'_t + \sum_{t=k+1}^{\infty} a_t y_t + \phi(0, 0, \dots) \\ &= W(z'_0, \dots, z'_k, y_{k+1}, \dots). \end{aligned}$$

Claim 2. If $0 < \phi(1, 1, \dots) < 1$, then W does not satisfy the Never-decisiveness of the present property.

To prove that, let $\varepsilon > 0$ satisfy $1 - \varepsilon > \phi(1, 1, \dots)$ and let $x_t = 1, \forall t$ and $y_t = 0, \forall t$. Hence $W(\mathbf{x}) > W(\mathbf{y})$. There exists N s.t. $\forall k \geq N, \sum_{t=0}^k a_t x_t = \sum_{t=0}^k a_t > \phi(1, 1, \dots) + \sum_{t=k+1}^{\infty} a_t$. Then, for any $k \geq N$, for any $(z_{k+1}, \dots) \in [0, 1]^{\infty}$, any $(z'_{k+1}, \dots) \in [0, 1]^{\infty}$ we have

$$\begin{aligned} W(y_0, \dots, y_k, z'_{k+1}, \dots) &= \sum_{t=k+1}^{\infty} a_t z'_t + \phi(y_0, \dots, y_k, z'_{k+1}, \dots) \\ &\leq \phi(1, 1, \dots) + \sum_{t=k+1}^{\infty} a_t \\ &\leq \sum_{t=0}^k a_t x_t + \sum_{t=k+1}^{\infty} a_t z_t + \phi(x_0, \dots, x_k, z_{k+1}, \dots) \\ &= W(x_0, \dots, x_k, z_{k+1}, \dots). \end{aligned}$$

The proof of the proposition is over. ■

We now extend the Never-decisiveness axioms to a SWF defined only on a subset of $[0, 1]^\infty$. In view of Proposition 2, we now assume $\sum_{t=0}^\infty a_t = +\infty$ and impose a domain restriction.

We consider a sequence \mathbf{a} which satisfies $a_t > 0, \forall t \geq 0, \sum_{t=0}^\infty a_t = +\infty, 0 < \underline{a} \leq a_t \leq \bar{a}, \forall t$. Let U be a subset of $[0, 1]^\infty$ and let $\bar{x} \in [0, 1]$. Define

$$X(\mathbf{a}, \bar{x}) = \left\{ \mathbf{x} \in U : \sum_{t=0}^\infty a_t(x_t - \bar{x}) \text{ exists in } \mathbb{R} \right\}$$

We make the following assumption on $X(\mathbf{a}, \bar{x})$:

Assumption 1 (\mathcal{T}_0): for any $\mathbf{x} \in X(\mathbf{a}, \bar{x})$, we have $x_t \rightarrow \bar{x}$ when $t \rightarrow +\infty$ and there exists a neighborhood $\mathcal{V}(\bar{x})$ of \bar{x} which satisfies:

(\mathcal{T}_1): there exists $x \in \mathcal{V}(\bar{x}), x \neq \bar{x}$ such that for any t , if $x_t \in \mathcal{V}(\bar{x}) \cap [0, 1]$ then there exist $s \in \mathcal{V}(\bar{x}) \cap [0, 1], s' \in \mathcal{V}(\bar{x}) \cap [0, 1], K$ such that for any $k \geq K$, the sequence \mathbf{x}' defined by $x'_i = x_i, i = 1, \dots, t, x'_{t+1} = s, x'_{t+i+1} = x, \forall i = 1, \dots, K, x'_{t+2+K} = s', x'_i = \bar{x}, i > t + 2 + K$ is in $X(\mathbf{a}, \bar{x})$.

(\mathcal{T}_2): there exist $x \in \mathcal{V}(\bar{x}) \cap [0, 1], x \neq \bar{x}, K$ such that for any $k \geq K$, there exists $s \in \mathcal{V}(\bar{x}) \cap [0, 1]$, and the sequence \mathbf{x}' defined by $x'_i = x, i = 1, \dots, k, x'_{k+1} = s, x'_{k+i+1} = x_{k+i+1}, \forall i > 0$, is in $X(\mathbf{a}, \bar{x})$.

Let $W(\mathbf{x}; \mathbf{a}, \bar{x}) = \sum_{t=0}^\infty a_t(x_t - \bar{x}) + \phi(\mathbf{x})$ for any $\mathbf{x} \in X(\mathbf{a}, \bar{x})$. Since $x_t \rightarrow \bar{x}$, we have $\phi(\mathbf{x}) = \phi(\bar{\mathbf{x}})$. Thus, we can take $W(\mathbf{x}; \mathbf{a}, \bar{x}) = \sum_{t=0}^\infty a_t(x_t - \bar{x})$.

Proposition 4 Suppose (\mathcal{T}_0), (\mathcal{T}_1), (\mathcal{T}_2) hold. Then the social welfare function $W(\cdot; \mathbf{a}, \bar{x})$ satisfies the two Never-decisiveness axioms on $X(\mathbf{a}, \bar{x})$.

Proof: (a) **Never-decisiveness of the present**

Let $W(\mathbf{x}; \mathbf{a}, \bar{x}) > W(\mathbf{y}; \mathbf{a}, \bar{x})$ with $\mathbf{x} \in X(\mathbf{a}, \bar{x}), \mathbf{y} \in X(\mathbf{a}, \bar{x})$. From (\mathcal{T}_0), (\mathcal{T}_1), since $x_t \rightarrow \bar{x}, y_t \rightarrow \bar{x}$ when $t \rightarrow +\infty$, one can find \hat{x}, s, s' , close to \bar{x} , and τ , s.t. for any $T \geq \tau$, for any k , when $\hat{x} > \bar{x}$, the sequence $\mathbf{y}^k = (y_0, y_1, \dots, y_T, s, z_{T+2}, \dots, z_{T+1+k}, s', \bar{x}, \bar{x}, \dots)$, with $z_{T+1+i} = \hat{x}, i = 1, \dots, k$, is in $X(\mathbf{a}, \bar{x})$ and when $\hat{x} < \bar{x}$, the sequence

$$\mathbf{x}^k = (x_0, x_1, \dots, x_T, s, z_{T+2}, \dots, z_{T+1+k}, s', \bar{x}, \bar{x}, \dots)$$

with $z_{T+1+i} = \hat{x}, i = 1, \dots, k$, is in $X(\mathbf{a}, \bar{x})$.

When $\hat{x} > \bar{x}$, we have

$$W(\mathbf{y}^k; \mathbf{a}, \bar{x}) = \sum_{t=0}^T a_t(y_t - \bar{x}) + a_{T+1}(s - \bar{x}) + (\hat{x} - \bar{x}) \sum_{T+2}^{T+1+k} a_t + a_{T+2+k}(s' - \bar{x})$$

Then $W(\mathbf{y}^k; \mathbf{a}, \bar{x}) \rightarrow +\infty$. Therefore, $W(\mathbf{y}^k; \mathbf{a}, \bar{x}) > W(\mathbf{x}; \mathbf{a}, \bar{x})$ for k large enough.

When $\hat{x} < \bar{x}$, we have

$$W(\mathbf{x}^k; \mathbf{a}, \bar{x}) = \sum_{t=0}^T a_t(x_t - \bar{x}) + a_{T+1}(s - \bar{x}) + (\hat{x} - \bar{x}) \sum_{T+2}^{T+1+k} a_t + a_{T+2+k}(s' - \bar{x})$$

then $W(\mathbf{x}^k; \mathbf{a}, \bar{x}) \rightarrow -\infty$. Therefore, $W(\mathbf{x}^k; \mathbf{a}, \bar{x}) < W(\mathbf{y}; \mathbf{a}, \bar{x})$ for k large enough.

(b) **Never-decisiveness of the future**

Again, let $W(\mathbf{x}; \mathbf{a}, \bar{x}) > W(\mathbf{y}; \mathbf{a}, \bar{x})$ with $\mathbf{x} \in X(\mathbf{a}, \bar{x})$, $\mathbf{y} \in X(\mathbf{a}, \bar{x})$. From (\mathcal{T}_0) , (\mathcal{T}_2) , since $x_t \rightarrow \bar{x}$, $y_t \rightarrow \bar{x}$ when $t \rightarrow +\infty$, one can find \hat{x} , close to \bar{x} , and τ , s.t. for any $T \geq \tau$, there exists s and when $\hat{x} > \bar{x}$, the sequence $\mathbf{y}^T = (\hat{x}, \hat{x}, \dots, \hat{x}, s, y_{T+2}, y_{T+3}, \dots)$ is in $X(\mathbf{a}, \bar{x})$ and when $\hat{x} < \bar{x}$, the sequence $\mathbf{x}^T = (\hat{x}, \hat{x}, \dots, \hat{x}, s, x_{T+2}, z_{T+3}, \dots)$ is in $X(\mathbf{a}, \bar{x})$. We have

$$W(\mathbf{y}^T; \mathbf{a}, \bar{x}) = \sum_{t=0}^T a_t(\hat{x} - \bar{x}) + a_{T+1}(s - \bar{x}) + \sum_{T+2}^{+\infty} a_t(y_t - \bar{x}).$$

When $T \rightarrow +\infty$, $\sum_{t=0}^T a_t(\hat{x} - \bar{x}) \rightarrow +\infty$, $|a_{T+1}(z - \bar{x})| \leq 2\bar{a}$ and $\sum_{T+2}^{+\infty} a_t(y_t - \bar{x}) \rightarrow 0$. Hence $W(\mathbf{y}^T; \mathbf{a}, \bar{x}) > W(\mathbf{x}; \mathbf{a}, \bar{x})$ for T large enough.

We also have

$$W(\mathbf{x}^T; \mathbf{a}, \bar{x}) = \sum_{t=0}^T a_t(\hat{x} - \bar{x}) + a_{T+1}(s - \bar{x}) + \sum_{T+2}^{+\infty} a_t(x_t - \bar{x}).$$

When $T \rightarrow +\infty$, $\sum_{t=0}^T a_t(\hat{x} - \bar{x}) + a_{T+1}(z - \bar{x}) + \sum_{T+2}^{+\infty} a_t(x_t - \bar{x}) \rightarrow -\infty$. Hence $W(\mathbf{x}^T; \mathbf{a}, \bar{x}) < W(\mathbf{y}; \mathbf{a}, \bar{x})$ for T large enough. ■

Corollary 1 *Let (a_t) satisfy*

(i) $\forall t, 0 < \underline{a} \leq a_t \leq \bar{a}$

(ii) $\sum_{t=0}^{\infty} a_t = +\infty$.

Let $X(\mathbf{a}, 0) = \{\mathbf{x} \in [0, 1]^{\infty} : \sum_{t=0}^{\infty} a_t x_t < +\infty\}$, and $W(\mathbf{x}; \mathbf{a}, 0) = \sum_{t=0}^{\infty} a_t x_t$ for $\mathbf{x} \in X(\mathbf{a}, 0)$.

Then $X(\mathbf{a}, 0)$ satisfies (\mathcal{T}_0) , (\mathcal{T}_1) , (\mathcal{T}_2) and $W(\cdot; \mathbf{a}, 0)$ satisfies the two Never-decisiveness axioms on X .

Proof: We just have to check that $X(\mathbf{a}, 0)$ satisfies (\mathcal{T}_0) , (\mathcal{T}_1) and (\mathcal{T}_2) .

\mathcal{T}_0 is obviously true since $x_t \rightarrow 0$ for any $\mathbf{x} \in X(\mathbf{a}, 0)$. \mathcal{T}_1 is satisfied with any $x > 0$, $s = 0$, $s' = 0$. \mathcal{T}_2 is satisfied with any $x > 0$, $s = 0$.

■

Definition 2 *A social welfare function V is anonymous if $V(\mathbf{x}) = V(\mathbf{y})$ when, for any i, j , $i \neq j$, $y_t = x_t, \forall t \neq i, j$, and $y_i = x_j, y_j = x_i$.*

It is immediate that there exists no affine SWF defined on $[0, 1]^\infty$ which is anonymous, excepted the purely finitely additive up to affine transformation. Unfortunately, no purely finitely additive SWF satisfies the Never-decisiveness of the future axiom. Indeed, let $V(x_0, x_1, \dots) > V(y_0, y_1, \dots)$. For any finite change $(x'_0, \dots, x'_T), (y'_0, \dots, y'_T)$ of the present we always have

$$V(x'_0, \dots, x'_T, x_{T+1}, \dots) = V(x_0, x_1, \dots) > V(y_0, y_1, \dots) = V(y'_0, \dots, y'_T, y_{T+1}, \dots)$$

For any $\bar{x} \in [0, 1]$, any $\bar{a} > 0$, define

$$X(\bar{a}, \bar{x}) = \{\mathbf{x} \in [0, 1]^\infty : \bar{a} \sum_{t=0}^{\infty} (x_t - \bar{x}) \in \mathbb{R}\}.$$

Let \mathcal{X} be the collection of $X(\bar{a}, \bar{x})$ which satisfy (\mathcal{T}_0) , (\mathcal{T}_1) and (\mathcal{T}_2) . We have the following Theorem

Theorem 1 *There is a set of domains in $2^{[0,1]^\infty} \setminus \{\emptyset\}$ such that, for each of these domains, there exists a SWF which is independent, anonymous and Pareto. Moreover, for each of these domains, the SWF satisfies the two Never-decisiveness properties.*

Proof: Consider \mathcal{X} . Define $W(.; \bar{a}, \bar{x})$ by: for any $X(\bar{a}, \bar{x}) \in \mathcal{X}$, any $\mathbf{x} \in X(\bar{a}, \bar{x})$,

$$W(\mathbf{x}; \bar{a}, \bar{x}) = \bar{a} \sum_{t=0}^{\infty} (x_t - \bar{x}).$$

From Definition 1, it is independent. Obviously, $W(.; \bar{a}, \bar{x})$ is anonymous and Pareto. From Proposition 4, it satisfies the two Never-decisiveness properties.

■

Corollary 2 *Let $\bar{a} > 0$ and*

$$X(\bar{a}, 0) = \{\mathbf{x} \in [0, 1]^\infty : \bar{a} \sum_{t=0}^{\infty} x_t < +\infty\}.$$

Then $X(\bar{a}, 0) \in \mathcal{X}$. The SWF $W(\cdot; \bar{a}, 0)$ defined by

$$\forall \mathbf{x} \in X(\bar{a}, 0), W(\mathbf{x}; \bar{a}, 0) = \bar{a} \sum_{t=0}^{\infty} x_t$$

is anonymous, independent, Pareto.

Proof: From Corollary 1, $X(\bar{a}, 0) \in \mathcal{X}$. Then apply Theorem 1.

■

3 The overtaking criterion as an anonymous and never-decisive criterion for optimal growth models

The difficulty is to exhibit a criterion which satisfies assumptions (\mathcal{T}_0) , (\mathcal{T}_1) , (\mathcal{T}_2) on some set X of sequences. We will show that the overtaking criterion introduced by Gale [1967] for optimal growth models actually induces an anonymous and never-decisive SWF on the set of utilities of good programmes of an intertemporal economy.

First, we construct the subset X on which the SWF (the overtaking criterion) is defined. It will be the set of the utilities yielded by good programmes. We have the property that any good programme converges, and therefore the utilities yielded by good programmes converge too (assumption (\mathcal{T}_0) above is satisfied). Second, we prove that under the assumptions on the economy presented below, the utilities of good programmes satisfy (\mathcal{T}_1) and (\mathcal{T}_2) .

We consider an intertemporal economy where the instantaneous utility of the representative consumer depends on κ_t , the capital stock on hand at date t , and on κ_{t+1} , the capital stock for date $t + 1$. Given κ_t , the set of feasible capital stocks for the next period $t + 1$ is $\Gamma(\kappa_t)$. We assume that at any period t the feasible capital stock on hand belongs to A , a subset of \mathbb{R}_+^n . More explicitly, we make the following assumptions⁸.

⁸This set-up is borrowed from Dana and Le Van [1990], but we do not assume here that $\text{graph}\Gamma$ is convex, i.e. we do not impose a technology with decreasing returns to scale.

Assumptions

H1: A is a compact, convex set of \mathbb{R}_+^n with non-empty interior.

H2: Γ is a continuous correspondence with non-empty images. It satisfies $\Gamma(A) \subseteq A$. Its graph is the set $\text{graph}\Gamma = \{(\kappa, \chi) \in A \times A : \chi \in \Gamma(\kappa)\}$.

H3: $\text{intgraph}\Gamma$, the interior of the graph of Γ , is non-empty.

We will denote by $\text{cograph}\Gamma$ the convex hull of $\text{graph}\Gamma$.

H4: The instantaneous utility function $u : \text{cograph}\Gamma \rightarrow \mathbb{R}_+$ is strictly concave and continuous. It is increasing with respect to the first variable and decreasing with respect to the second variable.

For the simplicity of our presentation we also assume the following.

H5: The function u is differentiable in the interior of $\text{cograph}\Gamma$.

H6: The set $I(A) = \{\kappa \in A : (\kappa, \kappa) \in \text{graph}\Gamma\}$ is non empty.

Remark 1 1. Dana and Le Van [1990] show that the following assumptions imply $\text{intgraph}\Gamma \neq \emptyset$.

- Free disposal assumption: If $\chi \in \Gamma(\kappa)$, $\kappa' \geq \kappa$, $\chi' \leq \chi$, $\kappa' \in A$, $\chi' \in A$, then $\chi' \in \Gamma(\kappa')$.

- Existence of expansible capital stocks: There exists $(\kappa, \chi) \in \text{graph}\Gamma$, with $\chi \gg \kappa$, i.e. $\chi_i > \kappa_i$, $\forall i \in \{1, \dots, n\}$.

2. If Γ is continuous and satisfies the condition:

There exists a number a such that $\{\|\kappa\| \geq a, \chi \in \Gamma(\kappa)\} \Rightarrow \|\chi\| < \|\kappa\|$, then $I(A)$ is compact.

3. An example where **H1–H2** and **H5** are satisfied. Let $\Gamma(\kappa) = \{\chi \in \mathbb{R}_+ : \chi \in [0, f(\kappa)]\}$, where f is a continuous increasing function which satisfies $f(0) = 0$, $f(1) = 1$, $f(\kappa) < \kappa$, $\forall \kappa > 1$. Take $A = [0, a]$ with $a > 1$.

A sequence κ is feasible from $\kappa_0 \in X$ if, $\forall t \geq 0$, $\kappa_{t+1} \in \Gamma(\kappa_t)$. We denote by $\Pi(\kappa_0)$ the set of feasible sequences from κ_0 . Following Gale [1967], a programme from κ_0 is a feasible sequence from κ_0 . The set of programmes is denoted by Π , i.e. $\Pi = \cup_{\kappa \in A} \Pi(\kappa)$.

Lemma 1 Assume **H1–H2**. Then $\Pi(\kappa_0)$ is compact for the product topology.

Proof: See e.g. Le Van and Dana [2003], chapter 4. ■

3.1 Good Programmes

Definition 3 A stationary point $(\bar{\kappa}, \bar{\chi})$ satisfies

$$\bar{x} = u(\bar{\kappa}, \bar{\chi}) = \max \{u(\kappa, \chi) : (\kappa, \chi) \in \text{graph}\Gamma\}.$$

Proposition 5 There exists a stationary point $(\bar{\kappa}, \bar{\chi})$ which satisfies

- (i) $\bar{x} = u(\bar{\kappa}, \bar{\chi}) = \max \{u(\kappa, \chi) : (\kappa, \chi) \in \text{graph}\Gamma\}$;
- (ii) if $(\bar{\kappa}, \bar{\chi}) \in \text{intgraph}\Gamma$, then

$$u(\kappa, \chi) + u_1(\bar{\kappa}, \bar{\chi})\chi - u_1(\bar{\kappa}, \bar{\chi})\kappa \leq u(\bar{\kappa}, \bar{\chi}), \quad \forall (\kappa, \chi) \in \text{graph}\Gamma,$$

where $u_1(\kappa, \chi) = \partial u(\kappa, \chi)/\partial \kappa$ and $u_2(\kappa, \chi) = \partial u(\kappa, \chi)/\partial \chi$.

Proof: (i) Since $I(A)$ is compact, the proof of the existence of $(\bar{\kappa}, \bar{\chi})$ is obvious.

(ii) If $(\bar{\kappa}, \bar{\chi})$ is interior then $u_1(\bar{\kappa}, \bar{\chi}) + u_2(\bar{\kappa}, \bar{\chi}) = 0$. Since u is concave, for any $(\kappa, \chi) \in \text{graph}\Gamma$, we have

$$u(\bar{\kappa}, \bar{\chi}) - u(\kappa, \chi) \geq u_1(\bar{\kappa}, \bar{\chi})(\bar{\kappa} - \kappa) + u_2(\bar{\kappa}, \bar{\chi})(\bar{\chi} - \chi) = u_1(\bar{\kappa}, \bar{\chi})\chi - u_1(\bar{\kappa}, \bar{\chi})\kappa.$$

■

We now add another assumption.

H7: $(\bar{\kappa}, \bar{\chi}) \in \text{intgraph}\Gamma$.

Assumption **H7** is not very restrictive, since in the one dimension Ramsey model it implies that consumption is strictly positive at the Golden Rule. Indeed, if $\Gamma(\kappa) = \{\chi : 0 \leq \chi \leq f(\kappa)\}$ where f is the production function, then $\bar{\kappa}$ satisfies $\bar{x} = \max_{\kappa} u(f(\kappa) - \kappa)$.

Let $\bar{p} = u_1(\bar{\kappa}, \bar{\chi})$.

Proposition 6 Under **H1-H7**, $(\bar{\kappa}, \bar{\chi})$ is the unique solution to problem

$$\max \{u(\kappa, \chi) + \bar{p}\chi - \bar{p}\kappa : (\kappa, \chi) \in \text{graph}\Gamma\}.$$

The proof is in Appendix 3.

Theorem 2 Assume **H1-H7**. For any programme κ , either

- (i) $\lim_{T \rightarrow +\infty} \sum_{t=0}^T [u(\kappa_t, \kappa_{t+1}) - \bar{x}]$ exists in \mathbb{R} and $\kappa_t \rightarrow \bar{\kappa}$, or
- (ii) $\lim_{T \rightarrow +\infty} \sum_{t=0}^T [u(\kappa_t, \kappa_{t+1}) - \bar{x}] = -\infty$.

The proof is in Appendix 3.

Definition 4 A programme $\kappa \in \Pi(\kappa_0)$ is good if $\lim_{T \rightarrow \infty} \sum_{t=0}^T [u(\kappa_t, \kappa_{t+1}) - u(\bar{\kappa}, \bar{\kappa})]$ exists in \mathbb{R} .

Definition 5 The set of the good programmes $\kappa \in \Pi(\kappa_0)$ is denoted $G(\kappa_0)$.

From Theorem 2 one gets:

Corollary 3 Assume **H1-H7**. If $\kappa \in G(\kappa_0)$, then $\kappa_t \rightarrow \bar{\kappa}$.

3.2 Never-decisiveness and anonymous criteria

Since A is compact, there exists a number $a > 0$ such that for any program κ we have $u(\kappa_t, \kappa_{t+1}) \in [0, a]$, $\forall t$. We define

$$U = \{(x_t)_t : \forall t, x_t = u(\kappa_t, \kappa_{t+1}), \kappa \in \Pi\}$$

and

$$X = \{\mathbf{x} \in U : \sum_{t=0}^{\infty} (x_t - \bar{x}) \in \mathbb{R}\}$$

Observe that we have $X = \cup_{\kappa} G(\kappa)$.

Let

$$W(\mathbf{x}) = \sum_{t=0}^{\infty} (x_t - \bar{x})$$

and

$$\mathcal{W}(\kappa) = \sum_{t=0}^{\infty} [u(\kappa_t, \kappa_{t+1}) - \bar{x}].$$

Proposition 7 The social welfare function W is anonymous and satisfies the Never-decisiveness properties on X .

Proof: Anonymity is obvious. To prove that this SWF satisfies the Never-decisiveness properties, we apply Proposition 4. We prove that assumptions (\mathcal{T}_0) , (\mathcal{T}_1) , (\mathcal{T}_2) are satisfied.

Assumption (\mathcal{T}_0) comes from Corollary 3.

To check (\mathcal{T}_1) : Since $(\bar{\kappa}, \bar{\kappa}) \in \text{intgraph}\Gamma$, there exists an open ball $B((\bar{\kappa}, \bar{\kappa}), \varepsilon) \subset \text{graph}\Gamma$ which satisfies the following properties:

- (i) There exists $(\hat{\kappa}, \hat{\kappa}) \in B((\bar{\kappa}, \bar{\kappa}), \varepsilon)$ with $\hat{\kappa} \neq \bar{\kappa}$,
- (ii) $(\hat{\kappa}, \bar{\kappa}) \in B((\bar{\kappa}, \bar{\kappa}), \varepsilon)$,
- (iii) For any y close enough to $\bar{\kappa}$, we have $(y, \hat{\kappa}) \in B((\bar{\kappa}, \bar{\kappa}), \varepsilon)$.

Since κ_t converges to $\bar{\kappa}$, there exists N such that for any $T \geq N$ we have $(\kappa_{T-1}, \kappa_T) \in B((\bar{\kappa}, \bar{\kappa}), \varepsilon)$ and $(\kappa_T, \hat{\kappa}) \in B((\bar{\kappa}, \bar{\kappa}), \varepsilon)$.

Define $s = u(\kappa_T, \hat{\kappa})$, $\hat{x} = u(\hat{\kappa}, \hat{\kappa})$, $s' = u(\hat{\kappa}, \bar{\kappa})$. Observe that $\hat{x} < \bar{x}$.

To check (\mathcal{T}_2) : take $\hat{x} = u(\hat{\kappa}, \hat{\kappa}) < \bar{x}$ and for any T large enough $s = u(\hat{\kappa}, \kappa_{T+1})$. ■

4 Solving optimal growth models with the overtaking criterion

We want to solve an optimal growth model with the SWF defined above. We first have:

Proposition 8 *Assume **H1–H7**. Then \mathcal{W} is upper semi-continuous for the product topology.*

Proof: See Dana and Le Van [1990]. ■

We then obtain

Theorem 3 *Assume **H1–H7**. If $G(\kappa_0) \neq \emptyset$, then there exists an optimal path. It is unique if $\text{graph}\Gamma$ is convex.*

Proof: From proposition 8 and lemma 1, for the product topology, \mathcal{W} is upper semi-continuous and $\Pi(\kappa_0)$ is compact. Therefore, an optimal path exists, since the problem is to maximize $\mathcal{W}(\kappa)$ with $\kappa \in \Pi(\kappa_0)$. This optimal path is unique when $\text{graph}\Gamma$ is convex since u is assumed strictly concave. ■

4.1 Applications

We now give examples of models which satisfy **H1–H7** and for which X is non-empty.

4.1.1 Convex technology

Assume $\text{graph}\Gamma$ is convex. Now, let $A = [0, 1]$, $\Gamma(\kappa) = [0, f(\kappa)]$ where f is strictly concave, increasing, differentiable with $f'(0) > 1$, $f(0) = 0$, $f(1) = 1$, and $u(\kappa, \chi) = v(f(\kappa) - \chi)$, where v is real-valued, defined on \mathbb{R}_+ , increasing, strictly concave, differentiable. It is easy to check **H1–H6**. Observe that the stationary point $\bar{\kappa}$ is defined by $f'(\bar{\kappa}) = 1$. It is therefore

unique. Obviously, $0 < \bar{\kappa} < f(\bar{\kappa}) < 1$. In other words, **H7** is satisfied. We claim that $G(\kappa_0) \neq \emptyset$, $\forall \kappa_0 > 0$. Indeed, since $f^t(\kappa_0) \rightarrow 1$ when $t \rightarrow +\infty$, there exists T such that $f^T(\kappa_0) \leq f(\bar{\kappa})$ and $f^{T+1}(\kappa_0) > f(\bar{\kappa})$. The sequence $(\kappa_0, f(\kappa_0), \dots, f^T(\kappa_0), \bar{\kappa}, \bar{\kappa}, \dots, \bar{\kappa}, \dots) \in G(\kappa_0)$.

4.1.2 Economy with a convex-concave production function

Let $A = [0, 1]$, $\Gamma(\kappa) = [0, f(\kappa)]$. Here we suppose that f is increasing, continuously differentiable, $f(0) = 0$, $f(1) = 1$, strictly convex between 0 and $\kappa_I < 1$ and strictly concave in $[\kappa_I, +\infty[$. The function u is as before $u(\kappa, \chi) = v(f(\kappa) - \chi)$, where v is real-valued, defined on \mathbb{R}_+ , increasing, strictly concave, differentiable.

Let $a = \max_{\kappa \in A} \left\{ \frac{f(\kappa)}{\kappa} \right\}$ and $\kappa_a \in A$ satisfy $a = \left\{ \frac{f(\kappa_a)}{\kappa_a} \right\}$. Let F be defined as follows: $F(\kappa) = a\kappa$, $\forall \kappa \leq \kappa_a$; $F(\kappa) = f(\kappa)$, $\kappa \geq \kappa_a$. Then F is concave and $\{(\kappa, \chi) \in A \times A : \chi \leq F(\kappa)\}$ is cograph Γ .

When $f'(0) > 1$, there exists a unique $\bar{\kappa} \in (\kappa_I, 1)$ such that $f'(\bar{\kappa}) = 1$. One can check that $0 < \bar{\kappa} < f(\bar{\kappa}) < 1$, i.e. $(\bar{\kappa}, \bar{\kappa}) \in \text{int graph}\Gamma$. Assumptions **H1–H7** are satisfied. Again $G(0) = \emptyset$ and $G(\kappa_0) \neq \emptyset$ if $\kappa_0 > 0$.

Now assume $f'(0) < 1 < \frac{f(\kappa_a)}{\kappa_a}$ and $f'(\kappa_I) > 1$ (see figure 1). There are two points $\underline{\kappa}, \bar{\kappa}$ such that $f'(\underline{\kappa}) = f'(\bar{\kappa}) = 1$. But $f(\underline{\kappa}) - \underline{\kappa} < 0$, i.e. $(\underline{\kappa}, \underline{\kappa}) \notin \text{graph}\Gamma$. We have a unique stationary point $\bar{\kappa}$. It satisfies $0 < \bar{\kappa} < f(\bar{\kappa}) < 1$, i.e. $(\bar{\kappa}, \bar{\kappa}) \in \text{int graph}\Gamma$. Again, **H1–H7** are satisfied.

Let $\tilde{\kappa}$ satisfy $f(\tilde{\kappa}) = \tilde{\kappa}$. Then $\tilde{\kappa} \in (0, 1)$. Moreover, $\kappa < \tilde{\kappa} \Rightarrow f(\kappa) < \kappa$ and $1 > \kappa > \tilde{\kappa} \Rightarrow f(\kappa) > \kappa$. That implies:

- (i) any feasible path from $\kappa_0 < \tilde{\kappa}$ will converge to 0. Hence, $G(\kappa_0) = \emptyset$ for any $\kappa_0 \leq \tilde{\kappa}$.
- (ii) for any $\kappa_0 > \tilde{\kappa}$, $f^t(\kappa_0) \rightarrow 1$. Hence, one can find T such that $f^T(\kappa_0) \leq \bar{\kappa}$ and $f^{T+1}(\kappa_0) > f(\bar{\kappa})$. The sequence $(\kappa_0, f(\kappa_0), \dots, f^T(\kappa_0), \bar{\kappa}, \bar{\kappa}, \dots, \bar{\kappa}, \dots) \in G(\kappa_0)$. Thus, $G(\kappa_0) \neq \emptyset$ for any $\kappa_0 > \tilde{\kappa}$.

One concludes that $\tilde{\kappa}$ is the poverty trap.

4.1.3 Growth and exhaustible resources

We consider the model in Le Van, Schubert and Nguyen [2010]. The country possesses a stock of a non-renewable natural resource S_0 . This resource is extracted at a rate R_t , and then sold abroad at a price P_t , in terms of the numeraire, which is the domestic single consumption good. The inverse demand function for the resource is $P(R_t)$. The revenue from the sale of the natural resource, $\phi(R_t) = P(R_t)R_t$, is used to buy a foreign good, which




Figure 1: The convex-concave technology

is supposed to be a perfect substitute of the domestic good, used for consumption and capital accumulation. The domestic production function is $F(k_t)$, where k_t is physical capital. The depreciation rate is δ . We define the function $f(k_t) = F(k_t) + (1 - \delta)k_t$, and we name it for simplicity the technology. The constraints of the economy are:

$$\begin{aligned} \forall t, c_t &\geq 0, k_t \geq 0, R_t \geq 0, \\ c_t + k_{t+1} &\leq f(k_t) + \phi(R_t), \\ \sum_{t=0}^{+\infty} R_t &\leq S_0, \\ S_0 &> 0, k_0 \geq 0 \text{ are given,} \end{aligned}$$

where c_t is consumption. We assume that f is strictly concave, $f(0) = 0$, $f'(+\infty) < 1$, $f'(0) > 1$. Let \tilde{k} satisfy $\tilde{k} = f(\tilde{k}) + \phi(S_0)$. Such a \tilde{k} is unique. Take $A = [0, \tilde{k}] \times [0, S_0]$. Let $\kappa = (k, R)$. Define $\Gamma(\kappa) = \{\chi = (y, R) : 0 \leq y \leq f(k) + \phi(R)\}$. One can check that Γ is continuous and maps A into A . Observe that any feasible sequence $\{R_t\}$ will converge to zero. We also assume $\phi'(0) < +\infty$. The representative consumer has an instantaneous utility function u which is increasing, strictly concave, satisfies

$u(0) = 0$. We show that the methodology of Sections 2–4 may be used for the present model.

Let \bar{k} satisfy

$$\begin{aligned} u(f(\bar{k}) - \bar{k}) &= \max \{u(f(k) - k) : 0 \leq k \leq f(k)\} \\ &= \max \{u(f(k) - k) : 0 \leq k \leq f(k) + \phi(S_0)\} \end{aligned}$$

Such a \bar{k} is unique and satisfies $0 < \bar{k} < f(\bar{k})$ and $f'(\bar{k}) = 1$. Let $\bar{p} = u'(f(\bar{k}) - \bar{k})$. We let to the reader to prove the following claim (see the proofs of Propositions 5, 6).

Claim 1

$(\bar{k}, \bar{k}, 0)$ is the unique solution to problem

$$\max \{u(f(k) - y + \phi(R)) + \bar{p}y - \bar{p}k - \bar{p}\phi'(0)R : ((k, R), (y, R)) \in \text{graph}\Gamma\}.$$

Let $\bar{x} = u(f(\bar{k}) - \bar{k})$.

Claim 2 For any program (\mathbf{k}, \mathbf{R}) , either

- (i) $\lim_{T \rightarrow +\infty} \sum_{t=0}^T [u(f(k_t) - k_{t+1} + \phi(R_t)) - \bar{x}]$ exists in \mathbb{R} and $k_t \rightarrow \bar{k}$, $R_t \rightarrow 0$, or
- (ii) $\lim_{T \rightarrow +\infty} \sum_{t=0}^T [u(f(k_t) - k_{t+1} + \phi(R_t)) - \bar{x}] = -\infty$

Proof of Claim 2

(i) and (ii): Observe that

$$\begin{aligned} &\sum_{t=0}^T [u(f(k_t) - k_{t+1} + \phi(R_t)) + \bar{p}k_{t+1} - \bar{p}k_t - \bar{p}\phi'(0)R_t - \bar{x}] \\ &= \sum_{t=0}^T [u(f(k_t) - k_{t+1} + \phi(R_t)) - \bar{x}] + \bar{p}(k_{T+1} - k_0) - \bar{p}\phi'(0) \sum_{t=0}^T R_t. \end{aligned}$$

Since $\sum_{t=0}^T R_t$ converges, use Claim 1 and the proof of Theorem 2 to conclude.

A programme $\{(\mathbf{k}, \mathbf{R})\}$ is *good* if $\sum_{t=0}^{+\infty} [u(f(k_t) - k_{t+1} + \phi(R_t)) - \bar{x}] \in \mathbb{R}$.

Now define

$$W(\mathbf{k}, \mathbf{R}) = \lim_{T \rightarrow +\infty} \sum_{t=0}^T [u(f(k_t) - k_{t+1} + \phi(R_t)) - \bar{x}].$$

W is upper semi-continuous and one can check that it is never-decisive. The optimal solution $(\mathbf{k}^*, \mathbf{R}^*)$ to

$$\max_{(\mathbf{k}, \mathbf{R})} W(\mathbf{k}, \mathbf{R})$$

under the constraints

$$\begin{aligned} \forall t, 0 \leq k_{t+1} &\leq f(k_t) + \phi(R_t), \quad k_0 \text{ is given} \\ \sum_{t=0}^{+\infty} R_t &\leq S_0, \end{aligned}$$

will converge to $(\bar{k}, 0)$.

4.1.4 Growth and renewable resources

We study the canonical model of growth with a renewable resource⁹. The economy possesses a stock of a renewable natural resource S_0 . This resource is extracted at a rate R_t . The domestic production function is $F(k_t, R_t)$. k_t is the stock of physical capital. The depreciation rate is δ . The natural growth of the renewable resource is $S_{t+1} - S_t = H(S_t)$. The function H is strictly concave, differentiable, satisfies $H(0) = 0$, $H(\hat{S}) = 0$, with $\hat{S} > 0$. There is a representative consumer with a utility function u depending on her consumption c . The technological constraints for each period t are as follows:

$$\begin{aligned} c_t + k_{t+1} &\leq F(k_t, R_t) + (1 - \delta)k_t \\ S_{t+1} &\leq S_t + H(S_t) - R_t. \end{aligned}$$

Let $\psi(S) = S + H(S)$. The function ψ is strictly concave, $\psi(0) = 0$, $\psi'(+\infty) < 1$, since $H'(+\infty) < 0$. We define the function $f(k_t, R_t) = F(k_t, R_t) + (1 - \delta)k_t$, and we name it for simplicity the technology. We assume that F is strictly concave, $F(0, R) = F(k, 0) = 0$, $F_1(+\infty, \psi(\hat{S})) < \delta$, $F_1(0, \psi(\hat{S})) > \delta$. The constraints become

$$\begin{aligned} 0 \leq k_{t+1} &\leq f(k_t, R_t) \\ 0 \leq S_{t+1} &\leq \psi(S_t) - R_t. \end{aligned}$$

Let \hat{k} satisfy $\hat{k} = f(\hat{k}, \psi(\hat{S}))$. Let $A = [0, \hat{k}] \times [0, \hat{S}]$ and for $\kappa = (k, S)$ define

$$\Gamma(\kappa) = \{\chi = (k', S') : 0 \leq k' \leq f(k, \psi(S) - S'), 0 \leq S' \leq \psi(S)\}$$

⁹See for instance Ayong Le Kama [2001].

Γ is continuous and maps A into A . Its graph is convex. There exists a unique $\bar{\kappa} = (\bar{k}, \bar{S})$ such that $(\bar{\kappa}, \bar{\kappa})$ is in the interior of $\text{graph}\Gamma$ and solves

$$\max \{u(f(k, \psi(S) - S) - k) : ((k, S), (k, S)) \in \text{graph}\Gamma\}.$$

Namely, $f_1(\bar{k}, \psi(\bar{S}) - \bar{S}) = 1$, $\psi'(\bar{S}) = 1$. This model satisfies assumptions **H1–H7**. Hence, one can define

$$W(\kappa_0, \kappa_1, \dots) = W(\mathbf{k}, \mathbf{S}) = \lim_{T \rightarrow +\infty} \sum_{t=0}^T [u(f(k_t, \psi(S_t) - S_{t+1}) - k_{t+1}) - \bar{x}]$$

where $\bar{x} = u(f(\bar{k}, \psi(\bar{S}) - \bar{S}) - \bar{k})$. The optimal solution to

$$\max_{\{\kappa_1, \kappa_2, \dots\}} W(\kappa_0, \kappa_1, \dots) = \max_{(\mathbf{k}, \mathbf{S})} W(\mathbf{k}, \mathbf{S})$$

under the constraints

$$\begin{aligned} \forall t, 0 \leq k_{t+1} &\leq f(k_t, \psi(S_t) - S_{t+1}), \\ 0 \leq S_{t+1} &\leq \psi(S_t), \quad (k_0, S_0) \text{ are given} \end{aligned}$$

will converges to (\bar{k}, \bar{S}) if there exists a good programme from (k_0, S_0) .

Appendix 1

We give a simple example where a SWF which satisfies our Never-decisiveness axioms does not satisfy the axiom of Hammond Equity for the Future. First, our criterion is not complete on $[0, 1]^\infty$. Now, consider \mathbf{x}, \mathbf{y} defined by $x_0 > 0$, $x_t = 0, \forall t \geq 1$, $y_0 = \frac{x_0}{2}, y_t = (\frac{1}{8} \times \frac{1}{2^{t-1}})x_0$, for $t \geq 1$. These sequences satisfy: $x_0 > y_0 > y_t > x_t$ for any $t \geq 1$. However we have $\sum_{t=0}^\infty x_t = x_0 > \sum_{t=0}^\infty y_t = \frac{3}{4}x_0$.

Appendix 2

Consider two problems:

$$\mathcal{R} : \max \sum_{t=0}^{\infty} \beta^t u(c_t)$$

$$\mathcal{C} : \max \sum_{t=0}^{\infty} \beta^t u(c_t) + \phi(u(c_1), u(c_2), \dots)$$

under the constraint $c_t + k_{t+1} \leq f(k_t)$ for all t , with $k_0 > 0$ given.

Let $\Pi(k_0)$ denote the set of feasible capital stocks. We make the following assumptions.

H0 $0 < \beta < 1$.

H1 The function $u : \mathbb{R} \rightarrow \mathbb{R}$ is twice continuously differentiable and satisfies $u(0) = 0$. Moreover, its derivatives satisfy $u' > 0$ (strictly increasing) and $u'' < 0$ (strictly concave).

H2 Inada condition $u'(0) = +\infty$.

H3 The function $f : \mathbb{R} \rightarrow \mathbb{R}$ is twice continuously differentiable and satisfies $f(0) = 0$. Its derivatives satisfy $f' > 0$ (strictly increasing), $f'' < 0$, strictly concave, $\lim_{x \rightarrow \infty} f'(x) < 1$.

H4 ϕ is purely finitely additive, continuous for the l_∞ -topology, homogeneous of degree 1, non decreasing, $\phi(0) = 0$, $\phi(\mathbf{1}) > 0$.

Lemma 2 Let $\mathbf{x} \in l_\infty$ be such that $\lim_{t \rightarrow \infty} x_t = \bar{x}$, then

$$\phi(\mathbf{x}) = \phi(\bar{x}, \bar{x}, \dots)$$

Proof: Denote $\bar{\mathbf{x}} = (\bar{x}, \bar{x}, \dots)$. Construct \mathbf{x}^n as:

$$\begin{aligned} x_t^n &= \bar{x} \text{ for } 0 \leq t \leq n \\ x_t^n &= x_t \text{ for } t > n \end{aligned}$$

Since ϕ is purely finitely additive, so for all n , $\phi(\mathbf{x}^n) = \phi(\mathbf{x})$. On the other hand, $\|\mathbf{x}^n - \bar{\mathbf{x}}\| = \sup_{t > n} |x_t - \bar{x}| \Rightarrow \lim_{n \rightarrow \infty} \|\mathbf{x}^n - \bar{\mathbf{x}}\| = 0$, then

$$\begin{aligned} \phi(\mathbf{x}) &= \lim_{n \rightarrow \infty} \phi(\mathbf{x}^n) \\ &= \phi(\bar{\mathbf{x}}) \end{aligned}$$

■

For each $\mathbf{k} \in \Pi(k_0)$, define:

$$\Phi(\mathbf{k}) = \sum_{t=0}^{\infty} \beta^t u(f(k_t) - k_{t+1}) + \phi(u(f(k_0) - k_1), u(f(k_1) - k_2), \dots),$$

$$W(\mathbf{k}) = \sum_{t=0}^{\infty} \beta^t u(f(k_t) - k_{t+1})$$

$$\varphi(\mathbf{k}) = \phi(u(f(k_0) - k_1), u(f(k_1) - k_2), \dots). \quad (2)$$

Lemma 3 *We have*

$$\sup_{\mathbf{k} \in \Pi(k_0)} \Phi(\mathbf{k}) \geq \sup_{\mathbf{k} \in \Pi(k_0)} W(\mathbf{k}) + \sup_{\mathbf{k} \in \Pi(k_0)} \varphi(\mathbf{k}). \quad (3)$$

Proof: Take any $\epsilon > 0$ satisfying $\epsilon < \sup_{\mathbf{k} \in \Pi(k_0)} \varphi(\mathbf{k})$. Fix T big enough such that for all $\mathbf{k} \in \Pi(k_0)$, we have

$$\sum_{t=T+1}^{\infty} \beta^t u(f(k_t) - k_{t+1}) < \epsilon.$$

Denote by \mathbf{k}^* the solution to Ramsey problem. Take \mathbf{k}' feasible, such that

$$\varphi(\mathbf{k}') > \sup_{\mathbf{k} \in \Pi(k_0)} \varphi(\mathbf{k}) - \epsilon > 0.$$

Define $f^t(k) = f(f(\dots(k)\dots))$, t times. We will prove that there exists τ satisfying $k'_{T+\tau} < f^\tau(k_T^*)$.

Indeed, suppose the contrary, $f^\tau(k_T^*) \leq k'_{T+\tau}$ for all τ . Denote by \bar{k} the solution to the equation $f(k) = k$. We can verify easily that $f^t(k) \rightarrow \bar{k}$ when $t \rightarrow \infty$, for all $k > 0$.

We have:

$$f^\tau(k_T^*) \leq k'_{T+\tau} \leq f^{T+\tau}(k_0).$$

From this inequality, we have $k'_t \rightarrow \bar{k}$ when t tends to infinity. Hence $f(k_t) - k_{t+1} \rightarrow 0$ when $t \rightarrow \infty$. This implies, by using the Lemma 2, $\varphi(\mathbf{k}') = 0$, a contradiction.

Now take τ such that $k'_{T+\tau} < f^\tau(k_T^*)$. Define \mathbf{k}'' as

$$\mathbf{k}'' = (k_0, k_1^*, k_2^*, \dots, k_T^*, f(k_T^*), f^2(k_T^*), \dots, f^{\tau-1}(k_T^*), k'_{T+\tau}, k'_{T+\tau+1}, \dots).$$

Since $k'_{T+\tau} < f^\tau(k_T^*)$, the sequence \mathbf{k}'' is feasible. Observe that $\varphi(\mathbf{k}'') = \varphi(\mathbf{k}')$. We have

$$\begin{aligned} \Phi(\mathbf{k}'') &= W(\mathbf{k}'') + \varphi(\mathbf{k}'') \\ &> \sum_{t=0}^T \beta^t u(f(k_t^*) - k_{t+1}^*) + \varphi(\mathbf{k}') \\ &> \sup_{\mathbf{k} \in \Pi(k_0)} W(\mathbf{k}) + \sup_{\mathbf{k} \in \Pi(k_0)} \varphi(\mathbf{k}) - 2\epsilon. \end{aligned}$$

This inequality is true for all ϵ small enough. Let $\epsilon \rightarrow 0$, we obtain (3). ■

Claim 1 (1) *Suppose that $f'(0) > 1$. Then problem \mathcal{C} has no solution.*

(2) *Suppose $f'(0) \leq 1$. Then problems \mathcal{C} and \mathcal{R} have the same solution.*

Proof: (1) Suppose that \mathcal{C} has a solution $\bar{\mathbf{k}}$. From the Lemma 3, we have $\bar{\mathbf{k}}$ is a solution to Ramsey problem and also a solution to problem

$$\max_{\mathbf{k} \in \Pi(k_0)} \varphi(\mathbf{k}).$$

Since $\bar{\mathbf{k}}$ is solution to Ramsey problem, we have $\bar{k}_t \rightarrow k^s$, solution to $f'(k) = \frac{1}{\beta}$, if $f'(0) > \frac{1}{\beta}$ and $k^s = 0$ if $f'(0) \leq \frac{1}{\beta}$. Denote by \tilde{k} the solution to $f'(k) = 1$. Observe that $f(k^s) - k^s < f(\tilde{k}) - \tilde{k}$.

Since $\tilde{k} < \bar{k}$, there exists T big enough such that $f^T(k_0) > \tilde{k}$. This implies the sequence

$$k''' = (k_0, f(k_0), \dots, f^T(k_0), \tilde{k}, \tilde{k}, \dots)$$

is feasible.

Since $f(k^s) - k^s < f(\tilde{k}) - \tilde{k}$, and by using the Lemma 2, we have

$$\varphi(\bar{\mathbf{k}}) < \varphi(k'''),$$

a contradiction.

(2) The proof is trivial since in this case any feasible sequence \mathbf{k} converges to 0 and hence $\varphi(\mathbf{k}) = 0$. ■

Appendix 3

Proof of Proposition 6:

Let $(\hat{\kappa}, \hat{\chi})$ be another solution. Then

$$u(\hat{\kappa}, \hat{\chi}) + \bar{p}\hat{\chi} - \bar{p}\hat{\kappa} = u(\bar{\kappa}, \bar{\kappa}). \quad (4)$$

Since $(\bar{\kappa}, \bar{\kappa}) \in \text{intgraph } \Gamma$, for $\lambda \in (0, 1)$, close enough to 1, we have

$$(\lambda\bar{\kappa} + (1-\lambda)\hat{\kappa}, \lambda\bar{\kappa} + (1-\lambda)\hat{\chi}) \in \text{graph}\Gamma.$$

Denote $\kappa_\lambda = \lambda\bar{\kappa} + (1-\lambda)\hat{\kappa}$, $\chi_\lambda = \lambda\bar{\kappa} + (1-\lambda)\hat{\chi}$. Then

$$\lambda u(\bar{\kappa}, \bar{\kappa}) + (1-\lambda)u(\hat{\kappa}, \hat{\chi}) + \bar{p}\chi_\lambda - \bar{p}\kappa_\lambda < u(\kappa_\lambda, \chi_\lambda) + \bar{p}\chi_\lambda - \bar{p}\kappa_\lambda \leq u(\bar{\kappa}, \bar{\kappa}),$$

and then

$$\lambda u(\bar{\kappa}, \bar{\kappa}) + (1-\lambda)[u(\hat{\kappa}, \hat{\chi}) + \bar{p}\hat{\chi} - \bar{p}\hat{\kappa}] < u(\bar{\kappa}, \bar{\kappa})$$

or equivalently

$$u(\hat{\kappa}, \hat{\chi}) + \bar{p}\hat{\chi} - \bar{p}\hat{\kappa} < u(\bar{\kappa}, \bar{\kappa}),$$

in contradiction with (4).

Proof of theorem 2:

Theorem 2 is identical to Proposition 1.3.1 and Corollary 1.3.2. in Le Van and Dana [2003]. But we have to modify the proof since $\text{graph}\Gamma$ is not assumed to be convex. Consider $\lim_{T \rightarrow +\infty} \sum_{t=0}^T [u(\kappa_t, \kappa_{t+1}) + \bar{p}\kappa_{t+1} - \bar{p}\kappa_t - \bar{x}]$ which exists in $\mathbb{R}_- \cup \{-\infty\}$ since $\forall t, u(\kappa_t, \kappa_{t+1}) + \bar{p}\kappa_{t+1} - \bar{p}\kappa_t - \bar{x} \leq 0$. We have

$$\sum_{t=0}^T [u(\kappa_t, \kappa_{t+1}) + \bar{p}\kappa_{t+1} - \bar{p}\kappa_t - \bar{x}] = \sum_{t=0}^T [u(\kappa_t, \kappa_{t+1}) - \bar{x}] - \bar{p}(\kappa_0 - \kappa_{T+1}). \quad (5)$$

If $\lim_{T \rightarrow +\infty} \sum_{t=0}^T [u(\kappa_t, \kappa_{t+1}) + \bar{p}\kappa_{t+1} - \bar{p}\kappa_t - \bar{x}] \in \mathbb{R}_-$ then $u(\kappa_t, \kappa_{t+1}) + \bar{p}\kappa_{t+1} - \bar{p}\kappa_t \rightarrow \bar{x}$. Let $(\hat{\kappa}, \hat{\chi})$ be a cluster point of $\{(\kappa_t, \kappa_{t+1})\}$. Then $u(\hat{\kappa}, \hat{\chi}) + \bar{p}\hat{\chi} - \bar{p}\hat{\kappa} = \bar{x}$. From Proposition 6, $(\hat{\kappa}, \hat{\chi}) = (\bar{\kappa}, \bar{\kappa})$. Therefore the sequence (κ_t) converges to $\bar{\kappa}$. From (5), $\sum_{t=0}^{\infty} [u(\kappa_t, \kappa_{t+1}) - \bar{x}] \in \mathbb{R}$.

Relation (5) implies

$$\sum_{t=0}^T [u(\kappa_t, \kappa_{t+1}) - \bar{x}] \leq \sum_{t=0}^T [u(\kappa_t, \kappa_{t+1}) + \bar{p}\kappa_{t+1} - \bar{p}\kappa_t - \bar{x}] + |\bar{p} \cdot \kappa_0| + |\bar{p} \cdot \kappa_{T+1}|.$$

Hence, if

$$\lim_{T \rightarrow +\infty} \sum_{t=0}^T [u(\kappa_t, \kappa_{t+1}) + \bar{p}\kappa_{t+1} - \bar{p}\kappa_t - \bar{x}] = -\infty,$$

then $\lim_{T \rightarrow +\infty} \sum_{t=0}^{\infty} [u(\kappa_t, \kappa_{t+1}) - \bar{x}] = -\infty$, since the sequence (κ_{T+1}) is bounded.

References

- [2009] Alvarez-Cuadrado, F. and N.V. Long, 2009, A Mixed Bentham–Rawls Criterion for Intergenerational Equity: Theory and Implications, *Journal of Environmental Economics and Management*, In Press.
- [1987] Arkin, V. I. and I. V. Evstigneev, 1987, *Stochastic Models of Control and Economic Dynamics*, Academic Press.
- [2010a] Asheim, G.B., d’Aspremont, C. and K. Banerjee, 2010, Generalized time-invariant overtaking, *Journal of Mathematical Economics*, 46, 519-533.

- [2010] Asheim, G. B. and T. Mitra, 2010, Sustainability and discounted utilitarianism in models of economic growth, *Mathematical Social Sciences* 59, 148-169.
- [2012] Asheim, G. B., T. Mitra and B. Tungodden, 2012, Sustainable recursive social welfare functions, *Economic Theory* 49, 267-292.
- [2004] Asheim G. B. and B. Tungodden, 2004, Do Koopmans postulates lead to discounted utilitarianism? Discussion Paper 32/04, Norwegian School of Economics and Business Administration.
- [1965] Atsumi, H., 1965, Neoclassical growth and the efficient program of capital accumulation, *Review of Economic Studies*, 32, 127–136.
- [2001] Ayong Le Kama, A., 2001, Sustainable growth, renewable resources and pollution, *Journal of Economic Dynamics and Control*, 25, 1911–1918.
- [2003] Basu, K., and T. Mitra, 2003, Aggregating infinite utility streams with intergenerational equity: the impossibility of being paretian, *Econometrica*, 71(5), 1557–1563.
- [2007] Basu, K., and T. Mitra, 2007, Utilitarianism for infinite utility streams: A new welfare criterion and its axiomatic characterization, *Journal of Economic Theory*, 133, 350–373.
- [1970a] Brock, W.A., 1970, On the existence of weakly maximal programmes in a multi-sector economy, *Review of Economic Studies*, 37, 275–280.
- [1970b] Brock, W.A., 1970, An axiomatic basis for the Ramsey-Weizsacker overtaking criterion, *Econometrica*, 38, 927–929.
- [1996] Chichilnisky, G., 1996, An axiomatic approach to sustainable development, *Social Choice and Welfare*, 13(2), 219–248.
- [1997] Chichilnisky, G., 1997, What is sustainable development?, *Land Economics*, 73(4), 467–491.
- [1990] Dana, R.A. and C. Le Van, 1990, On the Bellman equation of the overtaking criterion, *Journal of Optimization Theory and Applications*, 78(3), 605–612.

- [1993] Dana, R.A. and C. Le Van, 1993, On the Bellman equation of the overtaking criterion: Addendum, *Journal of Optimization Theory and Applications*, 67(3), 587–600.
- [1979] Dasgupta, P. and G.M. Heal, 1979, *Economic Theory and Exhaustible Resources*, Cambridge University Press, Cambridge.
- [1983] Dechert, W.D. and Nishimura, K., 1983, A complete characterization of optimal growth paths in an aggregate model with a non-concave production function, *Journal of Economic Theory*, 31, 332–354.
- [2012] Figuières, C. and M. Tidball, 2012, Sustainable exploitation of a natural resource: A satisfying use of Chichilnisky’s criterion, *Economic Theory* 49, 243-265.
- [2003] Fleurbaey, M. and P. Michel, 2003, Intertemporal equity and the extension of the Ramsey principle, *Journal of Mathematical Economics*, 39, 777–802.
- [1967] Gale, D., 1967, On optimal development in a multi-sector economy, *Review of Economic Studies*, XXXIV(1), 97, 1–18.
- [1998] Heal, G., 1998, *Valuing the Future, Economic Theory and Sustainability*, Columbia University Press, New York.
- [1960] Koopmans, T.C., 1960, Stationary Ordinal Utility and Impatience, *Econometrica* 28, 287–309.
- [1965] Koopmans, T., 1965, On the concept of optimal economic growth, *Pontificiae Academiae Scientiarum Varia*, 28, 225–300.
- [2012] Lauwers, L., 2012, Intergenerational equity, efficiency and constructibility, *Economic Theory* 49, 227-242.
- [2003] Le Van, C. and R.A. Dana, 2003, *Dynamic Programming in Economics*, Kluwer Academic Publishing.
- [2010] Le Van, C., K. Schubert and T-A. Nguyen, 2007, With exhaustible resources, can a developing country escape from the poverty trap?, *Journal of Economic Theory* 145, 2435–2447.
- [1928] Ramsey, F., 1928, A mathematical theory of saving, *Economic Journal*, 38, 543–559.

- [1971] Rawls, J., 1971, *A Theory of Justice*. Oxford, England: Clarendon.
- [1974] Solow, R., 1974, Intergenerational equity and exhaustible resources, *Review of Economic Studies* (symposium), 29–45.
- [1965] Von Weizsäcker, C.C., 1965, Existence of optimal programmes of accumulation for an infinite time horizon, *Review of Economic Studies*, 32, 85–104.
- [2007] Zaslavski, A., 2007, Turnpike results for discrete-time optimal control systems arising in economic dynamics, *Nonlinear Analysis* 67, 2024–2049.