



HAL
open science

Thermodynamic and transport properties of fluids: towards a LJ-SAFT molecular model valid for n-alkanes

Guillaume Galliero, C. Boned, Stephanie Delage Santacreu, Marc Odunlami,
F. Montel

► To cite this version:

Guillaume Galliero, C. Boned, Stephanie Delage Santacreu, Marc Odunlami, F. Montel. Thermodynamic and transport properties of fluids: towards a LJ-SAFT molecular model valid for n-alkanes. SAFT2011 discussion meeting. Pau, France, octobre 2011, 2011, pau, France. 2011. hal-01051603

HAL Id: hal-01051603

<https://hal.science/hal-01051603>

Submitted on 25 May 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

THERMODYNAMIC AND TRANSPORT PROPERTIES OF FLUIDS: TOWARDS A LJ-SAFT MOLECULAR MODEL VALID FOR N-ALKANES

Guillaume GALLIÉRO¹, Christian BONED¹, Stéphanie DELAGE², Marc ODUNLAMI² and François MONTEL³

¹Laboratory of Complex Fluids and their Reservoirs, UMR 5150 TOTAL-CNRS, Pau University, FRANCE.

²Laboratory of Applied Mathematics, UMR 5142 CNRS, Pau University, FRANCE.

³TOTAL E&P, CST, Avenue Larribau, Pau, FRANCE.

We study the limitations of the Lennard-Jones Chain fluid model (LJ-SAFT) to describe simultaneously all thermophysical properties (equilibrium and transport) of some n-alkanes along the vapor-liquid equilibrium line

MODELS AND THEORY

MOLECULAR DYNAMICS

Molecules movements numerically computed (Newton equations)

Spatial and temporal averages ($\sim 10^3$ particles, ~ 10 ns)

Exact thermophysical properties for a given molecular model

FLUID MODEL

Freely jointed Lennard-Jones spheres (Lennard-Jones Chain)

$$U_{LJ} = 4\epsilon \left(\left(\frac{\sigma}{r} \right)^{12} - \left(\frac{\sigma}{r} \right)^6 \right) + \text{internal bonds}$$

number of segments: N

Internal rigidity is included by $U_{flex} = k\epsilon(\theta - \theta_0)^2$

LJ-SAFT

No H-bonding, no polar and flexible $\Rightarrow A^{LJC} = A^{LJ} + A^{chain}$

A^{LJ} using the reference term of Kolafa & Nezbeda, (FPE 1994)

A^{chain} using the chain term of Johnson et al. (JPC 1994)

Accurate apart close to the critical point

VISCOSITY MODEL OF LJC

Based on MD results $\Rightarrow \eta(T, \rho) = \eta_0(T) + \eta_{res}(T, \rho)$

Galliero et al, IECR, 2005,
Galliero and Boned, PRE, 2009, PRE, 2010

Dilute gas Residual/Excess viscosity


Extended Chapman for η_0 and Rouse approximation for η_{res}

Valid for gas, liquid and supercritical states

RESULTS

VAPOR-LIQUID EQUILIBRIUM


$N=1$ for C1, $N=2$ for C4, $N=3$ for C7 and $N=4$ for C10, σ and ϵ have been adjusted



As well known this fluid model is correct for n-alkanes but $T_c^{SAFT} > T_c^{MD} > T_c$ (\nearrow with N)

Crossover approach is needed to deal with the critical region

SURFACE TENSION




Surface tensions are slightly overestimated

A simple correction of the critical temperature is sufficient


The coupling between DGT/DFT and LJ-SAFT yields good results

Galliero et al., JCP, 2008

SHEAR VISCOSITY



η is well estimated, except at low liquid T when $N \nearrow$
This weakness is related to the flexibility of the chain



η is highly affected by the rigidity while thermodynamic (fluid phase) only weakly

This degree of freedom (rigidity) can be used to improve simply the fluid model

Concerning thermal conductivity, λ_0 is out of range when $N \nearrow$ (internal degrees of freedom) but the λ_{res} of LJC is correct

Internal degrees of freedom (such as rigidity) should be included in a SAFT like approach