



HAL
open science

A note on Turing degree spectra of minimal subshifts

Mike Hochman, Pascal Vanier

► **To cite this version:**

Mike Hochman, Pascal Vanier. A note on Turing degree spectra of minimal subshifts. 2014. hal-01058198v2

HAL Id: hal-01058198

<https://hal.science/hal-01058198v2>

Preprint submitted on 27 Aug 2014 (v2), last revised 17 Sep 2014 (v4)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

A note on Turing degree spectra of minimal subshifts

Mike Hochman and Pascal Vanier

August 27, 2014

Abstract

Subshifts are shift invariant closed subsets of $\Sigma^{\mathbb{Z}^d}$, minimal subshifts are subshifts in which all points contain the same patterns. It has been proved by Jeandel and Vanier that the Turing degree spectra of non-periodic minimal subshifts always contain the cone of Turing degrees above any of its degree. It was however not known whether each minimal subshift's spectrum was formed of exactly one cone or not. We construct inductively a minimal subshift whose spectrum consists of an uncountable number of cones with disjoint base.

A \mathbb{Z}^d -subshift is a closed shift invariant subset of $\Sigma^{\mathbb{Z}^d}$. Subshifts may be seen as sets of colorings of \mathbb{Z}^d , with a finite number of colors, avoiding some set of forbidden patterns. Minimal subshifts are subshifts containing no proper subshift, or equivalently subshifts in which all configurations have the same patterns. They are fundamental in the sense that all subshifts contain at least a minimal subshift.

Degrees of unsolvability of subshifts have now been studied for a few years, Cenzer, Dashti, and King [CDK08] and Cenzer, Dashti, Toska, and Wyman [CDTW10; CDTW12] studied computability of one dimensional subshifts and proved some results about their Turing degree spectra: the *Turing degree spectrum* of a subshift is the set of Turing degrees of its points, see Kent and Lewis [KL10]. Simpson [Sim11], building on the work of Hanf [Han74] and Myers [Mye74], noticed that the Medvedev and Muchnik degrees of subshifts of finite type (SFTs) are the same as the Medvedev degrees of Π_1^0 classes: Π_1^0 classes are the subsets of $\{0, 1\}^{\mathbb{N}}$ for which there exists a Turing machine halting only on oracles not in the subset.

Subsequently Jeandel and Vanier [JV13] focused on Turing degree spectra of different classes of multidimensional subshifts: SFTs, sofic and effective subshifts. They proved in particular that the Turing degree spectra of SFTs are almost the same as the spectra of Π_1^0 classes: adding a computable point to the spectrum of any Π_1^0 class, one can construct an SFT with this spectrum. In order to prove that one cannot get a stronger statement, they proved that the spectrum of any non-periodic minimal subshift contains the cone above any of its degrees:

Theorem 1 (Jeandel and Vanier [JV13]). *Let X be a minimal non-finite subshift (i.e. non-periodic in at least one direction). For any point $x \in X$ and any degree $\mathbf{d} \geq_T \deg_T x$, there exists a point $y \in X$ such that $\mathbf{d} = \deg_T y$.*

Minimal subshifts are in particular interesting since any subshift contains a minimal subshift [Bir12]. Here we answer the followup question of whether a minimal subshift always corresponds to a single cone or if there exists one with at least two cones of disjoint base. It is quite easy to prove the following theorem:

Theorem 2. *For any Turing degree $\deg_T d$, there exists a minimal subshift X such that the set of Turing degrees of the points of X is a cone of base $\deg_T d$.*

For instance the spectrum of a Sturmian subshift [MH40] with an irrational angle is the cone whose base is the degree of the angle of the rotation. The theorem can also be seen as a corollary of Miller’s proof [Mil12][Proposition 3.1] of a result on Medvedev degrees.

In this paper, we prove the following result:

Theorem 3. *There exist a minimal subshift $X \subset \{0, 1\}^{\mathbb{Z}}$ and points $x_z \in X$ with $z \in \{0, 1\}^{\mathbb{N}}$ such that for any $z \neq z' \in \{0, 1\}^{\mathbb{N}}$, $\deg_T x_z$ and $\deg_T x_{z'}$ are incomparable and such that there exists no point $y \in X$ with $\deg_T y \leq_T \deg_T x_z$ and $\deg_T y \leq_T \deg_T x_{z'}$.*

The subshift constructed in this proof is not effective and cannot be “effectivized”, since minimal effective subshifts always contain a computable point and thus their spectra are the whole set of Turing degrees when they are non-periodic.

1 Preliminary definitions

We give here some standard definitions and facts about subshifts, one may consult the book of Lind and Marcus [LM95] for more details.

Let Σ be a finite alphabet, its elements are called *symbols*, the d -dimensional full shift on Σ is the set $\Sigma^{\mathbb{Z}^d}$ of all maps (colorings) from \mathbb{Z}^d to the Σ (the colors). For $v \in \mathbb{Z}^d$, the shift functions $\sigma_v : \Sigma^{\mathbb{Z}^d} \rightarrow \Sigma^{\mathbb{Z}^d}$, are defined locally by $\sigma_v(c_x) = c_{x+v}$. The full shift equipped with the distance $d(x, y) = 2^{-\min\{\|v\| \mid v \in \mathbb{Z}^d, x_v \neq y_v\}}$ is a compact metric space on which the shift functions act as homeomorphisms. An element of $\Sigma^{\mathbb{Z}^d}$ is called a *configuration*.

Every closed shift-invariant (invariant by application of any σ_v) subset X of $\Sigma^{\mathbb{Z}^d}$ is called a *subshift*. An element of a subshift is called a *point* of this subshift.

Alternatively, subshifts can be defined with the help of forbidden patterns. A *pattern* is a function $p : P \rightarrow \Sigma$, where P , the *support*, is a finite subset of \mathbb{Z}^d . We say that a configuration x contains a pattern $p : P \rightarrow \Sigma$, or equivalently that the pattern p appears in x , if there exists $z \in \mathbb{Z}^d$ such that $x|_{z+P} = p$.

Let \mathcal{F} be a collection of *forbidden* patterns, the subset $X_{\mathcal{F}}$ of $\Sigma^{\mathbb{Z}^d}$ containing the configurations having nowhere a pattern of \mathcal{F} . More formally, $X_{\mathcal{F}}$ is defined by

$$X_{\mathcal{F}} = \left\{ x \in \Sigma^{\mathbb{Z}^d} \mid \forall z \in \mathbb{Z}^d, \forall p \in \mathcal{F}, x|_{z+p} \neq p \right\}.$$

In particular, a subshift is said to be a *subshift of finite type* (SFT) when it can be defined by a collection of forbidden patterns that is finite. Similarly, an *effective subshift* is a subshift which can be defined by a recursively enumerable collection of forbidden patterns. A subshift is *sofic* if it is the image of an SFT by a letter by letter function.

Definition 1 (Minimal subshift). *A subshift X is called minimal if it verifies one of the following equivalent conditions:*

- *There is no subshift Y such that $Y \subsetneq X$.*
- *All the points of X contain the same patterns.*
- *It is the closure of the orbit of any of its points.*

We will use the two latter conditions.

For $x, y \in \{0, 1\}^{\mathbb{N}}$, we say that $x \leq_T y$ if there exists a Turing machine M such that M with oracle y computes x . Of course $x \equiv_T y$ when we have both $x \leq_T y$ and $y \leq_T x$. The Turing degree of x is the equivalence class of x with respect to \equiv_T . More details can be found in Rogers [Rog67].

For a possibly infinite word $w = w_0 \dots w_n$, we denote $w_{[i,j]} = w_i \dots w_j$.

2 Minimal subshifts with several cones

Lemma 2.1. *Let $\phi_1, \phi_2 : \{0, 1\}^{\mathbb{N}} \rightarrow \{0, 1\}^{\mathbb{N}}$ be two recursive operators and $L = \{w_1, w_2\} \subseteq \{0, 1\}^*$ two different words. There exists two words $w'_1, w'_2 \in L^*$ starting respectively with w_1 and w_2 such that we have one of the following:*

- (a) *either $\phi_1(w'_1 x) \neq \phi_2(w'_2 y)$ for any two sequences $x, y \in \{0, 1\}^{\mathbb{N}}$,*
- (b) *or $\phi_1(w'_1 x)$ and $\phi_2(w'_2 y)$ are recursive for all sequences $x, y \in \{0, 1\}^{\mathbb{N}}$,*

Proof. Let M_1, M_2 be the Turing machines computing the functions $x \mapsto \phi_1(w_1 x), x \mapsto \phi_2(w_2 x)$ respectively. It is quite clear that:

- either there exists some sequences $x, y \in L^*$ such that $M_1(x)$'s output differs from $M_2(y)$'s output at some step,
- or the outputs of both machines M_1, M_2 do not depend on their inputs, in this latter case, we are in case **b**, since in particular recursive inputs have recursive images.

In the former case, there exist prefixes w'_1 and w'_2 of x and y such that the partial outputs of M_1 once it has read w'_1 already differs from the partial output of M_2 once it has read w'_2 . \square

Theorem 4. *There exists a minimal subshift $X \subseteq \{0, 1\}^{\mathbb{N}}$ whose set of Turing degrees contains 2^{\aleph_0} disjoint cones of Turing degrees.*

Note that the following proof is in no way effective. As a matter of fact, all effective minimal subshifts contain some recursive point [BJ10], and their set of Turing degrees is the cone of *all* degrees.

Proof. We construct a sequence of sofic subshifts $(X_i)_{i \in \mathbb{N}}$ such that $X_{i+1} \subseteq X_i$ and such that the limit $X = \bigcap_{i \in \mathbb{N}} X_i$ is minimal. In the process of constructing the X_i , which will be formed of concatenations of allowed words x^i_1, \dots, x^i_k , we ensure that no extensions of two distinct words may compute an identical sequence with any of the first i Turing machines. At the same time, we make sure that all allowed words of level $i+1$ contain all words of level i , thus enforcing the minimality of the limit X . We also have to avoid that the limit X contains a computable point.

Let $(\mathcal{M}_i)_{i \in \mathbb{N}}$ be an enumeration of all minimal subshifts containing a recursive point. Such an enumeration exists since there are countably many computable sequences and minimal subshifts are the closure of any of their points. We do not know if such an enumeration is effective or not, but we do not need it to be for our purposes. We will also need an enumeration $(\phi_i)_{i \in \mathbb{N}}$ of the recursive operators from $\{0, 1\}^{\mathbb{N}}$ to $\{0, 1\}^{\mathbb{N}}$.

Now let us define the sequence of sofic shifts $(X_i)_{i \in \mathbb{N}}$. Each of these subshifts will be the shift invariant closure of the biinfinite words formed by concatenations of words of some language L_i which here will be finite languages. We define $X_0 = \{0, 1\}^{\mathbb{N}}$ that is to say X_0 is generated by $L_0 = \{w_0 = 0, w_1 = 1\}$. Let us now give the induction step. At each step, L_{i+1} will contain 2^{i+1} words $w_{0\dots 0}, \dots, w_{1\dots 1}$, the indices being binary words of length $i+1$, which will verify the following conditions:

1. The words w_{b_0}, w_{b_1} of L_{i+1} start with the word w_b of L_i .
2. The words w_b with $b \in \{0, 1\}^{i+1}$ of L_{i+1} each contain all the words of $w_{b'}$ with $b' \in \{0, 1\}^i$ of L_i .
3. For any two words $w_b, w_{b'}$ of L_{i+1} and for all $j, j' \leq i$:
 - Either for all $x, y \in L_i^\omega$, $\phi_j(w_b x) \neq \phi_{j'}(w_{b'} y)$,
 - Or for all $x, y \in L_i^\omega$, $\phi_j(w_b x), \phi_{j'}(w_{b'} y)$ are computable.
4. The words w_{b_0}, w_{b_1} do not appear in any configuration of \mathcal{M}_j , for all $j \leq i$.

Conditions 1 and 2 are easy to ensure: just make w_{b_a} start with $w' = w_b w_{0\dots 0} \dots w_{1\dots 1}$. We then use Lemma 2.1 to extend w' into a word w'' verifying condition 3, this is done several times, once for every quadruple w, w', j, j' . And

finally, since X_i is not minimal, we can extend w'' so that it contains a pattern appearing in none of the \mathcal{M}_j 's for $j \leq i$, to obtain condition . Now we can extend w'' with two different words thus obtaining w_{b0} and w_{b1} .

Now let's check that this leads to the desired result:

- $X = \bigcap X_i$ is a countable intersection of compact shift-invariant spaces, it is compact and shift-invariant, thus a subshift.
- Any pattern p appearing in some point of X is contained in a pattern w_b , with $b \in \{0, 1\}^i$ for some i , by construction (condition 2), all w_b with $b \in \{0, 1\}^{i+1}$ contain w_b . Therefore, all points of X , since they are contained in X_{i+1} , contain w_b and hence p . So X is minimal.
- For all $z \in \{0, 1\}^{\mathbb{N}}$, define the points $x_z = \lim_{i \rightarrow \infty} x_{z[0, i]}$, they are in X because they belong to each X_i . Condition 3 ensures that if two of them compute the same sequence $y \in \{0, 1\}^{\mathbb{N}}$, then this sequence is computable. Therefore they can only belong to the cone with base $\mathbf{0}$. And condition 2 ensures that no point of X belongs to a subshift containing a computable point.

□

It is quite straightforward to transform this proof in order to get a subshift on $\{0, 1\}^{\mathbb{Z}}$ instead of $\{0, 1\}^{\mathbb{N}}$ and obtain the following corollary:

Corollary 5. *For any dimension d , there exists a minimal subshift $X \subseteq \{0, 1\}^{\mathbb{Z}^d}$ whose set of Turing degrees contains 2^{\aleph_0} disjoint cones of Turing degrees.*

References

- [Bir12] M.-G. D. Birkhoff. “Quelques thormes sur le mouvement des systmes dynamiques”. In: *Bulletin de la SMF* 40 (1912), pp. 305–323.
- [BJ10] A. Ballier and E. Jeandel. “Computing (or not) quasi-periodicity functions of tilings”. In: *Second symposium on Cellular Automata (JAC 2010)*. 2010.
- [CDK08] D. Cenzer, A. Dashti, and J. L. F. King. “Computable symbolic dynamics”. In: *Mathematical Logic Quarterly* 54.5 (2008), pp. 460–469. DOI: [10.1002/malq.200710066](https://doi.org/10.1002/malq.200710066).
- [CDTW10] D. Cenzer, A. Dashti, F. Toska, and S. Wyman. “Computability of Countable Subshifts”. In: *Computability in Europe (CiE)*. Vol. 6158. Lecture Notes in Computer Science. 2010, pp. 88–97. DOI: [10.1007/978-3-642-13962-8_10](https://doi.org/10.1007/978-3-642-13962-8_10).
- [CDTW12] D. Cenzer, A. Dashti, F. Toska, and S. Wyman. “Computability of Countable Subshifts in One Dimension”. In: *Theory of Computing Systems* (2012). DOI: [10.1007/s00224-011-9358-z](https://doi.org/10.1007/s00224-011-9358-z).

- [Han74] W. Hanf. “Non Recursive Tilings of the Plane I”. In: *Journal of Symbolic Logic* 39.2 (June 1974), pp. 283–285.
- [JV13] E. Jeandel and P. Vanier. “Turing degrees of multidimensional SFTs”. In: *Theoretical Computer Science* 505.0 (2013). Theory and Applications of Models of Computation 2011, pp. 81–92. ISSN: 0304-3975. DOI: <http://dx.doi.org/10.1016/j.tcs.2012.08.027>. URL: <http://www.sciencedirect.com/science/article/pii/S0304397512008031>.
- [KL10] T. Kent and A. E. M. Lewis. “On the degree spectrum of a Π_1^0 class”. In: *Transactions of the American Mathematical Society* 362 (2010), pp. 5283–5319.
- [LM95] D. Lind and B. Marcus. *An Introduction to Symbolic Dynamics and Coding*. New York, NY, USA: Cambridge University Press, 1995. ISBN: 0-521-55900-6.
- [MH40] H. M. Morse and G. A. Hedlund. “Symbolic Dynamics II. Sturmian Trajectories”. In: *American Journal of Mathematics* 62.1 (1940), pp. 1–42.
- [Mil12] J. S. Miller. “Two Notes on subshifts”. In: *Proceedings of the American Mathematical Society* 140.5 (2012), pp. 1617–1622. DOI: [10.1090/S0002-9939-2011-11000-1](https://doi.org/10.1090/S0002-9939-2011-11000-1).
- [Mye74] D. Myers. “Non Recursive Tilings of the Plane II”. In: *Journal of Symbolic Logic* 39.2 (June 1974), pp. 286–294.
- [Rog67] H. Rogers. *Theory of recursive functions and effective computability*. McGraw-Hill series in higher mathematics. McGraw-Hill, 1967. URL: <http://books.google.fr/books?id=nAkzAAAAMAAJ>.
- [Sim11] S. G. Simpson. “Medvedev Degrees of 2-Dimensional Subshifts of Finite Type”. In: *Ergodic Theory and Dynamical Systems* (2011).